



# **King's Research Portal**

DOI: 10.12968/denu.2019.46.2.171

Document Version Peer reviewed version

Link to publication record in King's Research Portal

Citation for published version (APA):

O'Toole, S., Mylonas, P., & Bartlett, D. (2019). Practice-Based Risk Assessment- a Practical Guide for Oral Healthcare Teams: Tooth Wear. *Dental Update*, *46*(2), 171-178. https://doi.org/10.12968/denu.2019.46.2.171

### Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

#### General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

#### Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Generic Title: Erosive Tooth Wear Risk Assessment

Title: Practice-Based risk assessment – a practical guide for oral healthcare teams: Toothwear

O'Toole S<sup>1</sup>, BA B Dent Sc, MFD RCSI, FHEA, PhD Mylonas P<sup>1</sup>, BDS, DipMedEd, MFDS RCPS(Glasg), MJDF RCS (Eng), FHEA Bartlett D<sup>1</sup>, BDS, FDS RCS, MRD, FDS RCS (Restorative), CCST, FHEA, PhD

<sup>1</sup>Department of Prosthodontics, King's College London Dental Institute

Title: Practice-Based risk assessment – a practical guide for oral healthcare teams: Toothwear

Abstract: This paper discusses the presence of and interplay between erosion, abrasion and attrition as risk factors of tooth wear progression. The Basic Erosive Wear Examination is recommended as a simple practice-based screening tool to integrate into a risk assessment. Finally, the risk indicators are summarised and categorised using a high, medium or low risk classification system to facilitate practical risk assessment.

Clinical Relevance: This paper provides a practical guide for oral health providers to screen and risk assess tooth wear.

Brief Objectives Statement: This paper summarises the risk factors associated with erosive tooth wear and indicates how to perform a comprehensive risk assessment.

### Introduction

Tooth wear is a multi-factorial, complex process involving erosion, attrition, abrasion [1]. Erosive tooth wear is a term commonly used by European colleagues to represent that severe tooth wear rarely occurs without an underlying erosive aetiological component. Dental erosion is described as the loss of tooth tissue due to the effects of acid only, whilst attrition and abrasion describe the involvement of tooth-to-tooth and tooth-to-foreign object contact respectively. Current evidence indicates erosive tooth wear is common [2] and the prevalence is increasing, particularly in younger age groups [2,3].

Patients with erosive tooth wear can be difficult to risk assess as they are often unaware of their condition and may not see their dentist until it is significantly advanced. Being able to risk assess patients when they do present is necessary to minimise the progression of erosive tooth wear.

### **Risk Assessing Erosion**

This involves determining the potential origin of the underlying acids affecting patients'

teeth. These can be categorised into either intrinsic acids and/or extrinsic acids.

### Intrinsic acids (gastric contents)

Erosive tooth wear can result when stomach contents, containing gastric juice, frequently enter the oral cavity. The most common medical conditions resulting in erosive tooth wear from stomach acid are gastro-oesophageal reflux disease (GORD) and vomiting associated eating disorders (such as bulimia nervosa) affecting roughly 10% [4] and <1-2% [5] of the global population respectively. Most patients suffering from GORD are symptomatic and will most commonly complain of acid reflux and/or sternal pain and management from their GP should be encouraged to control the condition. The absence of symptoms does not necessarily mean that reflux is not involved. Silent gastro-oesophageal reflux can be a chronic long-term condition which remains symptomless, but the oral effects of the gastric acid reflux may be the only clinical sign. Other, more rare, causes of gastric contents entering the oral cavity include rumination habits, whereby the patient voluntarily regurgitates their food in order to rechew it and vomiting caused by other factors [6]. Pregnancy may also cause vomiting, particularly during the first trimester and hyperemesis gravidarum is a condition, affecting 0.3-2% of the population whereby vomiting starts early in the pregnancy and may last the duration of the entire pregnancy [7]. In addition, pregnant women are more prone to reflux.

Certain medications may have a central emetic effect such as chemotherapeutic drugs, opioids, digitalis and some oestrogens [8]. Alcoholism may also predispose to both vomiting and reflux disease [9]. With all these conditions, temporary periods of acid exposure are unlikely to result in severe pathological tooth wear, however, if they become chronic and uncontrolled, severe erosive tooth wear may result.

Eating disorders with an ongoing vomiting component can be challenging to assess from a patient history, given the unfortunate stigma attached to the associated mental health issues. Encouraging patients to discuss their current patterns regarding the frequency of vomiting, oral hygiene procedures before and after vomiting, and their diet will facilitate practical risk management while emphasising a supporting role.

Difficulty arises if an eating disorder is suspected but not diagnosed. Particularly as those with eating disorders may consume excessive amounts of diet drinks during bulimic phases, confounding the diagnosis between extrinsic and intrinsic erosive tooth wear. Erosive tooth wear may or may not be present in these patients. If present, it is often most severe on the palatal surfaces of the maxillary arch but generalised wear can be observed on all dental surfaces. Although there can be clinical indicators when the disease is active, such as enlargement of the parotid glands, soft tissue scarring on the backs of fingers where vomiting is forced with the hands and bruising or soft tissue damage on the hard and soft palate, it is difficult to risk assess and manage a patient without an open conversation about the severity of their condition.

When risk assessing erosive toothwear resulting from intrinsic acids, the frequency of acid entering the mouth is likely to be one of the key indicators of progression although there are few clinical studies to confirm this. Patients that are aware of weekly episodes of acid exposure can be classified as high risk of intrinsic erosive tooth wear progression. Those who experience acid exposure less than weekly may be considered as medium risk, whilst those that rarely experience acid exposure, due to good disease control, may be considered low risk.

### Extrinsic acids (diet)

The frequency of dietary acid intake has been shown as the most significant predictor of dietary erosive wear [10,11]. A recent case-control study reported that less-than-daily acid intake was associated with a negligible risk of wear [10]. The risk increased when dietary acids were consumed more than once daily and the further increased by 13 fold when three dietary acids or greater were consumed per day[10]. Common diet foods include fruits, fizzy drinks (excluding plain sparkling water), energy drinks, juices and smoothies. However, it also includes lesser known dietary acids such as fruit teas, fruit additions or flavourings in drinks eg. a slice of lemon or cordial/squash, sports drinks, fruit-flavoured lozenges or sweets, some medications, particularly effervescent vitamin C tablets. A patient who takes an effervescent multivitamin drink in the morning, has an apple as their mid-morning snack, takes a juice at lunchtime and then has a fruit tea that evening has had four acid attacks that day. The risk of developing erosion is reduced if acids are taken at meal times and this supports the current advice on balanced and healthy diets. Consuming fruit with meals showed no increased risk of erosive wear progression compared to those who snacked on fruit between meals. Similarly, those who drank acidic drinks with meals were half as likely to have severe erosive tooth wear than those who consumed the same frequency of acidic drinks between meals [10].

Drinking habits such as sipping, swishing or holding drinks in the mouth prior to swallowing increases the risk of having tooth wear [12]. There are rare occupational/recreational histories that might contribute to an increased risk of erosive wear such as wine tasters [13], athletes due to their need for hydration with erosive beverages containing electrolytes [14] or those with occupations or recreational habits associated with increased intake of

caffeinated drinks or soft drinks such as long-distance driving, night shift working or videogaming [15,16].

Risk assessment can be categorised according to the patient's daily habitual behaviour. If they consume three or greater dietary acids per day or greater than two dietary acids per day between meal, then they should be categorised as high risk. Daily acidic drink intake with meals would categorise them as medium risk and less than daily acid intake or daily fruit intake with meals places them in a low risk category. Patients should also be categorised as high risk if they have a habit of holding things in their mouth/cheeks prior to swallowing, sipping acidic drinks slowly or rinsing drinks around their mouth.

### **Risk Assessing Attrition**

Bruxism is defined as an oral habit consisting of involuntary rhythmic or spasmodic nonfunctional gnashing, grinding, or clenching of teeth, other than chewing movements of the mandible, which may lead to occlusal trauma [17].

Diagnosis of bruxism at early stages can be difficult as many patients are not aware of their habit. Jaw muscle discomfort, fatigue, pain, jaw lock, particularly upon awakening, have been reported by the American Academy of Sleep Medicine as being indicative of sleep bruxism. Often a sleep partner may be able to give evidence that the patient audibly or visibly bruxes at night. A comprehensive pain history involving the location, duration, precipitating factors, severity of the pain, relieving factors and where it radiates to can assist in the diagnosis of muscular or joint pain. The patient may give a history of chipping or fracturing restorations or cusps or both. In severe parafunction, occlusal loading can result in dental hypersensitivity. Occupation and lifestyle should be assessed for stress levels. Current or historical recreational drug use should be investigated if appropriate. The medical history should be checked for GORD symptoms, sleep apnoea or medications which may cause bruxism.

The risk assessment of bruxism starts with an extra-oral examination. Masseteric hypertrophy may be visible in chronic cases and the facial height may be reduced if the tooth wear has resulted in a reduced intraoral occlusal vertical dimension. There may be reduced or assymetric range of motion of the mandible, particularly in the morning. Clicking or crepitus of the joints may also indicate increased activity. Overactive muscles of mastication may or may not be tender on palpation and If symptoms are present they may assist in the diagnosis. The temporalis muscle can be particularly tender when a clenching habit is present as it is more actively involved in the final positioning of the mandible. However, the masseter may be tender for both grinding and clenching. An intraoral palpation of the medial pterygoid is not a particularly sensitive test as it can be painful even in cases of normal mandibular activity. Often though there are no clear symptoms and the diagnosis relies upon the clinical signs.

Intraorally, there can be soft tissue signs such as petechial haemorrhages, white lines of keratinisation on the buccal occlusal line, crenations on the tongue, broken or chipped restorations or cusps. Dental wear facets showing the location of interdigitation with the opposing arch, may also be visible. In severe bruxism cases the occlusal surfaces are flat.

In addition to history taking and clinical examination, specialised devices have been used to attempt to detect a bruxism habit. These include wear on occlusal splints or for research

purposes, sensors attached to occlusal splints and home muscle activity tests can be used. More advanced methods include electromyographic (EMG) recording of masticatory muscle activity in addition to the gold standard of polysomnography on sleep clinics.

Although the presence of attritional tooth wear is a confirmatory diagnosis, several studies have shown that the severity of bruxism is not related to the severity of tooth wear [18,19] which perhaps indicates that other tooth wear aetiological factors may be at play. However, if severe attritional wear is present, particularly at a young age, the patient should be categorised as high risk.

## **Risk Assessing Abrasion**

Abrasion is defined as an abnormal wearing away of the teeth by causes other than mastication [17]. Any foreign body misused or overused in the mouth has potential to be an aetiological factor. This includes chewing on pens, biting finger nails or any other foreign object, holding things between the teeth on a habitual basis, oral piercings and using items as toothpicks. A commonly quoted form of abrasion is from oral hygiene procedures [20]. Normal forces used to brush teeth with a low abrasivity toothpaste is unlikely to cause toothwear [21]. However, heavy pressure applied to the teeth from a brush has the potential to cause wear [21]. It remains unknown what force makes the abrasion risk severe and the diagnosis primarily relies on the clinical appearance of any lesions present.

Toothpaste abrasiveness is measured by a value known as the relative enamel abrasivity (REA) or relative dentine abrasivity (RDA). As dentine is more susceptible to abrasion due to its lower mineral content the RDA value is the widely used measure of abrasivity. An RDA over

100 is classified as high abrasivity and anything over 150 (predominantly found in whitening toothpastes) can be classified as harmful. Unfortunately, RDA testing is mandatory only in the US and not the UK/EU. Therefore, many toothpastes on the market, outside these geographical areas, do not have a known RDA value.

Studies have observed a relationship between the use of a hard toothbrush and an increase in tooth wear [11]. However, it is a difficult topic for clinical research. People who choose hard toothbrushes may be more likely to brush more aggressively and with a more abrasive toothpaste to get a clean feeling. Laboratory studies, have shown that increased tooth wear is associated with increased toothpaste abrasivity and increased force but not bristle stiffness. Several studies have shown that increased tooth wear was observed with a soft toothbrush. This was thought to be due to the increased flexing of the bristles which would hold more of the abrasive toothpaste. The combination of a medium bristled toothbrush and a low abrasivity toothpaste was observed to show the least tooth wear on both enamel and dentine in the laboratory [22].

A history of aggressive toothbrushing with a soft toothbrush may place patients at high risk. Dentine is more susceptible to mechanical forces than enamel and multiple studies have shown relationships between gingival recession and tooth wear [23,24]. Any exposed dentine will place patients in a higher risk category for a combined acid/mechanical wear challenge.

Not everybody uses a toothbrush and toothpaste to clean their teeth. Chewing on bark, sticks or using cloth with powders or salts is used in several countries to clean teeth and there is evidence of their efficacy for plaque removal. However, these can be very abrasive and have

also been associated with increased tooth wear [25]. It is important to recognise that a patient may be more comfortable using these oral hygiene procedures. In the absence of a daily acid source they may be categorised as medium risk category.

### BEWE - Screening wear already present

In today's society, it is difficult to justify not screening for tooth wear, not only to record it but also as part of a risk assessment. A useful tool for this is the Basic Erosive Wear Examination (BEWE), developed through international consensus, as a tool for screening erosive tooth wear in general practice. It grades the exposed surface based upon the percentage of the tooth surface affected and can be seen in Figure 1. A score between 0 - 3is given for each sextant (for the worst affected tooth), and a total BEWE score is given out of a maximum 18 as presented in Figure 2.

It was designed to be similar to a BPE (Basic Periodontal Examination) and therefore familiar to dentists. In common with the BPE, you do not need to record every surface but score the most severely worn surface in each sextant (Figure 2). An image of this can be seen in the below picture.

The sum score (in the above example 16 out of a total of 18) gives an overall representation of the tooth wear in the dentition. As the authors of the BEWE acknowledged, care must be taken with evaluating the total sum as the final measure. If severe, but localised wear is present in the upper anterior sextant, then this will present with an overall lower score and under the current classification system would not represent high risk despite the severe wear. A total wear score will also underestimate wear in a partially dentate arch. A useful method is to use both the maximum sextant score in conjunction with the total score in a risk assessment. Any mouth containing a score of 3 should be considered high risk, even if

elsewhere in the mouth the scores are lower. This recognises the distribution of tooth wear is often localised. A maximum score of 2 indicates moderate wear and a moderate risk whereas a maximum BEWE sextant score of 1 indicates low or no risk. Table 1 indicates a proposed method of risk assessment based upon tooth wear that is already present.

It is important to consider the age of the patient when risk assessing tooth wear. Physiological wear and tear is normal, and it would be unusual to see a sextant BEWE score of 0 on a patient over the age of 30. It would be equally as unusual to not see several sextant scores of 2 in a patient over the age of 55. The defining factor in risk assessment should be the presence of a BEWE score of 3. This is a sign of advanced wear at any age and the underlying aetiology needs to be diagnosed and managed. It is important to recognise that the need for restorative intervention is not related to the BEWE score.

### Summarising the Risk Assessment

A thorough history and examination is essential to risk assess tooth wear. Table 2 summarises the characteristics outlined in the previous chapter according to high (red), medium (amber) or low (green) characteristics. Behaviours change, and risk assessments should be repeated at regular intervals. Any positive behaviours can also be reinforced (Table 3).

## Conclusion

A comprehensive risk assessment for erosive tooth wear can be recorded at any stage and is important for the prevention of future wear. If the underlying aetiological factors are addressed there is evidence that tooth wear progression can reduce to physiological levels [26]. Active prevention methods such as effective preventive advice, fluoride measures and mouthguards should be a continuous intervention. Restorative intervention starts the patient on a lifelong treatment journey and should be undertaken with caution and care. Improved understanding of their progression through active monitoring for a period greater than 6 months will assist in your diagnosis. Monitoring the patient while you are determining the activity of the aetiological factors and risk assessing is not supervised neglect and may lead to an improved diagnosis.

## References

- [1] Carvalho TS, Colon P, Ganss C, Huysmans MC, Lussi A, Schlueter N, et al. Consensus report of the European Federation of Conservative Dentistry: erosive tooth wear—diagnosis and management. Clin Oral Investig 2015;19:1557–61. doi:10.1007/s00784-015-1511-7.
- [2] Bartlett DW, Lussi A, West NX, Bouchard P, Sanz M, Bourgeois D. Prevalence of tooth wear on buccal and lingual surfaces and possible risk factors in young European adults. J Dent 2013;41:1007–13. doi:10.1016/j.jdent.2013.08.018.
- [3] Jaeggi T, Lussi A. Prevalence, Incidence and Distribution of Erosion. In: Lussi A, Ganss
  C, editors. Erosive Tooth Wear From Diagnosis to Ther. 2nd ed., Basel Swizerland:
  Karger; 2014, p. 55–74.
- [4] El-Serag HB, Sweet S, Winchester CC, Dent J. Update on the epidemiology of gastrooesophageal reflux disease: a systematic review. Gut 2014;63:871–80.
   doi:10.1136/gutjnl-2012-304269.
- [5] Smink FRE, van Hoeken D, Hoek HW. Epidemiology of eating disorders: incidence, prevalence and mortality rates. Curr Psychiatry Rep 2012;14:406–14.
   doi:10.1007/s11920-012-0282-y.

- [6] Moazzez R, Bartlett D. Intrinsic causes of dental erosion. In: Lussi A, Ganss C, editors.
  Erosive Tooth Wear From Diagnosis to Ther. 2nd ed., Basel: Karger; 2014, p. 180–96.
- [7] Goodwin TM. Hyperemesis Gravidarum. Obstet Gynecol Clin North Am 2008;35:401–
  17. doi:10.1016/J.OGC.2008.04.002.
- [8] Scheutzel P. Etiology of dental erosion--intrinsic factors. Eur J Oral Sci 1996;104:178–90.
- [9] Teixeira L, Manso M-C, Manarte-Monteiro P. Erosive tooth wear status of institutionalized alcoholic patients under rehabilitation therapy in the north of Portugal. Clin Oral Investig 2016:epub ahead of print. doi:10.1007/s00784-016-1823-2.
- [10] O'Toole S, Bernabé E, Moazzez R, Bartlett D. Timing of dietary acid intake and erosive tooth wear: A case-control study. J Dent 2017;56. doi:10.1016/j.jdent.2016.11.005.
- [11] Lussi A, Schaffner M. Progression of and risk factors for dental erosion and wedgeshaped defects over a 6-year period. Caries Res 2000;34:182–7. doi:16587.
- [12] Hasselkvist A, Johansson A, Johansson AK. A 4 year prospective longitudinal study of progression of dental erosion associated to lifestyle in 13-14 year-old Swedish adolescents. J Dent 2016;47:55–62. doi:10.1016/j.jdent.2016.02.002.
- [13] Mulic A, Tveit AB, Hove LH, Skaare AB. Dental erosive wear among Norwegian wine tasters. Acta Odontol Scand 2011;69:21–6. doi:10.3109/00016357.2010.517554.
- [14] Ashley P, Di Iorio a, Cole E, Tanday a, Needleman I. Oral health of elite athletes and association with performance: a systematic review. Br J Sports Med 2014:1–7.
  doi:10.1136/bjsports-2014-093617.
- [15] Lowden A, Moreno C, Holmbäck U, Lennernäs M, Tucker P. Eating and shift workeffects on habits, metabolism, and performance. vol. 36. 2010.

- [16] Giammattei J, Blix G, Marshak HH, Wollitzer AO, Pettitt DJ. Television Watching and Soft Drink Consumption. Arch Pediatr Adolesc Med 2003;157:882.
   doi:10.1001/archpedi.157.9.882.
- [17] The Academy of Prosthodontics. The Glossary Of Prosthodontic Terms. J OfProsthetic Dent 2005;94:10–92. doi:10.1016/0168-8510(94)90003-5.
- [18] Jonsgar C, Hordvik PA, Berge ME, Johansson AK, Svensson P, Johansson A. Sleep bruxism in individuals with and without attrition-type tooth wear: An exploratory matched case-control electromyographic study. J Dent 2015;43:1504–10. doi:10.1016/j.jdent.2015.10.002.
- [19] Pergamalian A, Rudy TE, Zaki HS, Greco CM. The association between wear facets, bruxism, and severity of facial pain in patients with temporomandibular disorders. J Prosthet Dent 2003;90:194–200. doi:10.1016/S0022-3913(03)00332-9.
- [20] Hunter ML, Addy M, Pickles MJ, Joiner A. The role of toothpastes and toothbrushes in the aetiology of tooth wear. Int Dent J 2002;52:399–405. doi:10.1111/j.1875-595X.2002.tb00729.x.
- [21] Wiegand A, Köwing L, Attin T. Impact of brushing force on abrasion of acid-softened and sound enamel. Arch Oral Biol 2007;52:1043–7.
   doi:10.1016/j.archoralbio.2007.06.004.
- [22] Lippert F, Arrageg MA, Eckert GJ, Hara AT. Interaction between toothpaste abrasivity and toothbrush filament stiffness on the development of erosive/abrasive lesions *in vitro*. Int Dent J 2017:1–7. doi:10.1111/idj.12305.
- [23] West NX, Sanz M, Lussi A, Bartlett DW, Bouchard P, Bourgeois D. Prevalence of dentine hypersensitivity and study of associated factors: A European populationbased cross-sectional study. J Dent 2013;41:841–51. doi:10.1016/j.jdent.2013.07.017.

- [24] Fukumoto Y, Horibe M, Inagaki Y, Oishi K, Tamaki N, Ito HO, et al. Association of gingival recession and other factors with the presence of dentin hypersensitivity.
  Odontology 2014;102:42–9. doi:10.1007/s10266-012-0099-5.
- [25] Shah N, Mathur VP, Jain V, Logani A. Association between traditional oral hygiene methods with tooth wear, gingival bleeding, and recession: A descriptive crosssectional study. Indian J Dent Res 2018;29:150–4. doi:10.4103/ijdr.IJDR\_651\_16.
- [26] O'Toole S, Newton T, Moazzez R, Hasan A, Bartlett D. Randomised Controlled Clinical Trial Investigating the Impact of Implementation Planning on Behaviour Related to the Diet. Sci Rep 2018;8:8024. doi:10.1038/s41598-018-26418-0.



 $\frac{3}{3}$   $\frac{3}{2}$   $\frac{2}{2}$  Sum Score:16

Figure 2. BEWE score of clinical patient as would be recorded on clinical notes

	High Risk Characteristics	Medium Risk	Low Risk Characteristics
	(Red)	Characteristics (Amber)	(Green)
Maximum	3	2	1/0
BEWE Score			
in any			
Sextant			
Total BEWE	13 or greater	Between 7-13	6 or lower
Score		depending on age	

Table 1. Risk assessment based upon wear that is already present

High Risk Chara	cteristics	Medium Risk	Low Risk Characteristics
(Red)		Characteristics (Amber)	(Green)
Erosion Gastric Reflux Sympt	oms on a	Infrequent GORD	No history or symptoms of
weekly basis or	poorly	symptoms or well	GORD.
controlled GORD.		managed symptoms	
			No history of an eating
An active vomiting	g eating	Managing eating	disorder or previous
disorder (weekly bas	sis)	disorder and vomiting	history of an eating
		episodes occur	disorder but is inactive.
Total dietary acid i	ntake 3+	infrequently	
per day or 2+ pe	r day in		Less than daily acidic drink
between meals		Daily dietary acid intake	intake.
		but less than 2 per day	No sipping, swishing or
An occupation	which		holding drinks in mouth.
encourages	frequent		
consumption of acid	ic drinks		Fruit intake only with
			meals.
Drinking an acid dai	ly with a		
habit such as slow	sipping,		Consumes dietary acids
rinsing or swishing o	r holding		quickly in less than 10
the drink in the mo	uth prior		minutes.
to swallowing			
			No intrinsic sources of acid
Spending >10 minut	es eating		
a single portion of fr	uit every		
day			

Attrition	Flattened teeth already present and active soft tissue signs/symptoms.	Currently showing soft tissue signs/symptoms of bruxism but wearing mouthguard	No history of parafunction with no intraoral signs of parafunction
	High stress occupation and aware of grinding but unable to wear mouthguard		
Abrasion	Gingival recession and exposed dentine combined with aggressive brushing and interdental habits (3+ brushing per day and the use of a high RDA toothpaste)	Gingival recession and exposed dentine but non-aggressive brushing and interdental routine Oral piercing	Brushes twice a day with low abrasive toothpaste Non-aggressive interdental cleaning routine.
		Alternate form of oral hygiene procedures	No other sources of mechanical damage such as pen or nail biting.

Table 2. Risk assessment based on clinical presentation and daily behaviours

Overall risk characterisation for Erosive Tooth Wear				
RedReassessevery6-12monthsuntilconfidentthat the risk is nolonger present	Presence of one aetiological factor in the red category Daily exposure to a combination of acid and mechanical aetiological factors which for which management is not optimal			
Amber Reassess every 12 months until confident that the risk is no longer present	Presence of one aetiological factor in the amber category Multiple amber aetiological factors are present but are being managed to some degree of success.			
Green Reassess every 2-3 years until change in risk becomes evident	No daily risk of any component If characteristics are present, they are being managed effectively			

Table 3. Repeating Risk Assessments on Adults. Children and Adolescents should be risk assessed at the discretion of the practitioner.