



King's Research Portal

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Stevelink, S., Jones, M. M., Greenberg, N., Rona, R. J., & Fear, N. T. (2019). Mental health outcomes at the end of the British involvement in the Iraq and Afghanistan conflicts: a cohort study - Authors' reply. *British Journal of Psychiatry*.

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Correspondence

Edited by Kiriakos Xenitidis and Colin Campbell

Contents

- Threats to the validity of studies of post-traumatic stress disorder from unmeasured symptomatic exposure to mefloquine
- Author's reply

Threats to the validity of studies of post-traumatic stress disorder from unmeasured symptomatic exposure to mefloquine

I read with interest the recent study by Stevelink and colleagues¹ that studied the prevalence of various mental health outcomes, including post-traumatic stress disorder (PTSD), among UK military personnel following the conflicts in Iraq and Afghanistan. I am concerned that several of the authors' conclusions may be invalid, owing to a failure to measure and control for an important confounder in their study.

Participants in the authors' study were at risk of exposure to mefloquine, an antimalarial drug widely used by the UK military during the period, particularly during training missions in Africa and during deployments to Afghanistan. Mefloquine is known to cause psychiatric adverse effects, including nightmares, insomnia, depression and anxiety, which can last years after use. Individuals who experience psychiatric adverse effects during continued use of the drug, a condition known as symptomatic exposure, are at risk of these adverse effects becoming chronic. For example, based on a synthesis of recent data, abnormal dreams and nightmares lasting over 3 years after use of mefloquine may affect over 2% of those exposed to the drug.²

Various US military authors have cautioned that mefloquine use can 'confound the diagnosis and management of PTSD',³ and that given 'the overlapping symptoms of post-traumatic stress disorder and mefloquine toxicity, it can be challenging to distinguish between the two diagnoses'.⁴ The importance of identifying past mefloquine exposure is further underscored by the addition of criterion H to the PTSD diagnostic criteria in DSM-5, which requires that the disturbance not be the result of the effects of a medication.⁵ As the chronic adverse effects of mefloquine may mimic several symptoms of PTSD, including several symptoms assessed using the 17-item National Centre for PTSD Checklist (PCL-C), the authors use of this instrument without distinguishing which symptoms may have been the result of mefloquine risks the adverse effects of the drug having been misattributed to PTSD.

As mefloquine exposure is correlated with deployment, and as symptomatic mefloquine exposure creates a separate causal pathway for the development of several symptoms assessed by the PCL-C, symptomatic mefloquine exposure serves as a classic epidemiological confounder in the authors' study.²

In order to avoid potentially fatal threats to validity that result from such confounding, the authors are encouraged to measure symptomatic mefloquine exposure in future studies and to control for these effects during analysis. Our group has introduced the two-question White River Mefloquine Instrument (WRMI-2)⁶ for this purpose and encourages the use of this instrument in research and in the clinical evaluation of recent veterans at risk of mefloquine exposure.

Declaration of interest: R.N has been retained as consultant and expert witness in legal cases involving claims of adverse effects from antimalarial drugs, including mefloquine.

References

- 1 Stevelink SAM, Jones M, Hull L, Pernet D, MacCrimmon S, Goodwin L, et al. Mental health outcomes at the end of the British involvement in the Iraq and Afghanistan conflicts: a cohort study. *Br J Psychiatry* 2018; **213**: 690–7.
- 2 Nevin RL. Re: 'A decade of war: prospective trajectories of posttraumatic stress disorder symptoms among deployed US military personnel and the influence of combat exposure'. *Am J Epidemiol* 2018; **187**: 1573–4.
- 3 Magill A, Cersovsky S, DeFraités R. Special considerations for US military deployments. In *CDC Health Information for International Travel: The Yellow Book 2012* (ed GW Brunette): 561–5. Oxford University Press, 2012.
- 4 Livezey J, Oliver T, Cantilena L. Prolonged neuropsychiatric symptoms in a military service member exposed to mefloquine. *Drug Saf - Case Rep* 2016; **3**: 7.
- 5 Nevin RL. Mefloquine and posttraumatic stress disorder. In *Textbook of Military Medicine. Forensic and Ethical Issues in Military Behavioral Health* (ed EC Ritchie): 277–96. Borden Institute, 2015.
- 6 The Quinism Foundation. The Quinism Foundation Partners with the National Centre for Trauma to Screen U.K. Veterans for Symptomatic Mefloquine Exposure. July 30, 2018. Available at: <https://quinism.org/press-releases/the-quinism-foundation-partners-with-the-national-centre-for-trauma-to-screen-u-k-veterans-for-symptomatic-mefloquine-exposure>.

Remington Nevin, Executive Director, The Quinism Foundation, USA. Email: rnevin@quinism.org

doi:10.1192/bjp.2019.1

Author's reply

We appreciate the interest Dr Remington Nevin has shown in our article¹ and thank you for giving us the opportunity to respond. Dr Remington Nevin claims that mefloquine exposure is an important confounder that we have failed to control for and that our lack of adjustment for this variable may have invalidated our conclusions. We respectfully disagree.

First, there is no empirical evidence that mefloquine use confounds the association between deployment-related exposures and post-traumatic stress disorder (PTSD), nor that PTSD is a specific side-effect of mefloquine use. Adverse reactions that are reported in the National Institute for Health and Care Excellence guidelines include anxiety, depression, sleep disorders, abnormal dreams, dizziness, headache, vomiting, digestive problems and skin reactions.^{2–4} A recent Cochrane review found a low incidence of side-effects among mefloquine users.³ Further, mefloquine users do not report more frequent serious side-effects than more commonly used antimalarials such as atovaquone-proguanil or doxycycline.³

Second, data from the UK's Defence Medical Information Capability Programme indicated that between 1 April 2007 and 31 March 2015, approximately 120 000 UK armed forces personnel were prescribed an antimalarial drug, of which around 17 000 were prescribed mefloquine (14%).⁵ When looking specifically at personnel who deployed to Afghanistan between 1 April 2007 and 31 December 2014, only 536 (0.4%) of the 131 000 of personnel who deployed were prescribed mefloquine. A further 12 908 (10%) were prescribed an unknown malarial prophylaxis. The available data relating to both prescription and use, which is determined by service policy,⁶ suggest that mefloquine is not the first-choice malarial prophylactic for UK armed forces personnel other than in circumstances where resistance to other medication is a factor. Hence, it is highly unlikely that mefloquine was prescribed to this 10%. Therefore, its infrequent use would mean that it cannot account for any substantial proportion of the PTSD identified in our study.

Third, adherence to treatment during deployment varies substantially but is generally found to be low to moderate (range 38–61%), further weakening any potential confounding effect of mefloquine antimalarial medication.^{7,8} Although we could assess whether UK serving personnel were prescribed malarial prophylaxis as part of the health and well-being cohort study questionnaire, it is impossible to accurately assess through self-report whether personnel were compliant with the treatment regime and which medication was prescribed. Hence, even if prescription data were available, it would be difficult to identify the potentially marginal mental health and well-being impact of mefloquine particularly upon PTSD, as factors such as combat exposure have a direct effect.

Given the low prescription rate of mefloquine compounded with the low rates of adherence, it is implausible that mefloquine could have mimicked the pattern of associations of PTSD we observed with deployment, engagement type, serving status and combat role.

Declaration of interest: All authors are based at King's College London which, for the purpose of this study and other military-related studies, receives funding from the UK Ministry of Defence (MoD). S.A.M.S., M.J. and R.J.R.'s salaries were totally or partially paid by the UK MoD. The UK MoD provides support to the Academic Department of Military Mental Health, and the salaries of N.G. and N.T.F. are covered totally or partly by this contribution.

References

- 1 Stevelink SAM, Jones M, Hull L, Pernet D, MacCrimmon S, Goodwin L, et al. Mental health outcomes at the end of the British involvement in the Iraq and Afghanistan conflicts: a cohort study. *Br J Psychiatry* 2018; **213**: 690–7.
- 2 National Institute for Health and Care Excellence. Mefloquine. In *British National Formulary (BNF)*. NICE, no date (<https://bnf.nice.org.uk/drug/mefloquine.html#sideEffects>).
- 3 Tickell-Painter M, Maayan N, Saunders R, Pace C, Sinclair D. Mefloquine for preventing malaria during travel to endemic areas. *Cochrane Database Syst Rev* 2017; **10**: CD006491.
- 4 Tuck J, Williams J (2016). Malaria protection in Sierra Leone during the Ebola outbreak 2014/15; the UK military experience with malaria chemoprophylaxis Sep 14-Feb 15. *Travel Med Infect Dis* 2016; **14**: 471–4.
- 5 Ministry of Defence. Mefloquine hydrochloride prescribing in the UK Armed Forces, 1 April 2007–31 March 2015. *Ad Hoc Stat Bull* 2016; 12 January (https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/491134/20160112_Adhoc_Statistical_Bulletin_Mefloquine_prescribing_in_the_UK_Armed_Forces_-O.pdf).
- 6 Surgeon General. *Preventing Malaria in UK Armed Forces Personnel. Joint Service Publication 950, Part 1, Leaflet 3-3-1*. Ministry of Defence, 2017 (https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/650918/20170601-JSP_950_Lft_3-3-1_Prev_Mal_Mil_Pers_V2_01_Jun17_Endorsed-O.pdf).
- 7 Brisson M, Brisson P. Compliance with antimalaria chemoprophylaxis in a combat zone. *Am J Trop Med Hyg* 2012; **86**: 587–90.
- 8 Kotwal RS, Wenzel RB, Sterling RA, Porter WD, Jordan NN, Petruccioli BP. An outbreak of malaria in US Army Rangers returning from Afghanistan. *JAMA* 2005; **293**: 212–6.

Sharon A. M. Stevelink, Lecturer in Epidemiology, King's College London, UK;
Margaret Jones, Senior Research Associate, King's College London, UK;
Roberto J. Rona, Professor of Public Health Medicine, King's College London, UK;
Neil Greenberg, Professor of Defence Mental Health, King's College London, UK;
Nicola T. Fear, Professor of Epidemiology, King's College London, UK. Email: sharon.stevelink@kcl.ac.uk

doi:10.1192/bjp.2019.2