

King's Research Portal

DOI: [10.1007/s00222-018-0821-1](https://doi.org/10.1007/s00222-018-0821-1)

Document Version Peer reviewed version

[Link to publication record in King's Research Portal](https://kclpure.kcl.ac.uk/portal/en/publications/4be3ea1e-db48-4d14-a506-99f8845a4cd3)

Citation for published version (APA): Hadžić, M., & Jang, J. (2018). Expanding large global solutions of the equations of compressible fluid mechanics. INVENTIONES MATHEMATICAE, 214(3), 1205-1266. <https://doi.org/10.1007/s00222-018-0821-1>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Expanding large global solutions of the equations of compressible fluid mechanics

Mahir Hadžić * and Juhi Jang †

Abstract

Without any symmetry assumptions on the initial data we construct global-in-time unique solutions to the vacuum free boundary three-dimensional isentropic compressible Euler equations when the adiabatic exponent γ lies in the interval $(1, \frac{5}{3}]$. Our initial data lie sufficiently close to the expanding compactly supported affine motions recently constructed by Sideris and they satisfy the physical vacuum boundary condition.

1 Introduction

We consider the dynamics of moving gases in three dimensions as described by compressible isentropic Euler system. We are interested in fluids surrounded by vacuum and therefore the unknowns are the density ρ , fluid velocity vector-field u, and the free boundary of the support of ρ , denoted by $\partial\Omega(t)$. The resulting initial value problem takes the form:

$$
\partial_t \rho + \text{div}(\rho \mathbf{u}) = 0 \qquad \text{in } \Omega(t); \tag{1.1a}
$$

$$
\rho \left(\partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} \right) + \nabla p = 0 \qquad \text{in } \Omega(t) \, ; \tag{1.1b}
$$

$$
p = 0 \qquad \text{on } \partial\Omega(t) \, ; \tag{1.1c}
$$

$$
\mathcal{V}(\partial \Omega(t)) = \mathbf{u} \cdot \mathbf{n}(t) \quad \text{on} \quad \partial \Omega(t) \tag{1.1d}
$$

$$
(\rho(0,\cdot), \mathbf{u}(0,\cdot)) = (\rho_0, \mathbf{u}_0), \Omega(0) = \Omega_0.
$$
 (1.1e)

Here $V(\partial \Omega(t))$ denotes the normal velocity of $\partial \Omega(t)$ and $n(t)$ denotes the outward unit normal vector to $\partial\Omega(t)$. Equation (1.1a) is the well-known continuity equation, while (1.1b) expresses the conservation of momentum. Boundary condition (1.1c) is the vacuum boundary condition, while (1.1d) is the kinematic boundary condition stating that the boundary movement is tangential to the fluid particles.

In this article, we shall only consider ideal barotropic fluids, where the pressure depends only on the density, expressed through the following equation of state

$$
p = \rho^{\gamma}, \quad \gamma > 1,\tag{1.2}
$$

where we have set the entropy constant to be 1. We additionally demand that the initial density satisfies the *physical vacuum boundary condition* [16, 25]:

$$
-\infty < \frac{\partial c_s^2}{\partial n} \Big|_{\partial \Omega(0)} < 0 \tag{1.3}
$$

[∗]Department of Mathematics, King's College London, Strand, London WC2S 2LR, UK. Email: mahir.hadzic@kcl.ac.uk.

[†]Department of Mathematics, University of Southern California, Los Angeles, CA 90089, USA and Korea Institute for Advanced Study, Seoul, Korea. Email: juhijang@usc.edu.

where $c_s = \sqrt{\frac{d}{d\rho}p(\rho)}$ is the speed of the sound. We shall refer to the system of equations (1.1)–(1.3) as the *Euler system* and denote it by E_{γ} .

Due to the inherent lack of smoothness of the enthalpy c_s^2 at the vacuum boundary (implied by the assumption (1.3)), a rigorous understanding of the existence of physical vacuum states in compressible fluid dynamics has been a challenging problem. Only recently, a successful local-in-time well-posedness theory for the E_{γ} system was developed in [5, 18] using the Lagrangian formulation of the Euler system in the vacuum free boundary framework. The fundamental unknown is the flow map ζ defined as a solution of the ordinary differential equation (ODE)

$$
\partial_t \zeta = \mathbf{u} \circ \zeta.
$$

Very recently Sideris [44] constructed a special class of *affine* fluid motions¹ that solve (1.1) – (1.3) globally-in-time in the vacuum free boundary setting. The center of mass of such solutions is fixed to be at the origin. Here we consider a slightly larger class of affine motions by allowing the center of mass a to be time-dependent. The flow map of an affine solution with center of mass $a(t)$ by definition takes the form

$$
\zeta(t, y) = A(t)y + \mathbf{a}(t), \ \ A(t) \in \text{GL}^+(3), \ \ \mathbf{a}(t) \in \mathbb{R}^3. \tag{1.4}
$$

Plugging this ansatz into (1.1) one can effectively separate variables and discover a family of evolving vacuum states of the form

$$
\rho_{A,\mathbf{a}}(t,x) = \det A(t)^{-1} \left[\frac{(\gamma - 1)}{2\gamma} (1 - |A^{-1}(t)(x - \mathbf{a}(t))|^2) \right]^{\frac{1}{\gamma - 1}},\tag{1.5}
$$

$$
\mathbf{u}_{A,\mathbf{a}}(t,x) = \dot{A}(t)A^{-1}(t)(x - \mathbf{a}(t)) + \dot{\mathbf{a}}(t),
$$
\n(1.6)

whereby the flow matrix $t \to A(t)$ solves the following Cauchy problem for a system of ODEs:

$$
\ddot{A}(t) = \det A(t)^{1-\gamma} A(t)^{-\top},\tag{1.7}
$$

$$
(A(0), \dot{A}(0)) = (A_0, A_1) \in \text{GL}^+(3) \times \mathbb{M}^{3 \times 3}, \tag{1.8}
$$

and

$$
\mathbf{a}(t) = \mathbf{a}_0 + t\mathbf{a}_1, \ \mathbf{a}_0, \mathbf{a}_1 \in \mathbb{R}^3. \tag{1.9}
$$

The density and the velocity field are both supported on a moving ellipse $\Omega(t)$ of the form $a(t) + A(t)\Omega$, where $\Omega = B_1(0)$ is the unit ball in \mathbb{R}^3 . We denote the set of such affine motions by \mathscr{S} . Our main theorem states that the elements of $\mathscr S$ are stable under small perturbations if $\gamma \in (1, \frac{5}{3}]$.

 $¹$ Affine motions in the absence of free boundaries have been used before to understand qualitative behavior of solutions in</sup> fluid mechanics, for instance see Majda [31].

Elements of $\mathcal S$

Theorem 1.1 (Global existence in the vicinity of affine motions). Assume that $\gamma \in (1, \frac{5}{3}]$. Then small *perturbations of the expanding affine motions given by* (1.5)*–*(1.8) *give rise to unique globally-in-time defined solutions of the Euler system E*γ*. Moreover, their support expands at a linear rate and they remain close to the underlying set* S *of affine motions.*

To the best of our knowledge, Theorem 1.1 is the first global-in-time existence result to the physical vacuum free-boundary compressible Euler equations (1.1) – (1.2) without any symmetry assumptions on the initial data. These solutions are unique in the vacuum free boundary framework, and they are simultaneously weak solutions in \mathbb{R}^3 obtained by extending ρ to 0 in the vacuum region. Moreover, Theorem 1.1 is consistent with another result of Sideris [43]: *if the Euler system E*^γ *admits globalin-time compactly supported solutions then the diameter of the support has to grow at least linearly in time. In other words,* $\sup_{x \in \Omega(t)} |x| \gtrsim t$. The existence of such expanding global-in-time solutions has been shown only recently [44] in the context of a finite parameter family of affine motions $\mathscr S$ introduced above. One important novelty of our result is the introduction of a rigorous mathematical framework that allows us to describe the nonlinear stable expansion phenomenon of the Euler flows for all positive times around the set \mathscr{S} . Theorem 1.1 rigorously justifies the intuition that the expansion of a gas is a mechanism that prevents the development of shock singularities².

Our work brings to the fore a rich interaction of scaling symmetries of the compressible Euler system on one hand and the delicate well-posedness theory in the presence of vacuum boundaries on the other, see Section 1.2.

Remark 1.2. *A precise statement of Theorem 1.1 specifying the function spaces and the notions of "small" and "close" in the statement above is provided in Theorem 2.4.*

Remark 1.3. *Theorem 1.1 covers a range of adiabatic exponents of physical importance. The exponent* $\gamma = \frac{5}{3}$ is commonly used in the description of a monatomic gas, $\gamma = \frac{7}{5}$ corresponds to a diatomic gas, and $\gamma = \frac{4}{3}$ is often referred to as the radiative case.

Remark 1.4. As shown in [44], the affine motions exist even when $\gamma > \frac{5}{3}$. In the first version of *this manuscript we stated that it would be interesting to know whether one can go beyond the* $\gamma =$ 5 3 *threshold in our theorem. We have been made aware of a very recent preprint of Shkoller and Sideris [38], wherein the question raised above is answered to the positive - Sideris affine motions are stable in a suitable sense even if* $\gamma > \frac{5}{3}$ *.*

²In the absence of free boundaries, gas expansion also plays an important role in the global results in [39, 11]. The full nonlinear analysis of stabilizing effects of the fluid expansion in the context of general relativistic cosmological models was initiated by Rodnianski & Speck [36] and extended further in [41, 26, 13, 35, 10]

Remark 1.5. The theorem shows that in the forward time direction, in the vicinity of $\mathscr S$ the solution *naturally splits into an affine component (i.e. an element of S) and a remainder, which as we shall show, remains small due to the underlying expansion associated with the elements of* \mathscr{S} *.*

1.1 Existence theories for compressible Euler flows

We briefly review some existence theories of compressible flows, where we focus on the works most relevant to our problem. We will not attempt to provide an exhaustive overview, but for more references we refer the reader to [7].

It is well-known [7] that the system (1.1) is strictly hyperbolic if the density is bounded below away from zero. In the absence of vacuum, the theory of symmetric hyperbolic systems developed by Friedrichs-Lax-Kato applies and one can construct local-in-time smooth solutions, see for instance Majda's book [30]. These smooth solutions break down in a finite time for generic initial data, although global-in-time classical solutions can exist for special initial data with c_s sufficiently smooth and small, see Serre [39], Grassin [11], and Rozanova [37]. In particular, in two space dimensions there exist initial data leading to so-called eternal solutions for all $t \in \mathbb{R}$ [40, 11]. The first proof of formation of singularities for classical Euler flow with a nonzero constant density at infinity was given by Sideris [42]. A similar result was obtained by Makino-Ukai-Kawashima [32] for smooth compactly supported disturbance moving into vacuum. In an important work [3], Christodoulou gave a precise description of shock formation for irrotational relativistic fluids starting with small smooth initial data. We refer to the works by Luk-Speck [27] and Christodoulou-Miao [34] for more recent developments in this direction. We remark that the above results on singularity formation do not apply to the physical vacuum free boundary problem. In an early important work Liu-Smoller [23] showed that the shock waves vanish at the vacuum and the singular behavior is reminiscent of the centered rarefaction waves, which suggests that the vacuum has a regularizing effect in that setting, see Liu-Yang [25].

In the context of weak solutions it has been known since the work of DiPerna [8] and subsequently Chen [1], Lions-Perthame-Souganidis [21] that the one-dimensional isentropic Euler system allows for a globally defined notion of a weak solution. We mention that this theory requires the initial density ρ_0 to be strictly positive and supported uniformly away from zero, although a formation of vacuum regions is not necessarily dynamically precluded. We note that for multi-dimensional flows there is no entropy criterion that has so far been imposed so that the uniqueness of solutions is ensured and this question is still open. Recent results of Chiodaroli-DeLellis-Kreml [2] show that the two-dimensional compressible Euler system is in fact strongly ill-posed with respect to a commonly used notion of an admissible weak solution.

In the framework of the vacuum free boundary, the existence theory depends strongly on the behavior of initial data (containing vacuum). When the sound speed c_s is smooth across the vacuum boundary, Liu-Yang [24] constructed local-in-time solutions to one-dimensional Euler system with damping and showed that c_s^2 cannot be smooth across the vacuum after a finite time. Our physical vacuum assumption (1.3) corresponds precisely to the requirement that c_s is $\frac{1}{2}$ -Hölder continuous across the initial vacuum interface. Local-in-time existence theory for the Euler system with physical vacuum was developed by Coutand-Shkoller [4, 5] and Jang-Masmoudi [15, 18], where substantial new ideas with respect to prior works were introduced in order to handle the above mentioned vacuum degeneracy. See also a subsequent work by Luo-Xin-Zeng [28]. On the other hand, if c_s is Hölder continuous but does not satisfy the physical vacuum condition, some ill-posedness result can be found in [17]. In this regime, a satisfactory theory is still far from being complete. We also mention the work by Lindblad [20] on the vacuum free boundary problem for compressible liquids when the fluid is in contact with vacuum discontinuously, namely the density is positive at the vacuum boundary. For an elegant discussion on the expansion of a gas in vacuum and a motivation for the introduction of the physical vacuum condition (1.3) see also a recent work by Serre [40].

In the presence of damping, Liu [22] constructed explicit spherically symmetric self-similar solutions that satisfy the physical vacuum condition and asymptotically converge to Barenblatt solutions of the porous media equation. Liu conjectured that solutions of the Euler system with damping in vacuum converge to solutions of the porous media equation. Huang-Marcati-Pan [14] established the conjecture in the entropy solution framework, and in a recent work [29], Luo-Zheng justified the convergence for one-dimensional Euler system with damping in the physical vacuum free boundary setting.

1.2 Methodology and plan of the paper

The central conceptual challenge in this work is a "good" formulation of the stability question. Since the underlying affine motions are time-dependent it is a priori unclear how to achieve this. It turns out that there are (at least) two approaches to this question and they both, in one guise or another, make fundamental use of scaling properties of the system.

Approach 1. Motivated by the presence of the free boundary and the existing well-posedness theories, it is tempting to pull back the problem to a fixed domain via the Lagrangian coordinates. If ζ denotes the flow map, then the Euler system in the new variables can be written in the schematic form

$$
\zeta_{tt} + F[\zeta] = 0 \tag{1.10}
$$

for some nonlinearity F. The affine motions discovered in [44] can be obtained via a *separation-ofvariables* ansatz, by setting $\zeta_{\text{affine}}(t, y) = A(t)y$ in (1.10) wherein $\mathbf{a} \equiv 0$, for a matrix $t \mapsto A(t)$ to be found. Therefore, a natural way of studying the stability of thus obtained solutions is to pass to a *renormalized* unknown $\eta(t, y) = A(t)^{-1} \zeta(t, y)$. A related approach was developed by the authors in their study of stable supernovae expansion for the compressible Euler-Poisson system [12]. While quite natural from the point of view of Lagrangian coordinates, we choose to present the details of this approach in Appendix D; at the heart of our analysis is the second approach which we shall now endeavor to describe.

Approach 2. By contrast to Approach 1 above, we pursue here the idea that affine motions can be naturally interpreted in Eulerian variables only. This will help clarify the role that symmetries play in the existence of affine motions. It will also allow for a direct parallel to some of the ideas popularized in the study of dispersive equations, most notably the mass-critical nonlinear Schrödinger equation, see Remark 2.1. To achieve all this we require a good understanding of the symmetries of the Euler system.

We consider an almost invariant action of $GL^+(3)$ on the solutions of the Euler system E_γ: for any given (ρ, \mathbf{u}) and $(A, \mathbf{a}) \in GL^+(3) \times \mathbb{R}^3$ we consider the transformation

$$
(\rho, \mathbf{u}) \mapsto (\tilde{\rho}, \tilde{\mathbf{u}}) \tag{1.11}
$$

defined by

$$
\rho(t,x) = \det A^{-1} \tilde{\rho} \left(\det A^{\frac{1-3\gamma}{6}} t, A^{-1}(x - \mathbf{a}) \right)
$$
(1.12)

$$
\mathbf{u}(t,x) = \det A^{\frac{1-3\gamma}{6}} A \tilde{\mathbf{u}} \left(\det A^{\frac{1-3\gamma}{6}} t, A^{-1}(x - \mathbf{a}) \right)
$$
(1.13)

It is straightforward to check that if (ρ, \mathbf{u}) solve (1.1a)–(1.1b), then the pair $(\tilde{\rho}, \tilde{\mathbf{u}})$ solves the following generalized Euler system:

$$
\partial_s \tilde{\rho} + \text{div}(\tilde{\rho}\tilde{\mathbf{u}}) = 0 \text{ in } \tilde{\Omega}(s); \qquad (1.14a)
$$

$$
\tilde{\rho}\left(\partial_s\tilde{\mathbf{u}} + (\tilde{\mathbf{u}} \cdot \nabla)\tilde{\mathbf{u}}\right) + \Lambda \nabla(\tilde{\rho}^{\gamma}) = 0 \text{ in } \tilde{\Omega}(s); \tag{1.14b}
$$

where

$$
\Lambda := \det A^{\frac{2}{3}} A^{-1} A^{-\top}, \ \ \tilde{\Omega}(s) = A^{-1} (\Omega(s) - \mathbf{a}), \ \ s = \det A^{\frac{1 - 3\gamma}{6}} t,\tag{1.15}
$$

where $\Omega(s) - \mathbf{a} = \{x \in \mathbb{R}^3 \mid x + \mathbf{a} \in \Omega(s)\}\$. Matrix Λ is clearly symmetric, positive definite and belongs to SL(3). Here $A^{-\top}$ denotes the transpose of the inverse of A. With respect to the original momentum equation (1.1b), a structural novelty is the presence of the matrix Λ in (1.14b). We should stress here that a related similarity transformation appears in [39].

Remark 1.6 (Why is the transformation (1.12)–(1.13) natural?). *If one attempts to find all scaling symmetries of the Euler system, a simple algebraic calculation implies that there exists a one-parameter family of such invariant transformations! However, only one of them keeps the total mass preserved and the resulting exponents in the above similarity transformation appear naturally. The role of masscriticality in our case is fundamental to the existence of self-similar solutions, as we explain below.*

In the special case when A is a **conformal** matrix (i.e. $A(\det A)^{-\frac{1}{3}} \in SO(3)$) A is the identity matrix and the transformation (1.11) is an exact invariance. This invariance simply expresses the fact that the problem possesses both a scaling and a rotational symmetry. Conformal case corresponds to arbitrary compositions of these symmetries acting on the solution space of the Euler system *E*γ.

A rich scaling freedom of compressible Euler equations is responsible for the presence of a finite parameter family of global-in-time vacuum states discovered by Sideris [44]. We now proceed to explain this in a little more detail, while the rigorous arguments can be found in Section 2. Since the transformation (1.12)–(1.13) leaves the total mass $M(\rho)(t) = \int_{\mathbb{R}^3} \rho(t, x) dx$ unchanged we refer to it as *mass-critical*. It is thus plausible to look for paths $t \mapsto (A(t), \mathbf{a}(t)) \in GL^+(3) \times \mathbb{R}^3$ that yield special solutions of the E_{γ} -system of the form (1.12)–(1.13). After rescaling time and space according to the self-similar change of variables

$$
\frac{ds}{dt} = \det A(t)^{\frac{1-3\gamma}{6}}, \ \ y = A(t)^{-1}(x - \mathbf{a}(t))
$$
\n(1.16)

we discover that the unknowns $(\tilde{\rho}(s, y), \tilde{\mathbf{u}}(s, y))$ solve a new and more complicated looking system of equations. However, after introducing a modified velocity

$$
\mathbf{U} = \tilde{\mathbf{u}} - A^{-1}A_s y - A^{-1}\mathbf{a}_s,\tag{1.17}
$$

the unknowns $(\tilde{\rho}, \mathbf{U})$ in turn solve a system with a considerably more agreeable structure, reminiscent of the original E_{γ} system. The system takes a schematic form

$$
\partial_s \tilde{\rho} + \text{div} \, (\tilde{\rho} \mathbf{U}) = 0 \tag{1.18}
$$

$$
\partial_s \mathbf{U} + (\mathbf{U} \cdot \nabla) \mathbf{U} + \mathcal{F}_1(A, A_s) \mathbf{U} + \frac{\gamma}{\gamma - 1} \Lambda \nabla (\tilde{\rho}^{\gamma - 1}) = \mathcal{F}_2(A, A_s, A_{ss}) y + \mathcal{F}_3(A, A_s, \mathbf{a}_s, \mathbf{a}_{ss}),
$$
\n(1.19)

where \mathcal{F}_i , $i = 1, 2, 3$, are some explicitly given matrix-valued smooth functions (see Section 2.1) and $\Lambda = \Lambda(A)$ is defined in (1.15). For any $\delta > 0$ we can find a special solution of (1.18)–(1.19) by setting $\mathbf{U} = 0, \frac{\gamma}{\gamma - 1} \nabla (\tilde{\rho}^{\gamma - 1}) = -\delta y$ and

$$
\mathcal{F}_2(A, A_s, A_{ss}) = -\delta \Lambda,\tag{1.20}
$$

$$
\mathcal{F}_3(A, A_s, \mathbf{a}_s, \mathbf{a}_{ss}) = 0. \tag{1.21}
$$

A simple calculation shows that the equation (1.20) is just a restatement of the ODE (1.7) in the rescaled time variable s while (2.46) yields the ODE $a_{tt} = 0$. This way we rediscover all the affine motions from [44]! This situation exhibits a certain analogy to the work of Merle-Raphaël-Szeftel [33] wherein the stable self-similar behavior of solutions to the slightly supercritical Schrödinger equation is investigated, see Remark 2.1

The question of stability of a given element of $\mathscr S$ reduces to a problem of nonlinear stability of a particular steady state of (1.18)–(1.19) where \mathcal{F}_2 is replaced by $-\delta\Lambda$, \mathcal{F}_3 is zero, and \mathcal{F}_1 is a prescribed s-dependent matrix-valued function. However, there are several caveats. First, the solutions of (1.7) blow-up in finite s -time and therefore the self-similar time variable s is not well-suited for the study of stability. This is related to the fact that the affine motions from \mathscr{S} , even though of Type I^3 , are *not* self-similar with respect to the rescaling (1.16), but instead the associated solutions $A(t)$ grow linearly in t independently of $\gamma > 1$. We change to a new time scale τ

$$
\tau \sim_{t \to \infty} \log t,\tag{1.22}
$$

which resolves the above issue and simultaneously elucidates the stabilizing effect of the expansion of the background affine motion. In the new variables the τ -dependent coefficient \mathcal{F}_1 in (1.19) gives a manifestly dissipative energy contribution in the range $\gamma \in (1, \frac{5}{3})$ and suggests a possible nonlinear stability mechanism. The logarithmic change of time-scale (1.22) is often used in the study of Type I behavior and has been crucially exploited in authors' work on stable expanding solutions for the mass-critical Euler-Poisson system [12].

To turn the above strategy to a rigorous proof, we need a good well-posedness theory. Due to the physical vacuum condition (1.3) the initial enthalpy is not smooth across the vacuum interface and this makes the question of (even) local-in-time existence of solutions rather challenging. Only recently, the problem of local-in-time well-posedness in the presence of the physical vacuum condition was fully solved [5, 18]. At the core of both approaches is the use of Lagrangian coordinates which is particularly convenient as the problem is automatically pulled back to a fixed domain. Our work draws especially on the approach from [18] where a high-order energy method relying on only spatial differential operators is developed.

In our case we work with a Lagrangian formulation of (1.18) – (1.19) with respect to the modified velocity U, where the basic unknown is the flow map defined by the ODE

$$
\dot{\eta}(t, y) = \mathbf{U}(t, \eta(t, y)) \tag{1.23}
$$

such that $\eta \equiv y$ represents the expanding background solution. While the continuity equation (1.18) retains the same form as (1.1a), the presence of the matrix Λ encoding the geometry of the background expanding affine motion in the momentum equation (1.19) introduces two fundamental difficulties compared to the work [18]:

- (i) Since the curl of the modified pressure term $\Lambda \nabla (\tilde{\rho}^{\gamma-1})$ *does not* vanish, the curl of the velocity field *does not* satisfy a simple transport equation, a crucial tool in implementing both strategies in [5, 18];
- (ii) It is a priori not obvious how to extract a positive definite energy contribution from (1.19).

To handle the first issue we note that an adapted Λ -curl operator defined via

 $[\text{curl}_\Lambda \mathbf{F}]_i := \epsilon_{ijk} \Lambda_j^s \mathbf{F}^k$, $(\epsilon_{ijk}$ is the usual permutation symbol)

annihilates the Λ -gradient $\Lambda \nabla$.⁴ Our key observation is that the Λ -curl of U satisfies a transport equation which at the top order decouples from the rest of the dynamics, allowing us to obtain "good"

 $3By$ definition, Type I self-similar solutions approach an equilibrium exponentially fast in the logarithmic time variable (1.22). We refer to the review article [9] for a detailed discussion of this terminology.

⁴One could introduce a more geometric language so that curl_Λ and $\Lambda \nabla$ are "natural" operators with respect to a given metric structure, but for the sake of conciseness we choose not to.

estimates for the (Lagrangian pull-back of) Λ-curl of U. The perturbation around the steady state is given by

$$
\theta(\tau, y) := \eta(\tau, y) - y.
$$

A simplified schematic form of the equation for θ is

$$
w^{\alpha}e^{\mu(\gamma)\tau}(\partial_{\tau\tau}\theta + c\partial_{\tau}\theta) + (\Lambda\nabla)^{*}\mathcal{F}(w^{1+\alpha}D\theta) = 0 \text{ in } \Omega = B_1(\mathbf{0}),
$$
 (1.24)

where $(\Lambda \nabla)^*$ is the Lagrangian pull-back of $\Lambda \nabla$, $\mathcal{F} : \mathbb{M}^{3 \times 3} \to \mathbb{R}$ is a smooth function, $\mu(\gamma)$, $c, \alpha =$ $\alpha(\gamma) > 0$ given constants, and $w : \Omega \to \mathbb{R}$ is the equilibrium fluid enthalpy satisfying $w^{\alpha} = \rho_A(0)$ (initial density of the background expanding solution). See (2.64) for a precise version. Starting from (1.24) we derive a high-order energy by integrating the equation against spatial derivatives of $\Lambda^{-1}\theta_{\tau}$. To resolve the issue (ii) mentioned above we crucially use the fact that Λ is a real symmetric non-degenerate matrix depending only on τ . If ∂ denotes a generic high-order differential operator, we apply it to the Λ-gradient term in (1.24) and use the Leibniz rule to isolate the leading order term. After a suitable anti-symmetrization we can extract a coercive energy contribution at the expense of creating a new top order term containing $curl_A^*(\partial\theta)$ where $curl_A^*$ is the Lagrangian pull-back of the operator curl_Λ. This procedure relies on a delicate algebraic structure of the top-order term and requires diagonalizing the matrix Λ , see Lemma 4.3. The curl^{*}_{Λ} in turn satisfies a "good" transport equation and we are able to estimate it in terms of our natural high-order energy.

Going from a local well-posedness to a global theory for nonlinear equations is often nontrivial and it may not be true in general. For our problem, as hinted earlier, the background expanding affine motion stabilizes the dynamics for a certain range of γ in a well-designed energy space resolving the above difficulties (i) and (ii). The leading order energy contribution associated with ∂_{τ} -derivatives in (1.24) takes the following schematic form

$$
\frac{1}{2}\frac{d}{d\tau}\left(e^{\mu(\gamma)\tau}\|w^{\frac{\alpha}{2}}\sqrt{\Lambda^{-1}}\partial_{\tau}\theta\|_{L^{2}(\Omega)}^{2}\right)+(c-\frac{1}{2}\mu(\gamma))e^{\mu(\gamma)\tau}\|w^{\frac{\alpha}{2}}\sqrt{\Lambda^{-1}}\partial_{\tau}\theta\|_{L^{2}(\Omega)}^{2}.\tag{1.25}
$$

As shown in Proposition 4.6, the analogue of the requirement $c - \frac{1}{2}\mu(\gamma) \ge 0$ amounts to the condition $\gamma \leq \frac{5}{3}$ which is assumed in Theorem 1.1. An exponentially growing term $e^{\mu(\gamma)\tau}$ in (1.24) acts as a a medium for stabilization and we make crucial use of it in closing the high-order estimates.

After introducing a high-order weighted norm S^N , we prove an energy inequality of the form

$$
S^{N}(\tau) \leq C_0 + C \int_0^{\tau} e^{-\mu_* \tau'} S^{N}(\tau') d\tau' \text{ for some } \mu_* > 0.
$$
 (1.26)

The constant $C_0 > 0$ depends only on the initial data while the exponentially decaying factor inside the integral reflects the presence of exponential-in- τ weights in (1.25). The basic technical tool in the proof of (1.26) is the use of Hardy-Sobolev embeddings between weighted Sobolev spaces explained in Appendix C. The global existence for sufficiently small perturbed initial data follows from a standard continuity argument applied to (1.26).

We stress that our global-in-time result encompasses the case of the monatomic gas $\gamma = \frac{5}{3}$ when the effective "damping" coefficient $c - \frac{1}{2}\mu(\gamma)$ in (1.25) vanishes. This shows that "damping" is not the fundamental driving force behind stability. Instead, the important point is that there is no antidamping, which should be contrasted to the usual dissipation mechanism of a damped wave equation. The key stabilizing effect leading to the global-in-time result is the exponentially growing term $e^{\mu(\gamma)\tau}$ in (1.24). Its effect is felt through the exponentially decaying term inside the integral on the right-hand side of (1.26). Such a mechanism gives a new strategy to achieve the global theory.

Last but not least, it is important for our argument to use only spatial differential operators that do not destroy the structure of the time weight in (1.24) . If we use vector fields containing τ -derivatives, we encounter unpleasant commutator terms stemming from the exponential weights in (1.24) . By employing the approach of Jang-Masmoudi [18] with a significant refinement as explained above, we are able to control the high-order energy of spatial derivatives with powers of w as weights *uniformlyin-time*, see the definition of our high-order norm (2.77). The key weighting pattern [18] is that every normal derivative with respect to the boundary $\partial\Omega = \mathbb{S}^2$ requires adding a multiple of the fluid enthalpy w in the weight. If not for this weighting structure, it would be impossible to close the nonlinear energy estimates, see Remark 4.5. This is consistent with the equation (1.24) and such weighted spaces relate to the usual Sobolev spaces via Hardy-Sobolev embeddings, as laid out in Appendix C.

We conclude the introduction with the plan for the rest of the paper. In Section 2 we explain in detail how the affine motions relate to the symmetries of E_{γ} -system, Section 2.1, and we then formulate the stability problem and state the main result in Lagrangian coordinates, Sections 2.2–2.4. Vorticity bounds are explained in Section 3, main energy estimates in Section 4, and proof of the main theorem in Section 5. In Appendix A we summarize the properties of the background solution using the results from [44]. Appendix B contains the basic properties of commutators between various differential operators used in the paper, while Appendix C contains the statements of frequently used Hardy-Sobolev embeddings. In Appendix D, we give an alternative and equivalent formulation of the problem starting from the Lagrangian coordinates.

2 Formulation and Main result

2.1 A self-similar rescaling

Motivated by an (almost) invariant transformation (1.12) – (1.13) we are seeking for a path

$$
\mathbb{R}_{+} \ni t \mapsto (A(t), \mathbf{a}(t)) \in \mathrm{GL}^{+}(3) \times \mathbb{R}^{3}
$$

such that the unknowns $(\tilde{\rho}, \tilde{\mathbf{u}})$ defined by

$$
\rho(t,x) = \det A(s)^{-1} \tilde{\rho}(s,y),\tag{2.27}
$$

$$
\mathbf{u}(t,x) = \det A(s)^{\frac{1-3\gamma}{6}} A(s)\tilde{\mathbf{u}}(s,y),
$$
\n(2.28)

solve the Euler system E_{γ} . Here the new time and space coordinates s and y are given by

$$
\frac{ds}{dt} = \frac{1}{\det A(t)^{\frac{3\gamma - 1}{6}}}, \ \ y = A(t)^{-1}(x - \mathbf{a}(t)), \tag{2.29}
$$

motivated by the transformation (1.12)–(1.13). Introducing the notation

$$
\mu(s) := \det A(s)^{\frac{1}{3}},\tag{2.30}
$$

a simple application of the chain rule transforms the equations (1.1a)–(1.1b) into

$$
\tilde{\rho}_s - 3\frac{\mu_s}{\mu}\tilde{\rho} - \left(A^{-1}A_sy + A^{-1}\mathbf{a}_s\right) \cdot \nabla \tilde{\rho} + \text{div}(\tilde{\rho}\tilde{\mathbf{u}}) = 0,
$$
\n(2.31)

$$
\partial_s \tilde{\mathbf{u}} - \frac{3\gamma - 1}{2} \frac{\mu_s}{\mu} \tilde{\mathbf{u}} + A^{-1} A_s \tilde{\mathbf{u}} + (\tilde{\mathbf{u}} \cdot \nabla) \tilde{\mathbf{u}} - \left(A^{-1} A_s y + A^{-1} \mathbf{a}_s \right) \cdot \nabla \tilde{\mathbf{u}} + \frac{\gamma}{\gamma - 1} \Lambda \nabla (\tilde{\rho}^{\gamma - 1}) = 0,
$$
\n(2.32)

where we recall that $\Lambda(s) = \det A(s)^{\frac{2}{3}} A(s)^{-1} A(s)^{-\top}$. A structural miracle of the new system (2.31)– (2.32) is that it can be recast in a simpler form, close to the original Euler system E_y . To that end we introduce a modified velocity

$$
\mathbf{U}(s, y) := \tilde{\mathbf{u}}(s, y) + B(s)y + \mathbf{b}(s),
$$
\n(2.33)

where

$$
B(s) := -A^{-1}A_s, \ \mathbf{b}(s) := -A^{-1}\mathbf{a}_s,\tag{2.34}
$$

and note that

$$
\text{div}\,\left(A^{-1}A_s y\right) = \text{Tr}\,\left(A^{-1}A_s\right) = \partial_s \det A(s) \det A(s)^{-1} = \frac{3\mu_s}{\mu}.\tag{2.35}
$$

Then the system (2.31) – (2.32) can be rewritten as

$$
\partial_s \tilde{\rho} + \text{div} \left(\tilde{\rho} \mathbf{U} \right) = 0 \tag{2.36}
$$
\n
$$
\partial_s \mathbf{U} + (\mathbf{U} \cdot \nabla) \mathbf{U} + \left(-\frac{3\gamma - 1}{2} \frac{\mu_s}{\mu} \mathbf{Id} - 2B(s) \right) \mathbf{U} + \frac{\gamma}{\gamma - 1} \Lambda \nabla (\tilde{\rho}^{\gamma - 1})
$$
\n
$$
= \left[B_s - \left(\frac{3\gamma - 1}{2} \frac{\mu_s}{\mu} \mathbf{Id} + B \right) B \right] y + \mathbf{b}_s - \left(\frac{3\gamma - 1}{2} \frac{\mu_s}{\mu} \mathbf{Id} + B \right) \mathbf{b}, \tag{2.37}
$$

where we have used (2.35) in verifying (2.36).

For any given $\delta \in \mathbb{R}$ we look for solutions $(\mathbf{U}, \tilde{\rho}, B)$ of (2.36)–(2.37) of the form

$$
\mathbf{U} = \mathbf{0} \quad \text{in } \Omega,\tag{2.38}
$$

$$
B_s - \left(\frac{3\gamma - 1}{2}\frac{\mu_s}{\mu}\mathbf{Id} + B\right)B = -\delta\Lambda, \ \ s \ge 0,
$$
 (2.39)

$$
\frac{\gamma}{\gamma - 1} \nabla (\rho^{\gamma - 1}) = -\delta y \quad \text{in } \Omega,
$$
\n(2.40)

$$
\mathbf{b}_s - \left(\frac{3\gamma - 1}{2}\frac{\mu_s}{\mu}\,\mathbf{Id} + B\right)\mathbf{b} = 0, \ \ s \ge 0,\tag{2.41}
$$

where we recall that $\Omega = B_1(0)$ is the unit ball in \mathbb{R}^3 . Equation (2.40) is easy to solve and leads to the enthalpy profile $w := \tilde{\rho}^{\gamma - 1}$ given by

$$
w(y) = \frac{\delta(\gamma - 1)}{2\gamma} \left(1 - |y|^2\right), \quad y \in \Omega,
$$
\n(2.42)

where we consider only the case $\delta > 0$ to ensure that the physical vacuum condition (1.3) holds true. Equation (2.38) gives

$$
\tilde{\mathbf{u}}(s, y) = -By - \mathbf{b} = A^{-1}A_s y + A^{-1} \mathbf{a}_s.
$$
\n(2.43)

Interestingly, equation (2.39) does not posses globally defined solutions in the s-variable; instead we must rescale back to the original time variable t. Recalling that $\frac{dt}{ds} = \det A^{\frac{3\gamma - 1}{6}}$ and (2.34) we can re-express (2.39) as a differential equation for $A(t)$. Note that

$$
B_s = -\det A^{\frac{3\gamma - 1}{6}} \partial_t (A^{-1} A_t \det A^{\frac{3\gamma - 1}{6}})
$$

= $\det A^{\frac{3\gamma - 1}{3}} A^{-1} A_t A^{-1} A_t - \det A^{\frac{3\gamma - 1}{3}} A^{-1} A_{tt} + \frac{3\gamma - 1}{6} \det A^{\frac{3\gamma - 1}{3}} \frac{\partial_t \det A}{\det A}$
= $B^2 + \frac{3\gamma - 1}{2} \frac{\mu_s}{\mu} B - \det A^{\frac{3\gamma - 1}{3}} A_{tt}.$

Therefore, from (2.39) and the definition of Λ it follows that

$$
-\det A^{\frac{3\gamma-1}{3}} A^{-1} A_{tt} = -\delta \det A^{\frac{2}{3}} A^{-1} A^{-\top},
$$

which is equivalent to

$$
A_{tt} = \delta \det A^{1-\gamma} A^{-\top}.
$$
 (2.44)

We have recovered (1.7) (modulo a harmless parameter $\delta > 0$). This is exactly the ODE describing the affine motion of a fluid blob from [44]. By Theorem 6 [44] for any $\gamma > 1, \delta > 0$ there exists a global-in-t solution to (2.44) . By an analogous calculation equation (2.41) reduces to

$$
\mathbf{a}_{tt} = 0 \tag{2.45}
$$

in the original variables and therefore

$$
\mathbf{a}(t) = \mathbf{a}_0 + t\mathbf{a}_1,\tag{2.46}
$$

for any $a_0, a_1 \in \mathbb{R}^3$. The family of solutions $(U, \tilde{\rho}, B, a)$ with $a(t) = 0$ comprises all the affine motions constructed by Sideris in [44]. Using the translational invariance of the problem, we allow for the center $t \mapsto \mathbf{a}(t)$ of the expanding ellipsoidal solution of the original Euler system to move. We observe that the space of all such solutions denoted by $\mathscr S$ forms a finite-parameter family parametrized by the choice of the parameter δ and the initial conditions for the ODEs (2.45)–(2.46):

$$
(A_0, A_1, \delta, \mathbf{a}_0, \mathbf{a}_1) \in \mathrm{GL}^+(3) \times \mathbb{M}^{3 \times 3} \times \mathbb{R}_+ \times \mathbb{R}^3 \times \mathbb{R}^3.
$$

Here the parameter $\delta > 0$ dictates the total mass of the gas. From (2.42), one can easily see that the total mass can be viewed as a function of δ for any fixed $\gamma > 1$. In particular, since $\delta > 0$ is arbitrary, any positive finite total mass is allowed for the expansion of the gas. Vector a_1 is the total momentum per unit mass, while $a_0 \times a_1$ corresponds to the total angular momentum per unit mass, both being conserved quantities of the Euler flow E_{γ} .

Remark 2.1. *In [33] stable self-similar behavior of solutions to the slightly supercritical Schrodinger ¨ equation is investigated. There the analogue of the key transformation* (2.33) *leads to a reformulation of the nonlinear Schrodinger equation, analogous to ¨* (2.36)*–*(2.37)*. In the setting of [33] explicit formula for self-similar solutions is not available, but the new formulation is ideally suited for the study of their existence and stability. The modified velocity* **U** *defined by* (2.33) *differs from* \tilde{u} *by an affine transformation in analogy to the quantity* P_b *from [33] which arises by a suitable "conformal" change of variables honoring the symmetries of the nonlinear Schrodinger equation. ¨*

Remark 2.2. Although not explicitly used in this paper, we remark that in the special case $\gamma = \frac{5}{3}$ *the Euler system E*^γ *is pseudo-conformally invariant. To the best of our knowledge, the first explicit mention of this invariance was made by Serre [39] thus establishing an analogy to the mass-critical nonlinear Schrödinger equation, known to possess a pseudo-conformal invariance.*

2.2 Lagrangian formulation

Let us fix an element of $\mathscr S$ given by the quadruple $(\rho_A, \mathbf{u}_A, A, \mathbf{a})$, where (ρ_A, \mathbf{u}_A) are given by (1.5)– (1.6) and $(A, a) \in GL^+(3) \times \mathbb{R}^3$ are given as solutions of (1.7)–(1.9). Note that these solutions are fixed by a choice of free parameters $(A_0, A_1, \delta, a_0, a_1)$ as described above and they are in one-to-one correspondence with solutions to (2.36)–(2.37) given by the quadruple

$$
\mathbf{U} = 0, \ \ w(y) = \frac{\delta(\gamma - 1)}{2\gamma} (1 - |y|^2), \ \ B(s) = -A^{-1} A_s, \ \ \mathbf{a}(t) = \mathbf{a}_0 + t\mathbf{a}_1,\tag{2.47}
$$

where we recall (2.29) for the definition of s and the relationship

$$
\tilde{\rho}=w^{\alpha}, \ \alpha:=\frac{1}{\gamma-1}.
$$

Upon decomposing

$$
A = \mu O, \ \mu = \det A^{\frac{1}{3}}, \ O \in SL(3)
$$
 (2.48)

we see that

$$
B(s) = -\mu^{-1}O^{-1}(\mu_s O + \mu O_s) = -\frac{\mu_s}{\mu} \text{Id} - \Gamma, \ \Gamma := O^{-1}O_s.
$$
 (2.49)

To study the stability of the steady state given (2.47) we shall reformulate the problem in Lagrangian coordinates. Letting

$$
\Omega:=B_1(\mathbf{0})
$$

be the unit ball in \mathbb{R}^3 , we introduce $\eta : \Omega \to \tilde{\Omega}(s)$ as a flow map associated with the renormalized velocity field U :

$$
\eta_s(s, y) = \mathbf{U}(s, \eta(s, y)),\tag{2.50}
$$

$$
\eta(0, y) = \eta_0(y), \tag{2.51}
$$

where $\eta_0 : \Omega \to \tilde{\Omega}(0)$ is a sufficiently smooth diffeomorphism to be specified later. To pull-back (2.36)-(2.37) to the fixed domain Ω , we introduce the notation

$$
\mathscr{A} := [D\eta]^{-1} \quad \text{(Inverse of the Jacobian matrix)},\tag{2.52}
$$

$$
\mathscr{J} := \det[D\eta] \quad \text{(Jacobian determinant)}, \tag{2.53}
$$

$$
f := \tilde{\rho} \circ \eta \quad \text{(Lagrangian density)},\tag{2.54}
$$

$$
\mathbf{V} := \mathbf{U} \circ \eta \quad \text{(Lagrangian modified velocity)}, \tag{2.55}
$$

$$
a := \mathscr{J} \mathscr{A} \quad \text{(Cofactor matrix)}.
$$
\n
$$
(2.56)
$$

From $\mathscr{A}[D\eta] = \text{Id}$, one can obtain the differentiation formula for \mathscr{A} and \mathscr{J} :

$$
\partial \mathscr{A}_i^k = -\mathscr{A}_\ell^k \partial \eta^\ell,_{s} \mathscr{A}_i^s ; \quad \partial \mathscr{J} = \mathscr{J} \mathscr{A}_\ell^s \partial \eta^\ell,_{s}
$$

for $\partial = \partial_{\tau}$ or $\partial = \partial_i$, $i = 1, 2, 3$. Here we have used the Einstein summation convention and the notation F_{ik} to denote the k^{th} partial derivative of F. Both expressions will be used throughout the paper.

It is well-known [5, 18] that the continuity equation (2.36) reduces to the relationship

$$
f\mathscr{J}=f_0\mathscr{J}_0.
$$

We choose η_0 such that $w = (f_0 \mathscr{J}_0)^{\gamma-1}$ where w is given in (2.42). For given initial density function ρ_0 so that ρ_0/ρ_A is smooth, the existence of such η_0 follows from the Dacorogna-Moser theorem [6]. As a consequence the Lagrangian density can be expressed as

$$
f = w^{\alpha} \mathscr{J}^{-1} \tag{2.57}
$$

where we recall that $\alpha = \frac{1}{\gamma - 1}$. We remark that this specific choice of η_0 (gauge fixing) is important for our analysis; η_0 being the identity map corresponds to the background expanding solution, η_0 measures how the initial density is distributed with respect to the background equilibrium, and hence it provides a natural framework for stability analysis around the identity map.

Using (2.49) equation (2.37) takes the following form in Lagrangian coordinates:

$$
\eta_{ss} + \frac{5 - 3\gamma}{2} \frac{\mu_s}{\mu} \eta_s + 2\Gamma \eta_s + \delta \Lambda \eta + \frac{\gamma}{\gamma - 1} \Lambda \mathscr{A}^\top \nabla (f^{\gamma - 1}) = 0 \tag{2.58}
$$

which can be expressed in the component form [5, 18]

$$
w^{\alpha} \left(\partial_{ss} \eta_i - \frac{5 - 3\gamma}{2} b \partial_s \eta_i + 2 \Gamma_{ij} \partial_s \eta_j + \delta \Lambda_{ij} \eta_j \right) + \Lambda_{ij} (w^{1+\alpha} \mathscr{A}_j^k \mathscr{J}^{-\frac{1}{\alpha}}),_k = 0. \tag{2.59}
$$

Since in the s-time variable the matrix $B(s)$ blows up in finite time, we introduce a new time variable τ via

$$
\frac{d\tau}{ds} = \det A^{\frac{3\gamma - 3}{6}} = \mu^{\frac{3\gamma - 3}{2}} \text{ or equivalently } \frac{d\tau}{dt} = \frac{1}{\mu}.
$$
 (2.60)

Since μ grows linearly in t by results from [44] (see Lemma A.1), the new time τ grows like log t as $t \to \infty$ and therefore corresponds to a logarithmic time-scale with respect to the original time variable $t.$ Equation (2.58) takes the form

$$
\mu^{3\gamma - 4} \left(\mu \eta_{\tau \tau} + \mu_{\tau} \eta_{\tau} + 2\mu \Gamma^* \eta_{\tau} \right) + \delta \Lambda \eta + \frac{\gamma}{\gamma - 1} \Lambda \mathscr{A}^\top \nabla (f^{\gamma - 1}) = 0, \tag{2.61}
$$

where

$$
\Gamma^* = O^{-1}O_\tau.
$$

In coordinates, for any $i \in \{1, 2, 3\}$ it reads

$$
w^{\alpha}\mu^{3\gamma-4}\left(\mu\partial_{\tau\tau}\eta_i + \mu_{\tau}\partial_{\tau}\eta_i + 2\mu\Gamma^*_{ij}\partial_{\tau}\eta_j\right) + \delta w^{\alpha}\Lambda_{i\ell}\eta_{\ell} + \left(w^{1+\alpha}\Lambda_{ij}\mathscr{A}^k_j\mathscr{J}^{-\frac{1}{\alpha}}\right),_k = 0. \tag{2.62}
$$

Defining the perturbation

$$
\Theta(\tau, y) := \eta(\tau, y) - y \tag{2.63}
$$

equation (2.62) takes the form:

$$
w^{\alpha} \mu^{3\gamma - 3} \left(\partial_{\tau \tau} \theta_i + \frac{\mu_{\tau}}{\mu} \partial_{\tau} \theta_i + 2 \Gamma_{ij}^* \partial_{\tau} \theta_j \right) + \delta w^{\alpha} \Lambda_{i\ell} \theta_{\ell} + \left(w^{1 + \alpha} \Lambda_{ij} \left(\mathscr{A}_j^k \mathscr{J}^{-\frac{1}{\alpha}} - \delta_j^k \right) \right), k = 0,
$$
\n(2.64)

equipped with the initial conditions

$$
\theta(0, y) = \theta_0(y), \ \ \theta_\tau(0, y) = \mathbf{V}(0, y) = \mathbf{V}_0(y), \ \ (y \in \Omega = B_1(\mathbf{0})). \tag{2.65}
$$

We will use both $\theta_{\tau} = \partial_{\tau} \theta$ and V to denote the velocity field throughout the paper.

Remark 2.3. *The Lagrangian formulation* (2.50)*–*(2.51) *uses the modified velocity* U *to define the flow map* η ; this is a coherent strategy as we identified **U** as a natural perturbed velocity around the set $\mathscr S$ *of Sideris' affine motions in Eulerian coordinates. We remark that the order of this derivation can be reversed. For simplicity assume that the center of mass* a *is fixed to be* 0 *like in [44]. We can start with the Lagrangian formulation of the E*γ*-system using the original velocity* u *and a flow map* ζ*, wherein the affine motions are identified as solutions of the form* $\zeta(t, y) = A(t)y$ *as in* (1.4)*, wherein* $\mathbf{a} \equiv 0$ *. By suitably conjugating the flow map* ζ *by* $A(t)^{-1}$ *and introducing the time rescaling* (2.60) *we can arrive at* (2.61) *and subsequently* (2.64)*. See Appendix D for such a derivation. However, we have decided on presenting the formulation derived from Eulerian coordinates as it links the symmetries of the E*γ*-system to the existence of affine motions and provides a systematic approach to the question of global existence of solutions in their vicinity without any symmetry assumptions.*

2.3 Notation

Lie derivative of the flow map and modified vorticity. For vector-fields $\mathbf{F}: \Omega \to \mathbb{R}^3$, we introduce the Lie derivatives: full gradient along the flow map η

$$
[\nabla_{\eta} \mathbf{F}]_r^i := \mathscr{A}_r^s \mathbf{F}^i, \tag{2.66}
$$

the divergence

$$
\operatorname{div}_{\eta} \mathbf{F} := \mathscr{A}_{\ell}^{s} \mathbf{F}^{\ell}, \qquad (2.67)
$$

the modified curl-operator

$$
\left[\operatorname{curl}_{\Lambda \mathscr{A}} \mathbf{F}\right]^i := \epsilon_{ijk} (\Lambda \mathscr{A})^s_j \mathbf{F}^k, s = \epsilon_{ijk} \Lambda_{jm} \mathscr{A}^s_m \mathbf{F}^k, s \,, \tag{2.68}
$$

(where ϵ_{ijk} is the standard permutation symbol) and the anti-symmetric curl matrix

$$
\left[\text{Curl}_{\Lambda\mathscr{A}}\mathbf{F}\right]^{i}_{j} := \Lambda_{jm}\mathscr{A}^{s}_{m}\mathbf{F}^{i}_{\ ,s} - \Lambda_{im}\mathscr{A}^{s}_{m}\mathbf{F}^{j}_{\ ,s}\tag{2.69}
$$

We will also use $D \mathbf{F}$, div \mathbf{F} , curl \mathbf{F} to denote its full gradient, its divergence, and its curl:

$$
[D \mathbf{F}]^i_j = \mathbf{F}^i, j \; ; \; \text{ div } \mathbf{F} = \mathbf{F}^\ell, \ell \; ; \; \text{ [curl } \mathbf{F}]^i = \epsilon_{ijk} \mathbf{F}^k, j \; , \; i, j = 1, 2, 3.
$$

Function spaces. For any given integer $k \in \mathbb{N}$, a function $f : \Omega \to \mathbb{R}$ and any smooth non-negative function $\varphi : \Omega \to \mathbb{R}^+$ we introduce the notation

$$
||f||_{k,\varphi}^2 := \int_{\Omega} \varphi w^k |f(y)|^2 dy.
$$
 (2.70)

For vector-fields $\mathbf{F}: \Omega \to \mathbb{R}^3$ or 2-tensors $\mathbf{T}: \Omega \to \mathbb{M}^{3 \times 3}$ we analogously define

$$
\|\mathbf{F}\|_{k,\varphi}^2 := \sum_{i=1}^3 \|\mathbf{F}_i\|_{k,\varphi}^2, \ \ \|\mathbf{T}\|_{k,\varphi}^2 := \sum_{i,j=1}^3 \|\mathbf{T}_{ij}\|_{k,\varphi}^2. \tag{2.71}
$$

In this paper we will always choose the weight function φ to be either ψ or $1 - \psi$ where $\psi : B_1(\mathbf{0}) \to$ [0, 1] is a smooth cut-off function satisfying

$$
\psi = 0
$$
 on $B_{\frac{1}{4}}(\mathbf{0})$ and $\psi = 1$ on $B_1(\mathbf{0}) \setminus B_{\frac{3}{4}}(\mathbf{0})$. (2.72)

It will allow us to localize the estimates to two different regions of the domain Ω .

Angular gradient. For any $f \in H^1(\Omega)$ we define the *angular gradient* ∇ as the projection of the usual gradient onto the plane tangent to the sphere of constant (t, r) . In other words

$$
\nabla f(y) = \nabla f(y) - \frac{y}{r} \left(\frac{y}{r} \cdot \nabla\right) f(y) = \nabla f(y) - \frac{y}{r} \partial_r f(y),\tag{2.73}
$$

where $\partial_r := \frac{y}{r} \cdot \nabla$ and $r = |y|$. For $i \in \{1, 2, 3\}$ we can evaluate the components of ∇_i using (2.73):

$$
\nabla_i = \frac{y_j (y_j \partial_i - y_i \partial_j)}{r^2}.
$$
\n(2.74)

The angular gradients (2.74) are the natural, geometric tangential gradients near the boundary of our domain and they will be crucially used in the analysis. For the commutation rules between ∇ and other vector fields, see Lemma B.1.

For any multi-index $\beta = (\beta_1, \beta_2, \beta_3) \in \mathbb{Z}_{\geq 0}^3$ and any $a \in \mathbb{Z}_{\geq 0}$ we introduce the following differential operator

$$
\partial_r^a \nabla^\beta := \partial_r^a \nabla_1^{\beta_1} \nabla_2^{\beta_2} \nabla_3^{\beta_3}.
$$
\n(2.75)

Similarly for any multi-index $\nu = (\nu_1, \nu_2, \nu_3) \in \mathbb{Z}_{\geq 0}^3$ we denote

$$
\partial^{\nu} := \partial_{y_1}^{\nu_1} \partial_{y_2}^{\nu_2} \partial_{y_3}^{\nu_3},\tag{2.76}
$$

where ∂_{y_i} , $i = 1, 2, 3$ denote the usual Cartesian derivatives.

2.4 Main theorem in Lagrangian coordinates

To state the main theorem in our modified Lagrangian coordinates, we first introduce the high-order weighted Sobolev norm that measures the size of the deviation θ . For any $N \in \mathbb{N}$, let

$$
\mathcal{S}^{N}(\theta, \mathbf{V})(\tau) = \mathcal{S}^{N}(\tau)
$$
\n
$$
:= \sum_{a+|\beta| \leq N} \sup_{0 \leq \tau' \leq \tau} \left\{ \mu^{3\gamma - 3} \left\| \partial_{\tau}^a \nabla^{\beta} \mathbf{V} \right\|_{a+\alpha,\psi}^2 + \left\| \partial_{\tau}^a \nabla^{\beta} \theta \right\|_{a+\alpha,\psi}^2 + \left\| \nabla_{\eta} \partial_{\tau}^a \nabla^{\beta} \theta \right\|_{a+\alpha+1,\psi}^2 + \left\| \text{div}_{\eta} \partial_{\tau}^a \nabla^{\beta} \theta \right\|_{a+\alpha+1,\psi}^2 + \left\| \text{div}_{\eta} \partial_{\tau}^a \nabla^{\beta} \theta \right\|_{a+\alpha+1,\psi}^2 + \sum_{|\nu| \leq N} \sup_{0 \leq \tau' \leq \tau} \left\{ \mu^{3\gamma - 3} \left\| \partial^{\nu} \mathbf{V} \right\|_{\alpha,1-\psi}^2 + \left\| \partial^{\nu} \theta \right\|_{\alpha,1-\psi}^2 + \left\| \nabla_{\eta} \partial^{\nu} \theta \right\|_{\alpha+1,1-\psi}^2 + \left\| \text{div}_{\eta} \partial^{\nu} \theta \right\|_{\alpha+1,1-\psi}^2 \right\}
$$
\n(2.77)

Additionally we introduce another high-order quantity measuring the modified vorticity of V which is a priori not controlled by the norm $\mathcal{S}^N(\tau)$:

$$
\mathcal{B}^{N}[\mathbf{V}](\tau) := \sum_{a+|\beta| \leq N} \sup_{0 \leq \tau' \leq \tau} \left\| \operatorname{Curl}_{\Lambda \mathscr{A}} \partial_{\tau}^{a} \nabla^{\beta} \mathbf{V} \right\|_{a+\alpha+1,\psi}^{2} + \sum_{|\nu| \leq N} \sup_{0 \leq \tau' \leq \tau} \left\| \operatorname{Curl}_{\Lambda \mathscr{A}} \partial^{\nu} \mathbf{V} \right\|_{\alpha+1,1-\psi}^{2}.
$$
\n(2.78)

We also define the quantity $\mathcal{B}^{N}[\theta]$ analogously, with θ instead of V in the definition (2.78).

We are now ready to state our main result.

Theorem 2.4 (Nonlinear stability of affine motions). Assume that $\gamma \in (1, \frac{5}{3}]$ or equivalently $\alpha \in$ $\left[\frac{3}{2}, \infty\right)$ *. Let* $N \geq 2\lceil \alpha \rceil + 12$ *be fixed. Let* $(A_0, A_1, \delta, \mathbf{a}_0, \mathbf{a}_1) \in \mathrm{GL}^+(3) \times \mathbb{M}^{3 \times 3} \times \mathbb{R}_+ \times \mathbb{R}^3 \times \mathbb{R}^3$ *be a* given quintuple parametrizing an affine motion $A(\tau)$ from the set $\mathscr S$ and let

$$
\mu(\tau) = (\det A(\tau))^{\frac{1}{3}}, \ \Lambda(\tau) = \det A(\tau)^{\frac{2}{3}} A(\tau)^{-1} A(\tau)^{-\top}, \n\mu_0 := \frac{3\gamma - 3}{2} \mu_1, \ \mu_1 := \lim_{\tau \to \infty} \frac{\mu_\tau(\tau)}{\mu(\tau)}.
$$
\n(2.79)

Then there exists an $\varepsilon_0 > 0$ *such that for any* $0 \leq \varepsilon \leq \varepsilon_0$ and (θ_0, V_0) *satisfying* $S^N(\theta_0, V_0)$ + $\mathcal{B}^N(\mathbf{V}_0) \leq \varepsilon$, there exists a global-in-time solution to the initial value problem (2.64)–(2.65) and a *constant* C > 0 *such that*

$$
S^{N}(\mathbf{\theta}, \mathbf{V})(\tau) \le C\varepsilon, \ \ 0 \le \tau < \infty,\tag{2.80}
$$

and

$$
\mathcal{B}^{N}[\mathbf{V}](\tau) \leq \begin{cases} C\varepsilon e^{-2\mu_{0}\tau} & \text{if } 1 < \gamma < \frac{5}{3} \\ C\varepsilon(1+\tau^{2})e^{-2\mu_{0}\tau} & \text{if } \gamma = \frac{5}{3} \end{cases}, \quad 0 \leq \tau < \infty.
$$
 (2.81)

Furthermore, there exists a constant $C_* > 0$ *such that*

$$
\sum_{|\beta| \le N} \|\partial_r^a \nabla^\beta \mathbf{V}\|_{1+\alpha+a,\psi} + \sum_{|\nu| \le N} \|\partial^\nu \mathbf{V}\|_{1+\alpha,1-\psi} \le C_* \sqrt{\varepsilon} e^{-\mu_0 \tau},\tag{2.82}
$$

and a τ *-independent asymptotic attractor* θ[∞] *such that*

 \overline{a}

$$
\lim_{\tau \to \infty} \left(\sum_{a+|\beta| \le N} \left\| \partial_r^a \nabla^\beta \theta(\tau, \cdot) - \partial_r^a \nabla^\beta \theta_\infty(\cdot) \right\|_{a+\alpha,\psi} + \sum_{|\nu| \le N} \|\partial^\nu \theta(\tau, \cdot) - \partial^\nu \theta_\infty(\cdot) \|_{\alpha, 1-\psi} \right) = 0.
$$
\n(2.83)

Remark 2.5. *The exponential-in-*τ *decay stated in* (2.82) *implies an algebraic decay statement in* t*variable with an upper bound* $\frac{C_*}{(1+t)^{\mu_0}}$ *for some* $C_* > 0$ *. Note that the larger* $\gamma \in (1, \frac{5}{3}]$ *is, the faster the equilibration rate is. The same applies to the vorticity decay in* (2.81) *except for* $\gamma = \frac{5}{3}$ *where we* get $\frac{\ln(1+t)}{(1+t)}$ instead of $\frac{1}{1+t}$.

Remark 2.6. *The assumption* $N \ge 2\lceil \alpha \rceil + 12 = 2\lceil \frac{1}{\gamma - 1} \rceil + 12$ *is not an optimal lower bound for the number of derivatives necessary to prove Theorem 2.4. We have not tried to find an optimal* N*, as we are primarily interested in displaying the phenomenology and developing a robust approach to the construction of global-in-time solutions to the* E_γ *-system. We do observe that the larger* $\gamma \in (1, \frac{5}{3}]$ *is, the less smoothness is required in our norm.*

Remark 2.7 (Constants μ_0 and μ_1). As a simple consequence of the ODE analysis in [44], the limit *defining* µ¹ *in the statement of the theorem is well-defined, see Lemma A.1 for more details. Note that by definition* (2.77) *of* S ^N *, we have that*

$$
\sum_{|\beta| \le N} \|\partial_r^{\alpha} \nabla^{\beta} \mathbf{V}\|_{1+\alpha+a,\psi} + \sum_{|\nu| \le N} \|\partial^{\nu} \mathbf{V}\|_{1+\alpha,1-\psi} \lesssim e^{-\mu_0 \tau} \mathcal{S}^N(\tau)^{\frac{1}{2}}.
$$
 (2.84)

Bound (2.84) *will be commonly used in our analysis. We also observe that for any* $\gamma \in (1, \frac{5}{3}]$ *the inequality* $0 < \mu_0 = \mu_0(\gamma) \leq \mu_1$ *holds with equality if and only if* $\gamma = \frac{5}{3}$ *.*

Remark 2.8 (Initial density ρ_0). *Note that the density* ρ_0 *of the initial gas configuration relates to the density of the background affine motion via the composition*

$$
\rho_0(x) = \rho_{A,\mathbf{a}}((\zeta(0) \circ \eta_0)^{-1}(x)) \det[D(\zeta(0) \circ \eta_0)]^{-1},
$$

where the transformation $\eta_0 : \Omega \to \Omega(0)$ *is very close to the identity map and* ζ *is the flow map associated with the affine motion* (1.4)*.*

Local well-posedness

 \overline{a}

The local-well-posedness of (2.64)–(2.65) follows from an adaption of the argument in [18]. In fact, the method developed in [18] ensures the well-posedness upon having a good a priori estimates for the full energy and the curl energy of the Euler equations via a duality argument. Although (2.64) is different from the Euler equations due to the presence of the time weights such as Λ (note that these weights are prescribed functions independent of the solutions), the estimates in Section 3 and Section 4, which will be obtained through several new ideas, give rise to the desired bounds on the energy functionals S^N (2.77) and \mathcal{B}^N (2.78) that are sufficient to apply the methodology of [18] by using \mathcal{S}^N and \mathcal{B}^N :

Theorem 2.9 (Local well-posedness [18]). Let $N \ge 2\lceil \alpha \rceil + 12$ be given. Then for given initial data (θ_0, V_0) with $\mathcal{S}^N(\theta_0, V_0)+\mathcal{B}^N(V_0)<\infty$, there exists a $T>0$ and a unique solution $(\theta(\tau), V(\tau))$: Ω \rightarrow \mathbb{R}^3 \times \mathbb{R}^3 *for each* τ \in $[0,T]$ *to* (2.64)–(2.65) *such that* $\mathcal{S}^N(\theta,\mathbf{V})(\tau) + \mathcal{B}^N[\mathbf{V}](\tau)$ \le $2(S^N(\theta_0, \mathbf{V}_0) + \dot{B}^N(\mathbf{V}_0))$ for all $\tau \in [0, T]$. Moreover, the map $[0, T] \ni \tau \mapsto S^N(\tau) \in \mathbb{R}_+$ is *continuous.*

For the rest of the article, we will use this well-posedness result to guarantee the existence of a unique solution $(θ, V) : \Omega \to \mathbb{R}^3 \times \mathbb{R}^3$ with $\mathcal{S}^N(θ, V) + \mathcal{B}^N[V] < \infty$ on a finite interval $[0, T]$ where $T > 0$ is fixed.

A priori assumptions

It is convenient to make the following a priori assumptions:

$$
\mathcal{S}^N(\tau) < \frac{1}{3} \tag{2.85}
$$

as well as

$$
\|\mathscr{A} - \text{Id}\|_{W^{1,\infty}(\Omega)} < \frac{1}{3}, \quad \|D\theta\|_{W^{1,\infty}(\Omega)} < \frac{1}{3}, \quad \|\mathscr{J} - 1\|_{W^{1,\infty}(\Omega)} < \frac{1}{3} \tag{2.86}
$$

for each $\tau \in [0, T]$ where $T > 0$ is fixed. The latter assumption ensures the non-degeneracy of the flow map. For an $\varepsilon_0 \ll 1$ in Theorem 2.4 the assumptions (2.85) and (2.86) are initially true, and by the local well-posedness theory they remain true on some short time interval. By means of a continuity argument we will show in the proof of Theorem 2.4 (Section 5) that our a-priori assumptions are in fact *improved*, thereby justifying them in the first place.

3 Vorticity estimates

In this section we exploit an effective decoupling in the modified Euler system (2.64) which allows us to bound the modified vorticity tensors Curl $\Lambda_{\alpha\alpha}$ V and Curl $\Lambda_{\alpha\alpha}\theta$ in terms of the high-order norm S^N . The main result of this section is given in Proposition 3.3, but to achieve it we first prove a lemma which demonstrates the above mentioned decoupling and gives us a simplified formula for the modified vorticity.

Lemma 3.1. Let $(\theta, V) : \Omega \to \mathbb{R}^3 \times \mathbb{R}^3$ be a unique solution to (2.64)–(2.65) defined on some time *interval* [0, T]. For any $\tau \in [0, T]$ *the modified vorticity tensors* Curl_{ΛA} V *and* Curl_{ΛA} θ *satisfy*

$$
\text{Curl}_{\Lambda\mathscr{A}}\mathbf{V}(\tau) = \frac{\mu(0)\text{Curl}_{\Lambda\mathscr{A}}\mathbf{V}(0)}{\mu} + \frac{1}{\mu} \int_0^\tau \mu[\partial_\tau, \text{Curl}_{\Lambda\mathscr{A}}] \mathbf{V} d\tau' - \frac{2}{\mu} \int_0^\tau \mu \text{Curl}_{\Lambda\mathscr{A}} (\Gamma^*\mathbf{V}) d\tau' \qquad (3.87)
$$

and

$$
\text{Curl}_{\Lambda\mathscr{A}}\theta(\tau) = \text{Curl}_{\Lambda\mathscr{A}}\theta(0) + \mu(0)\text{Curl}_{\Lambda\mathscr{A}}\mathbf{V}(0)\int_0^\tau \frac{1}{\mu(\tau')}d\tau' + \int_0^\tau [\partial_\tau, \text{Curl}_{\Lambda\mathscr{A}}]\theta(\tau') d\tau' + \int_0^\tau \frac{1}{\mu(\tau')} \int_0^{\tau'} \mu(\tau'')[\partial_\tau, \text{Curl}_{\Lambda\mathscr{A}}]\mathbf{V} d\tau'' d\tau' - \int_0^\tau \frac{2}{\mu(\tau')} \int_0^{\tau'} \mu(\tau'')\text{Curl}_{\Lambda\mathscr{A}}(\Gamma^*\mathbf{V}) d\tau'' d\tau' (3.88)
$$

Proof. Recalling that $V = \theta_{\tau}$, equation (2.61) is best rewritten in the form

$$
\mu^{3\gamma - 3} \left(\mathbf{V}_{\tau} + \frac{\mu_{\tau}}{\mu} \mathbf{V} + 2\Gamma^* \mathbf{V} \right) + \frac{\gamma}{\gamma - 1} \Lambda \nabla_{\eta} (f^{\gamma - 1}) + \delta \Lambda \eta = 0 \tag{3.89}
$$

Recalling the definition (2.69), apply Curl $\Lambda_{\mathscr{A}}$ to (3.89) and obtain

$$
\mu^{3\gamma - 3} \left(\text{Curl}_{\Lambda \mathscr{A}} \mathbf{V}_{\tau} + \frac{\mu_{\tau}}{\mu} \text{Curl}_{\Lambda \mathscr{A}} \mathbf{V} + 2 \text{Curl}_{\Lambda \mathscr{A}} \left(\Gamma^* \mathbf{V} \right) \right) + \delta \text{Curl}_{\Lambda \mathscr{A}} \Lambda \eta = 0 \tag{3.90}
$$

Note that

$$
\mathrm{Curl}_{\Lambda\mathscr{A}}\Lambda\eta=0
$$

and therefore (3.90) gives

$$
\operatorname{Curl}_{\Lambda\mathscr{A}}\mathbf{V}_{\tau} + \frac{\mu_{\tau}}{\mu} \operatorname{Curl}_{\Lambda\mathscr{A}}\mathbf{V} + 2\operatorname{Curl}_{\Lambda\mathscr{A}}\left(\Gamma^*\mathbf{V}\right) = 0 \tag{3.91}
$$

To obtain the identities (3.87) and (3.88) we shall integrate (3.91) with respect to τ . We rewrite the first term as

$$
\text{Curl}_{\Lambda\mathscr{A}}\mathbf{V}_{\tau} = \partial_{\tau} \left(\text{Curl}_{\Lambda\mathscr{A}}\mathbf{V} \right) - \left[\partial_{\tau}, \text{Curl}_{\Lambda\mathscr{A}} \right] \mathbf{V} \tag{3.92}
$$

where

$$
[\partial_{\tau}, \text{Curl}_{\Lambda \mathscr{A}}]F_j^k = \partial_{\tau} \left(\Lambda_{jm} \mathscr{A}_m^s \right) F_{,s}^k - \partial_{\tau} \left(\Lambda_{km} \mathscr{A}_m^s \right) F_{,s}^j \tag{3.93}
$$

We may therefore rewrite (3.91) in the form

$$
\partial_{\tau} \left(\mu \mathrm{Curl}_{\Lambda \mathscr{A}} \mathbf{V} \right) = \mu [\partial_{\tau}, \mathrm{Curl}_{\Lambda \mathscr{A}}] \mathbf{V} - 2\mu \mathrm{Curl}_{\Lambda \mathscr{A}} \left(\Gamma^* \mathbf{V} \right). \tag{3.94}
$$

Integrating (3.94) in τ we obtain (3.87). Using the commutator formula again we may write (3.87) in the form

$$
\partial_{\tau} \text{Curl}_{\Lambda \mathscr{A}} \theta(\tau) = \frac{\mu(0) \text{Curl}_{\Lambda \mathscr{A}} \mathbf{V}(0)}{\mu} + [\partial_{\tau}, \text{Curl}_{\Lambda \mathscr{A}}] \theta + \frac{1}{\mu} \int_{0}^{\tau} \mu [\partial_{\tau}, \text{Curl}_{\Lambda \mathscr{A}}] \mathbf{V} d\tau' - \frac{2}{\mu} \int_{0}^{\tau} \mu \text{Curl}_{\Lambda \mathscr{A}} (\Gamma^* \mathbf{V}) d\tau' \qquad (3.95)
$$

and therefore (3.88) follows by integrating the previous identity in τ .

$$
\Box
$$

Lemma 3.2. Let $(\theta, V) : \Omega \to \mathbb{R}^3 \times \mathbb{R}^3$ be a unique solution to (2.64)–(2.65) defined on some time *interval* $[0, T]$ *and satisfying the a priori assumptions* (2.85)–(2.86)*. Then for any* $\tau \in [0, T]$

$$
\sum_{a+|\beta| \le N} \left\| \left[\partial_r^{\alpha} \nabla^{\beta}, \text{Curl}_{\Lambda \mathscr{A}} \right] \mathbf{V}(\tau) \right\|_{1+\alpha+a,\psi}^{2} + \sum_{|\nu| \le N} \left\| \left[\partial^{\nu}, \text{Curl}_{\Lambda \mathscr{A}} \right] \mathbf{V}(\tau) \right\|_{1+\alpha,1-\psi}^{2}
$$
\n
$$
\lesssim e^{-2\mu_0 \tau} \mathcal{S}^N(\tau) \left(1 + P(\mathcal{S}^N(\tau)) \right)
$$
\n
$$
\sum_{a+|\beta| \le N} \left\| \left[\partial_r^{\alpha} \nabla^{\beta}, \text{Curl}_{\Lambda \mathscr{A}} \right] \theta(\tau) \right\|_{1+\alpha+a,\psi}^{2} + \sum_{|\nu| \le N} \left\| \left[\partial^{\nu}, \text{Curl}_{\Lambda \mathscr{A}} \right] \theta(\tau) \right\|_{1+\alpha,1-\psi}^{2}
$$
\n
$$
\lesssim \mathcal{S}^N(0) + \kappa \mathcal{S}^N(\tau) + \left(1 + P(\mathcal{S}^N(\tau)) \right) \int_0^{\tau} e^{-\mu_0 \tau'} \mathcal{S}^N(\tau') d\tau' \tag{3.97}
$$

where the polynomial P *is of degree at least 1,* $0 < \kappa \ll 1$ *is a small constant, and* μ_0 *is defined in* (2.79)*.*

Proof. Proof of (3.96)*.* Recall that

$$
\begin{split}\n&\left[\partial_r^a \nabla^\beta, \text{Curl}_{\Lambda \mathscr{A}}\right] \mathbf{V}_i^k = \partial_r^a \nabla^\beta \left(\text{Curl}_{\Lambda \mathscr{A}} \mathbf{V}_i^k \right) - \text{Curl}_{\Lambda \mathscr{A}} (\partial_r^a \nabla^\beta \mathbf{V})_i^k \\
&= \partial_r^a \nabla^\beta \left(\Lambda_{im} \mathscr{A}_m^s \mathbf{V}^k, s - \Lambda_{km} \mathscr{A}_m^s \mathbf{V}^i, s \right) - \left(\Lambda_{im} \mathscr{A}_m^s (\partial_r^a \nabla^\beta \mathbf{V})^k, s - \Lambda_{km} \mathscr{A}_m^s (\partial_r^a \nabla^\beta \mathbf{V})^i, s \right) \\
&= \Lambda_{km} \left(\mathscr{A}_m^s (\partial_r^a \nabla^\beta \mathbf{V})^i, s - \partial_r^a \nabla^\beta (\mathscr{A}_m^s \mathbf{V})^i, s \right) - \Lambda_{im} \left(\mathscr{A}_m^s (\partial_r^a \nabla^\beta \mathbf{V})^k, s - \partial_r^a \nabla^\beta (\mathscr{A}_m^s \mathbf{V}^k, s) \right)\n\end{split} \tag{3.98}
$$

As the two summands appearing on the right-most side above behave identically from the point-of-view of derivative count, we will restrict our attention only to the second term. Applying the Leibniz rule,

$$
\mathcal{A}_{m}^{s}(\partial_{r}^{a}\nabla^{\beta}\mathbf{V})^{k},_{s}-\partial_{r}^{a}\nabla^{\beta}(\mathcal{A}_{m}^{s}\mathbf{V}^{k},_{s})=-\underbrace{\partial_{r}^{a}\nabla^{\beta}\mathcal{A}_{m}^{s}\mathbf{V}^{k}}_{=:A}
$$
\n
$$
-\sum_{1\leq a'+|\beta'|\leq N-1 \atop a'\leq a,\beta'\leq \beta}\partial_{r}^{a'}\nabla^{\beta'}\mathcal{A}_{m}^{s}\partial_{r}^{a-a'}\nabla^{\beta-\beta'}\mathbf{V}^{k},_{s}-\underbrace{\mathcal{A}_{m}^{s}[\partial_{r}^{a}\nabla^{\beta},\partial_{s}]\mathbf{V}^{k}}_{=:B}
$$
\n(3.99)

where $[\partial_r^a \vec{\nabla}^{\beta}, \partial_s]$ stands for the commutator defined in (B.177). Terms A and B are the two highestorder terms on the right-hand side of (3.99). To estimate A, we use formulas (B.175) (when β = $(0, 0, 0)$) and otherwise (B.176). Assume without loss of generality that $\beta \neq (0, 0, 0)$ since the other case can be done analogously. By (B.176) for some $\ell \in \{1, 2, 3\}$ we have

$$
A = -\underbrace{\mathscr{A}_{j}^{k}(\partial_{r}^{a}\nabla^{\beta}\theta^{j})_{,m}\mathscr{A}_{i}^{m}\mathbf{V}^{k}_{,s}}_{=:A_{1}} - \mathbf{V}^{k}_{,s}\Big(\sum_{\substack{0\n(3.100)
$$

Note that

$$
||A_1||_{1+\alpha+a,\psi}^2 \lesssim ||D\mathbf{V}||_{L^{\infty}(\Omega)}^2 ||\nabla_{\eta} \partial_r^a \nabla^{\beta} \theta||_{1+\alpha+a,\psi}^2
$$

\$\lesssim e^{-2\mu_0 \tau} (\mathcal{S}^N(\tau))^2 \left(1 + P(\mathcal{S}^N(\tau))\right), \qquad (3.101)\$

where we have used (C.182) to bound $||D{\bf V}||_{L^{\infty}(\Omega)}$ by a constant multiple of $e^{-\mu_0 \tau} \sqrt{S^N(\tau)}$ and the a priori bounds (2.86). The remaining terms on the right-hand side of (3.100) are estimated using the standard Moser type estimates in conjunction with Proposition C.2 and formula (B.177). The estimates for term B and the remaining terms in (3.99) rely on the formula $(B.177)$ and Proposition C.2. Namely, recalling (B.177) we have the following schematic formula for B :

$$
B=\mathscr{A}_m^s\sum_{\ell=0}^{a+|\beta|}\sum_{\substack{a'+|\beta'|=\ell\\ a'\leq a+1}}C_{a',\beta',\ell}^s\partial_r^{a'}{\not\!{\nabla}}^{\beta'}{\bf V}^k,
$$

for some universal functions $C^s_{\alpha',\beta',\ell}$, smooth away from the origin. Therefore

$$
||B||_{1+\alpha+a,\psi}^2 \lesssim e^{-2\mu_0 \tau} \mathcal{S}^N(\tau).
$$

Summing the above estimates we conclude that all the $\|\cdot\|_{1+\alpha+a,\psi}$ -norms on the left-hand side of (3.96) are bounded by the right-hand side. An entirely analogous strategy can be applied to $\|\cdot\|_{1+\alpha,1-\psi}$ -norms to conclude the proof.

Proof of (3.97). Using the formula (3.98) with θ instead of V, in analogy to (3.99) we reduce the problem to estimating the $\|\cdot\|_{1+\alpha+a,\psi}$ -norm of the expression

$$
\mathscr{A}_{m}^{s}(\partial_{r}^{a}\nabla^{\beta}\theta)^{k},_{s}-\partial_{r}^{a}\nabla^{\beta}(\mathscr{A}_{m}^{s}\theta^{k},_{s})=-\underbrace{\partial_{r}^{a}\nabla^{\beta}\mathscr{A}_{m}^{s}\theta^{k}}_{=:A_{\theta}}-\sum_{1\leq a'+|\beta'|\leq N-1}\partial_{r}^{a'}\nabla^{\beta'}\mathscr{A}_{m}^{s}\partial_{r}^{a-a'}\nabla^{\beta-\beta'}\theta^{k},_{s}-\underbrace{\mathscr{A}_{m}^{s}[\partial_{r}^{a}\nabla^{\beta},\partial_{s}]\theta^{k}}_{=:B_{\theta}},\tag{3.102}
$$

We illustrate the proof by estimating the term A_{θ} . Estimating θ^{k} , in L^{∞} norm and using the fundamental theorem of calculus we obtain

$$
\|A_{\theta}\|_{1+\alpha+a,\psi}^{2} \lesssim \|\partial_{r}^{\alpha}\nabla^{\beta}\mathscr{A}\|_{1+\alpha+a,\psi}^{2}\|D\theta\|_{L^{\infty}(\Omega)}^{2}
$$

$$
\lesssim \|\partial_{r}^{\alpha}\nabla^{\beta}\mathscr{A}\|_{1+\alpha+a,\psi}^{2}\left(\|D\theta_{0}\|_{L^{\infty}(\Omega)}+\int_{0}^{\tau}\|D\mathbf{V}\|_{L^{\infty}(\Omega)}\right)^{2}
$$

$$
\lesssim \mathcal{S}^{N}(\tau)\left(\mathcal{S}^{N}(0)+\left(\int_{0}^{\tau}e^{-\mu_{0}\tau'}\sqrt{\mathcal{S}^{N}(\tau')}\,d\tau'\right)^{2}\right)
$$

$$
\lesssim \mathcal{S}^{N}(\tau)\left(\mathcal{S}^{N}(0)+\int_{0}^{\tau}e^{-\mu_{0}\tau'}\mathcal{S}^{N}(\tau')\,d\tau'\right),
$$

where we have used the Hardy-Sobolev embedding $\|D\mathbf{V}\|_{L^{\infty}(\Omega)} \lesssim e^{-\mu_0 \tau} \sqrt{S^N(\tau)}$ in the secondto-last estimate, and the Cauchy-Schwarz inequality in the last line. We have also used the already established fact $\|\partial_r^a \psi^{\beta} \mathscr{A}\|_{1+\alpha+a,\psi}^2 \lesssim \mathcal{S}^N(\tau)$ which has been explained in the proof of (3.96). We leave out the details for the remaining terms in (3.102) as they follow similarly.

Proposition 3.3 (The vorticity estimates). Let $(\theta, \mathbf{V}) : \Omega \to \mathbb{R}^3 \times \mathbb{R}^3$ be a unique solution to (2.64)– (2.65) *defined on some time interval* [0, T] *and satisfying the a priori assumptions* (2.85)*–*(2.86)*. Then for any* $\tau \in [0, T]$

$$
\mathcal{B}^{N}[\mathbf{V}](\tau) \lesssim \begin{cases} e^{-2\mu_{0}\tau} \left(\mathcal{S}^{N}(0) + \mathcal{B}^{N}(0) \right) + e^{-2\mu_{0}\tau} \mathcal{S}^{N}(\tau) & \text{if } 1 < \gamma < \frac{5}{3} \\ e^{-2\mu_{0}\tau} \left(\mathcal{S}^{N}(0) + \mathcal{B}^{N}(0) \right) + (1 + \tau^{2})e^{-2\mu_{0}\tau} \mathcal{S}^{N}(\tau) & \text{if } \gamma = \frac{5}{3} \end{cases} (3.103)
$$

$$
\mathcal{B}^{N}[\theta](\tau) \lesssim \mathcal{S}^{N}(0) + \mathcal{B}^{N}(0) + \kappa \mathcal{S}^{N}(\tau) + \int_{0}^{\tau} e^{-\mu_{0}\tau'} \mathcal{S}^{N}(\tau') d\tau'
$$
\n(3.104)

where $0 < \kappa \ll 1$ is a small constant, μ_0 is defined in (2.79), and ${\cal B}^N({\bf V}),$ ${\cal B}^N[\theta]$ are defined in (2.78).

Proof. Proof of (3.103). Applying $\partial_r^{\alpha} \overline{\mathbf{\mathcal{P}}}^{\beta}$ to (3.87) we obtain

$$
\text{Curl}_{\Lambda\mathscr{A}}\partial_r^{\alpha}\nabla^{\beta}\mathbf{V} = \frac{\mu(0)\partial_r^{\alpha}\nabla^{\beta}\text{Curl}_{\Lambda\mathscr{A}}\mathbf{V}(0)}{\mu} + \left[\partial_r^{\alpha}\nabla^{\beta}, \text{Curl}_{\Lambda\mathscr{A}}\right]\mathbf{V} + \frac{1}{\mu}\int_0^{\tau} \mu \partial_r^{\alpha}\nabla^{\beta}[\partial_\tau, \text{Curl}_{\Lambda\mathscr{A}}]\mathbf{V}d\tau' - \frac{2}{\mu}\int_0^{\tau} \mu \partial_r^{\alpha}\nabla^{\beta}\text{Curl}_{\Lambda\mathscr{A}}(\Gamma^*\mathbf{V})d\tau'. \tag{3.105}
$$

We shall exploit the exponential-in- τ growth of μ to show that for any pair of indices $a + |\beta| \leq N$ the left-hand side satisfies a "good" bound in L^2 -sense. Note that by definition (2.77) of \mathcal{S}^N it immediately follows that

$$
\left\| \frac{\mu(0)\partial_r^a \nabla^\beta \text{Curl}_{\Lambda \mathscr{A}} \mathbf{V}(0)}{\mu} \right\|_{1+\alpha+a,\psi}^2 \lesssim e^{-2\mu_1 \tau} (\mathcal{S}^N(0) + \mathcal{B}^N(0)),\tag{3.106}
$$

where we used the exponential growth (A.162) of μ . The second term on the right-hand side of (3.105) is bounded by Lemma 3.2. Concerning the third term on the right-hand side of (3.105), by (3.93) we have

$$
\partial_r^a \nabla^{\beta} [\partial_{\tau}, \text{Curl}_{\Lambda \mathscr{A}}] \mathbf{V}_{j}^{k} = \partial_r^a \nabla^{\beta} (\partial_{\tau} (\Lambda_{jm} \mathscr{A}_{m}^{s}) \mathbf{V}_{,s}^{k} - \partial_{\tau} (\Lambda_{km} \mathscr{A}_{m}^{s}) \mathbf{V}_{,s}^{j}).
$$

Schematically speaking both terms inside the parenthesis on the right-hand side behave the same way from the point of view of derivative count. It is therefore sufficient to restrict our attention to one term only. Note that

$$
\partial_r^{\alpha} \nabla^{\beta} \left(\partial_{\tau} \left(\Lambda_{jm} \mathscr{A}_{m}^{s} \right) \mathbf{V}_{,s}^{k} \right)
$$
\n
$$
= \partial_r^{\alpha} \nabla^{\beta} \left[\left(\partial_{\tau} \Lambda_{jm} \mathscr{A}_{m}^{s} + \Lambda_{jm} \partial_{\tau} \mathscr{A}_{m}^{s} \right) \mathbf{V}_{,s}^{k} \right]
$$
\n
$$
= \partial_{\tau} \Lambda_{jm} \left(\partial_r^{\alpha} \nabla^{\beta} \mathscr{A}_{m}^{s} \mathbf{V}_{,s}^{k} + \mathscr{A}_{m}^{s} \partial_r^{\alpha} \nabla^{\beta} \mathbf{V}_{,s}^{k} \right) + \Lambda_{jm} \left(\partial_r^{\alpha} \nabla^{\beta} \partial_{\tau} \mathscr{A}_{m}^{s} \mathbf{V}_{,s}^{k} + \partial_{\tau} \mathscr{A}_{m}^{s} \partial_r^{\alpha} \nabla^{\beta} \mathbf{V}_{,s}^{k} \right)
$$
\n
$$
+ \sum_{1 \leq \alpha' + |\beta'| \leq N-1} C_{\alpha',\beta'} \left(\partial_{\tau} \Lambda_{jm} \partial_r^{\alpha'} \nabla^{\beta'} \mathscr{A}_{m}^{s} + \Lambda_{jm} \partial_r^{\alpha'} \nabla^{\beta'} \partial_{\tau} \mathscr{A}_{m}^{s} \right) \partial_r^{\alpha - \alpha'} \nabla^{\beta - \beta'} \mathbf{V}_{,s}^{k} \tag{3.107}
$$

From the point of view of derivative count we may schematically rewrite the first two terms on the right-most side in the form

$$
\underbrace{\partial_{\tau} \Lambda \partial_r^a \nabla^{\beta} D \theta D \mathbf{V}}_{=:C_1} + \underbrace{\partial_{\tau} \Lambda D \eta \partial_r^a \nabla^{\beta} D \mathbf{V}}_{=:C_2} + \underbrace{\Lambda D \mathbf{V} \partial_r^a \nabla^{\beta} D \mathbf{V}}_{=:C_3}
$$
(3.108)

For the term C_1 , observe that

$$
\int_{\Omega} \psi w^{1+\alpha+a} \frac{1}{\mu^2} \Big| \int_0^{\tau} \mu \partial_{\tau} \Lambda \partial_{r}^{\alpha} \overline{\psi}^{\beta} D\theta D\mathbf{V} d\tau' \Big|^2 dx
$$
\n
$$
\lesssim e^{-2\mu_1 \tau} \int_{\Omega} \psi w^{1+\alpha+a} \Big| \int_0^{\tau} \Big| e^{\mu_1 \tau'} \partial_{\tau} \Lambda \partial_{r}^{\alpha} \overline{\psi}^{\beta} D\theta D\mathbf{V} \Big| d\tau' \Big|^2 dx
$$
\n
$$
\lesssim e^{-2\mu_1 \tau} \sup_{0 \le \tau' \le \tau} \int_{\Omega} \psi w^{1+\alpha+a} |\partial_{r}^{\alpha} \overline{\psi}^{\beta} D\theta|^{2} \Big| \int_0^{\tau} |e^{\mu_1 \tau'} |\partial_{\tau} \Lambda D\mathbf{V}| d\tau' \Big|^2 dx
$$
\n
$$
\lesssim e^{-2\mu_1 \tau} \sup_{0 \le \tau' \le \tau} \Big(||\partial_{r}^{\alpha} \overline{\psi}^{\beta} D\theta||_{1+\alpha+a,\psi}^{2} \Big) \Big| \int_0^{\tau} e^{\mu_1 \tau'} |\partial_{\tau} \Lambda| e^{-\mu_0 \tau'} (\mathcal{S}^N(\tau'))^{\frac{1}{2}} d\tau' \Big|^2
$$
\n
$$
\lesssim e^{-2\mu_1 \tau} (\mathcal{S}^N(\tau))^2
$$
\n(3.109)

where we used Proposition C.2 to bound $||D{\bf V}||_{L^{\infty}(\Omega)}$ by $e^{-\mu_0 \tau'}(S^N(\tau'))^{\frac{1}{2}}$, the sharp exponential behavior (A.162) of $\mu(\tau)$, and the exponential decay (A.164) of $|\Lambda_{\tau}|$.

To estimate the $\|\cdot\|_{1+\alpha+a,\psi}$ -norm of term C_2 in (3.108) we first note that the expression $\partial_r^a \nabla^\beta D{\bf V}$ contains one too many derivative to belong to our energy space. To resolve this difficulty, we must integrate-by-parts in τ first. Observe that

$$
\frac{1}{\mu} \int_0^{\tau} \mu \partial_{\tau} \Lambda D \eta \, \partial_{r}^a \nabla^{\beta} D \mathbf{V} d\tau' = \frac{1}{\mu} \left(\mu \partial_{\tau} \Lambda D \eta \, \partial_{r}^a \nabla^{\beta} D \theta \right) \Big|_0^{\tau} \n- \frac{1}{\mu} \int_0^{\tau} \mu \left(\frac{\mu_{\tau}}{\mu} \partial_{\tau} \Lambda D \eta + \partial_{\tau \tau} \Lambda D \eta + \partial_{\tau} \Lambda D \mathbf{V} \right) \partial_{r}^a \nabla^{\beta} D \theta d\tau' .
$$
\n(3.110)

Applying now the same line of reasoning as in the case of term C_1 , using thereby the exponential decay of $\partial_{\tau} \Lambda$ and $\partial_{\tau} \Lambda$ as stated in Lemma A.1 we arrive at

$$
\|\frac{1}{\mu}\int_0^\tau \mu \partial_\tau \Lambda D\eta \,\partial_r^a \nabla^\beta D\mathbf{V} d\tau'\|_{1+\alpha+a,\psi}^2 \lesssim e^{-2\mu_1 \tau} \mathcal{S}^N(0) + \left((1+\tau^2)e^{-2\mu_1 \tau} + Q(\tau)\right)\mathcal{S}^N(\tau)
$$
\n(3.111)

where

$$
Q(\tau) = \begin{cases} e^{-4\mu_0 \tau} & \text{if } 1 < \gamma < \frac{4}{3} \\ \tau^2 e^{-4\mu_0 \tau} & \text{if } \gamma = \frac{4}{3} \\ e^{-2\mu_1 \tau} & \text{if } \frac{4}{3} < \gamma \le \frac{5}{3} \end{cases}.
$$

The first τ^2 on the right-hand side of (3.111) comes from estimating $\int_0^{\tau} \mu_{\tau} \partial_{\tau} \Lambda d\tau'$ and $Q(\tau)$ from estimating $\int_0^{\tau} \mu \partial_{\tau \tau} \Lambda d\tau'$ ² corresponding to the first two terms in the τ integral in the right-hand side of (3.110).

The estimate for the error term C_3 from (3.108) is more subtle. Quantity $|\Lambda|$ does *not* decay exponentially but is merely bounded. Therefore when estimating $\frac{1}{\mu(\tau)} \int_0^{\tau} \mu(\tau') C_3(\tau') d\tau'$ in L^{∞}_{τ} -norm, we must proceed with more caution. As in the above, since the expression $\partial_r^a \nabla^{\beta} D \mathbf{V}$ contains one too many derivative to belong to our energy space, we integrate-by-parts. As a result

$$
\frac{1}{\mu} \int_0^{\tau} \mu \Lambda D \mathbf{V} \, \partial_r^{\alpha} \overline{\mathbf{\mathbf{\nabla}}}^{\beta} D \mathbf{V} d\tau' = \frac{1}{\mu} \left[\mu \Lambda D \mathbf{V} \, \partial_r^{\alpha} \overline{\mathbf{\mathbf{\nabla}}}^{\beta} D \theta \right]_0^{\tau} \n- \frac{1}{\mu} \int_0^{\tau} \mu \left(\frac{\mu \tau}{\mu} \Lambda D \mathbf{V} + \partial_\tau \Lambda D \mathbf{V} + \Lambda D \mathbf{V}_\tau \right) \partial_r^{\alpha} \overline{\mathbf{\mathbf{\nabla}}}^{\beta} D \theta \, d\tau'. \quad (3.112)
$$

The $\|\cdot\|_{1+\alpha+a,\psi}$ -norm of the first term on the right-hand side of (3.112) is obviously bounded by

$$
\mu^{-1}S^{N}(0) + \|D\mathbf{V}\|_{L^{\infty}(\Omega)}(\mathcal{S}^{N}(\tau))^{\frac{1}{2}} \lesssim e^{-\mu_{1}\tau} \mathcal{S}^{N}(0) + e^{-\mu_{0}\tau} \mathcal{S}^{N}(\tau)
$$

where we used the Hardy-Sobolev embeddings of Appendix C to obtain $||D\textbf{V}||_{L^{\infty}(\Omega)} \leq$ $e^{-\mu_0 \tau} (\mathcal{S}^N(\tau))^{\frac{1}{2}}$. By the sharp exponential behavior (A.162) of μ we have

$$
\left| \frac{1}{\mu} \int_0^{\tau} \mu_{\tau} \Lambda D \mathbf{V} \partial_{\tau}^a \overline{\mathbf{V}}^{\beta} D \theta \, d\tau' \right| \lesssim \sup_{0 \le \tau' \le \tau} |\partial_{\tau}^a \overline{\mathbf{V}}^{\beta} D \theta| e^{-\mu_1 \tau} \int_0^{\tau} e^{\mu_1 \tau'} ||D \mathbf{V}||_{L^{\infty}(\Omega)} \, d\tau'
$$

$$
\lesssim \sup_{0 \le \tau' \le \tau} |\partial_{\tau}^a \overline{\mathbf{V}}^{\beta} D \theta| (\mathcal{S}^N(\tau))^{\frac{1}{2}} e^{-\mu_1 \tau} \int_0^t e^{(\mu_1 - \mu_0) \tau'} d\tau'
$$

=:L

Note that

$$
L \lesssim \begin{cases} e^{-\mu_0 \tau} & \text{if } 1 < \gamma < \frac{5}{3} \\ \tau e^{-\mu_0 \tau} & \text{if } \gamma = \frac{5}{3} \end{cases}
$$

because $\mu_0 = \mu_1$ when $\gamma = \frac{5}{3}$. Therefore

$$
\|\frac{1}{\mu}\int_0^\tau \mu_\tau \Lambda D\theta \partial_r^a \nabla^\beta D\theta \, d\tau'\|_{1+\alpha+a,\psi}^2 \lesssim \begin{cases} e^{-2\mu_0\tau} (\mathcal{S}^N(\tau))^2 & \text{if } 1 < \gamma < \frac{5}{3} \\ \tau^2 e^{-2\mu_0\tau} (\mathcal{S}^N(\tau))^2 & \text{if } \gamma = \frac{5}{3} \end{cases}
$$

The remaining two terms inside the integral on the right-hand side of (3.112) can be estimated similarly. The only difference is that after integrating-by-parts in τ one τ -derivative will fall on DV in the third term on the right-hand side of (3.112). Since $D{\bf V}_\tau$ is a priori not in our energy space, we must use equation (2.64) to express V_τ in terms of V and purely spatial derivatives of θ. The rest of the proof is then analogous and we obtain

$$
\|\frac{1}{\mu}\int_0^\tau \mu(\tau')C_3(\tau') d\tau'\|_{1+\alpha+a,\psi}^2 \lesssim \begin{cases} e^{-2\mu_0\tau} \mathcal{S}^N(\tau) & \text{if } 1 < \gamma < \frac{5}{3} \\ (1+\tau^2)e^{-2\mu_0\tau} \mathcal{S}^N(\tau) & \text{if } \gamma = \frac{5}{3} \end{cases} (3.113)
$$

The lower-order remainder appearing as the last term on the right-hand side of (3.107) is straightforward to estimate using the above ideas, Moser-type estimates, Proposition C.2, Lemma A.1, and the definition of S^N :

$$
\left\| \sum_{1 \le a'+|\beta'| \le N-1} C_{a',\beta'} \left(\partial_{\tau} \Lambda_{jm} \partial_{r}^{a'} \nabla^{\beta'} \mathscr{A}_{m}^{s} + \Lambda_{jm} \partial_{r}^{a'} \nabla^{\beta'} \partial_{\tau} \mathscr{A}_{m}^{s} \right) \partial_{r}^{a-a'} \nabla^{\beta-\beta'} \mathbf{V}^{k},_{s} \right\|_{1+\alpha+a,\psi}^{2} \n\lesssim e^{-2\mu_{0}\tau} \mathcal{S}^{N}(\tau) \left(1 + P(\mathcal{S}^{N}(\tau)) \right)
$$
\n(3.114)

where P is a polynomial of degree at least 1. By (2.85), it is in turn bounded by $e^{-2\mu_0 \tau} S^N(\tau)$.

From (3.109)–(3.114) and by using $\mu_0 \leq \mu_1$ and (2.85), we conclude that

$$
\|\frac{1}{\mu}\int_0^\tau \mu \partial_r^a \nabla^\beta [\partial_\tau, \text{Curl}_{\Lambda \varnothing}] \mathbf{V} d\tau' \|^2_{1+\alpha+a,\psi} \lesssim \begin{cases} e^{-2\mu_0 \tau} \mathcal{S}^N(\tau) & \text{if } 1 < \gamma < \frac{5}{3} \\ (1+\tau^2)e^{-2\mu_0 \tau} \mathcal{S}^N(\tau) & \text{if } \gamma = \frac{5}{3} \end{cases} \tag{3.115}
$$

In an entirely analogous fashion we can estimate the last term on the right-hand side of (3.105) to conclude

$$
\|\frac{2}{\mu}\int_0^\tau \mu \partial_\tau^a \overline{\mathbf{\mathbf{\nabla}}}^{\beta} \text{Curl}_{\Lambda \mathscr{A}} \left(\Gamma^* \mathbf{V} \right) d\tau' \|^2_{1+\alpha+a,\psi} \lesssim \begin{cases} e^{-2\mu_0 \tau} \mathcal{S}^N(\tau) & \text{if } 1 < \gamma < \frac{5}{3} \\ (1+\tau^2)e^{-2\mu_0 \tau} \mathcal{S}^N(\tau) & \text{if } \gamma = \frac{5}{3} \end{cases} \tag{3.116}
$$

Combining (3.105) – (3.106) , Lemma 3.2, bounds (3.115) – (3.116) , and Proposition 4.2 we obtain (3.103).

Proof of (3.104). Applying $\partial_r^a \overline{\psi}^\beta$ to (3.88) we obtain

$$
\text{Curl}_{\Lambda\mathscr{A}}\partial_r^{\alpha}\nabla^{\beta}\theta = \partial_r^{\alpha}\nabla^{\beta}\text{Curl}_{\Lambda\mathscr{A}}\theta(0) + \mu(0)\partial_r^{\alpha}\nabla^{\beta}\text{Curl}_{\Lambda\mathscr{A}}\mathbf{V}(0)\int_0^{\tau} \frac{1}{\mu(\tau')}d\tau'
$$

+
$$
\left[\partial_r^{\alpha}\nabla^{\beta}, \text{Curl}_{\Lambda\mathscr{A}}\right]\theta + \int_0^{\tau} \frac{1}{\mu(\tau')} \int_0^{\tau'} \mu(\tau'')\partial_r^{\alpha}\nabla^{\beta}[\partial_\tau, \text{Curl}_{\Lambda\mathscr{A}}]\mathbf{V}d\tau'' d\tau'
$$

-
$$
\int_0^{\tau} \frac{2}{\mu(\tau')} \int_0^{\tau'} \mu(\tau'')\partial_r^{\alpha}\nabla^{\beta}\text{Curl}_{\Lambda\mathscr{A}}(\Gamma^*\mathbf{V}) d\tau'' d\tau'.
$$
(3.117)

The $\|\cdot\|_{1+\alpha+a,\psi}$ norm of the first two terms on the right-hand side of (3.117) is bounded by a constant multiple of $S^N(0) + B^N(0)$ since $\int_0^{\tau} \frac{1}{\mu(\tau)} d\tau'$ is bounded uniformly-in- τ due to the exponential growth of $\mu(\tau)$. The term $\|\left[\partial_r^a \nabla^\beta, \text{Curl}_{\Lambda \mathscr{A}}\right] \theta\|_{1+\alpha+a,\psi}$ is bounded by Lemma 3.2. To bound the fourth term on the right-hand side of (3.117) we use the Cauchy-Schwarz inequality and (3.115) to obtain

$$
\begin{split}\n&\left\|\int_{0}^{\tau} \frac{1}{\mu(\tau')}\int_{0}^{\tau'} \mu(\tau'') \partial_{r}^{a} \overline{\mathbf{\mathbf{\mathbf{\mathcal{V}}}}}^{\beta} [\partial_{\tau}, \text{Curl}_{\Lambda \mathscr{A}}] \mathbf{V} d\tau'' d\tau'\right\|_{1+\alpha+a,\psi}^{2} \\
&\lesssim \int_{\Omega} \left[\int_{0}^{\tau} \frac{(1+\tau')^{2}}{\mu(\tau')} d\tau' \int_{0}^{\tau} \frac{1}{(1+\tau')^{2} \mu(\tau')} \left(\int_{0}^{\tau'} \mu(\tau'') \partial_{r}^{a} \overline{\mathbf{\mathbf{\mathbf{\mathcal{V}}}}}^{\beta} [\partial_{\tau}, \text{Curl}_{\Lambda \mathscr{A}}] \mathbf{V} d\tau''\right)^{2} d\tau'\right] w^{1+\alpha+a} \psi d\mathbf{x} \\
&\lesssim \int_{0}^{\tau} \frac{1}{(1+\tau')^{2} \mu(\tau')} \left\|\int_{0}^{\tau'} \mu(\tau'') \partial_{r}^{a} \overline{\mathbf{\mathbf{\mathbf{\mathcal{V}}}}}^{\beta} [\partial_{\tau}, \text{Curl}_{\Lambda \mathscr{A}}] \mathbf{V} d\tau''\right\|_{1+\alpha+a,\psi}^{2} d\tau' \\
&\lesssim \int_{0}^{\tau} e^{-\mu_{0}\tau} \mathbf{S}^{N}(\tau') d\tau',\n\end{split}
$$

where the last bound follows from a minor adaptation of the argument leading to (3.115). Using the Cauchy-Schwarz inequality and (3.116) instead, we analogously show that the last term on the righthand side of (3.117) satisfies

$$
\Big\| \int_0^{\tau} \frac{2}{\mu(\tau')}\int_0^{\tau'} \mu(\tau'') \partial_{\tau}^a \nabla^{\beta} \text{Curl}_{\Lambda \mathscr{A}}\left(\Gamma^* \mathbf{V}\right) d\tau'' d\tau' \Big\|_{1+\alpha+a,\psi}^2 \lesssim \int_0^{\tau} e^{-\mu_0 \tau'} \mathcal{S}^N(\tau') d\tau'
$$

From the above estimates, the desired bound for $\sup_{\tau \in [0,T]} \sum_{a+|\beta| \le N} ||\text{Curl}_{\Lambda \mathscr{A}} \partial_r^a \nabla^\beta \theta||_{1+\alpha+a,\psi}^2$
in (3.104) is obtained. A completely analogous argument yields the remaining bound on $\sum_{|\nu| \le N} ||\partial^{\nu}, \text{Curl}_{\Lambda \mathscr{A}}] \Theta ||_{1+\alpha+a,1-\psi}^2$ and this concludes the proof of the proposition. \Box

4 Energy estimates

4.1 High-order energies

The high-order *energy* that emerges naturally from the problem is intricate and a priori different from the high-order norm S^N . Nevertheless, they are equivalent as we shall establish shortly. Before we give the precise expression for the energy functional, we first identify some important quantities.

Recall that Λ (1.15) is a positive symmetric matrix with the determinant 1 and therefore it is diagonalisable:

$$
\Lambda = P^{\top}QP, \ P \in SO(3), \ Q = \text{diag}(d_1, d_2, d_3) \tag{4.118}
$$

where $P \in SO(3)$ is the orthogonal matrix that diagonalizes Λ and $d_i > 0$ are the eigenvalues of Λ . We then define $\mathcal{M}_{a,\beta}$ and \mathcal{N}_{ν} as the P−conjugates of $\nabla_{\eta}\partial_{r}^{\alpha}\overline{\psi}^{\beta}$ θ and $\nabla_{\eta}\partial^{\nu}\theta$:

$$
\mathcal{M}_{\alpha,\beta} := P \, \nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta \, P^{\top} \text{ and } \mathcal{N}_{\nu} := P \, \nabla_{\eta} \partial^{\nu} \theta \, P^{\top} \tag{4.119}
$$

for any $(a, \beta), 0 \le a + |\beta| \le N$ and $\nu, 0 \le |\nu| \le N$ respectively. By denoting the standard inner product of two vectors in \mathbb{R}^3 by $\langle \cdot, \cdot \rangle$:

$$
\langle u, v \rangle = u^{\top} v = v^{\top} u \text{ for any } u, v \in \mathbb{R}^{3}
$$

we define the high-order energy functional:

$$
\mathcal{E}^{N}(\theta, \mathbf{V}) = \mathcal{E}^{N}(\tau)
$$
\n
$$
= \frac{1}{2} \sum_{a+|\beta| \leq N} \int_{\Omega} \psi \left[\mu^{3\gamma - 3} \left\langle \Lambda^{-1} \partial_{r}^{a} \nabla^{\beta} \mathbf{V}, \partial_{r}^{a} \nabla^{\beta} \mathbf{V} \right\rangle + \delta \left\langle \Lambda^{-1} \partial_{r}^{a} \nabla^{\beta} \theta, \partial_{r}^{a} \nabla^{\beta} \theta \right\rangle \right] w^{a+\alpha}
$$
\n
$$
+ \psi \mathcal{J}^{-\frac{1}{\alpha}} \Big[\sum_{i,j=1}^{3} d_{i} d_{j}^{-1} \left((\mathcal{M}_{a,\beta})_{i}^{j} \right)^{2} + \frac{1}{\alpha} \left(\text{div}_{\eta} \partial_{r}^{a} \nabla^{\beta} \theta \right)^{2} \Big] w^{a+\alpha+1} dy
$$
\n
$$
+ \frac{1}{2} \sum_{|\nu| \leq N} \int_{\Omega} (1 - \psi) \Big[\mu^{3\gamma - 3} \left\langle \Lambda^{-1} \partial^{\nu} \mathbf{V}, \partial^{\nu} \mathbf{V} \right\rangle + \delta \left\langle \Lambda^{-1} \partial^{\nu} \mathbf{V}, \partial^{\nu} \mathbf{V} \right\rangle \Big] w^{\alpha}
$$
\n
$$
+ (1 - \psi) \mathcal{J}^{-\frac{1}{\alpha}} \Big[\sum_{i,j=1}^{3} d_{i} d_{j}^{-1} \left((\mathcal{N}_{\nu})_{i}^{j} \right)^{2} + \frac{1}{\alpha} \left(\text{div}_{\eta} \partial^{\nu} \theta \right)^{2} \Big] w^{1+\alpha} dy \qquad (4.120)
$$

Note that when $a + |\beta| = 0$ and $|\nu| = 0$, the sum of the two integrals above produces the zeroth order energy that does not depend on the cutoff function ψ .

Remark 4.1. *Since the eigenvalues* $d_i(\tau)$, $i = 1, 2, 3$ *are strictly positive for all* $\tau \geq 0$ *and bounded from above and below uniformly-in-* τ *by Lemma A.1 the ratios* $\frac{d_i}{d_j}$ *are strictly positive and uniformly bounded from above and below for all* $\tau \geq 0$ *.*

We also introduce the *dissipation* functional:

$$
\begin{split} \mathcal{D}^N(\mathbf{V})=\mathcal{D}^N(\tau) =\frac{5-3\gamma}{2}\mu^{3\gamma-3}\frac{\mu_\tau}{\mu}\int_{\Omega}\Big[\psi\sum_{a+|\beta|\leq N}\left\langle\Lambda^{-1}\partial_r^a\vec{\nabla}^{\beta}\mathbf{V},\,\partial_r^a\vec{\nabla}^{\beta}\mathbf{V}\right\rangle w^{a+\alpha}\\ &\qquad\qquad+(1-\psi)\sum_{|\nu|\leq N}\left\langle\Lambda^{-1}\partial^\nu\mathbf{V},\,\partial^\nu\mathbf{V}\right\rangle w^{\alpha}\Big]\,dy \end{split}
$$

Note that $\mathcal{D}^N(\mathbf{V}) \geq 0$ for $\gamma \leq \frac{5}{3}$.

Proposition 4.2 (Equivalence of norm and energy). Let $(\theta, \mathbf{V}) : \Omega \to \mathbb{R}^3 \times \mathbb{R}^3$ be a unique solution *to* (2.64)*–*(2.65) *defined on some time interval* [0, T] *and satisfying the a priori assumptions* (2.85)*–* (2.86)*. Then there exist constants* $C_1, C_2 > 0$ *depending only on* A_0, A_1, δ *such that*

$$
C_1 \mathcal{S}^N(\tau) \le \sup_{0 \le \tau' \le \tau} \mathcal{E}^N(\tau') \le C_2 \mathcal{S}^N(\tau). \tag{4.121}
$$

Proof. The proof is a simple consequence of the definitions of S^N , \mathcal{E}^N and uniform boundedness of the matrix Λ and its eigenvalues, see Lemma A.1. \Box

4.2 Energy identities

Before we derive the fundamental energy identity, we shall need the following lemma, which captures the key algebraic structure necessary for the extraction of a positive-definite energy.

Lemma 4.3. Let $(\theta, V) : \Omega \to \mathbb{R}^3 \times \mathbb{R}^3$ be a unique solution to (2.64)–(2.65) defined on some time *interval* [0, T]. Then for any (a, β) , $0 \le a + |\beta| \le N$, and a multi-index ν , $|\nu| \le N$, the following *identities hold:*

$$
\Lambda_{\ell j} (\nabla_{\eta} \partial_r^a \nabla^{\beta} \theta)^i_j \Lambda_{im}^{-1} (\nabla_{\eta} \partial_r^a \nabla^{\beta} \theta)^m_{\ell, \tau} = \frac{1}{2} \frac{d}{d\tau} \left(\sum_{i,j=1}^3 d_i d_j^{-1} (\mathcal{M}_{a,\beta})^j_i)^2 \right) + \mathcal{T}_{a,\beta}
$$
(4.122)

$$
\Lambda_{\ell j} (\nabla_{\eta} \partial^{\nu} \theta)^{i}_{j} \Lambda_{im}^{-1} (\nabla_{\eta} \partial^{\nu} \theta)^{m}_{\ell, \tau} = \frac{1}{2} \frac{d}{d\tau} \left(\sum_{i,j=1}^{3} d_{i} d_{j}^{-1} (\mathcal{N}_{\nu})^{j}_{i})^{2} \right) + \mathcal{T}_{\nu}
$$
\n(4.123)

where $\mathcal{M}_{a,\beta}$ *and* \mathcal{N}_{ν} *are defined in* (4.119) *and the error terms* $\mathcal{T}_{a,\beta}$ *and* \mathcal{T}_{ν} *are given by*

$$
\mathcal{T}_{a,\beta} = -\frac{1}{2} \sum_{i,j=1}^{3} \frac{d}{d\tau} \left(d_i d_j^{-1} \right) \left(\left(\mathcal{M}_{a,\beta} \right)_i^j \right)^2 - \text{Tr} \left(Q \mathcal{M}_{a,\beta} Q^{-1} \left(\partial_\tau P P^\top \mathcal{M}_{a,\beta}^\top + \mathcal{M}_{a,\beta}^\top P \partial_\tau P^\top \right) \right)
$$
\n(4.124)

$$
\mathcal{T}_{\nu} = -\frac{1}{2} \sum_{i,j=1}^{3} \frac{d}{d\tau} \left(d_i d_j^{-1} \right) \left((\mathcal{N}_{\nu})_i^j \right)^2 - \text{Tr} \left(Q \mathcal{N}_{\nu} Q^{-1} \left(\partial_{\tau} P P^{\top} \mathcal{N}_{\nu}^{\top} + \mathcal{N}_{\nu}^{\top} P \partial_{\tau} P^{\top} \right) \right). \tag{4.125}
$$

Proof. For any matrix $\tau \mapsto M(\tau)$ let $\tilde{M} = PMP^{\top}$ where $P \in SO(\mathbb{R}^3)$ is the orthogonal matrix introduced in (4.118). We then have the following identity

$$
\operatorname{Tr}\left(\Lambda M\Lambda^{-1}M_{\tau}^{\top}\right) = \operatorname{Tr}\left(Q(PMP^{\top})Q^{-1}PM_{\tau}^{\top}P^{\top}\right)
$$
\n
$$
= \operatorname{Tr}\left(Q\tilde{M}Q^{-1}\partial_{\tau}\tilde{M}^{\top}\right) - \operatorname{Tr}\left(Q\tilde{M}Q^{-1}\left(\partial_{\tau}PP^{\top}\tilde{M}^{\top} + \tilde{M}^{\top}P\partial_{\tau}P^{\top}\right)\right)
$$
\n
$$
= \frac{1}{2}\frac{d}{d\tau}\sum_{k,\ell=1}^{3} \left(d_{k}d_{\ell}^{-1}\tilde{M}_{k\ell}^{2}\right) - \frac{1}{2}\sum_{k,\ell=1}^{3}\frac{d}{d\tau}\left(d_{k}d_{\ell}^{-1}\right)\tilde{M}_{k\ell}^{2}
$$
\n
$$
- \operatorname{Tr}\left(Q\tilde{M}Q^{-1}\left(\partial_{\tau}PP^{\top}\tilde{M}^{\top} + \tilde{M}^{\top}P\partial_{\tau}P^{\top}\right)\right). \tag{4.126}
$$

The matrix identity (4.126) is a simple algebraic manipulation, where we used the identity

$$
PM_{\tau}^{\top} P^{\top} = \partial_{\tau} (PM^{\top} P^{\top}) - \partial_{\tau} PM^{\top} P^{\top} - PM^{\top} \partial_{\tau} P^{\top}
$$

$$
= \partial_{\tau} (\tilde{M}^{\top}) - \partial_{\tau} PP^{\top} \tilde{M}^{\top} - \tilde{M}^{\top} P \partial_{\tau} P^{\top}
$$

but it allows us to extract a positive-definite energy quantity modulo a small error term. Applying (4.126) to $M=\nabla_{\eta}\partial_r^a\nabla^{\beta}\theta,$ we have

$$
\Lambda_{\ell j} \Lambda_{im}^{-1} (\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta)^{i}_{j} (\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta)^{m}_{\ell, \tau} = (\Lambda (\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta))_{\ell i} (\Lambda^{-1} \partial_{\tau} (\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta)^{\top})_{i\ell}
$$
\n
$$
= \frac{1}{2} \frac{d}{d\tau} \left(\sum_{i,j=1}^{3} d_{i} d_{j}^{-1} ((\mathcal{M}_{a,\beta})^{j}_{i})^{2} \right) - \frac{1}{2} \sum_{ij} \frac{d}{d\tau} (d_{i} d_{j}^{-1}) ((\mathcal{M}_{a,\beta})^{j}_{i})^{2}
$$
\n
$$
- \text{Tr} (Q \mathcal{M}_{a,\beta} Q^{-1} (\partial_{\tau} P P^{\top} \mathcal{M}_{a,\beta}^{\top} + \mathcal{M}_{a,\beta}^{\top} P \partial_{\tau} P^{\top})) \tag{4.127}
$$

where we recall the definitions of $\mathcal{M}_{a,\beta}$ and \mathcal{N}_{ν} given in (4.119). This proves (4.122) and the proof of (4.123) is completely analogous. \Box

The next lemma reveals how vector fields ∂_r and ∇_i , $i = 1, 2, 3$, commute with the degenerate (spatial) differential operator coming from the pressure gradient term in the momentum equation. This commutation lemma in particular provides a clue as to why the strength of the weights used in our norm scales proportionally to the number of spatial derivatives used.

Lemma 4.4. *Let* $q \in \mathbb{R}_+$ *be given. Then for a given* 2*-tensor* $\mathbf{T} : \Omega \to \mathbb{M}^{3 \times 3}$ *and any* $i, j \in \{1, 2, 3\}$ *the following commutation identities hold*

$$
\partial_r \left[\frac{1}{w^q} (w^{1+q} \mathbf{T}_i^k)_{,k} \right] = \frac{1}{w^{1+q}} (w^{2+q} \partial_r \mathbf{T}_i^k)_{,k} + C_i^{q+1} [\mathbf{T}], \tag{4.128}
$$

$$
\nabla_j \left[\frac{1}{w^q} (w^{1+q} \mathbf{T}_i^k)_{,k} \right] = \frac{1}{w^q} (w^{1+q} \nabla_j \mathbf{T}_i^k)_{,k} + C_{ij}^q [\mathbf{T}], \tag{4.129}
$$

where the commutators $\mathcal{C}_i^{q+1}[\mathbf{T}]$ and $\mathcal{C}_{ij}^q[\mathbf{T}]$ are given as follows:

$$
\mathcal{C}_i^{q+1}[\mathbf{T}] = \left(\partial_r w - \frac{w}{r}\right) \nabla_k \mathbf{T}_i^k + (1+q)\partial_r w_{,k} \mathbf{T}_i^k \tag{4.130}
$$

$$
\mathcal{C}_{ij}^q[\mathbf{T}] = w \left(\frac{y_j \nabla_k}{r^2} + \frac{\delta_{kj} r^2 - y_k y_j}{r^3} \partial_r \right) \mathbf{T}_i^k + (1+q) \nabla_j w_{,k} \mathbf{T}_i^k \text{ for } j = 1, 2, 3. \tag{4.131}
$$

Proof. Note that

$$
\partial_r \left[\frac{1}{w^q} (w^{1+q} \mathbf{T}_i^k)_{,k} \right] = \partial_r \left[w \mathbf{T}_i^k,_k + (1+q) w,_k \mathbf{T}_i^k \right] \n= w \partial_r \mathbf{T}_i^k,_k + \partial_r w \mathbf{T}_i^k,_k + (1+q) w,_k \partial_r \mathbf{T}_i^k + (1+q) \partial_r w,_k \mathbf{T}_i^k \n= w (\partial_r \mathbf{T}_i^k)_, _k + (2+q) w,_k \partial_r \mathbf{T}_i^k + w [\partial_r, \partial_k] \mathbf{T}_i^k + \partial_r w \mathbf{T}_i^k,_k - w,_k \partial_r \mathbf{T}_i^k + (1+q) \partial_r w,_k \mathbf{T}_i^k \n= \frac{1}{w^{1+q}} (w^{2+q} \partial_r \mathbf{T}_i^k)_, _k - \frac{w}{r} \nabla_k \mathbf{T}_i^k + \partial_r w \mathbf{T}_i^k,_k - w,_k \partial_r \mathbf{T}_i^k + (1+q) \partial_r w,_k \mathbf{T}_i^k
$$

Since $\partial_r = \frac{y}{r} \cdot \nabla$ and $w = \frac{\delta}{2(1+\alpha)}(1-|y|^2) = \frac{\delta}{2(1+\alpha)}(1-r^2)$, we deduce

$$
\partial_r w \mathbf{T}^k_{i,k} - w_{,k} \partial_r \mathbf{T}^k_i = \partial_r w \nabla_k \mathbf{T}^k_i
$$

which finishes the proof of (4.128). Using $\nabla_i w = 0$, we obtain

$$
\begin{split} &\nabla_j\left[\frac{1}{w^q}(w^{1+q}\mathbf{T}^k_i)_{,\mathbf{k}}\right] = \nabla_j\left[w\mathbf{T}^k_i, k + (1+q)w_{,\mathbf{k}}\mathbf{T}^k_i\right] \\ & = w\nabla_j\mathbf{T}^k_i, k + (1+q)w_{,\mathbf{k}}\nabla_j\mathbf{T}^k_i + (1+q)\nabla_j w_{,\mathbf{k}}\mathbf{T}^k_i \\ & = w(\nabla_j\mathbf{T}^k_i), k + w[\nabla_j, \partial_k]\mathbf{T}^k_i + (1+q)w_{,\mathbf{k}}\nabla_j\mathbf{T}^k_i + (1+q)\nabla_j w_{,\mathbf{k}}\mathbf{T}^k_i \\ & = \frac{1}{w^q}(w^{1+q}\nabla_j\mathbf{T}^k_i), k + \mathcal{C}^q_{ij}[\mathbf{T}] \end{split}
$$

where we used the commutator formula (B.172) and the definition (4.131).

 \Box

Remark 4.5. *The commutators appearing in the previous lemma consist of lower order terms with respect to both the number of derivatives and the strength of weights in* w*. Even though the term* $\partial_r w \nabla_k \mathbf{T}^k_i$ appearing in the expression \mathcal{C}^{q+1}_i (4.130) is potentially very dangerous as it looses a factor of w , it is nevertheless of lower order because it contains the term \mathbf{T}^k_i with purely tangential derivatives. *Additionally all the coefficients appearing in the commutators in Lemma 4.4 are smooth in the annulus* $\frac{1}{4} \leq r \leq 1$ where the commutation formulas (4.128)–(4.129) will be actually applied.

Proposition 4.6. Let $(\theta, \mathbf{V}) : \Omega \to \mathbb{R}^3 \times \mathbb{R}^3$ be a unique solution to (2.64)–(2.65) defined on some *time interval* $[0, T]$ *and satisfying the a priori assumptions* (2.85)–(2.86)*. Then for any* $\tau \in [0, T]$ *the* *following energy inequality holds:*

$$
\mathcal{E}^{N}(\tau) + \int_{0}^{\tau} \mathcal{D}^{N}(\tau') d\tau' \lesssim \mathcal{S}^{N}(0) + \mathcal{B}^{N}[\theta](\tau) + \int_{0}^{\tau} (\mathcal{S}^{N}(\tau'))^{\frac{1}{2}} (\mathcal{B}^{N}[\mathbf{V}](\tau'))^{\frac{1}{2}} d\tau'
$$

+ $\kappa \mathcal{S}^{N}(\tau) + \int_{0}^{\tau} e^{-\mu_{0}\tau'} \mathcal{S}^{N}(\tau') d\tau'$ (4.132)

where $0 < \kappa \ll 1$ *is a small constant and the constant* μ_0 *is defined in* (2.79).

Proof. To simplify notations, we shall denote by R any lower order error term satisfying an estimate of the form

$$
\Big|\int_0^{\tau} \int_{\Omega} \mathcal{R} \, dx \, d\tau' \Big| \lesssim \mathcal{S}^N(0) + \kappa \mathcal{S}^N(\tau) + \int_0^{\tau} e^{-\mu_0 \tau'} \mathcal{S}^N(\tau') d\tau'. \tag{4.133}
$$

Step 1. Zeroth order estimate. We start with $a + |\beta| = 0$ and $|\nu| = 0$. In this case, there is no need to obtain two separate estimates by using the cutoff function ψ . Multiply (2.64) by $\Lambda_{im}^{-1}\partial_{\tau}\theta^m$ and integrate over Ω to obtain

$$
\int_{\Omega} w^{\alpha} \left(\mu^{3\gamma - 3} \partial_{\tau \tau} \theta_i + \mu^{3\gamma - 4} \mu_{\tau} \partial_{\tau} \theta_i + 2 \mu^{3\gamma - 3} \Gamma^*_{ij} \partial_{\tau} \theta_j + \delta \Lambda_{i\ell} \theta_{\ell} \right) \Lambda^{-1}_{im} \partial_{\tau} \theta^m dy + \int_{\Omega} \left(w^{1 + \alpha} \Lambda_{ij} \left(\mathscr{A}^k_j \mathscr{J}^{-\frac{1}{\alpha}} - \delta^k_j \right) \right), k \Lambda^{-1}_{im} \partial_{\tau} \theta^m dy = 0.
$$

The first integral can be rewritten as

$$
\frac{1}{2}\frac{d}{d\tau}\left(\mu^{3\gamma-3}\int_{\Omega}w^{\alpha}\langle\Lambda^{-1}\partial_{\tau}\theta,\partial_{\tau}\theta\rangle dy+\delta\int_{\Omega}w^{\alpha}|\theta|^{2}dy\right)+\frac{5-3\gamma}{2}\mu^{3\gamma-4}\mu_{\tau}\int_{\Omega}w^{\alpha}\langle\Lambda^{-1}\partial_{\tau}\theta,\partial_{\tau}\theta\rangle dy\\-\frac{\mu^{3\gamma-3}}{2}\int_{\Omega}w^{\alpha}\langle\partial_{\tau}\Lambda^{-1}\partial_{\tau}\theta,\partial_{\tau}\theta\rangle dy+2\mu^{3\gamma-3}\int_{\Omega}w^{\alpha}\langle\Lambda^{-1}\partial_{\tau}\theta,\Gamma^{*}\partial_{\tau}\theta\rangle dy
$$

For the second integral, by divergence theorem (integration by parts) and by using the following identities

$$
\mathscr{A}_j^k \mathscr{J}^{-\frac{1}{\alpha}} - \delta_j^k = (\mathscr{A}_j^k - \delta_j^k) \mathscr{J}^{-\frac{1}{\alpha}} + \delta_j^k (\mathscr{J}^{-\frac{1}{\alpha}} - 1)
$$

$$
\mathscr{A}_j^k - \delta_j^k = \mathscr{A}_l^k \delta_j^l - \mathscr{A}_l^k [D\eta]_j^l = \mathscr{A}_l^k (\delta_j^l - [D\eta]_j^l - [D\theta]_j^l) = -\mathscr{A}_l^k [D\theta]_j^l
$$

we obtain

$$
- \int_{\Omega} w^{1+\alpha} \Lambda_{ij} \left(\mathcal{A}_{j}^{k} \mathcal{J}^{-\frac{1}{\alpha}} - \delta_{j}^{k} \right) \Lambda_{im}^{-1} \partial_{\tau} \theta^{m}, k \, dy
$$

=
$$
\int_{\Omega} w^{1+\alpha} \mathcal{J}^{-\frac{1}{\alpha}} \Lambda_{ij} \mathcal{A}_{l}^{k} \theta^{l}, j \Lambda_{im}^{-1} \partial_{\tau} \theta^{m}, k \, dy - \int_{\Omega} w^{1+\alpha} (\mathcal{J}^{-\frac{1}{\alpha}} - 1) \partial_{\tau} \theta^{k}, k \, dy
$$

=: $(i) + (ii)$

Now for (*i*), by rewriting $\Lambda_{ij} = \Lambda_{ip} \delta_p^j = \Lambda_{ip} \mathscr{A}_l^j \eta^l_{\, ,p} = \Lambda_{ip} (\mathscr{A}_p^j + \mathscr{A}_l^j \theta^l_{\, ,p})$, we have

$$
(i) = \int_{\Omega} w^{1+\alpha} \mathcal{J}^{-\frac{1}{\alpha}} \Lambda_{ip} \mathcal{A}_{p}^{j} \theta^{\ell}, j \Lambda_{im}^{-1} \mathcal{A}_{\ell}^{k} \partial_{\tau} \theta^{m}, k \, dy + \int_{\Omega} w^{1+\alpha} \mathcal{J}^{-\frac{1}{\alpha}} \Lambda_{ip} \mathcal{A}_{l}^{j} \theta^{l}, p \, \mathcal{A}_{\ell}^{k} \theta^{\ell}, j \Lambda_{im}^{-1} \partial_{\tau} \theta^{m}, k \, dy
$$

\n
$$
= \int_{\Omega} w^{1+\alpha} \mathcal{J}^{-\frac{1}{\alpha}} \Lambda_{\ell j} [\nabla_{\eta} \theta]_{j}^{i} \Lambda_{im}^{-1} [\nabla_{\eta} \partial_{\tau} \theta]_{\ell}^{m} dy + \int_{\Omega} w^{1+\alpha} \mathcal{J}^{-\frac{1}{\alpha}} [\text{Curl}_{\Lambda \mathcal{A}} \theta]_{i}^{\ell} \Lambda_{im}^{-1} [\nabla_{\eta} \partial_{\tau} \theta]_{\ell}^{m} dy
$$

\n
$$
+ \int_{\Omega} w^{1+\alpha} \mathcal{J}^{-\frac{1}{\alpha}} \mathcal{A}_{l}^{j} \theta^{l}, m \, \mathcal{A}_{\ell}^{k} \theta^{\ell}, j \, \partial_{\tau} \theta^{m}, k \, dy
$$

\n
$$
= : (i)_{1} + (i)_{2} + (i)_{3}.
$$

By Lemma 4.3, we rewrite the first term

$$
(i)_1 = \frac{1}{2} \frac{d}{d\tau} \int_{\Omega} w^{1+\alpha} \mathcal{J}^{-\frac{1}{\alpha}} \sum_{i,j=1}^{3} d_i d_j^{-1} ((\mathcal{M}_{0,0})_i^j)^2 dy + \int_{\Omega} w^{1+\alpha} \mathcal{J}^{-\frac{1}{\alpha}} \mathcal{T}_{0,0} dy
$$

$$
+ \frac{1}{2\alpha} \int_{\Omega} w^{1+\alpha} \mathcal{J}^{-\frac{1}{\alpha}-1} \mathcal{J}_{\tau} \sum_{i,j=1}^{3} d_i d_j^{-1} ((\mathcal{M}_{0,0})_i^j)^2 dy
$$

The first term contributes to the energy. The second term contains τ −derivative of P or d_i from $\mathcal{T}_{0,0}$ decaying exponentially fast and the third term contains $D\theta_\tau$ from \mathscr{J}_τ , and therefore they are bounded by $e^{-\mu_0 \tau} S^N(\tau)$. Now the term $(i)_2$ contributes to the second right-hand side term in (4.132). The term $(i)_3$ is bounded by $e^{-\mu_0 \tau} S^N(\tau)$.

For (ii) , we first note that

$$
\mathscr{J} = \det[D\eta] = \det[\mathbf{Id} + D\theta] = 1 + \text{Tr}[D\theta] + O(|D\theta|^2).
$$

By writing $1 - \mathscr{J}^{-\frac{1}{\alpha}} = \frac{1}{\alpha} \text{Tr}[D\theta] + R_J$ where $R_J = O(|D\theta|^2)$, we deduce that

 $|(ii)| \lesssim e^{-\mu_0 \tau} \mathcal{S}^N(\tau)$

To be consistent with the definition of our energy \mathcal{E}^N , we may recover the zeroth order divergence energy in \mathcal{E}^N by adding the following identity

$$
\frac{1}{2}\frac{d}{d\tau}\frac{1}{\alpha}\int_{\Omega}w^{1+\alpha}\mathscr{J}^{-\frac{1}{\alpha}}|\text{div}_{\eta}\theta|^{2}dy = \frac{1}{2\alpha}\int_{\Omega}w^{1+\alpha}\left(\partial_{\tau}(\mathscr{J}^{-\frac{1}{\alpha}})|\text{div}_{\eta}\theta|^{2} + \mathscr{J}^{-\frac{1}{\alpha}}\partial_{\tau}(|\text{div}_{\eta}\theta|^{2})\right)dy
$$

where the right-hand side is clearly bounded by $e^{-\mu_0 \tau} S^N(\tau)$ due to the presence of τ -derivatives of θ. This complete the zeroth order estimate.

Step 2. High order estimates. In order to obtain high order estimates, we will need to deal with degeneracy caused by vacuum and having correct weights and commuting with right vector fields near the boundary is important. This is why we invoke the cutoff function ψ to localize the difficulty. We focus on getting the estimates of \mathcal{E}^N involving ψ . Here we will crucially use Lemma 4.4.

Let (a, β) where $a + |\beta| \ge 1$ be given. We first rewrite (2.64) as

$$
\mu^{3\gamma-4} \left(\mu \partial_{\tau \tau} \theta_i + \mu_\tau \partial_\tau \theta_i + 2\mu \Gamma_{ij}^* \partial_\tau \theta_j \right) + \delta \Lambda_{i\ell} \theta_\ell + \frac{1}{w^\alpha} \left(w^{1+\alpha} \Lambda_{ij} \left(\mathscr{A}_j^k \mathscr{J}^{-\frac{1}{\alpha}} - \delta_j^k \right) \right), k = 0. \tag{4.134}
$$

We apply $\partial_r^a \nabla^{\beta}$ to (4.134) by using (4.128) and (4.129) and multiply the resulting equation by the cutoff function ψ :

$$
\psi \mu^{3\gamma-4} \left(\mu \partial_{\tau \tau} \partial_r^{\alpha} \nabla^{\beta} \theta_i + \mu_{\tau} \partial_{\tau} \partial_r^{\alpha} \nabla^{\beta} \theta_i + 2 \mu \Gamma_{ij}^* \partial_{\tau} \partial_r^{\alpha} \nabla^{\beta} \theta_j \right) + \delta \psi \Lambda_{i\ell} \partial_r^{\alpha} \nabla^{\beta} \theta_{\ell} + \psi \frac{1}{w^{\alpha+\alpha}} \left(w^{1+\alpha+\alpha} \Lambda_{ij} \partial_r^{\alpha} \nabla^{\beta} \left(\mathcal{A}_j^k \mathcal{J}^{-\frac{1}{\alpha}} - \delta_j^k \right) \right),_k = -\psi \mathcal{R}_i^{a,\beta},
$$
(4.135)

where the error terms $\mathcal{R}_i^{a,\beta}$ can be written inductively as follows. For any given index β with $|\beta|=n$ we represent it as $\beta = b^n := (b_n, ..., b_1), b_i \in \{1, 2, 3\}$ so that $b^n = (b_n, b^{n-1})$ and $b^1 = (b_1)$. Then for $a = 0$ we have the following inductive formula for $\mathcal{R}_i^{a,\beta}$

$$
\mathcal{R}_{i}^{0,b^{1}} = \mathcal{C}_{ib_{1}}^{\alpha} [\mathscr{A} \mathscr{J}^{-\frac{1}{\alpha}} - \mathrm{Id}]
$$

$$
\mathcal{R}_{i}^{0,b^{n}} = \mathcal{C}_{ib_{n}}^{\alpha} [\overline{\mathbf{v}}^{b^{n-1}} (\mathscr{A} \mathscr{J}^{-\frac{1}{\alpha}} - \mathrm{Id})] + \overline{\mathbf{v}}_{b_{n}} \mathcal{R}_{i}^{0,b^{n-1}} \text{ for } n \ge 2
$$

where $\mathcal{C}_{ij}^{\alpha}[\cdot]$ is defined in (4.131) and we recall that $\alpha = \frac{1}{\gamma - 1}$. Now in general for $a \ge 1$, by setting $\mathcal{R}^{0,0} = 0$, we have

$$
\mathcal{R}^{a,\beta}_i = \mathcal{C}^{\alpha+a}_i [\partial^{a-1}_r \overline{\mathbb{V}}^\beta(\mathscr{A} \mathscr{J}^{-\frac{1}{\alpha}} - \mathrm{Id})] + \partial_r \mathcal{R}^{a-1,\beta}_i
$$

where $\mathcal{C}_i^{\alpha+1}[\cdot]$ is defined in (4.130). Note that $\mathcal{R}_i^{\alpha,\beta}$ is a lower order term in the sense that it satisfies the error estimate

$$
\|\mathcal{R}_i^{a,\beta}\|_{a+\alpha,\psi}^2 \lesssim \mathcal{S}^N\left((1+P(\mathcal{S}^N)\right),\tag{4.136}
$$

where P is a polynomial of order at least 1. The proof follows from the Proposition C.2. See also Remark 4.5.

We will now derive the energy estimates from (4.135) . Multiplying (4.135) by $w^{\alpha+a} \Lambda_{im}^{-1} \partial_\tau \partial_r^a \overline{\mathbf{\nabla}}^{\beta} \mathbf{\theta}^m$ and integrating over Ω we arrive at

$$
\frac{1}{2} \frac{d}{d\tau} \left(\mu^{3\gamma - 3} \int_{\Omega} \psi w^{\alpha + a} \langle \Lambda^{-1} \partial_r^a \nabla^\beta \mathbf{V}, \partial_r^a \nabla^\beta \mathbf{V} \rangle dy + \delta \int_{\Omega} \psi w^{\alpha + a} |\partial_r^a \nabla^\beta \theta|^2 dy \right) \n+ \frac{5 - 3\gamma}{2} \mu^{3\gamma - 3} \frac{\mu \tau}{\mu} \int_{\Omega} \psi w^{\alpha + a} \langle \Lambda^{-1} \partial_r^a \nabla^\beta \mathbf{V}, \partial_r^a \nabla^\beta \mathbf{V} \rangle dy \n- \frac{1}{2} \mu^{3\gamma - 3} \int_{\Omega} \psi w^{\alpha + a} \langle [\partial_\tau \Lambda^{-1} - 4\Lambda^{-1} \Gamma^*] \partial_r^a \nabla^\beta \mathbf{V}, \partial_r^a \nabla^\beta \mathbf{V} \rangle dy \n+ \int_{\Omega} \psi \left(w^{1 + \alpha + a} \Lambda_{ij} \partial_r^a \nabla^\beta \left(\mathcal{A}_j^k \mathcal{J}^{-\frac{1}{\alpha}} - \delta_j^k \right) \right), k \Lambda_{im}^{-1} \partial_\tau \partial_r^a \nabla^\beta \theta^m dy \n= \int_{\Omega} \psi w^{\alpha + a} \mathcal{R}_{a,\beta}^i \Lambda_{im}^{-1} \partial_r^a \nabla^\beta \theta^m_{\tau} dy.
$$
\n(4.137)

Note that

$$
\partial_{\tau} \Lambda^{-1} - 4 \Lambda^{-1} \Gamma^* = \partial_{\tau} (O^{\top} O) - 4 O^{\top} O O^{-1} O_{\tau} = -\partial_{\tau} \Lambda^{-1} + 2(O_{\tau}^{\top} O - O^{\top} O_{\tau}),
$$

where the orthogonal matrix O is defined in (2.48). Since $(O_\tau^\top O - O^\top O_\tau)^\top = -(O_\tau^\top O - O^\top O_\tau)$, the third line of (4.137) reduces to

$$
\frac{1}{2}\mu^{3\gamma-3}\int_{\Omega}\psi w^{\alpha+a}\left\langle \partial_{\tau}\Lambda^{-1}\partial_{r}^{a}\overline{\mathbf{\mathbb{V}}}^{\beta}\mathbf{V},\partial_{r}^{a}\overline{\mathbf{\mathbb{V}}}^{\beta}\mathbf{V}\right\rangle dy.
$$

Hence, all lines except the fourth one in the left-hand side of (4.137) are written in desirable forms. The fourth line will give rise to the full gradient energy at the expense of a top order term, which will contain the favorable Curl $\Lambda_{\mathcal{A}}$ -component. To see this, we start with the following key formula [18]:

$$
\partial(\mathscr{A}_j^k \mathscr{J}^{-\frac{1}{\alpha}}) = -\mathscr{J}^{-\frac{1}{\alpha}} \mathscr{A}_\ell^k \mathscr{A}_j^s \partial \eta^\ell,_{s} - \frac{1}{\alpha} \mathscr{J}^{-\frac{1}{\alpha}} \mathscr{A}_j^k \mathscr{A}_\ell^s \partial \eta^\ell,_{s}
$$
(4.138)

where ∂ is either ∂_i , $i = 1, 2, 3$, or $\partial = \partial_{\tau}$. This formula is still valid for ∂_r or ∇_i if we write the formula with more care:

$$
X(\mathscr{A}_j^k \mathscr{J}^{-\frac{1}{\alpha}}) = -\mathscr{J}^{-\frac{1}{\alpha}} \mathscr{A}_\ell^k \mathscr{A}_j^s X(\eta^\ell, s) - \frac{1}{\alpha} \mathscr{J}^{-\frac{1}{\alpha}} \mathscr{A}_j^k \mathscr{A}_\ell^s X(\eta^\ell, s)
$$

for $X \in \mathcal{X}$ where $\mathcal{X} = \{\partial_i, \vec{\nabla}_j, \partial_r\}_{i,j=1,2,3}$. We also note that

$$
X(\eta^{\ell},s) = X(y^{\ell},s + \theta^{\ell},s) = X(\theta^{\ell},s) = (X\theta^{\ell}),s + [X,\partial_{s}]\theta^{\ell}, \quad X \in \mathcal{X},\tag{4.139}
$$

where $[X, \partial_s]$ is given in Lemma B.1. Hence, for any multi-index (a, β) so that $a + |\beta| > 0$, we have the identity

$$
\Lambda_{ij}\partial_r^{\alpha}\nabla^{\beta}(\mathscr{A}_j^k \mathscr{J}^{-\frac{1}{\alpha}} - \delta_j^k)
$$
\n
$$
= -\Lambda_{ij}\left(\mathscr{J}^{-\frac{1}{\alpha}}\mathscr{A}_\ell^k \mathscr{A}_j^s \partial_r^{\alpha}\nabla^{\beta}\theta^{\ell}, s + \frac{1}{\alpha}\mathscr{J}^{-\frac{1}{\alpha}}\mathscr{A}_j^k \mathscr{A}_\ell^s \partial_r^{\alpha}\nabla^{\beta}\theta^{\ell}, s\right) + C_i^{a,\beta,k}(\theta)
$$
\n
$$
= -\mathscr{J}^{-\frac{1}{\alpha}}\mathscr{A}_\ell^k \left(\Lambda_{ij}\mathscr{A}_j^s \partial_r^{\alpha}\nabla^{\beta}\theta^{\ell}, s - \Lambda_{\ell j}\mathscr{A}_j^s \partial_r^{\alpha}\nabla^{\beta}\theta^i, s\right) - \mathscr{J}^{-\frac{1}{\alpha}}\mathscr{A}_\ell^k \Lambda_{\ell j}\mathscr{A}_j^s \partial_r^{\alpha}\nabla^{\beta}\theta^i, s - \frac{1}{\alpha}\mathscr{J}^{-\frac{1}{\alpha}}\Lambda_{ij}\mathscr{A}_j^k \mathscr{A}_\ell^s \partial_r^{\alpha}\nabla^{\beta}\theta^{\ell}, s + C_i^{a,\beta,k}(\theta)
$$
\n
$$
= -\mathscr{J}^{-\frac{1}{\alpha}}\mathscr{A}_\ell^k \left[\text{Curl}_{\Lambda\mathscr{A}}\partial_r^{\alpha}\nabla^{\beta}\theta\right]_i^{\ell} - \mathscr{J}^{-\frac{1}{\alpha}}\mathscr{A}_\ell^k \Lambda_{\ell j} \left[\nabla_\eta \partial_r^{\alpha}\nabla^{\beta}\theta\right]_j^i
$$
\n
$$
- \frac{1}{\alpha}\mathscr{J}^{-\frac{1}{\alpha}}\Lambda_{ij}\mathscr{A}_j^k \text{div}_{\eta}(\partial_r^{\alpha}\nabla^{\beta}\theta) + C_i^{a,\beta,k}(\theta).
$$
\n(4.140)

All that matters in our determination of the lower order commutator $\mathcal{C}_i^{a,\beta,k}$ is that for any choice of $i, k \in \{1, 2, 3\}$ its schematic form is given by

$$
\mathcal{C}_i^{a,\beta,k}(\theta) = \sum_{g \in G_{a,\beta}} C_g(\partial \theta) \prod_{\{(a_i,\beta_i)\} \in G_{a,\beta}} \partial_r^{a_i} \nabla^{\beta_i} \theta,
$$
\n(4.141)

where the set $G_{a,\beta}$ denotes the set of all decompositions into finite sequences of pairs $\{(a_i,\beta_i)\}_{i=1,\dots,m}$ such that $\sum_{i=1}^{m} (a_i + |\beta_i|) < a + |\beta|$, $\beta_i \leq \beta$ for all $i = 1, ..., m$, and $C_g(\cdot)$ are some smooth universal functions on the exterior of any ball around the origin $r = 0$. Such a decomposition is a simple consequence of the Leibniz rule and the commutator properties given in Lemma B.1. Using Proposition C.2 and the standard Moser inequalities we obtain the bound

$$
\|\mathcal{C}_i^{a,\beta,k}(\theta)\|_{\alpha+a,\psi}^2 + \|D\mathcal{C}_i^{a,\beta,k}(\theta)\|_{1+\alpha+a,\psi}^2 \lesssim \mathcal{S}^N(\tau). \tag{4.142}
$$

For the sake of completeness we briefly explain the proof of the above estimate. Without loss of generality assume that $a_1 + |\beta_1| = \max_{i=1,...,m} \{a_i + |\beta_i|\}$. Then for any $j = 2, ..., m, a_j + |\beta_j| \leq \lfloor \frac{N}{2} \rfloor$ and therefore by Proposition C.2, estimate (C.181), we have the bound

$$
\prod_{j=2}^{m} \|w^{a_j} \partial_r^{a_j} \nabla^{\beta_j} \theta\|_{L^{\infty}(\Omega)} \lesssim \mathcal{S}^N(\tau)^{\frac{m-1}{2}}.
$$
\n(4.143)

Therefore, using the Cauchy-Schwarz inequality, the definition of S^N , decomposition (4.141) we conclude that

$$
\|\mathcal{C}_i^{a,\beta,k}(\theta)\|_{\alpha+a,\psi}^2 \lesssim \mathcal{S}^N(\tau)
$$

as claimed. A similar bound using (C.182) leads to the second inequality in (4.142). We now plug (4.140) into the last term on the left-hand side of (4.137) and after integration by parts we ob-

$$
\int_{\Omega} \psi \left(w^{1+\alpha+a} \Lambda_{ij} \partial_r^{\alpha} \nabla^{\beta} \left(\mathcal{A}_j^k \mathcal{J}^{-\frac{1}{\alpha}} - \delta_j^k \right) \right) , k \Lambda_{im}^{-1} \partial_\tau \partial_r^{\alpha} \nabla^{\beta} \theta^m dy
$$
\n
$$
= \int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\mathcal{A}_\ell^k [\text{Curl}_{\Lambda \mathcal{A}} \partial_r^{\alpha} \nabla^{\beta} \theta]_{i}^{\ell} + \mathcal{A}_\ell^k \Lambda_{\ell j} [\nabla_\eta \partial_r^{\alpha} \nabla^{\beta} \theta]_{j}^{\ell} + \frac{1}{\alpha} \Lambda_{ij} \mathcal{A}_j^k \text{div}_{\eta} (\partial_r^{\alpha} \nabla^{\beta} \theta) \right)
$$
\n
$$
\Lambda_{im}^{-1} \partial_\tau \left(\partial_r^{\alpha} \nabla^{\beta} \theta^m \right) , k dy
$$
\n
$$
+ \int_{\Omega} \psi , k w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\mathcal{A}_\ell^k [\text{Curl}_{\Lambda \mathcal{A}} \partial_r^{\alpha} \nabla^{\beta} \theta]_{i}^{\ell} + \mathcal{A}_\ell^k \Lambda_{\ell j} [\nabla_\eta \partial_r^{\alpha} \nabla^{\beta} \theta]_{j}^{\ell} + \frac{1}{\alpha} \Lambda_{ij} \mathcal{A}_j^k \text{div}_{\eta} (\partial_r^{\alpha} \nabla^{\beta} \theta) \right)
$$
\n
$$
\Lambda_{im}^{-1} \partial_\tau \partial_r^{\alpha} \nabla^{\beta} \theta^m dy
$$
\n
$$
+ \int_{\Omega} \psi \left(w^{1+\alpha+a} \mathcal{C}_i^{\alpha,\beta,k} \right) , k \Lambda_{im}^{-1} \partial_\tau \partial_r^{\alpha} \nabla^{\beta} \theta^m dy
$$
\n
$$
=: I_1 + I_2 + I_3.
$$
\n(4.144)

The rest of the proof will be devoted to the estimation of I_1 , I_2 and I_3 .

Estimation of I_3 in (4.144). By the Cauchy-Schwarz inequality and (A.164), the last integral I_3 above satisfies the bound

$$
\left| \int_0^{\tau} \int_{\Omega} \psi \left(w^{1+\alpha+a} C_i^{a,\beta,k} \right) \kappa \Lambda_{im}^{-1} \partial_{\tau} \partial_{\tau}^{\alpha} \overline{\mathbf{\mathbb{V}}}^{\beta} \mathbf{\Theta}^m \, dy d\tau' \right|
$$

\n
$$
\lesssim \int_0^{\tau} \|\partial_{\tau}^{\alpha} \overline{\mathbf{\mathbb{V}}}^{\beta} \mathbf{V}^m \|_{1+\alpha+a,\psi} \left(\|C_i^{a,\beta,k}(\theta)\|_{\alpha+a,\psi} + \|D C_i^{a,\beta,k} \|_{1+\alpha+a,\psi} \right) d\tau'
$$

\n
$$
\lesssim \int_0^{\tau} e^{-\mu_0 \tau'} \mathcal{S}^N(\tau') d\tau', \tag{4.145}
$$

where we used the definition of S^N and (4.142).

Estimation of I_2 in (4.144). We now explain the second-to-last integral I_2 on the right-hand side of (4.144). Note that $\psi_{,k}$ is supported in $B_{\frac{3}{4}}(0) \setminus B_{\frac{1}{4}}(0)$ and it is not bounded by ψ , and therefore the integral cannot be controlled by using only ψ norms. On $B_{\frac{3}{4}}(0) \setminus B_{\frac{1}{4}}(0)$, however, $\partial_r^a \overline{\psi}^\beta \theta$ and $\partial_r^a \nabla^{\beta} \mathbf{V}$ can be expressed in terms of cartesian derivatives ∂^ν of θ and \mathbf{V} for $|\nu| \le a + |\beta|$. Hence we use both norms involving ψ and $1 - \psi$ to control the integral:

$$
\begin{split}\n&\left|\int_{0}^{\tau} \int_{B_{\frac{3}{4}}(0) \setminus B_{\frac{1}{4}}(0)} \psi_{,k} \, w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\mathcal{A}_{\ell}^{k} \left[\text{Curl}_{\Lambda \mathcal{A}} \partial_{r}^{\alpha} \nabla^{\beta} \theta \right]_{i}^{\ell} + \mathcal{A}_{\ell}^{k} \Lambda_{\ell j} \left[\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta \right]_{j}^{\ell} + \frac{1}{\alpha} \Lambda_{ij} \mathcal{A}_{j}^{k} \text{div}_{\eta} (\partial_{r}^{\alpha} \nabla^{\beta} \theta) \right) \\
&\Lambda_{im}^{-1} \partial_{\tau} \partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \, dy \, d\tau'\n\end{split}
$$
\n
$$
\lesssim \int_{0}^{\tau} \|D\psi\|_{L^{\infty}(\Omega)} \left(\|\partial_{r}^{\alpha} \nabla^{\beta} \mathbf{V}^{m}\|_{\alpha+a,\psi} \|\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta\|_{1+\alpha+a,\psi} + \sum_{|\nu| \leq \alpha+|\beta|} \|\partial^{\nu} \mathbf{V}\|_{\alpha,1-\psi} \sum_{|\nu| \leq \alpha+|\beta|} \|\nabla_{\eta} \partial^{\nu} \theta\|_{1+\alpha,1-\psi} \right) d\tau' \\
\lesssim \int_{0}^{\tau} e^{-\mu_{0}\tau'} \mathcal{S}^{N}(\tau') \, d\tau'.\n\tag{4.146}
$$

The last two integrals I_2 and I_3 on the right-hand side of (4.144) therefore contribute to the error term R satisfying the bound (4.133).

tain

Estimation of I_1 in (4.144). We evaluate now the first integral on the right-hand side of (4.144):

$$
\int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\mathcal{A}_{\ell}^{k} \left[\text{Curl}_{\Lambda \mathcal{A}} \partial_{r}^{\alpha} \nabla^{\beta} \theta \right]_{i}^{\ell} + \mathcal{A}_{\ell}^{k} \Lambda_{\ell j} \left[\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta \right]_{j}^{\ell} + \frac{1}{\alpha} \Lambda_{ij} \mathcal{A}_{j}^{k} \text{div}_{\eta} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta \right) \right) \n\Lambda_{im}^{-1} \partial_{\tau} \partial_{r}^{\alpha} \nabla^{\beta} \theta^{m}, \quad dy \n= \int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left[\text{Curl}_{\Lambda \mathcal{A}} \partial_{r}^{\alpha} \nabla^{\beta} \theta \right]_{i}^{\ell} \mathcal{A}_{\ell}^{k} \Lambda_{im}^{-1} \partial_{\tau} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \right), \quad dy \n+ \int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\mathcal{A}_{\ell}^{k} \Lambda_{\ell j} \left[\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta \right]_{j}^{\ell} + \frac{1}{\alpha} \Lambda_{ij} \mathcal{A}_{j}^{k} \text{div}_{\eta} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta \right) \right) \Lambda_{im}^{-1} \partial_{\tau} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \right), \quad dy
$$
\n(4.147)

We first focus on the first integral on the right-hand side of (4.147) . We integrate-by-parts with respect to τ and obtain

$$
\int_{0}^{\tau} \int_{\Omega} w^{1+\alpha+a} \mathcal{J}^{\frac{1}{\alpha}} \mathcal{A}_{\ell}^{k} \Lambda_{im}^{-1} \psi \left[\text{Curl}_{\Lambda \mathcal{A}} \partial_{r}^{\alpha} \nabla^{\beta} \theta \right]_{i}^{\ell} \partial_{\tau} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \right),_{k} dx d\tau'
$$
\n
$$
= \int_{\Omega} w^{1+\alpha+a} \mathcal{J}^{\frac{1}{\alpha}} \mathcal{A}_{\ell}^{k} \Lambda_{im}^{-1} \psi \left[\text{Curl}_{\Lambda \mathcal{A}} \partial_{r}^{\alpha} \nabla^{\beta} \theta \right]_{i}^{\ell} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \right),_{k} dx \Big|_{0}^{\tau}
$$
\n
$$
- \int_{0}^{\tau} \int_{\Omega} w^{1+\alpha+a} \mathcal{J}^{\frac{1}{\alpha}} \mathcal{A}_{\ell}^{k} \Lambda_{im}^{-1} \psi \left[\text{Curl}_{\Lambda \mathcal{A}} \partial_{r}^{\alpha} \nabla^{\beta} \mathbf{V} \right]_{i}^{\ell} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \right),_{k} dx d\tau'
$$
\n
$$
- \int_{0}^{\tau} \int_{\Omega} w^{1+\alpha+a} \partial_{\tau} \left(\mathcal{J}^{\frac{1}{\alpha}} \mathcal{A}_{\ell}^{k} \Lambda_{im}^{-1} \right) \psi \left[\text{Curl}_{\Lambda \mathcal{A}} \partial_{r}^{\alpha} \nabla^{\beta} \theta \right]_{i}^{\ell} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \right),_{k} dx d\tau'
$$
\n
$$
- \int_{0}^{\tau} \int_{\Omega} w^{1+\alpha+a} \mathcal{J}^{\frac{1}{\alpha}} \mathcal{A}_{\ell}^{k} \Lambda_{im}^{-1} \psi \left[\text{Curl}_{\Lambda \tau \mathcal{A}} \partial_{r}^{\alpha} \nabla^{\beta} \theta \right]_{i}^{\ell} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \right),_{k
$$

The first term on the right-hand side of (4.148) is estimated by using the Young inequality and Proposition 3.3:

$$
\int_{\Omega} w^{1+\alpha+a} \mathcal{J}^{\frac{1}{\alpha}} \mathcal{A}_{\ell}^{k} \Lambda_{im}^{-1} \psi \sum_{a+|\beta| \leq N} [\text{Curl}_{\Lambda \mathcal{A}} \partial_{r}^{a} \nabla^{\beta} \theta]_{i}^{\ell} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \right)_{,k} dx \Big|_{0}^{\tau}
$$
\n
$$
\lesssim \mathcal{S}^{N}(0) + \kappa \|\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta\|_{1+\alpha+a,\psi}^{2} + \frac{1}{\kappa} \|\text{Curl}_{\Lambda \mathcal{A}} \partial_{r}^{\alpha} \nabla^{\beta} \theta\|_{1+\alpha+a,\psi}^{2}
$$
\n
$$
\lesssim \mathcal{S}^{N}(0) + \kappa \mathcal{S}^{N}(\tau) + \mathcal{B}^{N}[\theta](\tau) \tag{4.149}
$$

where we used the Young inequality in the first estimate. To bound the second term on the right-hand side of (4.148) we use the Cauchy-Schwarz inequality and the a priori bounds (2.86),

$$
\left| \int_0^{\tau} \int_{\Omega} w^{1+\alpha+a} \mathcal{J}^{\frac{1}{\alpha}} \mathcal{A}_{\ell}^{k} \Lambda_{im}^{-1} \psi \left[\text{Curl}_{\Lambda \mathscr{A}} \partial_r^{\alpha} \nabla^{\beta} \mathbf{V} \right]_{i}^{\ell} \left(\partial_r^{\alpha} \nabla^{\beta} \theta^{m} \right) ,_{k} dx d\tau' \right|
$$

\n
$$
\lesssim \int_0^{\tau} \left\| \nabla_{\eta} \partial_r^{\alpha} \nabla^{\beta} \theta \right\|_{1+\alpha+a,\psi} \left\| \text{Curl}_{\Lambda \mathscr{A}} \partial_r^{\alpha} \nabla^{\beta} \mathbf{V} \right\|_{1+\alpha+a,\psi} d\tau'
$$

\n
$$
\lesssim \int_0^{\tau} (\mathcal{S}^N(\tau'))^{\frac{1}{2}} (\mathcal{B}^N[\mathbf{V}](\tau'))^{\frac{1}{2}} d\tau'. \tag{4.150}
$$

The last two integrals on the right-hand side of (4.148) are easily estimated by using the Young inequality, definition of S^N , and Lemma A.1 and they therefore contribute to the error term R satisfying the bound (4.133).

We next move onto the second integral on the right-hand side of (4.147) , which will give rise to our energy. Note that

$$
\int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\mathcal{A}_{\ell}^{k} \Lambda_{\ell j} [\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta]_{j}^{i} + \frac{1}{\alpha} \Lambda_{ij} \mathcal{A}_{j}^{k} \text{div}_{\eta} (\partial_{r}^{\alpha} \nabla^{\beta} \theta) \right) \Lambda_{im}^{-1} \partial_{\tau} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \right), k \, dy
$$
\n
$$
= \int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\Lambda_{\ell j} [\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta]_{j}^{i} \Lambda_{im}^{-1} [\nabla_{\eta} \partial_{\tau} \partial_{r}^{\alpha} \nabla^{\beta} \theta]_{\ell}^{m} + \frac{1}{\alpha} \text{div}_{\eta} (\partial_{r}^{\alpha} \nabla^{\beta} \theta) \text{div}_{\eta} (\partial_{\tau} \partial_{r}^{\alpha} \nabla^{\beta} \theta) \right) dy
$$
\n
$$
= \int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\Lambda_{\ell j} [\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta]_{j}^{i} \Lambda_{im}^{-1} \partial_{\tau} [\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta]_{\ell}^{m} + \frac{1}{\alpha} \text{div}_{\eta} (\partial_{r}^{\alpha} \nabla^{\beta} \theta) \partial_{\tau} \text{div}_{\eta} (\partial_{r}^{\alpha} \nabla^{\beta} \theta) \right) dy
$$
\n
$$
- \int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\Lambda_{\ell j} [\nabla_{\eta} \partial_{r}^{\alpha} \nabla^{\beta} \theta]_{j}^{i} \Lambda_{im}^{-1} \partial_{\tau} \mathcal{A}_{\ell}^{k} \left(\partial_{r}^{\alpha} \nabla^{\beta} \theta^{m} \right), k + \frac{1}{\alpha} \text{div}_{
$$

The last integral consists of lower order terms and they will contribute to the error term R satisfying (4.133). The source of the exponential-in- τ decay are the terms of the form $\partial_{\tau}\mathscr{A}$ which at the top order behave like DV. By Lemma 4.3, the first integral in the right hand side of (4.151) can be rewritten in the form

$$
\int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\Lambda_{rj} [\nabla_{\eta} \partial_r^{\alpha} \nabla^{\beta} \theta]_{j}^{i} \Lambda_{im}^{-1} \partial_{\tau} [\nabla_{\eta} \partial_r^{\alpha} \nabla^{\beta} \theta]_{\ell}^{m} + \frac{1}{\alpha} \text{div}_{\eta} (\partial_r^{\alpha} \nabla^{\beta} \theta) \partial_{\tau} \text{div}_{\eta} (\partial_r^{\alpha} \nabla^{\beta} \theta) \right) dy
$$
\n
$$
= \frac{1}{2} \frac{d}{d\tau} \left\{ \int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \left(\sum_{i,j=1}^{3} d_i d_j^{-1} (\mathcal{M}_{\alpha,\beta})_i^j)^2 + \frac{1}{\alpha} \left(\text{div}_{\eta} \partial_r^{\alpha} \nabla^{\beta} \theta \right)^2 \right) dy \right\}
$$
\n
$$
+ \int_{\Omega} \psi w^{1+\alpha+a} \mathcal{J}^{-\frac{1}{\alpha}} \mathcal{T}_{\alpha,\beta} dy
$$
\n
$$
- \frac{1}{2} \int_{\Omega} \psi w^{1+\alpha+a} \partial_{\tau} \left(\mathcal{J}^{-\frac{1}{\alpha}} \right) \left(\sum_{i,j=1}^{3} d_i d_j^{-1} (\mathcal{M}_{\alpha,\beta})_i^j)^2 + \frac{1}{\alpha} \left(\text{div}_{\eta} \partial_r^{\alpha} \nabla^{\beta} \theta \right)^2 \right) dy, \qquad (4.152)
$$

where we recall that $\mathcal{M}_{a,\beta}$ is defined in (4.119). Using the a priori bounds (2.86), definition (4.124), and (A.166) it follows that

$$
\Big|\int_{\Omega} \psi w^{1+\alpha+a}\mathscr{J}^{-\frac{1}{\alpha}}\mathcal{T}_{a,\beta}dy\Big|\lesssim e^{-\mu_0\tau}\mathcal{S}^N(\tau).
$$

Similar argument applies to the last term on the right-hand side of (4.152), where the source of an exponentially decaying term is ∂_{τ} \mathscr{J} and we rely on Lemma A.1. Therefore, the last two terms on the right-hand side of (4.152) will contribute to error term R and are bounded by $e^{-\mu_0 \tau} S^N(\tau)$.

From (4.137), (4.144), (4.147), (4.151), (4.149), (4.150), and (4.152) we infer that

$$
\frac{1}{2}\left(\mu^{3\gamma-3}\int_{\Omega}\psi w^{\alpha+a}\langle\Lambda^{-1}\partial_{r}^{a}\nabla^{\beta}\mathbf{V},\partial_{r}^{a}\nabla^{\beta}\mathbf{V}\rangle dy+\delta\int_{\Omega}\psi w^{\alpha+a}\left|\partial_{r}^{a}\nabla^{\beta}\theta\right|^{2}dy\right) \n+\frac{1}{2}\left(\int_{\Omega}\psi w^{1+\alpha+a}\mathscr{J}^{-\frac{1}{\alpha}}\left(\sum_{i,j=1}^{3}d_{i}d_{j}^{-1}(\mathscr{M}_{a,\beta})_{i}^{j}\right)^{2}+\frac{1}{\alpha}\left(\text{div}_{\eta}\partial_{r}^{\alpha}\nabla^{\beta}\theta\right)^{2}\right)dy\right) \n+\int_{0}^{\tau}\frac{5-3\gamma}{2}\mu^{3\gamma-3}\frac{\mu_{\tau}}{\mu}\left(\int_{\Omega}\psi w^{\alpha+a}\langle\Lambda^{-1}\partial_{r}^{\alpha}\nabla^{\beta}\mathbf{V},\partial_{r}^{\alpha}\nabla^{\beta}\mathbf{V}\rangle dy\right)d\tau'\n\lesssim\mathcal{S}^{N}(0)+\mathcal{B}^{N}[\theta](\tau)+\int_{0}^{\tau}\mathcal{S}^{N}(\tau')^{\frac{1}{2}}\mathcal{B}^{N}[\mathbf{V}](\tau')^{\frac{1}{2}}d\tau'+\kappa\mathcal{S}^{N}(\tau)+\int_{0}^{\tau}e^{-\mu_{0}\tau'}\mathcal{S}^{N}(\tau')d\tau'.\tag{4.153}
$$

For any index $\nu = (\nu_1, \nu_2, \nu_3) \in \mathbb{Z}_{\geq 0}^3$ we now commute (2.64) with the operator $(1 - \psi)\partial^{\nu}$, where we recall that the cut-off function ψ is defined in (2.72). Here the estimates will be obtained away from the boundary and therefore, the weight w has a positive lower bound in $B_{\frac{3}{2}}(0)$. Hence the change of the weights is not critical to closing the estimates. A similar computation yields the following energy inequality:

$$
\frac{1}{2}\left(\mu^{3\gamma-3}\int_{\Omega}(1-\psi)w^{\alpha}\langle\Lambda^{-1}\partial^{\nu}\mathbf{V},\partial^{\nu}\mathbf{V}\rangle dy+\delta\int_{\Omega}(1-\psi)w^{\alpha}|\partial^{\nu}\theta|^{2}dy\right) \n+\frac{1}{2}\left(\int_{\Omega}(1-\psi)w^{1+\alpha}\mathscr{J}^{-\frac{1}{\alpha}}\left(\sum_{i,j=1}^{3}d_{i}d_{j}^{-1}(\mathscr{N}_{\nu})_{i}^{j})^{2}+\frac{1}{\alpha}\left(\text{div}_{\eta}\partial^{\nu}\theta\right)^{2}\right)dy\right) \n+\int_{0}^{\tau}\frac{5-3\gamma}{2}\mu^{3\gamma-3}\frac{\mu_{\tau}}{\mu}\left(\int_{\Omega}(1-\psi)w^{\alpha}\langle\Lambda^{-1}\partial^{\nu}\mathbf{V},\partial^{\nu}\mathbf{V}\rangle dy\right)d\tau' \n\lesssim\mathcal{S}^{N}(0)+\mathcal{B}^{N}[\theta](\tau)+\int_{0}^{\tau}\mathcal{S}^{N}(\tau')^{\frac{1}{2}}\mathcal{B}^{N}[\mathbf{V}](\tau')^{\frac{1}{2}}d\tau'+\kappa\mathcal{S}^{N}(\tau)+\int_{0}^{\tau}e^{-\mu_{0}\tau'}\mathcal{S}^{N}(\tau')d\tau'.
$$
\n(4.154)

 \Box

Summing (4.153) and (4.154) we arrive at (4.132).

σ≤τ

5 Nonlinear energy inequality and proof of the main theorem

By the local-in-time well-posedness theory from [18] there exists a unique solution to the initial value problem (2.64)–(2.65) on some time interval $[0, T]$, $T > 0$. See Theorem 2.9. From Propositions 3.3 and 4.6 with $\kappa > 0$ chosen small enough and by using the equivalence of the norm and energy in Proposition 4.2, we conclude that there exist universal constants $c_1, c_2, c_3 \geq 1$ such that for any $0 \leq$ $\sigma \leq \tau \leq T$

$$
\sup_{0\leq\tau'\leq\tau}\mathcal{E}^N(\tau')\leq c_1\mathcal{E}^N(\sigma)+c_2\left(\mathcal{S}^N(0)+\mathcal{B}^N[\mathbf{V}](0)\right)+c_3\int_{\sigma}^{\tau}e^{-\frac{\mu_0}{2}\tau'}\mathcal{S}^N(\tau';\sigma)d\tau'.\tag{5.155}
$$

Here $S^N(\tau;\sigma)$ denotes the sliced norm of S^N from σ to τ with $\sup_{0\leq \tau'\leq \tau}$ replaced by $\sup_{\sigma\leq \tau'\leq \tau}$ in the definition of S^N (2.77). Note that $S^N(\tau;0) = S^N(\tau)$ and $S^N(\tau) = \max\{S^N(\sigma), S^N(\tau;\sigma)\}\$ for any $\sigma \in [0, \tau]$. When $\sigma = 0$, the estimate (5.155) is a direct consequence of Propositions 3.3 and 4.6, while for general $\sigma > 0$ the estimate follows by an identical argument, applying our analysis to the interval $[\sigma, \tau]$.

By a standard well-posedness estimate, we deduce that the time of existence T is inversely proportional to the size of the initial data, i.e.: $T \sim (S^N(0) + B^N[\mathbf{V}](0))$. Choose $\varepsilon > 0$ so small that the time of existence T satisfies

$$
e^{-\mu_0 T/4} \le \frac{\kappa \mu_0}{c_3}, \ \ \sup_{\tau \le T} \mathcal{S}^N(\tau) \le C \left(\mathcal{S}^N(0) + \mathcal{B}^N[\mathbf{V}](0) \right) \tag{5.156}
$$

where $C > 0$ is a universal constant provided by the local-in-time well-posedness theory. Let

$$
C_* = \frac{3(c_1C_2C + c_2)}{C_1}
$$

where $C_1, C_2, (C_1 \leq C_2)$ are the constants appearing in the equivalence of the energy and the norm given in Proposition 4.2. Define

$$
\mathcal{T} := \sup_{\tau \geq 0} \{ \text{ solution to (2.64)} - (2.65) \text{ exists on } [0, \tau) \text{ and } \mathcal{S}^N(\tau) \leq C_* \left(\mathcal{S}^N(0) + \mathcal{B}^N[\mathbf{V}](0) \right) \}.
$$

Observe that $\mathcal{T} \geq T$. Letting $\sigma = \frac{T}{2}$ in (5.155) for any $\tau \in [\frac{T}{2}, \mathcal{T}]$ we obtain

$$
\sup_{\tau' \in [\frac{T}{2},\tau]} \mathcal{E}^N(\tau') \le c_1 \mathcal{E}^N(\frac{T}{2}) + c_2 \left(\mathcal{S}^N(0) + \mathcal{B}^N[\mathbf{V}](0) \right) + c_3 \int_{\frac{T}{2}}^{\tau} e^{-\frac{\mu_0}{2}\tau'} \mathcal{S}^N(\tau';\frac{T}{2}) d\tau'.
$$

Therefore, using (5.156) we conclude that for any $\tau \in [\frac{T}{2}, \mathcal{T}]$

$$
\sup_{\tau' \in [\frac{T}{2}, \tau]} \mathcal{E}^{N}(\tau') \le c_1 \mathcal{E}^{N}(\frac{T}{2}) + c_2 \left(\mathcal{S}^{N}(0) + \mathcal{B}^{N}[\mathbf{V}](0) \right) + \frac{c_3}{\mu_0} e^{-\mu_0 T/4} \mathcal{S}^{N}(\tau; \frac{T}{2})
$$

$$
\le c_1 C_2 \mathcal{S}^{N}(\frac{T}{2}) + c_2 \left(\mathcal{S}^{N}(0) + \mathcal{B}^{N}[\mathbf{V}](0) \right) + \kappa \mathcal{S}^{N}(\tau; \frac{T}{2}). \tag{5.157}
$$

Since by (5.156), $S^{N}(\frac{T}{2}) \leq C(S^{N}(0) + \mathcal{B}^{N}[\mathbf{V}](0))$ we conclude from (5.157) that

$$
C_1\mathcal{S}^N(\tau;\frac{T}{2})\leq (c_1C_2C+c_2)\left(\mathcal{S}^N(0)+\mathcal{B}^N[\mathbf{V}](0)\right)+\kappa\mathcal{S}^N(\tau;\frac{T}{2}),
$$

which for sufficiently small κ gives

$$
S^{N}(\tau; \frac{T}{2}) \leq \frac{2(c_{1}C_{2}C + c_{2})}{C_{1}} (\mathcal{S}^{N}(0) + \mathcal{B}^{N}[\mathbf{V}](0)) < C_{*} (\mathcal{S}^{N}(0) + \mathcal{B}^{N}[\mathbf{V}](0))
$$

and hence $S^N(\tau) < C_* \left(S^N(0) + \mathcal{B}^N[\mathbf{V}](0) \right)$. It is now easy to check the a priori bounds in (2.85) and (2.86) are in fact improved. For instance, by the fundamental theorem of calculus

$$
||D\theta||_{W^{1,\infty}} = ||\int_0^{\tau} D\mathbf{V}||_{W^{1,\infty}} \leq \int_0^{\tau} e^{-\mu_0 \tau'} \mathcal{S}^N(\tau') d\tau' \lesssim \varepsilon < \frac{1}{6}, \ \ \tau \in [0, \mathcal{T})
$$

for $\varepsilon > 0$ small enough. Similar arguments apply to the remaining a priori assumptions.

From the continuity of the map $\tau \mapsto \mathcal{S}^N(\tau')$ and the definition of $\mathcal T$ we conclude that $\mathcal T = \infty$ and the solution to (2.64)–(2.65) exists globally-in-time. Together with Proposition 3.3 the global bound (2.80) follows.

Note that by Lemma A.1 $\mu(\tau) \sim_{\tau \to \infty} e^{\mu_1 \tau}$ where $\mu_1 > 0$ is defined in (2.79), see also Lemma A.1. Statement (2.82) follows from the already established global bound $S^N(\tau) \leq C\varepsilon$ and the presence of the exponentially growing term $\mu^{3\gamma-3}$ in the definition (2.77) of \mathcal{S}^N . Estimate (2.81) is a simple consequence of (2.80) and Proposition 3.3.

From the global-in-time boundedness of \mathcal{S}^N , there exists a weak limit θ_{∞} independent of τ such that $\mathcal{S}^N(\theta_\infty, 0) \lesssim \mathcal{S}^N(\tau) \leq C \varepsilon$. Observe that for any $0 < \tau_2 < \tau_1$,

$$
\begin{split}\n&\|\partial_r^a \nabla^{\beta} \theta(\tau_1) - \partial_r^a \nabla^{\beta} \theta(\tau_2)\|_{a+\alpha,\psi}^2 = \int_{\Omega} \psi w^{\alpha+a} \left| \int_{\tau_2}^{\tau_1} \partial_r^a \nabla^{\beta} \theta_\tau d\tau \right|^2 dy \\
&\leq \left(\int_{\tau_2}^{\tau_1} \mu^{\frac{3-3\gamma}{2}} d\tau \right) \left(\int_{\tau_2}^{\tau_1} \mu^{\frac{3\gamma-3}{2}} \int_{\Omega} \psi w^{\alpha+a} \left| \partial_r^a \nabla^{\beta} \mathbf{V} \right|^2 dy d\tau \right) \\
&\lesssim (e^{-\frac{3\gamma-3}{2}\mu_0 \tau_2} - e^{-\frac{3\gamma-3}{2}\mu_0 \tau_1}) \int_{\tau_2}^{\tau_1} \mu^{-\frac{3\gamma-3}{2}} \mu^{3\gamma-3} \int_{\Omega} \psi w^{\alpha+a} \left| \partial_r^a \nabla^{\beta} \mathbf{V} \right|^2 dy d\tau \\
&\lesssim \varepsilon (e^{-\frac{3\gamma-3}{2}\mu_0 \tau_2} - e^{-\frac{3\gamma-3}{2}\mu_0 \tau_1}),\n\end{split}
$$

where the last line follows from the integrability of $\int_{\tau_2}^{\tau_1} \mu^{-\frac{3\gamma-3}{2}} d\tau$ and the fact that $\sup_{0\leq \tau'\leq \tau}\left(\mu^{3\gamma-3}\|\partial_r^a\nabla^{\beta}\mathbf{V}(\tau')\|_{L^2_{\alpha+a,\psi}}^2\right)\lesssim \mathcal{S}^N(\tau)\lesssim \varepsilon\text{ for any }\tau\geq 0.$ Therefore, given a strictly increasing sequence $\tau_n \to \infty$, the sequence $\{\partial_n^{\alpha} \nabla^{\beta} \theta(\tau_n)\}_{n=1}^{\infty}$ is Cauchy in $L^2_{a+\alpha,\psi}$ for any (a,β) satisfying $a + |\beta| \leq N$. By an analogous argument we can show that the sequence $\{\partial^{\nu} \theta(\tau_n)\}_{n=1}^{\infty}$ is Cauchy in $L^2_{\alpha,1-\psi}$. This completes the proof of (2.83).

Acknowledgements

The authors express their gratitude to P. Raphaël for fruitful discussions and for pointing out connections to the treatment of self-similar singular behavior for nonlinear Schrödinger equations. They also thank C. Dafermos for his feedback and pointing out important references. JJ is supported in part by NSF grants DMS-1608492 and DMS-1608494 and a von Neumann fellowship of the Institute for Advanced Study through the NSF grant DMS-1128155. MH acknowledges the support of the EPSRC Grant EP/N016777/1.

A Asymptotic-in- τ behavior of affine solutions

In this section we collect some of the basic properties of affine motions that are used at many places in our estimates. Their proofs are rather straightforward and follow directly from the description of the asymptotic behavior of the solutions of (2.44) from [44].

We remind the reader that for any $M \in \mathbb{M}^{3 \times 3}$ we denote by $||M||$ the Hilbert-Schmidt norm of the matrix M.

Lemma A.1 (Asymptotic behavior of $A, \Gamma^* = O^{-1}O_\tau$, and $\Lambda = O^{-1}O^{-\top}$). For any $\gamma \in (1, \frac{5}{3}]$ and any pair of initial conditions $(A(0),\dot{A}(0))\in$ $\rm GL^+(3)\times\mathbb{M}^{3\times 3}$ there exist matrices $A_0,A_1,M(t)$ such *that the unique solution* A(t) *to the Cauchy problem*

$$
A_{tt} = \delta \det A^{1-\gamma} A^{-\top} \tag{A.158}
$$

$$
A(0) = A(0), At(0) = \dot{A}(0)
$$
 (A.159)

can be written in the form

$$
A(t) = A_0 + tA_1 + M(t), \quad t \ge 0,
$$
\n(A.160)

where A_0 , A_1 *are both time-independent and* $M(t)$ *satisfies the bounds*

$$
||M(t)|| = o_{t \to \infty}(1+t), ||\partial_t M(t)|| \lesssim (1+t)^{3-3\gamma}.
$$
 (A.161)

Moreover

$$
e^{\mu_1 \tau} \lesssim \mu(\tau) \lesssim e^{\mu_1 \tau}, \ \ \tau \ge 0,
$$
\n(A.162)

$$
\lim_{\tau \to \infty} \|\Gamma^* - e^{-\mu_1 \tau} \mu_1 A_0 A_1^{-1}\| = 0,
$$
\n(A.163)

where

$$
\Gamma^* = O^{-1}O_\tau, \quad \mu_1 = (\det A_1)^{\frac{1}{3}} > 0.
$$

Furthermore there exists a constant C > 0 *such that*

$$
\|\Lambda_{\tau}\| \le Ce^{-\mu_1 \tau}, \ \|\Lambda_{\tau\tau}\| \le Ce^{-2\mu_0 \tau}, \ \|\Lambda\| + \|\Lambda^{-1}\| \le C,
$$
\n(A.164)

$$
\sum_{i=1}^{3} \left(d_i + \frac{1}{d_i} \right) \le C \tag{A.165}
$$

$$
\sum_{i=1}^{3} |\partial_{\tau} d_i| + ||\partial_{\tau} P|| \leq Ce^{-\mu_1 \tau}
$$
\n(A.166)

$$
\frac{1}{C}|\mathbf{w}|^2 \le \langle \Lambda^{-1}\mathbf{w}, \mathbf{w} \rangle \le C|\mathbf{w}|^2, \ \mathbf{w} \in \mathbb{R}^3. \tag{A.167}
$$

where d_i , $i = 1, 2, 3$, are the eigenvalues of the matrix Λ and $P \in SO(3)$ satisfies $\Lambda = P^{\top}QP$, $Q = \text{diag}(d_i)$.

Proof. Asymptotic behavior (A.160)–(A.161) and bound (A.162) are a consequence of Theorem 3 and Lemma 6 from [44].

Proof of (A.163). Since $O = \frac{A}{\mu}$ we have

$$
O_t = \frac{A_t}{\mu} - \frac{A\mu_t}{\mu^2} = \frac{A_t}{\mu} - \frac{A \text{Tr}(A^{-1}A_t)}{3\mu},
$$
 (A.168)

where we used the formula

$$
\mu_t = \partial_t \left((\det A)^{\frac{1}{3}} \right) = \frac{1}{3} \mu^{-2} \partial_t \det A = \frac{1}{3} \mu^{-2} \mu^3 \text{Tr}(A^{-1} A_t) = \frac{1}{3} \mu \text{Tr}(A^{-1} A_t).
$$

Therefore, from (A.160) and (A.161) it is easy to obtain the following asymptotics:

$$
\operatorname{Tr}(A^{-1}A_t) \sim \operatorname{Tr}((A_0 + tA_1 + M(t))^{-1}(A_1 + \partial_t M(t))) = t^{-1}\operatorname{Tr}((\frac{A_0}{t} + A_1)^{-1}(A_1 + O(t^{3-3\gamma}))
$$

$$
\sim t^{-1}\operatorname{Tr}(A_1^{-1}(A_1 + O(t^{3-3\gamma})) = \frac{3}{t} + O(t^{2-3\gamma}).
$$
\n(A.169)

This implies that

$$
O_t \sim \frac{A_1}{\mu} - \frac{(A_0 + tA_1)\frac{3}{t}}{3\mu} = \frac{A_0}{t\mu},
$$

where we made use of (A.161) again. Recalling that $\frac{d\tau}{dt} = \frac{1}{\mu}$ we obtain the following asymptotic behavior $O_{\tau} \sim_{\tau \to \infty} \frac{A_0}{t(\tau)}$. Using (A.160) again it follows that

$$
\det A \sim \det(A_0 + tA_1) = t^3 \det(\frac{A_0}{t} + A_1) \sim t^3 \det A_1, \ \mu \sim t (\det A_1)^{\frac{1}{3}}.
$$
 (A.170)

Since

$$
t = t(\tau) \sim_{\tau \to \infty} e^{(\det A_1)^{\frac{1}{3}} \tau}.
$$

and $O \sim \frac{A_0 + tA_1}{\mu}$ we conclude that

$$
O^{-1} \sim \frac{\mu}{t} \left(\frac{A_0}{t} + A_1\right)^{-1} \sim \frac{\mu}{t} A_1^{-1} \sim (\det A_1)^{\frac{1}{3}} A_1^{-1},
$$

where we used (A.170). Therefore

$$
\Gamma^* = O^{-1}O_\tau \sim (\det A_1)^{\frac{1}{3}} A_1^{-1} \frac{A_0}{t(\tau)} = e^{-\mu_1 \tau} \mu_1 A_0 A_1^{-1},
$$

where $\mu_1 = (\det A_1)^{\frac{1}{3}}$, and this completes the proof of (A.163). Proof of (A.164) is similar.

Proof of (A.164)–(A.166). From the definition of Λ we have

$$
\Lambda_{\tau} = -O^{-1}O_{\tau}O^{-1}O^{-\top} - O^{-1}O^{-\top}O_{\tau}^{\top}O^{-\top} = -\Gamma^{*}\Lambda - \Lambda(\Gamma^{*})^{\top} = -2\Lambda(\Gamma^{*})^{\top}, \quad (A.171)
$$

where we used the symmetry of Λ in the last equality. Since $\|\Lambda\| \leq 1$ it follows by part (i) that $\|\Lambda_{\tau}\| \lesssim e^{-\mu_1 \tau}$. To bound $\Lambda_{\tau\tau}$ we note that by (A.171) $\Lambda_{\tau\tau} = -2\Lambda_{\tau}(\Gamma^*)^{\top} - 2\Lambda(\Gamma^*)^{\top}$. Since both

 $\|\Lambda_{\tau}\|$ and $\|\Gamma^*\|$ decay exponentially, it remains to prove the decay of $\|\Gamma_{\tau}^*\|$. From $\Gamma^* = O^{-1}O_{\tau}$ it follows that $\Gamma_\tau^* = -(\Gamma^*)^2 + O^{-1}O_{\tau\tau}$, and therefore it remains to prove the decay of $||O_{\tau\tau}||$. A simple calculation shows that

$$
O_{\tau\tau} = \mu \left(A_{tt} - \frac{A_t \text{Tr}(A^{-1}A_t) + A\partial_t \text{Tr}(A^{-1}A_t)}{3} \right).
$$

Using the asymptotic behavior (A.169), (2.44), we can refine the asymptotics (A.169) to show that A_t Tr $(A^{-1}A_t) + A\partial_t$ Tr $(A^{-1}A_t) = O(t^{2-3\gamma})$ and therefore from the above equation it follows that

$$
||O_{\tau\tau}|| \lesssim (1+t)^{3-3\gamma} \lesssim e^{-2\mu_0\tau}.
$$

This yields the second bound in (A.164). Bounds (A.165) (A.166) follow by similar arguments using (A.160), while (A.167) is a direct consequence of (A.165). \Box

B Commutators

In order to evaluate various commutator terms that arise from commuting differential operators $\partial_r^a \vec{\nabla}^\beta$ with the usual Cartesian derivatives or apply the Leibniz rule we shall rely on the fact that the high-order Sobolev norms expressed in polar coordinates are equivalent to the usual high-order Sobolev norms on the support of function ψ .

Lemma B.1. *Let* $\mathcal{X} := \{\partial_i, \nabla_j, \partial_r\}_{i,j=1,2,3}$ *be a collection of the standard Cartesian, normal, and tangential vector-fields. For any two vector fields* $X_k, X_\ell \in \mathcal{X}, k, \ell = 1, \ldots, 7$ *there commutator satisfies the following relationship*

$$
[X_k, X_\ell] = \sum_{m=1}^3 c_{k\ell}^m \nabla_m + c_{k\ell} \partial_r, \ \ k, \ell = 1, \ldots 7,
$$

where the functions $c_{k\ell}^m$, $c_{k\ell}$, are C^{∞} on the exterior of any ball around the origin $r = 0$.

Proof. The proof is a simple consequence of the following direct calculations. For any $i, j \in \{1, 2, 3\}$ we have

$$
[\partial_r, \partial_i] = [\partial_r, \nabla_i] = -\frac{1}{r} \nabla_i, \quad [\partial_i, \partial_j] = 0,
$$

\n
$$
[\nabla_i, \nabla_j] = \frac{y_i \nabla_j - y_j \nabla_i}{r^2},
$$

\n
$$
[\partial_i, \nabla_j] = -\frac{y_j \nabla_i}{r^2} + \Delta_{ij} \partial_r, \quad \Delta_{ij} := \frac{y_i y_j - \delta_{ij} r^2}{r^3}.
$$
 (B.172)

 \Box

Lemma B.1 is a technical tool allowing us to bound the lower order commutators in our energy estimates.

Using the product rule and the relationship $\mathscr{A} = [D\eta]^{-1}$ it is easy to see that the following formulas hold

$$
\partial_r \mathscr{A}_i^k = -\mathscr{A}_s^k \partial_r \partial_m \eta^s \mathscr{A}_i^m = -\mathscr{A}_s^k \partial_r (\partial_m \theta^s + \delta_m^s) \mathscr{A}_i^m
$$

= $-\mathscr{A}_s^k (\partial_r \theta^s),_m \mathscr{A}_i^m + \mathscr{A}_s^k [\partial_m, \partial_r] \theta^s \mathscr{A}_i^m.$ (B.173)

Similarly, for any $j = 1, 2, 3$ we have

$$
\nabla_j \mathscr{A}_i^k = -\mathscr{A}_s^k (\nabla_j \theta^s),_m \mathscr{A}_i^m + \mathscr{A}_s^k [\partial_m, \nabla_j] \theta^s \mathscr{A}_i^m.
$$
 (B.174)

Therefore, if $\beta = (0, 0, 0)$ we have the formula

$$
\partial_r^a \nabla^{\beta} \mathscr{A}_i^k = \partial_r^a \mathscr{A}_i^k = -\mathscr{A}_s^k (\partial_r^a \theta^s),_m \mathscr{A}_i^m - \mathscr{A}_r^k [\partial_r^{a-1}, \partial_m] \partial_r \theta^s \mathscr{A}_i^m
$$

$$
- \sum_{1 \le a' \le a} c_{a'} \partial_r^{a'} (\mathscr{A}_s^k \mathscr{A}_i^m) \partial_r^{a-a'} (\partial_r \theta^s),_m + \partial_r^{a-1} (\mathscr{A}_s^k [\partial_m, \partial_r] \theta^s_{,j} \mathscr{A}_i^m).
$$

(B.175)

If $|\beta| > 0$, then with $e_1 = (1, 0, 0), e_2 = (0, 1, 0),$ and $e_3 = (0, 0, 1)$ we obtain

$$
\partial_r^{\alpha} \nabla^{\beta} \mathcal{A}_i^k = -\mathcal{A}_s^k (\partial_r^{\alpha} \nabla^{\beta} \theta^s),_m \mathcal{A}_i^m - \mathcal{A}_r^k [\partial_r^{\alpha} \nabla^{\beta - e_j}, \partial_m] \nabla_j \theta^s \mathcal{A}_i^m
$$

$$
- \sum_{0 < a' + |\beta'| \leq a + |\beta| \atop \beta' \leq \beta - e_j} c_{a' \beta'} \partial_r^{\alpha'} \nabla^{\beta'} (\mathcal{A}_s^k \mathcal{A}_i^m) \partial_r^{\alpha - a'} \nabla^{\beta - e_j - \beta'} (\nabla_j \theta^s),_m
$$

$$
- \partial_r^{\alpha} \nabla^{\beta - e_j} (\mathcal{A}_s^k [\partial_m, \nabla_j] \theta^s \mathcal{A}_i^m), \qquad (B.176)
$$

where $c_{a' \beta'}$, $c_{\beta'}$ are positive universal constants and the Einstein summation convention does not apply to the index j . Finally, the high-order commutators appearing on the right-hand side in the identities (B.175)–(B.176) can be expressed as a linear combination of the elements of X with smooth coefficients away from zero. In other words, for any $j = 1, 2, 3$ the following commutator identity holds:

$$
[\partial_r^a \nabla^\beta, \partial_j] = \sum_{\ell=0}^{a+|\beta|} \sum_{\substack{a'+|\beta'|=\ell\\ a'\leq a+1}} C_{a',\beta',\ell}^j \partial_r^{a'} \nabla^{\beta'}
$$
(B.177)

for some universal coefficients $C^j_{a',\beta',\ell}$ which are smooth on the exterior of any ball around the origin $r = 0$. Formula (B.177) is a direct consequence of Lemma B.1.

C Weighted spaces, Hardy inequalities, and Sobolev embeddings

In this section, we recall Hardy inequalities and embedding results of weighted function spaces. First of all, we state the Hardy inequality near $r = 1$.

Lemma C.1. *(Hardy inequality [19]) Let* k *be a real number and g a function satisfying* $\int_0^1 (1 (r)^{k+2}(g^2+g^2)dr < \infty$.

If $k > -1$, then we have $\int_0^1 (1 - r)^k g^2 dr \leq C \int_0^1 (1 - r)^{k+2} (g^2 + |g'|^2) dr$. *If* $k < -1$, then g has a trace at $r = 1$ and $\int_0^1 (1 - r)^k (g - g(1))^2 dr \le C \int_0^1 (1 - r)^{k+2} |g'|^2 dr$.

Since w depends only on r and w behaves like a distance function $1 - r$, using Lemma C.1, we in particular get

$$
\int_{B_1(0)\setminus B_{\frac{1}{4}}(0)} \psi w^k |u|^2 dy \lesssim \int_{B_1(0)\setminus B_{\frac{1}{4}}(0)} \psi w^{k+2} \left(|\partial_r u|^2 + |u|^2 \right) dy \tag{C.178}
$$

for any nonnegative real number $k > -1$ and for any $u \in C^{\infty}(B_1(\mathbf{0}) \setminus B_{\frac{1}{4}}(\mathbf{0}))$. We can apply (C.178) to $\partial_r^a \nabla^{\beta} u$ and thereafter apply (C.178) repeatedly to the right-hand side. As a consequence for any $-1 < k < \alpha + a$,

$$
\int_{B_1(\mathbf{0})\backslash B_{\frac{1}{4}}(\mathbf{0})} \psi w^k |\partial_r^{\alpha} \nabla^{\beta} u|^2 dy \lesssim \int_{B_1(\mathbf{0})\backslash B_{\frac{1}{4}}(\mathbf{0})} \psi w^{k+2} \left(|\partial_r \partial_r^{\alpha} \nabla^{\beta} u|^2 + |\partial_r^{\alpha} \nabla^{\beta} u|^2 \right) dy
$$
\n
$$
\lesssim \sum_{j=0}^m \int_{B_1(\mathbf{0})\backslash B_{\frac{1}{4}}(\mathbf{0})} \psi w^{k+2j} |\partial_r^j \partial_r^{\alpha} \nabla^{\beta} u|^2 dy.
$$
\n(C.179)

Upon choosing $m = [a + \alpha - k]$ where $\lceil \cdot \rceil$ is the ceiling function, we obtain

$$
\|\partial_r^{\alpha}\nabla^{\beta}u\|_{k,\psi}^2 \lesssim \sum_{j=0}^{\lceil a+\alpha-k\rceil} \|\partial_r^j \partial_r^{\alpha}\nabla^{\beta}u\|_{\alpha+a+j,\psi}^2
$$
 (C.180)

for any $u \in C^{\infty}(B_1(\mathbf{0}) \setminus B_{\frac{1}{2}}(\mathbf{0})).$

As a consequence of $(C.\overline{180})$, we obtain the weighted Sobolev-Hardy inequality:

Proposition C.2. *For any* $u \in C^{\infty}(B_1(\mathbf{0}))$ *, we have*

$$
\sup_{B_1(0)\setminus B_{\frac{1}{4}}(0)} \left| w^{\frac{a_1}{2}} \partial_r^{a_1} \nabla^{\beta_1} u \right|
$$
\n
$$
\lesssim \sum_{a+|\beta| \le a_1 + |\beta_1| + \lceil \alpha \rceil + 6} \|\partial_r^a \nabla^\beta u\|_{a+\alpha,\psi} + \sum_{|\nu| \le a_1 + |\beta_1| + 2} \|\partial^\nu u\|_{\alpha,1-\psi}, \tag{C.181}
$$
\n
$$
\sup_{B_1(0)\setminus B_{\frac{1}{4}}(0)} \left| w^{\frac{a_1}{2}} D \partial_r^{a_1} \nabla^{\beta_1} u \right|
$$
\n
$$
\lesssim \sum_{a+|\beta| \le a_1 + |\beta_1| + \lceil \alpha \rceil + 6} \|\nabla_\eta \partial_r^a \nabla^\beta u\|_{a+\alpha+1,\psi} + \sum_{|\nu| \le a_1 + |\beta_1| + 2} \|\nabla_\eta \partial^\nu u\|_{\alpha+1,1-\psi}. \tag{C.182}
$$

We omit the technical details of the proof, as it follows from standard estimates relying on the $H^2(\Omega) \hookrightarrow L^{\infty}(\Omega)$ continuous embedding and the Hardy inequality (C.180).

D Starting from Lagrangian formulation

Consider the Lagrangian formulation of the E_{γ} -system [5, 18]:

$$
\zeta_{tt} + \frac{\gamma}{\gamma - 1} \mathscr{A}_{\zeta}^T \nabla (w \mathscr{J}_{\zeta}^{1 - \gamma}) = 0
$$
 (D.183)

where \mathcal{A}_{ζ} and \mathcal{J}_{ζ} are induced by the flow map $\partial_t \zeta(t, y) = \mathbf{u}(t, \zeta(t, y))$ (**u** is the original fluid velocity appearing in (1.1)) and $w = (\rho_0 \mathcal{J}_\zeta(0))^{\gamma-1} \geq 0$ is an enthalpy function that depends only on the initial data. To find affine motions (with center of mass at 0) one makes the ansatz $\zeta(t,y) = A(t)y$ [44]. With $\mathscr{A}_{\zeta}^{\top} = [D\zeta]^{-\top} = A(t)^{-\top}$ and $\mathscr{J}_{\zeta} = \det A$, by plugging this ansatz into (D.183), we obtain

$$
A_{tt}y + \frac{\gamma}{\gamma - 1} (\det A)^{1 - \gamma} A^{-\top} \nabla w = 0
$$
 (D.184)

Since w is independent of t , (D.184) will be satisfied if we demand

$$
A_{tt} = \delta(\det A)^{1-\gamma} A^{-\top}, \quad \delta y = -\frac{\gamma}{\gamma - 1} \nabla w, \quad \delta > 0,
$$
 (D.185)

which precisely yields the affine motions (1.5)–(1.6) (discovered by Sideris when $a \equiv 0$) which form the set $\mathscr S$ with the center of mass fixed at the origin.

In order to study the stability of elements of $\mathscr S$ we want to realize them as time-independent background solutions. Given such an affine motion A we modify the flow map ζ and define $\eta := A^{-1}\zeta$. Since $\mathscr{A}_{\zeta}^{\top} = A^{-\top} \mathscr{A}_{\eta}^{\top}$ and $\mathscr{J}_{\zeta} = (\det A) \mathscr{J}_{\eta}$, we obtain

$$
\eta_{tt} + 2A^{-1}A_t \eta_t + A^{-1}A_{tt}\eta + \frac{\gamma}{\gamma - 1} (\det A)^{1 - \gamma} A^{-1} A^{-\top} \mathscr{A}_{\eta}^{\top} \nabla (w \mathscr{J}_{\eta}^{1 - \gamma}) = 0
$$

By using (D.185) we can rewrite the previous equation in the form

$$
(\det A)^{\gamma - \frac{1}{3}} \eta_{tt} + 2(\det A)^{\gamma - \frac{1}{3}} A^{-1} A_t \eta_t + \delta \Lambda \eta + \frac{\gamma}{\gamma - 1} \Lambda \mathscr{A}_{\eta}^{\top} \nabla (w \mathscr{J}_{\eta}^{1 - \gamma}) = 0 \tag{D.186}
$$

where $\Lambda = \det A^{\frac{2}{3}} A^{-1} A^{-T}$.

We rescale the time variable t so that $1 + t \sim e^{\mu_1 \tau}$ by setting $\frac{d\tau}{dt} = (\det A)^{-\frac{1}{3}}$. Then (D.186) can be written as

$$
(\det A)^{\gamma - 1} \eta_{\tau \tau} - \frac{1}{3} (\det A)^{\gamma - 2} (\det A)_{\tau} \eta_{\tau} + 2 (\det A)^{\gamma - 1} A^{-1} A_{\tau} \eta_{\tau} + \delta \Lambda \eta + \frac{\gamma}{\gamma - 1} \Lambda \mathscr{A}_{\eta}^{\top} \nabla (w \mathscr{J}_{\eta}^{1 - \gamma}) = 0
$$
\n(D.187)

We now recall $A^{-1}A_{\tau} = \mu^{-1}\mu_{\tau}I + O^{-1}O_{\tau}$ where $A = \mu O$ and $\mu = (\det A)^{\frac{1}{3}}$ (see Section 2.1). The equation for η reads

$$
\mu^{3\gamma - 3} \eta_{\tau \tau} + \mu^{3\gamma - 4} \mu_{\tau} \eta_{\tau} + 2\mu^{3\gamma - 3} \Gamma^* \eta_{\tau} + \delta \Lambda \eta + \frac{\gamma}{\gamma - 1} \Lambda \mathscr{A}_{\eta}^{\top} \nabla (w \mathscr{J}_{\eta}^{1 - \gamma}) = 0 \tag{D.188}
$$

where $\Gamma^* = O^{-1}O_{\tau}$. It is clear that $\eta(y) \equiv y$ corresponds to Sideris' affine motions, and equation (D.188) is nothing but (2.61). By considering $\theta = \eta - y$, we obtain the θ -equation (2.64).

References

- [1] CHEN, G.-Q.: Remarks on R. J. DiPerna's paper: "Convergence of the viscosity method for isentropic gas dynamics" [Comm. Math. Phys. 91 (1983), no. 1, 1–30; MR0719807 (85i:35118)]. *Proc. Amer. Math. Soc.*, 125(10):2981–2986, 1997.
- [2] CHIODAROLI, E., DE LELLIS, C., KREML, O. Global ill-posedness of the isentropic system of gas dynamics. *Comm. Pure Appl. Math.* 68, no. 7, 1157–1190 (2015)
- [3] CHRISTODOULOU, D.: The Formation of Shocks in 3-Dimensional Fluids. *EMS Monographs in Mathematics, EMS Publishing House* 2007.
- [4] COUTAND, D., SHKOLLER, S.: Well-posedness in smooth function spaces for the movingboundary 1-D compressible Euler equations in physical vacuum, *Comm. Pure Appl. Math.* 64, no. 3, 328–366 (2011)
- [5] COUTAND, D., SHKOLLER, S.: Well-posedness in smooth function spaces for the moving boundary three-dimensional compressible Euler equations in physical vacuum. *Arch. Ration. Mech. Anal.* 206, no. 2, 515–616 (2012)
- [6] DACOROGNA B., MOSER, J.: On a partial differential equation involving the Jacobian determinant, *Ann. Inst. H. Poincare Anal. Non Lin ´ eaire ´* 7 (1990), no. 1, 1-26
- [7] DAFERMOS, C.: *Hyperbolic conservation laws in continuum physics*. Second edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 325. Springer-Verlag, Berlin, 2005.
- [8] DIPERNA, R. J. Convergence of the viscosity method for isentropic gas dynamics. *Comm. Math. Phys.* 91, 1–30 (1983)
- [9] EGGERS J., FONTELOS, A. M.: The role of self-similarity in singularities of partial differential equations. *Nonlinearity* 22 R1–R44 (2009)
- [10] FRIEDRICH, H.: Sharp asymptotics for Einstein-λ-dust flows. *Available online at https://arxiv.org/abs/1601.04506*
- [11] GRASSIN, M.: Global smooth solutions to Euler equations for a perfect gas. *Indiana Univ. Math. J.* 47, 1397-1432 (1998)
- [12] HADŽIĆ, M., JANG, J.: Nonlinear stability of expanding star solutions in the radially-symmetric mass-critical Euler-Poisson system. To appear in *Comm. Pure Appl. Math.*
- [13] HADŽIĆ, M., SPECK, J.: The Global Future Stability of the FLRW Solutions to the Dust-Einstein System with a Positive Cosmological Constant. *Jour. Hyp. Diff. Eqns.,* 12, no.1, 87–188 (2015)
- [14] HUANG, H., MARCATI, P., PAN, R.: Convergence to the Barenblatt solution for the compressible Euler equations with damping and vacuum, *Arch. Ration. Mech. Anal.* 176, 1-24 (2005)
- [15] JANG, J., MASMOUDI, N. Well-posedness for compressible Euler equations with physical vacuum singularity, *Comm. Pure Appl. Math.* 62 (2009), 1327–1385
- [16] JANG, J., MASMOUDI, N.: Vacuum in Gas and Fluid dynamics, *Proceedings of the IMA summer school on Nonlinear Conservation Laws and Applications*, Springer (2011), 315-329
- [17] JANG, J., MASMOUDI, N.: Well and ill-posedness for compressible Euler equations with vacuum. *J. Math. Phys.* 53 (11) (2012) 115625
- [18] JANG, J., MASMOUDI, N. Well-posedness of compressible Euler equations in a physical vacuum. *Communications on Pure and Applied Mathematics* 68, no. 1, 61–111 (2015)
- [19] KUFNER, A., MALGRANDA, L., PERSSON, L.-E. *The Hardy inequality,* Vydavatelsky Servis, ´ Plzen, 2007.
- [20] LINDBLAD, H.: Well posedness for the motion of a compressible liquid with free surface boundary. *Comm. Math. Phys.* 260, 319–392 (2005)
- [21] LIONS, P. L., PERTHAME, B., SOUGANIDIS, P. E. Existence and stability of entropy solutions for the hyperbolic systems of isentropic gas dynamics in Eulerian and Lagrangian coordinates. *Comm. Pure Appl. Math.*, 49(6):599–638, 1996.
- [22] LIU, T.-P.: Compressible flow with damping and vacuum, *Japan J. Appl. Math* 13, 25-32 (1996)
- [23] LIU, T.-P., SMOLLER, J.: On the vacuum state for isentropic gas dynamics equations, *Advances in Math.* 1, 345-359 (1980)
- [24] LIU, T.-P., YANG, T.: Compressible Euler equations with vacuum, *J. Differential Equations* 140, 223-237 (1997)
- [25] LIU, T.-P., YANG, T.: Compressible flow with vacuum and physical singularity, *Methods Appl. Anal.* 7, 495-509 (2000)
- [26] LÜBBE, C., VALIENTE-KROON, J. A.: A conformal approach for the analysis of the nonlinear stability of pure radiation cosmologies. *Ann. Phys.* 328, 1–25, (2013)
- [27] LUK, J., SPECK, J.: Shock formation in solutions to the $2D$ compressible Euler equations in the presence of non-zero vorticity *Available online at https://arxiv.org/abs/1610.00737*
- [28] LUO, T., XIN, Z., ZENG, H.: Well-posedness for the motion of physical vacuum of the threedimensional compressible Euler equations with or without self-gravitation. *Arch. Ration. Mech. Anal.* 213, no. 3, 763-831 (2014)
- [29] LUO, T., ZENG, H.: Global Existence of Smooth Solutions and Convergence to Barenblatt Solutions for the Physical Vacuum Free Boundary Problem of Compressible Euler Equations with Damping, *Comm. Pure Appl. Math.* 69 no.7, 1354 –1396 (2016)
- [30] MAJDA, A.: *Compressible fluid flow and systems of conservation laws in several space variables*, volume 53 of *Applied Mathematical Sciences*. Springer-Verlag, New York (1984)
- [31] MAJDA, A.: Vorticity and the mathematical theory of incompressible fluid flow. *Comm. Pure Appl. Math.*, 39(S, suppl.):S187–S220 (1986)
- [32] MAKINO, T., UKAI, S., KAWASHIMA, S.: Sur la solution à support compact de l'équations d'Euler compressible. *Japan J. Appl. Math.* 3, 249-257 (1986)
- [33] MERLE, F., RAPHAËL, P., SZEFTEL, J. Stable self similar blow up dynamics for L^2 -supercritical NLS equations. *Geom. Funct. Anal.* 20 no 4, 1028–1071 (2010)
- [34] CHRISTODOULOU, D., MIAO, S.: *Compressible Flow and Euler's Equations*, Surveys in Modern Mathematics Vol. 9, International Press (2014)
- [35] OLIYNYK, T.: Future stability of the FLRW fluid solutions in the presence of a positive cosmological constant. *To appear in Commun. Math. Phys.*
- [36] RODNIANSKI, I., SPECK, J.: The Nonlinear Future Stability of the FLRW Family of Solutions to the Irrotational Euler-Einstein System with a Positive Cosmological Constant. *Journal of European Math. Soc.* 15 (6), 2369–2462 (2013)
- [37] ROZANOVA, O.: Solutions with linear profile of velocity to the Euler equations in several dimensions. *Hyperbolic problems: theory, numerics, applications, 861-870, Springer, Berlin,* 2003
- [38] Shkoller, S., Sideris, T. C. Global existence of near-affine solutions to the compressible Euler equations. Preprint, arXiv:1710.08368
- [39] SERRE, D. Solutions classiques globales des equations d'Euler pour un fluide parfait compress- ´ ible. *Annales de l'Institut Fourier* 47, 139–153 (1997)
- [40] SERRE, D.: Expansion of a compressible gas in vacuum. *Bulletin of the Institute of Mathematics, Academia Sinica, Taiwan.* 10, 695–716 (2015)
- [41] SPECK, J.: The nonlinear future stability of the FLRW family of solutions to the Euler-Einstein system with a positive cosmological constant. *Selecta Mathematica*, 18, no. 3, 633–715 (2012)
- [42] SIDERIS, T. C.: Formation of singularities in three-dimensional compressible fluids. *Comm. Math. Phys.* 101(4), 475–485 (1985)
- [43] SIDERIS, T. C.: Spreading of the free boundary of an ideal fluid in a vacuum. *J. Differential Equations*, 257(1), 1–14 (2014)
- [44] SIDERIS, T., C.: Global existence and asymptotic behavior of affine motion of 3D ideal fluids surrounded by vacuum, *Arch. Ration. Mech. Anal.* 225, no. 1, 141–176 (2017)