



King's Research Portal

Document Version

Early version, also known as pre-print

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Pushnitski, A., & Miheisi, N. (in press). Restriction theorems for Hankel operators. *Studia Mathematica*.

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

RESTRICTION THEOREMS FOR HANKEL OPERATORS

NAZAR MIHEISI AND ALEXANDER PUSHNITSKI

ABSTRACT. We consider a class of maps from integral Hankel operators to Hankel matrices, which we call restriction maps. In the simplest case, such a map is simply a restriction of the integral kernel onto integers. More generally, it is given by an averaging of the kernel with a sufficiently regular weight function. We study the boundedness of restriction maps with respect to the operator norm and the Schatten norms.

1. INTRODUCTION

1.1. Hankel operators. Let $\alpha = \{\alpha(j)\}_{j \geq 0}$ be a sequence of complex numbers. The *Hankel matrix* $H(\alpha)$ is the “infinite matrix” $\{\alpha(j+k)\}_{j,k \geq 0}$, considered as a linear operator on $\ell^2(\mathbb{Z}_+)$, $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$, so that

$$(H(\alpha)x)(k) = \sum_{j \geq 0} \alpha(j+k)x(j), \quad k \geq 0, \quad x = \{x(j)\}_{j \geq 0} \in \ell^2(\mathbb{Z}_+).$$

Similarly, for a *kernel function* $\mathbf{a} \in L^1_{\text{loc}}(0, \infty)$, the *integral Hankel operator* on $L^2(0, \infty)$ is defined by the formula

$$(\mathbf{H}(\mathbf{a})\mathbf{f})(t) = \int_0^\infty \mathbf{a}(t+s)\mathbf{f}(s) ds, \quad t > 0, \quad \mathbf{f} \in L^2(0, \infty).$$

In order to distinguish between these two classes of operators, we use boldface font for objects associated with integral Hankel operators.

For general background on Hankel operators, see [7, 9]. In what follows, we will only consider bounded Hankel matrices and bounded integral Hankel operators.

1.2. Restrictions. The purpose of this paper is to examine the linear map, which we call *the restriction map*, between the set of integral Hankel operators and the set of Hankel matrices. To set the scene, let us consider the *pointwise restriction* of integral kernels to integers. For a given kernel function \mathbf{a} , define the sequence

$$\alpha(j) := \mathbf{a}(j+1), \quad j \geq 0. \tag{1.1}$$

Date: May 28, 2019.

2010 Mathematics Subject Classification. 47B35, 47B10.

Key words and phrases. Hankel operator, integral kernel, Schatten class.

Of course, for this operation to make sense, the kernel function \mathbf{a} has to be continuous. Here is our first result; we denote by \mathbf{S}_p , $0 < p < \infty$, the standard Schatten class of compact operators (see Section 2).

Theorem 1.1. *Let $\mathbf{H}(\mathbf{a}) \in \mathbf{S}_p$ for some $0 < p \leq 1$. Then the kernel function $\mathbf{a}(t)$ is continuous in $t > 0$, so the restriction (1.1) is well defined. The operator $H(\alpha)$ is in \mathbf{S}_p and we have the estimate*

$$\|H(\alpha)\|_{\mathbf{S}_p} \leq C_p \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p}. \quad (1.2)$$

The continuity of the kernel function \mathbf{a} for trace class integral Hankel operators is well known (see e.g. [8, Corollary 7.10]); the main point here is the estimate (1.2). In Section 3 we give a slightly more precise version of Theorem 1.1 and show that it does not extend to $p > 1$. Further, we show that if we restrict the map $\mathbf{H}(\mathbf{a}) \mapsto H(\alpha)$ to *non-negative* integral Hankel operators, then it is bounded in \mathbf{S}_p norm for all $0 < p < \infty$ (and also in the operator norm).

Further, along with the pointwise restriction (1.1), we consider the following *restrictions by averaging*. For a suitably regular function φ on \mathbb{R} and for a kernel function \mathbf{a} , we define the restriction $\mathcal{R}_\varphi \mathbf{a}$ to be the sequence

$$\mathcal{R}_\varphi \mathbf{a}(j) = \int_0^\infty \mathbf{a}(t) \varphi(t-j) dt, \quad j \geq 0.$$

In particular, formally taking $\varphi(t) = \delta(t-1)$, where δ is the Dirac δ -function, we recover the pointwise restrictions (1.1). In Section 5 we prove that, under suitable regularity conditions on φ , the map

$$\mathbf{H}(\mathbf{a}) \mapsto H(\mathcal{R}_\varphi \mathbf{a})$$

is bounded in \mathbf{S}_p norm for all $0 < p < \infty$ (and also in the operator norm). We also relate this result to the well known unitary equivalence between Hankel matrices and integral Hankel operators.

The results of this paper seem to parallel some restriction theorems for Fourier multipliers; see e.g. [3, 5, 2]. However, this connection is not completely understood (at least by the authors).

We note in passing that one can consider a converse operation, an *extension* of a Hankel matrix to an integral kernel. For a suitably regular function φ and a sequence $\alpha = \{\alpha(j)\}_{j \geq 0}$, one can define the *extension* $\mathcal{E}_\varphi \alpha$ to be the function

$$\mathcal{E}_\varphi \alpha(t) = \sum_{k \geq 0} \alpha(k) \varphi(t-k), \quad t > 0,$$

and one can consider the map

$$H(\alpha) \mapsto \mathbf{H}(\mathcal{E}_\varphi \alpha).$$

Although some Schatten norm boundedness results for this map are not difficult to prove, we have not succeeded in finding a coherent set of estimates for it and therefore we do not discuss extensions here.

1.3. Helson operators. In order to put this paper into context, here we say a few words about Helson operators, although they are not mentioned in the rest of the paper. *Helson matrices* (also known as multiplicative Hankel matrices) are infinite matrices of the form $M(\alpha) = \{\alpha(jk)\}_{j,k \geq 1}$. The “continuous” analogues of these are integral operators on $L^2(1, \infty)$ defined by

$$\mathbf{M}(\mathbf{a})\mathbf{f}(t) = \int_1^\infty \mathbf{a}(ts)\mathbf{f}(s) ds, \quad t > 1.$$

We refer to these as *integral Helson operators*. It is easily seen that, by an exponential change of variables, each integral Helson operator is unitarily equivalent to an integral Hankel operator; this is not the case for Helson matrices.

The analogue of Theorem 1.1 holds for Helson operators. To be precise, if $\mathbf{M}(\mathbf{a}) \in \mathbf{S}_p$, $0 < p \leq 1$, then $\mathbf{a}(t)$ is continuous in $t > 1$. If we set $\alpha(j) = \mathbf{a}(j+1)$, $j \geq 1$, the following bound holds [6, Theorem 3.2]:

$$\|M(\alpha)\|_{\mathbf{S}_p} \leq C_p \|\mathbf{M}(\mathbf{a})\|_{\mathbf{S}_p}. \tag{1.3}$$

The estimate (1.3) was a key component in determining explicit spectral asymptotics for a family of compact modifications of the multiplicative Hilbert matrix [6] (see [1] for background on the multiplicative Hilbert matrix). The present paper appeared as an attempt to consider this technical ingredient on a more systematic basis.

In proving Theorem 1.1 (and its extensions) we will make use of V. Peller’s characterization of Hankel operators of class \mathbf{S}_p (Proposition 2.1). While this also gives a description of the integral Helson operators of class \mathbf{S}_p , no such characterization is known for Helson matrices and so the proof of [6, Theorem 3.2] is more involved.

1.4. Symbols. For a bounded Hankel matrix $H(\alpha)$, its analytic symbol is the function

$$\check{\alpha}(z) = \sum_{m \geq 0} \alpha_m z^m, \quad |z| < 1.$$

Similarly, for a bounded integral Hankel operator $\mathbf{H}(\mathbf{a})$, its analytic symbol is the function

$$\check{\mathbf{a}}(\xi) = \int_0^\infty \mathbf{a}(t)e^{2\pi i t \xi} dt, \quad \text{Im } \xi > 0.$$

It is instructive to view restriction maps on Hankel operators in terms of the symbols. If $\alpha = \mathcal{R}_\varphi \mathbf{a}$, then for the symbols we have

$$\check{\alpha}(z) = \int_{\mathbb{R}} \frac{\check{\mathbf{a}}(\xi + i0)\check{\varphi}(-\xi + i0)}{1 - ze^{-2\pi i\xi}} d\xi, \quad |z| < 1. \quad (1.4)$$

In particular, for the pointwise restriction (1.1) we have (see Section 2.3)

$$\check{\alpha}(e^{2\pi i\xi}) = e^{-2\pi i\xi} \sum_{j \in \mathbb{Z}} \check{\mathbf{a}}(\xi - j), \quad \text{Im } \xi > 0. \quad (1.5)$$

Since Schatten norms of Hankel operators correspond to Besov norms of the symbols (see Section 2), one can view the topic of this paper as the study of the map induced by (1.4) between Besov classes. We prefer to use an operator theoretic viewpoint whenever possible, although sometimes we have to resort to proofs in terms of Besov classes.

2. PRELIMINARIES

Throughout this paper, the symbol ‘ C ’ with a (possibly empty) set of subscripts will denote a positive constant, depending only on the subscripts, whose precise value may change with each occurrence. Moreover, we write $X \asymp Y$ for two expressions X and Y if $X \leq CY$ and $Y \leq CX$.

2.1. Operator theory, Schatten classes. For a bounded linear operator A in a Hilbert space, we denote by $\|A\|_{\mathcal{B}}$ the operator norm of A .

For a compact operator A in a Hilbert space, let $\{s_n(A)\}_{n=1}^{\infty}$ be the sequence of singular values of A , enumerated with multiplicities taken into account. For $0 < p < \infty$, the standard Schatten class \mathbf{S}_p of compact operators is defined by the condition

$$A \in \mathbf{S}_p \quad \Leftrightarrow \quad \|A\|_{\mathbf{S}_p}^p := \sum_{n \geq 1} s_n(A)^p < \infty.$$

$\|\cdot\|_{\mathbf{S}_p}$ is a norm on \mathbf{S}_p for $p \geq 1$ and a quasinorm for $0 < p < 1$. We denote by \mathbf{S}_{∞} the class of compact operators and we take $\|\cdot\|_{\mathbf{S}_{\infty}}$ to be the operator norm on \mathbf{S}_{∞} .

2.2. Characterisation of Schatten class Hankel operators. Let \mathbb{T} denote the unit circle. We consider the Fourier transform \mathcal{F} as the unitary map from $L^2(\mathbb{T})$ to $\ell^2(\mathbb{Z})$,

$$(\mathcal{F}f)(j) = \hat{f}(j) = \int_0^1 f(e^{2\pi it}) e^{-2\pi ijt} dt, \quad j \in \mathbb{Z}.$$

We also use its inverse $\mathcal{F}^{-1} : \ell^2(\mathbb{Z}) \rightarrow L^2(\mathbb{T})$ and denote $\check{\alpha} = \mathcal{F}^{-1}\alpha$. Similarly, we use the Fourier integral transform \mathcal{F} in $L^2(\mathbb{R})$ and its inverse

$$(\mathcal{F}^{-1}\mathbf{f})(\xi) = \check{\mathbf{f}}(\xi) = \int_{\mathbb{R}} \mathbf{f}(t)e^{2\pi i\xi t} dt, \quad \xi \in \mathbb{R}.$$

Let $\mathbf{w} \in C^\infty(\mathbb{R})$ be a non-negative function such that $\text{supp } \mathbf{w} \subset [1/2, 2]$ and

$$\sum_{m \in \mathbb{Z}} \mathbf{w}(t/2^m) = 1, \quad t > 0.$$

We set $\mathbf{w}_m(t) = \mathbf{w}(t/2^m)$. For $m \geq 0$, we denote by w_m the restriction of the function \mathbf{w}_m onto \mathbb{Z}_+ , i.e. $w_m(j) = \mathbf{w}_m(j)$, $j \geq 0$.

Proposition 2.1. *Let $0 < p < \infty$.*

(i) [9, Theorem 6.1.1, 6.2.1, 6.3.1] *For a bounded Hankel matrix $H(\alpha)$, one has*

$$\|H(\alpha)\|_{\mathfrak{S}_p}^p \asymp |\alpha(0)|^p + \sum_{m \geq 0} 2^m \|\mathcal{F}^{-1}(\alpha w_m)\|_{L^p(\mathbb{T})}^p.$$

(ii) [9, Theorem 6.7.4] *For a bounded integral Hankel operator $\mathbf{H}(\mathbf{a})$ one has*

$$\|\mathbf{H}(\mathbf{a})\|_{\mathfrak{S}_p}^p \asymp \sum_{m \in \mathbb{Z}} 2^m \|\mathcal{F}^{-1}(\mathbf{a} w_m)\|_{L^p(\mathbb{R})}^p.$$

The expressions in the right side here are exactly the (quasi-)norms of the symbols in the Besov class $B_p^{1/p}$.

2.3. Periodization operator. Here we discuss the map induced by (1.5). For a compactly supported function $\mathbf{f} \in C(\mathbb{R})$, we define the *periodization* of \mathbf{f} as the function on the unit circle given by

$$\mathcal{P}\mathbf{f}(e^{2\pi it}) = \sum_{j \in \mathbb{Z}} \mathbf{f}(t - j), \quad e^{2\pi it} \in \mathbb{T}. \quad (2.1)$$

We call \mathcal{P} the *periodization operator*. Applying the ‘‘triangle inequality’’ $|a+b|^p \leq |a|^p + |b|^p$ for $0 < p \leq 1$ to (2.1) and then integrating over t we see that

$$\|\mathcal{P}\mathbf{f}\|_{L^p(\mathbb{T})} \leq \|\mathbf{f}\|_{L^p(\mathbb{R})}, \quad 0 < p \leq 1.$$

This allows one to extend \mathcal{P} to a map from $L^1(\mathbb{R})$ to $L^1(\mathbb{T})$. For $\mathbf{f} \in L^1(\mathbb{R})$ it is straightforward to see that

$$\widehat{\mathcal{P}\mathbf{f}}(j) = \widehat{\mathbf{f}}(j), \quad j \in \mathbb{Z},$$

i.e. the map $\widehat{\mathbf{f}} \mapsto \widehat{\mathcal{P}\mathbf{f}}$ is exactly the restriction onto integers. Thus we have the estimate

$$\left\| \sum_{j \in \mathbb{Z}} \widehat{\mathbf{f}}(j) z^j \right\|_{L^p(\mathbb{T})} \leq \|\mathbf{f}\|_{L^p(\mathbb{R})}, \quad 0 < p \leq 1. \quad (2.2)$$

3. POINTWISE RESTRICTIONS

3.1. Pointwise restrictions for operators of class \mathbf{S}_p . For $\lambda > 0$, let $\delta_\lambda(t) = \delta(t - \lambda)$, where $\delta(t)$ is the Dirac delta function, so that if $\mathbf{a} \in C(0, \infty)$, then

$$(\mathcal{R}_{\delta_\lambda} \mathbf{a})(j) = \mathbf{a}(j + \lambda), \quad j \geq 0.$$

If \mathbf{a} is the kernel function of an integral Hankel operator of class \mathbf{S}_1 , then \mathbf{a} is almost everywhere equal to a continuous function on $(0, \infty)$ [8, Corollary 7.10], and the estimate

$$|\mathbf{a}(t)| \leq C \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_1} / t, \quad t > 0, \quad (3.1)$$

holds true with some absolute constant C . Thus, the definition of $\mathcal{R}_{\delta_\lambda} \mathbf{a}$ makes sense without any further assumptions on \mathbf{a} .

The aim of this section is to prove the following.

Theorem 3.1. *Let $0 < p \leq 1$, $\lambda > 0$. If $\mathbf{H}(\mathbf{a}) \in \mathbf{S}_p$ then $H(\mathcal{R}_{\delta_\lambda} \mathbf{a}) \in \mathbf{S}_p$ and*

$$\|H(\mathcal{R}_{\delta_\lambda} \mathbf{a})\|_{\mathbf{S}_p}^p \leq C_p (|\mathbf{a}(\lambda)|^p + \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p}^p) \leq C_p (1 + 1/\lambda^p) \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p}^p. \quad (3.2)$$

The main component in the proof of Theorem 3.1 is the estimate (2.2).

Proof. Denote $\mathbf{b}(t) = \mathbf{a}(t + \lambda)$. By Proposition 2.1(i), we have

$$\|H(\mathcal{R}_{\delta_\lambda} \mathbf{a})\|_{\mathbf{S}_p}^p \leq C_p |\mathbf{b}(0)|^p + C_p \sum_{m \geq 0} 2^m \left\| \sum_{j \geq 0} \mathbf{b}(j) w_m(j) z^j \right\|_{L^p(\mathbb{T})}^p. \quad (3.3)$$

Let us first estimate the series in the right hand side of (3.3). Applying (2.2) to $\mathbf{f} = \mathcal{F}^{-1}(\mathbf{b} w_m)$, we obtain

$$\left\| \sum_{j \geq 0} \mathbf{b}(j) w_m(j) z^j \right\|_{L^p(\mathbb{T})}^p \leq \|\mathcal{F}^{-1}(\mathbf{b} w_m)\|_{L^p(\mathbb{R})}^p$$

for every $m \geq 0$. By Proposition 2.1(ii), this yields

$$\sum_{m \geq 0} 2^m \left\| \sum_{j \geq 0} \mathbf{b}(j) w_m(j) z^j \right\|_{L^p(\mathbb{T})}^p \leq \sum_{m \in \mathbb{Z}} 2^m \|\mathcal{F}^{-1}(\mathbf{b} w_m)\|_{L^p(\mathbb{R})}^p \leq C_p \|\mathbf{H}(\mathbf{b})\|_{\mathbf{S}_p}^p.$$

Let us relate the norm of $\mathbf{H}(\mathbf{b})$ to the norm of $\mathbf{H}(\mathbf{a})$. Writing

$$\int_0^\infty \int_0^\infty \mathbf{b}(t+s) \mathbf{f}(t) \overline{\mathbf{g}(s)} dt ds = \int_{\lambda/2}^\infty \int_{\lambda/2}^\infty \mathbf{a}(t+s) \mathbf{f}(t - \lambda/2) \overline{\mathbf{g}(s - \lambda/2)} dt ds,$$

we see that $\mathbf{H}(\mathbf{b})$ is unitarily equivalent to the compression of $\mathbf{H}(\mathbf{a})$ onto the subspace $L^2(\lambda/2, \infty) \subset L^2(0, \infty)$. It follows that

$$\|\mathbf{H}(\mathbf{b})\|_{\mathbf{S}_p} \leq \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p} \quad (3.4)$$

for all $p > 0$. Finally, consider the first term in the right hand side of (3.3). By (3.1) we have

$$|\mathbf{b}(0)| = |\mathbf{a}(\lambda)| \leq C\|\mathbf{H}(\mathbf{a})\|_{\mathbf{s}_1}/\lambda \leq C\|\mathbf{H}(\mathbf{a})\|_{\mathbf{s}_p}/\lambda.$$

Combining the above estimates, we arrive at the required statement. \square

Remark. One can also consider restrictions of \mathbf{a} to the scaled lattice $\{\gamma j + \lambda\}_{j \geq 0}$ for some $\gamma > 0$. For $\gamma > 0$, let $\mathbf{a}_\gamma(t) = \mathbf{a}(\gamma t)$ and let $V_\gamma : L^2(0, \infty) \rightarrow L^2(0, \infty)$ be the unitary operator

$$V_\gamma \mathbf{f}(t) = \sqrt{\gamma} \mathbf{f}(\gamma t), \quad t > 0.$$

Then $\gamma \mathbf{H}(\mathbf{a}_\gamma) = V_\gamma \mathbf{H}(\mathbf{a}) V_\gamma^*$ and so $\gamma \|\mathbf{H}(\mathbf{a}_\gamma)\|_{\mathbf{s}_p} = \|\mathbf{H}(\mathbf{a})\|_{\mathbf{s}_p}$ for all $0 < p < \infty$. It follows from this and Theorem 3.1 that if $0 < p \leq 1$, then $\gamma \|H(\mathcal{R}_{\delta_\lambda} \mathbf{a}_\gamma)\|_{\mathbf{s}_p} \leq C_p \|\mathbf{H}(\mathbf{a})\|_{\mathbf{s}_p}$, and thus

$$\sup_{\gamma > 0} \gamma \|H(\mathcal{R}_{\delta_\lambda} \mathbf{a}_\gamma)\|_{\mathbf{s}_p} \leq C_p \|\mathbf{H}(\mathbf{a})\|_{\mathbf{s}_p}. \quad (3.5)$$

3.2. Counterexample for $p > 1$. For $p > 1$, is it no longer the case that the kernel of an integral Hankel operator of class \mathbf{S}_p is necessarily continuous. However, even if we only consider operators with continuous kernels, the conclusions of Theorem 3.1 still fail and thus the condition $0 < p \leq 1$ is sharp.

To show this, we fix a smooth kernel function \mathbf{a} with $\text{supp } \mathbf{a} \subset [1/2, 2]$ and $\mathbf{a}(1) = 1$ and let $\mathbf{a}^{(N)}(t) = \mathbf{a}(1 + N(t - 1))$ for $N \in \mathbb{N}$. Then for each N we have

$$\begin{aligned} \mathcal{R}_{\delta_1} \mathbf{a}^{(N)}(0) &= \mathbf{a}^{(N)}(1) = 1, \\ \mathcal{R}_{\delta_1} \mathbf{a}^{(N)}(j) &= \mathbf{a}^{(N)}(1 + j) = 0, \quad j \geq 1. \end{aligned}$$

It follows that

$$\|H(\mathcal{R}_{\delta_1} \mathbf{a}^{(N)})\|_{\mathbf{s}_p} = 1$$

for all $p \geq 1$ and $N \in \mathbb{N}$. On the other hand, it is not difficult to show that

$$\|\mathbf{H}(\mathbf{a}^{(N)})\|_{\mathbf{s}_p}^p \leq CN^{1-p}$$

which tends to zero as $N \rightarrow \infty$ whenever $p > 1$. Indeed, by the assumption on the support of \mathbf{a} we have

$$\mathbf{a}^{(N)} = \mathbf{a}^{(N)} \mathbf{w}_{-1} + \mathbf{a}^{(N)} \mathbf{w}_0 + \mathbf{a}^{(N)} \mathbf{w}_1$$

for all N , where \mathbf{w}_m are defined in Section 2.2. It is easy to conclude that

$$\sum_{m=-1,0,1} 2^m \|\mathcal{F}^{-1}(\mathbf{a}^{(N)} \mathbf{w}_m)\|_{L^p(\mathbb{R})}^p \leq C_p \|\mathcal{F}^{-1}(\mathbf{a}^{(N)})\|_{L^p(\mathbb{R})}^p = CN^{1-p}.$$

3.3. Partial converse of Theorem 3.1. It is clear that one cannot bound $\mathbf{H}(\mathbf{a})$ by $H(\mathcal{R}_{\delta_\lambda} \mathbf{a})$ in any norm. However, one can achieve a partial converse if we vary our restriction operators in an appropriate sense and take a supremum over all restrictions in the right side. Here we briefly sketch a sample argument of this nature. Fix $0 < p \leq 1$; we use “continuous” counterparts of the expressions in Proposition 2.1; see e.g. [16, Section 2.3.3, p.99]:

$$\begin{aligned} \|H(\alpha)\|_{\mathbf{S}_p}^p &\asymp |\alpha(0)|^p + \int_0^2 \|\mathcal{F}^{-1}(\alpha w^\tau)\|_{L^p(\mathbb{T})}^p \frac{d\tau}{\tau^2}, \\ \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p}^p &\asymp \int_0^\infty \|\mathcal{F}^{-1}(\mathbf{a} w^\tau)\|_{L^p(\mathbb{R})}^p \frac{d\tau}{\tau^2}, \end{aligned} \quad (3.6)$$

where $\mathbf{w}^\tau(t) = \mathbf{w}(\tau t)$ and $w^\tau = \{\mathbf{w}(\tau j)\}_{j \geq 0}$.

Let \mathbf{a} be a continuous function on $(0, \infty)$ and, for $\gamma > 0$, let $\mathbf{a}_\gamma(t) = \mathbf{a}(\gamma t)$. Observe that, by a change of variable,

$$\begin{aligned} \|\mathcal{F}^{-1}(w^\tau \mathcal{R}_{\delta_\lambda} \mathbf{a}_\gamma)\|_{L^p(\mathbb{T})}^p &= \int_{-1/2}^{1/2} \left| \sum_{j \geq 0} \mathbf{w}(\tau j) \mathbf{a}(\gamma(j + \lambda)) e^{2\pi i j s} \right|^p ds \\ &= \gamma^{1-p} \int_{-1/2\gamma}^{1/2\gamma} \left| \gamma \sum_{j \geq 0} \mathbf{w}(\tau j) \mathbf{a}(\gamma(j + \lambda)) e^{2\pi i j \gamma s} \right|^p ds. \end{aligned}$$

By another change of variable, it then follows from (3.6) that

$$\begin{aligned} \gamma^p \|H(\mathcal{R}_{\delta_\lambda} \mathbf{a}_\gamma)\|_{\mathbf{S}_p}^p &\geq C_p \gamma \int_0^2 \int_{-1/2\gamma}^{1/2\gamma} \left| \gamma \sum_{j \geq 0} \mathbf{w}(\tau j) \mathbf{a}(\gamma(j + \lambda)) e^{2\pi i j \gamma s} \right|^p ds \frac{d\tau}{\tau^2} \\ &= C_p \int_0^{2/\gamma} \int_{-1/2\gamma}^{1/2\gamma} \left| \gamma \sum_{j \geq 0} \mathbf{w}(\tau \gamma j) \mathbf{a}(\gamma(j + \lambda)) e^{2\pi i j \gamma s} \right|^p ds \frac{d\tau}{\tau^2}. \end{aligned} \quad (3.7)$$

Since \mathbf{a} is continuous, for each $s \in \mathbb{R}$ and $\tau > 0$ the integrand in (3.7) converges to $|\mathcal{F}^{-1}(\mathbf{a} w^\tau)(s)|^p$ as $\gamma \rightarrow 0$. Then by Fatou’s Lemma we see that

$$\begin{aligned} \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p}^p &\asymp \int_0^\infty \|\mathcal{F}^{-1}(\mathbf{a} w^\tau)\|_{L^p(\mathbb{R})}^p \frac{d\tau}{\tau^2} \\ &\leq \lim_{\gamma \rightarrow 0} \int_0^{2/\gamma} \int_{-1/2\gamma}^{1/2\gamma} \left| \gamma \sum_{j \geq 0} \mathbf{w}(\tau \gamma j) \mathbf{a}(\gamma(j + \lambda)) e^{2\pi i j \gamma s} \right|^p ds \frac{d\tau}{\tau^2} \\ &\leq C_p \lim_{\gamma \rightarrow 0} \gamma^p \|H(\mathcal{R}_{\delta_\lambda} \mathbf{a}_\gamma)\|_{\mathbf{S}_p}^p. \end{aligned} \quad (3.8)$$

This gives an analogue of Igari’s theorem for Fourier multipliers [5]. Combining (3.8) with (3.5) gives the estimate

$$\|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p} \asymp \sup_{\gamma > 0} \gamma \|H(\mathcal{R}_{\delta_\lambda} \mathbf{a}_\gamma)\|_{\mathbf{S}_p}, \quad 0 < p \leq 1.$$

4. POINTWISE RESTRICTION FOR NON-NEGATIVE OPERATORS

4.1. **Statement of the result.** Although Theorem 3.1 fails for $p > 1$, the estimate (3.2) remains valid for all $0 < p < \infty$ if we restrict to the class of non-negative operators (in the usual quadratic form sense). Before stating this precisely we recall (see e.g. [17, page 22]) that a bounded integral Hankel operator $\mathbf{H}(\mathbf{a})$ is non-negative if and only if the kernel function \mathbf{a} can be represented as

$$\mathbf{a}(t) = \int_0^\infty e^{-t\eta} d\mu(\eta), \quad (4.1)$$

where the measure μ satisfies

$$\mu((0, \eta)) \leq C\eta, \quad \eta > 0.$$

In particular, it follows that the kernel function $\mathbf{a}(t)$ is continuous in $t > 0$, and therefore the restriction $\mathcal{R}_{\delta_\lambda} \mathbf{a}$ is well defined for all $\lambda > 0$.

We have the following theorem.

Theorem 4.1. *Let $\mathbf{H}(\mathbf{a}) \geq 0$ be a bounded integral Hankel operator and $\lambda > 0$. Then the following hold:*

(i) $H(\mathcal{R}_{\delta_\lambda} \mathbf{a})$ is bounded and

$$\|H(\mathcal{R}_{\delta_\lambda} \mathbf{a})\|_{\mathcal{B}} \leq C(1 + 1/\lambda) \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}. \quad (4.2)$$

(ii) If $\mathbf{H}(\mathbf{a}) \in \mathbf{S}_p$ for some $0 < p \leq \infty$, then $H(\mathcal{R}_{\delta_\lambda} \mathbf{a}) \in \mathbf{S}_p$ and

$$\|H(\mathcal{R}_{\delta_\lambda} \mathbf{a})\|_{\mathbf{S}_p} \leq C_p(1 + 1/\lambda) \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p}. \quad (4.3)$$

Remark 4.2. (1) We note for clarity that part (ii) of the theorem includes the statement that if $\mathbf{H}(\mathbf{a}) \geq 0$ is compact, then $H(\mathcal{R}_{\delta_\lambda} \mathbf{a})$ is compact.

(2) Observe that by (4.1), the kernel function \mathbf{a} is necessarily positive, monotone decreasing and continuous on $(0, \infty)$. In fact, the proof of Theorem 4.1 depends only on these properties of \mathbf{a} .

(3) If $\lambda \geq 2$, one can slightly improve the statement of Theorem 4.1. In this case one gets

$$\begin{aligned} \|H(\mathcal{R}_{\delta_\lambda} \mathbf{a})\|_{\mathcal{B}} &\leq \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}, \quad \lambda \geq 2, \\ \|H(\mathcal{R}_{\delta_\lambda} \mathbf{a})\|_{\mathbf{S}_p} &\leq \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p}, \quad \lambda \geq 2, \quad p \in 2\mathbb{N}, \end{aligned}$$

i.e. the constants in the estimates are equal to one in these cases.

In the rest of this section we prove Theorem 4.1. Observe that we only need to consider the case $p > 1$, as for $0 < p \leq 1$ the required result follows from Theorem 3.1.

Our proof consists of two different parts. The first one is a short operator theoretic argument based on pointwise domination which however works only for

$p \in 2\mathbb{N}$ or $p = \infty$. The second one is a direct calculation based on Proposition 2.1 which applies to all $p \geq 1$.

4.2. **Proof for $p \in 2\mathbb{N} \cup \{\infty\}$.** First we need a version of (3.1) for non-negative operators.

Lemma 4.3. *Let $\mathbf{H}(\mathbf{a}) \geq 0$ be a bounded integral Hankel operator and $\lambda > 0$. Then*

$$\lambda \mathbf{a}(\lambda) \leq 2\|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}.$$

Proof. Take $\mathbf{f}(t) = e^{-t/\lambda}$; then $\|\mathbf{f}\|_{L^2(0,\infty)}^2 = \lambda/2$ and using the monotonicity of $\mathbf{a}(t)$,

$$\begin{aligned} (\mathbf{H}(\mathbf{a})\mathbf{f}, \mathbf{f}) &= \int_0^\infty \int_0^\infty \mathbf{a}(t+s)e^{-(t+s)/\lambda} dt ds = \int_0^\infty \mathbf{a}(t)e^{-t/\lambda} t dt \\ &\geq \int_0^\lambda \mathbf{a}(t)e^{-t/\lambda} t dt \geq \mathbf{a}(\lambda) \int_0^\lambda e^{-t/\lambda} t dt = (1 - 2e^{-1})\lambda^2 \mathbf{a}(\lambda). \end{aligned}$$

On the other hand,

$$(\mathbf{H}(\mathbf{a})\mathbf{f}, \mathbf{f}) \leq \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}} \|\mathbf{f}\|_{L^2(0,\infty)}^2 = (\lambda/2)\|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}.$$

Combining these two estimates, we obtain

$$\lambda \mathbf{a}(\lambda) \leq \frac{1}{2(1 - 2e^{-1})} \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}} \leq 2\|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}},$$

as required. □

Proof of Theorem 4.1 for $p \in 2\mathbb{N} \cup \{\infty\}$. First let us **assume that** $\lambda \geq 2$. Let K be the integral operator in $L^2(0, \infty)$ with the integral kernel

$$K(t, s) = \mathbf{a}(\lambda + [t] + [s]),$$

where $[t]$ is the largest integer less than or equal to t . Since

$$\lambda + [t] + [s] \geq \lambda + (t - 1) + (s - 1) \geq t + s,$$

by monotonicity of \mathbf{a} we have

$$K(t, s) \leq \mathbf{a}(t + s).$$

In the terminology of [13, Chapter 2], this means that K is *pointwise dominated* by $\mathbf{H}(\mathbf{a})$. By [13, Theorem 2.13], it follows that

$$\|K\| \leq \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}} \quad \text{and} \quad \|K\|_{\mathbf{s}_p} \leq \|\mathbf{H}(\mathbf{a})\|_{\mathbf{s}_p}$$

for all $p \in 2\mathbb{N}$. (This implication does not extend to $p \notin 2\mathbb{N}$; see e.g. [10, 14].) It is also true (see [11, 4]) that the compactness of $\mathbf{H}(\mathbf{a})$ implies the compactness of K .

Next, let us relate K to $H(\mathcal{R}_{\delta_\lambda} \mathbf{a})$. For $\mathbf{f} \in L^2(0, \infty)$ let us write the quadratic form of K as

$$(K\mathbf{f}, \mathbf{f}) = \sum_{j,k \geq 0} \mathbf{a}(\lambda + j + k) f_j \overline{f_k}, \quad f_j = \int_j^{j+1} \mathbf{f}(t) dt.$$

This means that, writing $L^2(0, \infty) = \ell^2(\mathbb{Z}_+) \otimes L^2(0, 1)$, the operator K can be represented as

$$K = H(\mathcal{R}_{\delta_\lambda} \mathbf{a}) \otimes (\cdot, \mathbb{1}) \mathbb{1},$$

where $(\cdot, \mathbb{1}) \mathbb{1}$ is the rank one operator in $L^2(0, 1)$ acting as

$$\mathbf{f} \mapsto \int_0^1 \mathbf{f}(t) dt.$$

It follows that

$$\|K\|_{\mathcal{B}} = \|H(\mathcal{R}_{\delta_\lambda} \mathbf{a})\|_{\mathcal{B}} \quad \text{and} \quad \|K\|_{\mathbf{s}_p} = \|H(\mathcal{R}_{\delta_\lambda} \mathbf{a})\|_{\mathbf{s}_p}$$

for all $p > 0$. This completes the proof for $\lambda \geq 2$ and $p \in 2\mathbb{N} \cup \{\infty\}$.

Let us consider the case $0 < \lambda < 2$. Let P_2 be the projection onto $\ell^2(\{2, 3, \dots\})$ in $\ell^2(\mathbb{Z}_+)$. Write

$$H(\mathcal{R}_{\delta_\lambda} \mathbf{a}) = P_2 H(\mathcal{R}_{\delta_\lambda} \mathbf{a}) P_2 + \tilde{H}.$$

Inspecting the matrix elements of \tilde{H} and using Lemma 4.3, we see that

$$\|\tilde{H}\|_{\mathbf{s}_2}^2 \leq 4 \sum_{j \geq 0} |\mathbf{a}(\lambda + j)|^2 \leq 16 \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}^2 \sum_{j \geq 0} \frac{1}{|\lambda + j|^2} \leq C(1 + 1/\lambda)^2 \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}^2.$$

On the other hand, the operator $P_2 H(\mathcal{R}_{\delta_\lambda} \mathbf{a}) P_2$ is unitarily equivalent to $H(\mathcal{R}_{\delta_{\lambda+2}} \mathbf{a})$. Thus, applying the previous step of the proof, we obtain

$$\|P_2 H(\mathcal{R}_{\delta_\lambda} \mathbf{a}) P_2\|_{\mathcal{B}} \leq \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}} \quad \text{and} \quad \|P_2 H(\mathcal{R}_{\delta_\lambda} \mathbf{a}) P_2\|_{\mathbf{s}_p} \leq \|\mathbf{H}(\mathbf{a})\|_{\mathbf{s}_p}$$

for $p \in 2\mathbb{N}$. Combining these estimates, we arrive at (4.2) and (4.3) for $p \in 2\mathbb{N}$ and $p = \infty$. \square

As already mentioned, this proof does not extend to $p \notin 2\mathbb{N}$; see e.g. [10, 14]. Below we give a different proof which works for all $1 \leq p < \infty$, but does not give precise information about the constants in the estimates.

4.3. Proof of Theorem 4.1 for $1 \leq p < \infty$. In order to simplify our notation, we set $\mathbf{b}(t) = \mathbf{a}(t + \lambda)$, $b(k) = \mathbf{a}(k + \lambda)$, and

$$\begin{aligned} \check{b}_m(z) &= \sum_{k \geq 0} b(k) w_m(k) z^k, \quad m \in \mathbb{Z}_+, \quad z \in \mathbb{T}, \\ \check{\mathbf{b}}_m(\xi) &= \int_0^\infty \mathbf{b}(t) \mathbf{w}_m(t) e^{2\pi i \xi t} dt, \quad m \in \mathbb{Z}, \quad \xi \in \mathbb{R}. \end{aligned}$$

The core of the proof is the bound

$$\sum_{m \geq 1} 2^m \|\check{b}_m\|_{L^p(\mathbb{T})}^p \leq C_p \sum_{m \in \mathbb{Z}} 2^m \|\check{\mathbf{b}}_m\|_{L^p(\mathbb{R})}^p, \quad (4.4)$$

which we prove below. Throughout the proof, we use the property that \mathbf{b} and b are positive and monotone decreasing.

First step: upper bound for $\|\check{b}_m\|_{L^p(\mathbb{T})}$. Fix $m \geq 1$. First we prepare two pointwise bounds for $\check{b}_m(z)$. The first one is trivial:

$$|\check{b}_m(z)| \leq \sum_{k \geq 0} b(k) w_m(k) \leq 2^{m+1} b(2^{m-1}). \quad (4.5)$$

The second one is obtained through a discrete version of integration by parts (Abel summation). We have

$$\begin{aligned} \check{b}_m(z) &= \frac{1}{z-1} \sum_{k \geq 0} b(k) w_m(k) (z^{k+1} - z^k) \\ &= \frac{1}{z-1} \sum_{k \geq 0} (b(k) w_m(k) - b(k+1) w_m(k+1)) z^{k+1} \\ &= \frac{1}{z-1} \sum_{k \geq 0} ((b(k) - b(k+1)) w_m(k) + b(k+1) (w_m(k) - w_m(k+1))) z^{k+1}, \end{aligned}$$

and therefore

$$|\check{b}_m(z)| \leq \frac{1}{|z-1|} \sum_{k=2^{m-1}}^{2^{m+1}} (b(k) - b(k+1)) + \frac{1}{|z-1|} \sum_{k=2^{m-1}}^{1+2^{m+1}} b(k) |w_m(k-1) - w_m(k)|.$$

Clearly, the first sum here is telescoping. For the second sum, we use the estimate

$$|w_m(k-1) - w_m(k)| \leq C 2^{-m}.$$

Putting this together, we obtain

$$\begin{aligned} |\check{b}_m(z)| &\leq \frac{1}{|z-1|} (b(2^{m-1}) - b(1 + 2^{m+1})) + \frac{C}{|z-1|} 2^{-m} \sum_{k=2^{m-1}}^{1+2^{m+1}} b(k) \\ &\leq \frac{C}{|z-1|} b(2^{m-1}), \end{aligned} \quad (4.6)$$

which is our second bound for $\check{b}_m(z)$.

Now we can estimate the norm $\|\check{b}_m\|_{L^p(\mathbb{T})}$. We split the integral over the unit circle into two parts and estimate them separately. Using (4.5), we obtain

$$2^m \int_{|t| < 2^{-m}} |\check{b}_m(e^{2\pi i t})|^p dt \leq C 2^{pm} b(2^{m-1})^p.$$

Using (4.6), we get

$$\begin{aligned} 2^m \int_{|t|>2^{-m}} |\check{b}_m(e^{2\pi it})|^p dt &\leq C2^m \int_{|t|>2^{-m}} \frac{dt}{|e^{2\pi it} - 1|^p} b(2^{m-1})^p \\ &\leq C2^m \int_{2^{-m}}^1 \frac{dt}{t^p} b(2^{m-1})^p \leq C2^{pm} b(2^{m-1})^p. \end{aligned}$$

Combining the estimates for two integrals above, we obtain

$$2^m \|\check{b}_m\|_{L^p(\mathbb{T})}^p \leq C2^{pm} b(2^{m-1})^p.$$

Second step: lower bound for $\|\check{\mathbf{b}}_m\|_{L^p(\mathbb{R})}$. For the derivative of \mathbf{b}_m we have

$$\check{\mathbf{b}}'_m(\xi) = 2\pi i \int_0^\infty \mathbf{b}(t) \mathbf{w}_m(t) t e^{2\pi i t \xi} dt,$$

and therefore

$$|\check{\mathbf{b}}'_m(\xi)| \leq 2\pi \int_0^\infty \mathbf{b}(t) \mathbf{w}_m(t) t dt \leq 2^{m+2} \pi \int_0^\infty \mathbf{b}(t) \mathbf{w}_m(t) dt = 2^{m+2} \pi \check{\mathbf{b}}_m(0).$$

It follows that

$$|\check{\mathbf{b}}_m(\xi) - \check{\mathbf{b}}_m(0)| \leq |\xi| 2^{m+2} \pi \check{\mathbf{b}}_m(0),$$

and therefore for $|\xi| < 2^{-m-5}$ we have

$$|\check{\mathbf{b}}_m(\xi)| \geq \check{\mathbf{b}}_m(0)/2.$$

We use this to obtain a lower bound for the integral of $|\check{\mathbf{b}}_m|^p$:

$$2^m \int_{\mathbb{R}} |\check{\mathbf{b}}_m(\xi)|^p d\xi \geq 2^m \int_{|\xi| < 2^{-m-5}} |\check{\mathbf{b}}_m(\xi)|^p d\xi \geq 2^{-5} (\check{\mathbf{b}}_m(0)/2)^p = C\check{\mathbf{b}}_m(0)^p.$$

Finally,

$$\check{\mathbf{b}}_m(0) = \int_0^\infty \mathbf{b}(t) \mathbf{w}_m(t) dt \geq \mathbf{b}(2^{m+1}) \int_0^\infty \mathbf{w}_m(t) dt = C2^m \mathbf{b}(2^{m+1}),$$

and so we obtain

$$2^m \|\check{\mathbf{b}}_m\|_{L^p(\mathbb{R})}^p \geq C2^{mp} b(2^{m+1})^p.$$

Combining the two steps and completing the proof. Combining the upper bound for $\|\check{b}_m\|_{L^p(\mathbb{T})}$ and the lower bound for $\|\check{\mathbf{b}}_m\|_{L^p(\mathbb{R})}$, we obtain

$$2^m \|\check{b}_m\|_{L^p(\mathbb{T})}^p \leq C2^{p(m-1)} b(2^{m-1})^p \leq C2^{m-2} \|\check{\mathbf{b}}_{m-2}\|_{L^p(\mathbb{R})}^p, \quad m \geq 1.$$

Summing over m , we obtain the bound (4.4).

By Proposition 2.1(i), we have

$$\|H(b)\|_{\mathbf{S}_p}^p \leq C_p |b(0)|^p + C_p \sum_{m \geq 0} 2^m \|\check{b}_m\|_{L^p(\mathbb{T})}^p. \quad (4.7)$$

By Lemma 4.3, we have

$$|b(0)|^p = |\mathbf{a}(\lambda)|^p \leq 2^p \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}^p / \lambda^p.$$

Similarly, the $m = 0$ term in the series in (4.7) can be estimated as follows:

$$\|\check{b}_0\|_{L^p(\mathbb{T})}^p = |b(1)|^p = |\mathbf{a}(\lambda + 1)|^p \leq 2^p \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}^p / (1 + \lambda)^p \leq 2^p \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}^p / \lambda^p.$$

Combining this with (4.4) and using Proposition 2.1(ii), we obtain

$$\|H(b)\|_{\mathbf{S}_p}^p \leq C_p \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}^p / \lambda^p + C_p \sum_{m \in \mathbb{Z}} 2^m \|\check{\mathbf{b}}_m\|_{L^p(\mathbb{R})}^p \leq C_p \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}}^p / \lambda^p + C_p \|\mathbf{H}(\mathbf{b})\|_{\mathbf{S}_p}^p.$$

Finally, as in (3.4), we have $\|\mathbf{H}(\mathbf{b})\|_{\mathbf{S}_p} \leq \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p}$, and we arrive at the required estimate (4.3).

5. RESTRICTION BY AVERAGING

5.1. Boundedness of restrictions by averaging. The main result of this section says that if the function φ is sufficiently regular, then the map $\mathbf{H}(\mathbf{a}) \mapsto H(\mathcal{R}_\varphi \mathbf{a})$ is bounded with respect to all Schatten norms. We will make use of the periodisation operator \mathcal{P} from Section 2.3.

Theorem 5.1. *Let $\varphi \in C(\mathbb{R})$ be such that $\text{supp } \varphi \subset [0, \infty)$ and $\mathcal{P}(|\check{\varphi}|) \in L^\infty(\mathbb{T})$. Then there exist bounded operators Φ_1 and Φ_2 acting from $\ell^2(\mathbb{Z}_+)$ to $L^2(0, \infty)$ such that*

$$\Phi_2^* \mathbf{H}(\mathbf{a}) \Phi_1 = H(\mathcal{R}_\varphi \mathbf{a}) \tag{5.1}$$

and $\|\Phi_1\|_{\mathcal{B}} = \|\Phi_2\|_{\mathcal{B}} = \sqrt{A}$, where $A = \|\mathcal{P}(|\check{\varphi}|)\|_{L^\infty(\mathbb{T})}$. Consequently, we have

$$\|H(\mathcal{R}_\varphi \mathbf{a})\|_{\mathcal{B}} \leq A \|\mathbf{H}(\mathbf{a})\|_{\mathcal{B}} \quad \text{and} \quad \|H(\mathcal{R}_\varphi \mathbf{a})\|_{\mathbf{S}_p} \leq A \|\mathbf{H}(\mathbf{a})\|_{\mathbf{S}_p}$$

for every $0 < p < \infty$.

A close inspection of the proof of Theorem 5.1 will reveal that the condition $\mathcal{P}(|\check{\varphi}|) \in L^\infty(\mathbb{T})$ is necessary, in the sense that if there exist bounded operators $\Phi_1, \Phi_2 : \ell^2(\mathbb{Z}_+) \rightarrow L^2(0, \infty)$ such that (5.1) holds, then $\mathcal{P}(|\check{\varphi}|) \in L^\infty(\mathbb{T})$.

It will be convenient to separate the statement related to the boundedness of the maps Φ_1 and Φ_2 .

Lemma 5.2. *Let $\psi \in L^2(\mathbb{R})$ with $\mathcal{P}(|\check{\psi}|^2) \in L^\infty(\mathbb{T})$. Then the map*

$$\Phi : x = \{x(j)\}_{j \geq 0} \mapsto \sum_{j \geq 0} x(j) \psi(t - j), \quad t \in \mathbb{R},$$

is bounded from $\ell^2(\mathbb{Z}_+)$ to $L^2(\mathbb{R})$, with $\|\Phi\|_{\mathcal{B}}^2 = \|\mathcal{P}(|\check{\psi}|^2)\|_{L^\infty(\mathbb{T})}$.

Proof. Let x be a finitely supported sequence. We have, using Parseval's theorem,

$$\begin{aligned}
 \|\Phi x\|_{L^2(\mathbb{R})}^2 &= \left\| \sum_{j \geq 0} x(j) \psi(\cdot - j) \right\|_{L^2(\mathbb{R})}^2 = \int_{\mathbb{R}} \left| \sum_{j \geq 0} x(j) \check{\psi}(\xi) e^{2\pi i j \xi} \right|^2 d\xi \\
 &= \int_{\mathbb{R}} |\check{x}(e^{2\pi i \xi})|^2 |\check{\psi}(\xi)|^2 d\xi = \sum_{j \in \mathbb{Z}} \int_0^1 |\check{x}(e^{2\pi i \xi})|^2 |\check{\psi}(\xi - j)|^2 d\xi \\
 &= \int_0^1 |\check{x}(e^{2\pi i \xi})|^2 \mathcal{P}(|\check{\psi}|^2)(\xi) d\xi \leq \|\mathcal{P}(|\check{\psi}|^2)\|_{L^\infty(\mathbb{T})} \|\check{x}\|_{L^2(\mathbb{T})}^2 \\
 &= \|\mathcal{P}(|\check{\psi}|^2)\|_{L^\infty(\mathbb{T})} \|x\|_{\ell^2(\mathbb{Z}_+)}^2.
 \end{aligned}$$

It is also clear that the inequality here is sharp in the sense that

$$\sup_{\|\check{x}\|_{L^2(\mathbb{T})}=1} \int_{\mathbb{R}} |\check{x}(e^{2\pi i \xi})|^2 |\check{\psi}(\xi)|^2 d\xi = \|\mathcal{P}(|\check{\psi}|^2)\|_{L^\infty(\mathbb{T})}.$$

This proves the claim. \square

Below \mathbb{C}_+ will denote the upper half-plane; $H^\infty(\mathbb{C}_+)$, $H^2(\mathbb{C}_+)$ etc. are the standard Hardy classes. We will sometimes identify functions in these Hardy classes with their boundary values on \mathbb{R} .

Proof of Theorem 5.1. By assumption, we have $\mathcal{P}(|\check{\varphi}|) \in L^\infty(\mathbb{T}) \subset L^1(\mathbb{T})$; it follows that $\check{\varphi} \in L^1(\mathbb{R})$. Recalling that $\text{supp } \varphi \subset [0, \infty)$, we obtain that $\check{\varphi} \in H^1(\mathbb{C}_+)$. Thus, we can factorise $\check{\varphi}$ into a product of two $H^2(\mathbb{C}_+)$ -functions. More precisely, there exist $\varphi_1, \varphi_2 \in L^2(0, \infty)$ such that

$$\check{\varphi}(\xi) = \check{\varphi}_1(\xi) \overline{\check{\varphi}_2(-\xi)} \quad \text{and} \quad |\check{\varphi}_1(\xi)| = |\check{\varphi}_2(-\xi)|, \quad \forall \xi \in \mathbb{R}.$$

Then $\varphi = \varphi_1 * \overline{\varphi_2}$ and

$$\|\mathcal{P}(|\check{\varphi}_1|^2)\|_{L^\infty(\mathbb{T})} = \|\mathcal{P}(|\check{\varphi}_2|^2)\|_{L^\infty(\mathbb{T})} = \|\mathcal{P}(|\check{\varphi}|)\|_{L^\infty(\mathbb{T})}.$$

Next, for $i = 1, 2$, let us define the map $\Phi_i : \ell^2(\mathbb{Z}_+) \rightarrow L^2(0, \infty)$ by

$$\Phi_i : x = \{x(j)\}_{j \geq 0} \mapsto \sum_{j \geq 0} x(j) \varphi_i(\cdot - j).$$

By Lemma 5.2, both Φ_1 and Φ_2 are bounded with norms equal to $\sqrt{\|\mathcal{P}(|\check{\varphi}|)\|_{L^\infty(\mathbb{T})}}$.

In order to prove (5.1), let us first rearrange the definition of $\mathcal{R}_\varphi \mathbf{a}$. For each $j, k \geq 0$

$$\begin{aligned}
\mathcal{R}_\varphi \mathbf{a}(j+k) &= \int_0^\infty \mathbf{a}(t) \varphi(t-j-k) dt \\
&= \int_0^\infty \mathbf{a}(t) \int_{\mathbb{R}} \varphi_1(t-s-j-k) \overline{\varphi_2(s)} ds dt \\
&= \int \int_{s+t>0} \mathbf{a}(s+t) \varphi_1(t-j) \overline{\varphi_2(s-k)} dt ds.
\end{aligned}$$

Since both φ_1 and φ_2 are supported on $(0, \infty)$, we can rewrite this as

$$\mathcal{R}_\varphi \mathbf{a}(j+k) = \int_0^\infty \int_0^\infty \mathbf{a}(s+t) \varphi_1(t-j) \overline{\varphi_2(s-k)} ds dt.$$

Now for $x = \{x(j)\}_{j \geq 0} \in \ell^2(\mathbb{Z}_+)$, let us compute the quadratic form

$$\begin{aligned}
(\mathbf{H}(\mathbf{a})\Phi_1 x, \Phi_2 x) &= \sum_{j,k \geq 0} \int_0^\infty \int_0^\infty \mathbf{a}(t+s) x(j) \overline{x(k)} \varphi_1(t-j) \overline{\varphi_2(s-k)} dt ds \\
&= \sum_{j,k \geq 0} \mathcal{R}_\varphi \mathbf{a}(j+k) x(j) \overline{x(k)},
\end{aligned}$$

which yields (5.1). □

5.2. Unitary equivalence and restrictions associated to general convolutions. Let $L_n = L_n^{(0)}$ be the n 'th Laguerre polynomial (see [15, Ch. V] for the definition) and let

$$\mathbf{u}_n(t) = -2i\sqrt{\pi} L_n(4\pi t) e^{-2\pi t}, \quad t > 0. \quad (5.2)$$

Then $\{\mathbf{u}_n\}_{n \geq 0}$ is an orthonormal basis of $L^2(0, \infty)$. It is well known that the matrix of an integral Hankel operator is a Hankel matrix in the basis $\{\mathbf{u}_n\}_{n \geq 0}$ and hence the classes of Hankel matrices and integral Hankel operators are unitarily equivalent [9, Ch. 1, Thm 8.9].

In this subsection we discuss how this unitary equivalence fits into our ‘‘restriction by averaging’’ framework. This requires looking at restrictions by averaging of a more general type than considered above. To a given integral Hankel operator $\mathbf{H}(\mathbf{a})$ we associate the Hankel matrix $H(\alpha)$ with

$$\alpha_j = \int_0^\infty \mathbf{a}(t) \varphi_j(t) dt, \quad j \geq 0,$$

where φ_j is a certain sequence of smooth functions, a more general one than just translations of a single function. Our sequence φ_j will be given by the multiple convolution of the form

$$\varphi_j = \varphi * \underbrace{\nu * \nu * \cdots * \nu}_{j \text{ terms}}, \quad j \geq 0,$$

where φ is a sufficiently regular function supported on $[0, \infty)$, and ν is a positive finite measure supported on $[0, \infty)$. Observe that if $d\nu(t) = \delta(t-1)dt$, then $\varphi_j(t) = \varphi(t-j)$, so we recover the definition of \mathcal{R}_φ .

To make the multiple convolution notation more readable, we introduce the (formal) convolution with ν operator

$$T_\nu \mathbf{f} = \mathbf{f} * \nu;$$

then $\varphi_j = T_\nu^j \varphi$.

Theorem 5.3. *Let ν be a complex valued measure on $[0, \infty)$ with $\|\check{\nu}\|_{H^\infty(\mathbb{C}_+)} \leq 1$, and let $\varphi \in C(\mathbb{R})$ satisfy $\text{supp } \varphi \subset [0, \infty)$ and*

$$|\check{\varphi}(\xi)| \leq \frac{C}{1 + \xi^2}, \quad \xi \in \mathbb{R}. \quad (5.3)$$

For $j \geq 0$, set $\varphi_j = T_\nu^j \varphi$ and consider the map

$$\mathbf{a}(t) \mapsto \alpha = \{\alpha(j)\}_{j=0}^\infty, \quad \alpha(j) = \int_0^\infty \mathbf{a}(t) \varphi_j(t) dt.$$

Then there exist bounded operators Φ_1 and Φ_2 acting from $\ell^2(\mathbb{Z}_+)$ to $L^2(0, \infty)$ such that

$$\Phi_2^* \mathbf{H}(\mathbf{a}) \Phi_1 = H(\alpha). \quad (5.4)$$

Consequently,

$$\|H(\alpha)\| \leq A \|\mathbf{H}(\mathbf{a})\| \quad \text{and} \quad \|H(\alpha)\|_{\mathbf{s}_p} \leq A \|\mathbf{H}(\mathbf{a})\|_{\mathbf{s}_p}$$

for all $0 < p < \infty$, where $A = \|\Phi_1\| \|\Phi_2\|$.

It will again be convenient to separate the boundedness of Φ_1 and Φ_2 into a lemma.

Lemma 5.4. *Let $\omega \in H^\infty(\mathbb{C}_+)$ with $\|\omega\|_{H^\infty} \leq 1$. Then the map*

$$x = \{x(j)\}_{j=0}^\infty \mapsto \sum_{j \geq 0} \frac{x(j) \omega(\xi)^j}{\xi + i}, \quad \xi \in \mathbb{C}_+, \quad (5.5)$$

is bounded from $\ell^2(\mathbb{Z}_+)$ to $H^2(\mathbb{C}_+)$.

Proof. Consider the conformal map

$$\mathbb{D} \ni \zeta \mapsto \xi = i \frac{1 + \zeta}{1 - \zeta} \in \mathbb{C}_+$$

and the corresponding unitary operator $U : H^2(\mathbb{C}_+) \rightarrow H^2(\mathbb{D})$,

$$(U\mathbf{f})(\zeta) = \frac{2\sqrt{\pi}}{1 - \zeta} \mathbf{f} \left(i \frac{1 + \zeta}{1 - \zeta} \right).$$

We have

$$U : \frac{\omega(\xi)^j}{\xi + i} \mapsto -i\sqrt{\pi}\psi(\zeta)^j, \quad \psi(\zeta) = \omega\left(i\frac{1+\zeta}{1-\zeta}\right).$$

It follows that U maps the right hand side of (5.5) to the function

$$-i\sqrt{\pi} \sum_{j \geq 0} x(j)\psi(\zeta)^j.$$

Since $|\psi(\zeta)| \leq 1$, by the Littlewood subordination theorem [12, Chap. 1.3], we have

$$\left\| \sum_{j \geq 0} x(j)\psi(\zeta)^j \right\|_{H^2(\mathbb{D})} \leq C \left\| \sum_{j \geq 0} x(j)\zeta^j \right\|_{H^2(\mathbb{D})} = C\|x\|_{\ell^2}.$$

Putting this together, we obtain the required statement. \square

Proof of Theorem 5.3. Let us write

$$\check{\varphi}(\xi) = \check{\varphi}_1(\xi)\overline{\check{\varphi}_2(-\xi)}, \quad \check{\varphi}_1(\xi) = \check{\varphi}(\xi)(\xi + i), \quad \check{\varphi}_2(\xi) = -\frac{1}{\xi + i},$$

so that $\varphi = \varphi_1 * \overline{\varphi_2}$. By (5.3) combined with the condition on the support of φ , we have $\check{\varphi}_1, \check{\varphi}_2 \in H^2(\mathbb{C}_+)$ and so $\varphi_1, \varphi_2 \in L^2(0, \infty)$.

For $i = 1, 2$, let $\Phi_i : \ell^2(\mathbb{Z}_+) \rightarrow L^2(0, \infty)$ be the map

$$\Phi_i : x = \{x(j)\}_{j \geq 0} \mapsto \sum_{j \geq 0} x(j)T_\nu^j \varphi_i.$$

Observe that

$$\overline{T_\nu^j \varphi_i(\xi)} = \check{\nu}(\xi)^j \check{\varphi}_i(\xi).$$

Let us first show that Φ_2 is bounded. By applying the inverse Fourier transform, it suffices to check that the map

$$x = \{x(j)\}_{j \geq 0} \mapsto \left(\sum_{j \geq 0} x(j)\check{\nu}(\xi)^j \right) \check{\varphi}_2(\xi)$$

is bounded from ℓ^2 to $H^2(\mathbb{C}_+)$. Recalling the definition of $\check{\varphi}_2$, we see that this immediately follows from Lemma 5.4.

To prove that Φ_1 is bounded, we write

$$\check{\varphi}_1(\xi) = \frac{\mathbf{h}(\xi)}{\xi + i}, \quad \mathbf{h}(\xi) = \check{\varphi}(\xi)(\xi + i)^2.$$

By (5.3), we have $\mathbf{h} \in H^\infty(\mathbb{C}_+)$, and so the boundedness of Φ_1 again follows by an application of Lemma 5.4.

It remains to check formula (5.4). This is the same argument as the one in the proof of Theorem 5.1. Indeed, we have

$$T_\nu^{j+k} \varphi = T_\nu^{j+k}(\varphi_1 * \overline{\varphi_2}) = (T_\nu^j \varphi_1) * \overline{(T_\nu^k \varphi_2)},$$

and therefore

$$\alpha(j+k) = \int_0^\infty \mathbf{a}(t) \int_{\mathbb{R}} (T_\nu^j \varphi_1)(t-s) \overline{(T_\nu^k \varphi_2)}(s) ds dt.$$

Since $\text{supp } T_\nu^j \varphi_i \subset [0, \infty)$, by a change of variable this can be rewritten as

$$\alpha(j+k) = \int_0^\infty \int_0^\infty \mathbf{a}(t+s) (T_\nu^j \varphi_1)(t) \overline{(T_\nu^k \varphi_2)}(s) ds dt.$$

Now we see that

$$\begin{aligned} (\mathbf{H}(\mathbf{a})\Phi_1 x, \Phi_2 x) &= \sum_{j,k \geq 0} x(j) \overline{x(k)} \int_0^\infty \int_0^\infty \mathbf{a}(t+s) (T_\nu^j \varphi_1)(t) \overline{(T_\nu^k \varphi_2)}(s) ds dt \\ &= (H(\alpha)x, x). \end{aligned}$$

□

Example 5.5. For $t \geq 0$, let

$$\varphi(t) = -4\pi t e^{-2\pi t} \quad \text{and} \quad \nu(t) = \delta(t) - 4\pi e^{-2\pi t}.$$

Then

$$\check{\varphi}(\xi) = \frac{1}{\pi(\xi+i)^2} \quad \text{and} \quad \check{\nu}(\xi) = \frac{\xi-i}{\xi+i} \in L^\infty(\mathbb{R}).$$

Hence the conclusions of Theorem 5.3 hold. However, we can say more in this case. We also have that $\varphi = \psi * \psi$, with $\psi(t) = -2i\sqrt{\pi}e^{-2\pi t}$, $t \geq 0$, and so we can take

$$\Phi_1 x = \Phi_2 x = \sum_{j \geq 0} x(j) T_\nu^j \psi$$

in the proof of Theorem 5.3. It can be shown that $T_\nu^j \psi = \mathbf{u}_j$, $j \geq 0$, where $\{\mathbf{u}_j\}_{j \geq 0}$ is the orthonormal basis given by (5.2). Thus Φ_1 (and hence Φ_2) is unitary. Consequently, this choice of φ and ν produces the well-known unitary equivalence between Hankel matrices and integral Hankel operators.

REFERENCES

- [1] O. F. BREVIG, K.-M. PERFECT, K. SEIP, A. G. SISKAKIS, D. VUKOTIĆ, *The multiplicative Hilbert matrix*, Adv. Math. **302** (2016), 410–432.
- [2] R. COIFMAN, G. WEISS, *Transference methods in analysis*, American Mathematical Society, 1976.
- [3] K. DE LEEUW, *On L^p multipliers*, Ann. of Math. **81** (1965), 364–379.
- [4] P. G. DODDS, D. H. FREMLIN, *Compact operators in Banach lattices*, Israel J. Math. **34** (1979), no. 4, 287–320.
- [5] S. IGARI, *Functions of L^p -multipliers*, Tôhoku Math. J. **21** (1969), 304–320.
- [6] N. MIHEISI, A. PUSHNITSKI, *A Helson matrix with explicit eigenvalue asymptotics*, J. Funct. Anal. **275** (2018), no.4, 967–987.
- [7] N. NIKOLSKI, *Operators, Functions and Systems: An Easy Reading. Volume 1*, American Mathematical Society, 2002.

- [8] J. R. PARTINGTON, *An Introduction to Hankel Operators*, Cambridge University Press, 1988.
- [9] V. V. PELLER, *Hankel operators and their applications*, Springer, 2003.
- [10] V. V. PELLER, *Hankel operators of class \mathfrak{S}_p and their applications (rational approximation, Gaussian processes, the problem of majorization of operators)*, Math. USSR Sbornik **41** (1982), 443–479.
- [11] L. D. PITT, *A compactness condition for linear operators of function spaces*, J. Operator Theory **1** (1979), 49–54.
- [12] J. H. SHAPIRO, *Composition operators and classical function theory*, Springer, 1993.
- [13] B. SIMON, *Trace Ideals and Their Applications*, Second Edition, American Mathematical Society, 2005.
- [14] B. SIMON, *Pointwise domination of matrices and comparison of \mathcal{I}_p norms*, Pacific J. Math. **97** (1981), no.2, 471–475.
- [15] G. SZEGŐ, *Orthogonal Polynomials*, American Mathematical Society, 1939.
- [16] H. TRIEBEL, *Theory of function spaces II*, Springer, 1992.
- [17] H. WIDOM, *Hankel matrices*, Trans. Amer. Math. Soc. **121** (1966), 1–35.

DEPARTMENT OF MATHEMATICS, KING'S COLLEGE LONDON, STRAND, LONDON WC2R 2LS, UNITED KINGDOM

E-mail address: nazar.miheisi@kcl.ac.uk

E-mail address: alexander.pushnitski@kcl.ac.uk