



## King's Research Portal

DOI:

[10.1126/scitranslmed.aam7828](https://doi.org/10.1126/scitranslmed.aam7828)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Galleu, A., Riffo-Vasquez, Y., Trento, C., Lomas, C., Dolcetti, L., Cheung, T. S., von Bonin, M., Barbieri, L., Halai, K., Ward, S., Weng, L., Chakraverty, R., Lombardi, G., Watt, F. M., Orchard, K., Marks, D. I., Apperley, J., Bornhauser, M., Walczak, H., ... Dazzi, F. (2017). Apoptosis in mesenchymal stromal cells induces in vivo recipient-mediated immunomodulation. *Science Translational Medicine*, 9(416), Article eaam7828. <https://doi.org/10.1126/scitranslmed.aam7828>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Title: Apoptosis in mesenchymal stromal cells is required to initiate *in vivo* recipient-mediated immunomodulation**

**Single sentence summary:** Apoptosis in mesenchymal stromal cells is required for immunosuppression, is induced by recipient cytotoxic cells and can predict clinical responses in graft-versus-host disease

**Authors:** Antonio Galleu<sup>1</sup>, Yanira Riffo-Vasquez<sup>2</sup>, Cristina Trento<sup>1</sup>, Cara Lomas<sup>3,4</sup>, Luigi Dolcetti<sup>1</sup>, Tik Shing Cheung<sup>1</sup>, Malte von Bonin<sup>5</sup>, Laura Barbieri<sup>1</sup>, Krishma Halai<sup>1</sup>, Sophie Ward<sup>3,4</sup>, Ling Weng<sup>1</sup>, Ronjon Chakraverty<sup>3,4</sup>, Giovanna Lombardi<sup>6</sup>, Fiona M. Watt<sup>7</sup>, Kim Orchard<sup>8</sup>, David I. Marks<sup>9</sup>, Jane Apperley<sup>10</sup>, Martin Bornhauser<sup>1,5</sup>, Henning Walczak<sup>\*4</sup>, Clare Bennett<sup>\*3,4</sup>, and Francesco Dazzi<sup>1,10</sup>

\* Contributed equally

**Affiliations:** <sup>1</sup>Regenerative Medicine, Division of Cancer Studies, King's College London, London, UK. <sup>2</sup>Institute of Pharmaceutical Science, King's College London, London, UK. <sup>3</sup>UCL Institute of Immunity and Transplantation, London, UK. <sup>4</sup>UCL Cancer Institute, London, UK. <sup>5</sup>University Hospital Carl Gustav Carus, Dresden, Germany. <sup>6</sup>MRC Centre for Transplantation King's College London, London, UK. <sup>7</sup>Centre for Regenerative Medicine, King's College London, London, UK. <sup>8</sup>Southampton University Hospitals Trust, Southampton, UK. <sup>9</sup>Haematology and Oncology Centre, Bristol, UK. <sup>10</sup>Centre for Haematology, Imperial College London, London, UK.

## Abstract

The immunosuppressive activity of mesenchymal stromal cells (MSC) is well documented. However, the therapeutic benefit is completely unpredictable, thus raising concerns about MSC efficacy. One of the affecting factors is the unresolved conundrum that, despite being immunosuppressive, MSC are undetectable following injection. Therefore, understanding the fate of infused MSC could help to predict clinical responses. Using a murine model of graft-versus-host disease (GvHD) we demonstrate that MSC are actively induced to undergo perforin-dependent apoptosis by recipient cytotoxic cells and that this process is essential to initiate MSC-induced immunosuppression. When examining patients with GvHD who received MSC we found a striking parallel, whereby only those with high cytotoxic activity against MSC responded to MSC infusion whereas those with low activity did not. Importantly, the need for recipient cytotoxic cell activity could be replaced by the infusion of apoptotic MSC generated *ex vivo*. Recipient phagocytes engulf apoptotic MSC and produce indoleamine 2,3-dioxygenase (IDO) that is ultimately necessary for effecting immunosuppression. Therefore, we propose the innovative concept that patients should be stratified for MSC treatment according to their ability to kill MSC or that all patients could be treated with *ex vivo* apoptotic MSC.

## Introduction

Mesenchymal stromal cells (MSC) exhibit potent immunosuppressive and anti-inflammatory activities (1) that have been extensively tested in several clinical conditions (2–5). However, the results have often been controversial and proof of efficacy inconclusive. Two major unresolved challenges undermine progress in the field. The first is that, only a proportion of patients, although affected by the same disease, responds to MSC infusions and this response cannot be predicted. The second is that, to be efficacious, MSC are not required to engraft. The vast majority of infused MSC resides transiently in the lungs before becoming undetectable within a few hours (6). Since our current knowledge cannot provide an explanation to this paradox (7–10) a better understanding of the mechanisms underlying MSC therapeutic activity would be highly desirable.

We selected to address these challenges in Graft-versus-Host Disease (GvHD) because there is proof of principle that MSC are efficacious (4, 11). Thereby, we demonstrate that the activated cytotoxic cells harbored in GvHD mice rapidly induce extensive *in vivo* caspase activation in infused MSC. The presence of activated cytotoxic cells in MSC recipients is required for inducing MSC apoptosis and, as a consequence, for triggering the MSC immunosuppressive effect. These findings do not only explain the rapid clearance of infused MSC but also mechanistically reconcile their disappearance with their immunomodulatory activity. The link between cytotoxic cell activity, MSC apoptosis and immunosuppression is confirmed by the correlation in GvHD patients between high levels of cytotoxic activity against MSC and clinical responses. Recipient-mediated killing of MSC can therefore be regarded as the first

mechanism-based biomarker for patient stratification. Strikingly, apoptotic MSC generated *ex vivo* exhibit a potent immunosuppressive effect that is dependent on phagocyte-derived indoleamine 2,3-dioxygenase (IDO) activity and bypasses the need for cytotoxic cells in the recipient. Eventually, we propose that novel treatment concepts may be devised based on the application of apoptotic MSC.

## Results

### *MSC undergo apoptosis in recipient GvHD animals.*

In order to explain the mechanism by which MSC are rapidly cleared after injection (6, 12), we tested the hypothesis that MSC undergo apoptosis. We injected human MSC in a mouse model of GvHD in which lethally irradiated C57BL/6 male mice were transplanted with bone marrow (BM) and polyclonal purified CD4<sup>+</sup> cells from female syngeneic donors, and purified CD8<sup>+</sup> cells transgenic for a T-cell receptor specific for the male HY antigen Uty (Matahari, Mh) as GvHD effectors (13) (Fig. S1A). In this model the expansion of the T cells effecting GvHD (CD8<sup>+</sup>Vβ8.3<sup>+</sup>) can be precisely enumerated.

*In vivo* MSC caspase activation was evaluated using MSC that were transfected with the pGL3 control vector for the expression of firefly luciferase (*luc+*) (luc-MS). Caspase activation was measured as luciferase activity using DEVD-aminoluciferin. In this system, caspase 3 activation could be quantified on the basis of emitted light since DEVD is cleaved upon activation of caspase 3, leading to release of aminoluciferin which in turn can be metabolized by the firefly luciferase expressed in MSC. Luc-MS were injected into recipients of BM transplant with CD8<sup>+</sup> Mh T cells

(GvHD group) and one hour later caspase activity was measured as total luminescence signal (TLS). Control mice consisted of naïve males (naïve group) and a group of mice which were irradiated and received CD4<sup>+</sup> and BM cells (BM group) without the transgenic T cells to reproduce the condition of MSC infusion in the absence of activated cytotoxic T cells (Fig. S1A). We observed high caspase activity only in MSC injected into GvHD mice (Fig. 1, A and B). High signal could be detected from the lungs of all animals when the control D-luciferin (firefly luciferase substrate) was used (Fig. S1, B and C), thus confirming that luc-MSCs can be tracked in the lungs also when caspase activity could not be detected.

The evidence that MSCs undergo apoptosis after infusion prompted the question of whether they are still capable of suppressing antigen-driven T cell expansion. Therefore, we analyzed their immunosuppressive effect by enumerating CD8<sup>+</sup>Vβ8.3<sup>+</sup> Mh T cells (GvHD effector cells) in MSC-treated or -untreated GvHD mice. MSCs produced a significant reduction in GvHD effector cell infiltration in both spleen and lungs (Fig. 1, C and D). These results indicate that, despite the presence of MSC apoptosis after infusion (Fig. 1, A and B), MSC immunosuppression still occurs.

We can exclude the possibility that the observed immunosuppressive activity could be the consequence of the recipient inflammatory cytokines because in our xenogeneic combination murine inflammatory cytokines will not cross-react with the corresponding human receptors and will not activate immunosuppressive molecules in human MSCs (14–16), whilst retaining the ability to expand murine effector cells mediating GvHD (17). Accordingly, human MSCs were not able to inhibit concanavalin-A (ConA) induced proliferation of murine splenocytes (mSpl) unless pre-activated by human cytokines

(Fig. S2, A, B and C). Furthermore, exposure of human MSC to murine inflammatory cytokines did not upregulate *IDO*, TNF-stimulated gene 6 protein (*TSG-6*) or prostaglandin-endoperoxide synthase-2 (*PTSG2*), considered major effectors of human MSC-mediated *in vitro* immunosuppression (8) (Fig. S2 D).

#### *In vivo MSC apoptosis depends on activated recipient GvHD effector cells*

Our results show that MSC rapidly undergo apoptosis after infusion, providing the long-sought after explanation for the rapid clearance of transplanted MSC in the recipient. The absence of *in vivo* MSC apoptosis in naïve and BM mice clearly demonstrates that MSC apoptosis is not the result of xenogeneic recognition of human MSC, because it is detected only in GvHD mice. When we enumerated GvHD effector cell infiltrate (CD8<sup>+</sup>Vβ8.3<sup>+</sup>) in the lungs of mice, where MSC apoptosis occurs, we found that only the lungs of GvHD but not naïve and BM mice contained a large proportion of CD8<sup>+</sup>Vβ8.3<sup>+</sup> cells (Fig. 2A), thus confirming the correlation between caspase activation in MSC and the presence of GvHD effector cells.

To test the hypothesis that GvHD effector cells were responsible for MSC apoptosis, MSC were cultivated with CD8<sup>+</sup> T cells purified from the lungs or spleens of GvHD (*in vivo* activated) or naïve Mh (*in vivo* resting) mice. Activated, but not resting, Mh CD8<sup>+</sup> cells induced MSC apoptosis (Fig. 2B). In further support of the lack of antigen-specificity in the induction of the killing activity, high levels of cytotoxic activity could be elicited by naïve Mh CD8<sup>+</sup> cells stimulated *in vitro* by CD3/CD28 beads (Fig. S3A). The requirement of cytotoxic cells in the induction of MSC apoptosis and the consequent immunosuppression was evaluated using Mh/Perforin Knock-Out mice (Mh/Perf<sup>-/-</sup>) as donors of cytotoxic defective GvHD effector cells (GvHDPerf<sup>-/-</sup> group).

Luc-MSCs were infused into GvHDPerf<sup>-/-</sup> or control GvHD mice which had received Mh CD8<sup>+</sup> T cells. Mice were imaged 1 hour later and caspase activation measured as described before. We observed much lower caspase activity in GvHDPerf<sup>-/-</sup> mice compared to GvHD controls (Fig. 2, C and D). High signal was detected in the lungs of all animals when the control D-luciferin was used (Fig. S3, B and C), thus confirming that luc-MSCs were in the lungs also when caspase activity could not be detected. Importantly, the infiltration of GvHD effector cells in the spleen and lungs of mice receiving MSC was not reduced in GvHDPerf<sup>-/-</sup> receiving MSC (Fig. 2, E and F). We conclude that MSC apoptosis is indispensable for immunosuppression and requires functionally activated cytotoxic cells in the recipient.

*Cytotoxic activity against MSC is a biomarker predictive of clinical response to MSC in GvHD patients*

Based on these findings, we inferred that the presence of cytotoxic cells in the recipient could be predictive of MSC therapeutic activity. Sixteen patients, mean age 40.5 years (10-69), with severe steroid resistant grade 3-4 GvHD received a total of 17 doses. Patient characteristics are summarized in Table S1.

Clinical responses to MSC were defined by an improvement of at least 50% in at least 1 organ affected by GvHD as previously described (4, 11, 18). Five patients obtained a clinical response. Peripheral blood mononuclear cells (PBMC) were freshly collected within the 24 hours preceding the MSC infusion and tested directly for their ability to induce MSC apoptosis *ex vivo* in a 4-hour cytotoxic assay. One patient received two doses of MSC and the cytotoxic assay was performed before each dose independently. MSC were sourced from the same donor used for the infusion (N=8)



or from a different donor ( $N=9$ ). At the time of performing the assay and cytofluorimetric analysis the operator was blind to patients' clinical details. PBMC from healthy donors ( $N=5$ ) were used as controls.

Overall, PBMC from GvHD patients exhibited an average cytotoxic activity against MSC higher than that detected in control PBMC but without reaching a significant difference (mean $\pm$ SD were: 10.63 $\pm$ 8.76% and 3.82 $\pm$ 2.50%, respectively;  $p=.10$ ). However, the level of cytotoxicity between clinical responders and non-responders to MSC was markedly different, with the proportion of apoptotic MSC (annexin-V<sup>+</sup>/7AAD<sup>-</sup>) exhibiting a four-fold difference (Fig. 3, A and B). The discrimination threshold of apoptotic MSC between responders and non-responders calculated using the receiver-operating characteristic curve revealed that a 14.85% cut-off was predictive of clinical response with the highest sensitivity and specificity. The level of cytotoxicity did not vary amongst MSC preparations, because when we tested patients' PBMC against the MSC used for the infusion as compared to another preparation obtained from an unrelated donor, no difference in apoptosis induction could be detected (Fig. S4, A). To further confirm the irrelevance of the specific MSC preparation, we evaluated the susceptibility of MSC sourced from different unrelated donors to be killed by 4 different mixed lymphocyte reaction (MLR) combinations. The proportion of apoptotic MSC was similar amongst the different MSC preparations when the same MLR was tested. Conversely, the cytotoxic activity against the same MSC varied amongst different MLR (Fig. S4, B).

Finally, we ruled out the possibility that different proportions of CD8<sup>+</sup> and CD56<sup>+</sup> cells could account for the differing cytotoxic activity because the average frequency in the PBMC of responders and non-responders was similar (Fig. S4, C and D). Therefore,

we conclude that the presence of activated cytotoxic cells in the recipient is predictive of MSC therapeutic activity.

*MSC apoptosis induced by cytotoxic cells is the result of a bystander effect*

To define the mechanisms that drive apoptosis in MSC, we used *in-vitro*-activated PBMC as effector cells. We found that activated but not resting PBMC induced extensive early apoptosis (annexinV<sup>+</sup>/7AAD<sup>-</sup>) in MSC (Fig. 4A), which peaked at 4 hours and shifted towards late apoptosis (annexinV<sup>+</sup>/7AAD<sup>+</sup>) by 24 hours (Fig. S5A). In accord with our *in vivo* observations (Fig. 1, A and B), only activated PBMC induced caspase activation in MSC with a peak at 90 minutes (Movie S1, S2 and S3), and this was completely abrogated by the pan-caspase inhibitor Z-VAD-FMK (Fig. 4B, Fig. S5B and movie S4).

In order to identify the cells inducing apoptosis in MSC, we performed selective enrichment and depletion experiments amongst activated PBMC. We found that CD56<sup>+</sup> natural killer (NK) and CD8<sup>+</sup> populations were the only cells responsible for initiating MSC apoptosis (Fig. 4, C and D). To characterize the mechanisms mediating MSC apoptosis induced by activated cytotoxic cells, we studied potential factors involved in caspase 3 activation. Inhibition of either Granzyme B (GrB) or perforin completely abolished the ability of activated PBMC to kill MSC (Fig. 4E) and activate caspase 3 (Fig. S5C, movie S5 and S6). We also observed reduced PBMC mediated cytotoxicity when CD95 ligand (CD95L, also known as FasL or APO-1L) was neutralized (Fig. 4F), but not when TNF- $\alpha$  or TNF-related apoptosis-inducing ligand (TRAIL) were inhibited, even in the presence of very high concentrations of their respective inhibitors (Fig. S5D).

We then interrogated the nature of the MSC-cytotoxic cell interaction. We observed that apoptosis was not affected by the presence of anti-HLA class I- or anti-HLA class II neutralizing antibodies. Consistently, the cytotoxic activity of activated PBMC on autologous or allogeneic MSC did not differ (Fig. 4G). However, although PBMC required physical contact with MSC to induce apoptosis (Fig. 4H), blocking immunological synapse formation by inhibiting the polarization of microtubule organizing center (19) (Fig. 4I), had no effect. These results demonstrate that MSC killing by activated cytotoxic cells is a bystander effect that does not involve the immunological synapse.

*MSC apoptosis does not interfere with the recognition of the specific target of cytotoxic cells*

Having determined that the MSC apoptosis induced by cytotoxic cells is MHC-independent and not antigen-specific, we asked whether MSC could exert their immunosuppressive effects by competing with and antagonizing antigen-specific recognition. NY-ESO1-specific CD8<sup>+</sup> T cell clone (4D8) or IL2-activated polyclonal CD56<sup>+</sup> purified NK cells were used as effector cells against NY-ESO-1 peptide pulsed T2 or K562 cells, respectively. Two different sets of experiments were performed. In the first set, 4D8 or NK cells were tested against fixed numbers of putative (*susceptible*) target cells in the presence of escalating numbers of MSC used as a *cold* target. The alternative condition consisted of escalating the numbers of the putative target cells – now used as *cold* targets – in the presence of a fixed number of MSC then considered as the *susceptible* target. MSC did not compete with antigen-specific T cell cytotoxicity, since the killing of peptide-pulsed T2 cells was not affected by the presence of MSC (Fig. 5A). The same results were obtained using NK cells

(Fig. 5B). In contrast, the presence of the putative target cells markedly reduced MSC killing in a dose dependent manner in both systems (Fig. 5, C and D). Our data show that MSC killing does not interfere with the primary recognition of the cognate antigen.

*Apoptotic MSC are immunosuppressive in a Th2-type inflammation model*

Our data imply that, since MSC killing does not interfere with the primary recognition of the cognate antigen, induction of apoptosis must be prominently involved in the immunosuppressive activity. Accordingly, in the GvHD model described, MSC apoptosis produced by recipient cytotoxic cells is required for immunosuppression. Therefore, we asked whether this causative relationship remains valid in a different disease model associated with non-cytotoxic Th2-type inflammation. We selected the model of ovalbumin (OVA)-induced allergic airway inflammation (20) summarized in Fig. S6A. Although cytotoxic immune cells have been implicated in the induction of this condition (21, 22), CD8<sup>+</sup> and NK1.1<sup>+</sup> cells infiltrating broncho-alveolar lavage (BAL) and lung tissues were less than 2% one hour after the last challenge, when MSC were infused (Fig. S6, B, C, D and E). To confirm the absence of MSC killing, mice received luc-MSK to assess caspase activation after infusion and imaged one hour later. No caspase activation was detected in any of the mice (Fig. 6, A and B). High signal could be detected in all animals receiving control D-luciferin (Fig. S6, F and G).

The therapeutic activity, assessed by quantitating the eosinophil infiltration in the BAL showed no difference between MSC-treated and untreated mice (Fig. 6C). Together, these results indicate that also in this model MSC immunosuppression relies on the presence of recipient cytotoxic cells that mediate MSC apoptosis.

Therefore, we decided to test whether *in vitro* generated apoptotic MSC (apoMSC) could bypass the need of cytotoxic cells and ameliorate eosinophil infiltration. When apoMSC were administered to recipient mice we observed that the eosinophil infiltrate in BAL was much reduced (Fig. 6D).

*Apoptotic MSC infused in GvHD are immunosuppressive and induce IDO production in recipient phagocytes*

We subsequently investigated whether apoMSC could be immunosuppressive also in the GvHD model. ApoMSC were administered either intravenously (i.v.) or intraperitoneally (i.p.) and the infiltration of CD8<sup>+</sup>Vβ8.3<sup>+</sup> Mh T cells was assessed and compared to untreated GvHD mice. ApoMSC produced a significant reduction in GvHD effector cell infiltration in both spleen and lungs, but this could only be observed in those mice treated with ApoMSC infused i.p. (Fig. 7, A, B, C and D).

It has been reported that the injection of irradiated thymocytes into animals results in their phagocytosis by recipient macrophages and induction of IDO (23). We therefore tested whether apoMSC followed the same destiny by eliciting *in vivo* efferocytosis by recipient phagocytes and inducing IDO production. For this purpose, labelled apoMSC were traced in recipient phagocytes after injection. Following i.p. administration, apoMSC were largely identified in CD11b<sup>+</sup> (Fig. 7, E) and CD11c<sup>+</sup> (Fig. 7, F) phagocytes in the peritoneal draining lymph nodes (24) (Fig. 7, E) but absent when searched for in the lungs and spleen. When the i.v. route was used, amongst the several phagocytic populations investigated (25), CD11b<sup>high</sup>CD11c<sup>int</sup>, CD11b<sup>high</sup>CD11c<sup>-</sup> and CD11b<sup>-</sup>CD11c<sup>+</sup> were detected as engulfing apoMSC in lungs

(Fig. 7, G, H and I, respectively). The analysis of IDO expression in the phagocytes engulfing ApoMSC both in the i.v. and i.p. groups revealed that only the phagocytes in the i.p. group were able to increase IDO expression at a significantly higher level in comparison with their counterparts in untreated GvHD mice (Fig. 7, J and K). These findings strongly suggest that the immunosuppressive effect of apoMSC involve recipient phagocytes and IDO as a crucial effector mechanisms.

*Recipient derived IDO-producing phagocytes are indispensable for MSC immunosuppression in GvHD*

To directly test the importance of recipient-derived phagocytes and recipient-produced IDO in MSC immunosuppressive activity, we depleted phagocytes and inhibited IDO activity in GvHD mice before MSC treatment and evaluated the effect of live MSC on the expansion of GvHD effectors.

To deplete phagocytes liposome clodronate was given to mice 72 hours before MSC injection. The treatment, dramatically impaired the ability of MSC to suppress Mh T cell infiltration (Fig. 8, A and B).

Finally, animals were given the IDO inhibitor 1-methyl-D-tryptophan (1-DMT) (26) before MSC injection. Also in this case, the beneficial effect of MSC on Mh T cell infiltration was much reduced in mice receiving 1-DMT compared to controls (Fig. 8, C and D). We therefore conclude that the immunosuppressive effect of MSC requires the presence of recipient phagocytic cells or IDO production.

## Discussion

This study finally sheds light on the controversial topic of MSC therapeutics by identifying a crucial mechanism that explains several unresolved issues in the field. The first striking piece of information provides the resolution to the until now paradox that MSC are therapeutically efficacious despite the lack of engraftment (27–29). We demonstrate that MSC undergo extensive caspase activation and apoptosis after infusion in the presence of cytotoxic cells, and that this is a requirement for their immunosuppressive function. Although other recipient dependent reactions have been described as mediating MSC lysis *in vitro* (30) and MSC clearance *in vivo* (28, 29, 31, 32) only our study has unveiled the instrumental role of *in vivo* MSC apoptosis in delivering immunosuppression after infusion. Furthermore, although several studies have reported the ability of apoptotic cells to modulate immune responses, here we provide evidence that *in vivo* naturally occurring cell death drives immunosuppression.

MSC apoptosis requires and is effected by cytotoxic granules contained in recipient cytotoxic cells that also mediate GvHD in recipient mice (Fig. 1, A and B, fig. 2, C and D). Importantly, the cytotoxic activity against MSC can also be detected in the PBMC of GvHD patients and it is predictive of clinical responses. Patients displaying high cytotoxicity respond to MSC, whilst those with low or absent cytotoxic activity do not improve following MSC infusion (Fig. 3, A and B). Therefore, the ability of the recipient to generate apoptotic MSC appears to be a requirement for the therapeutic efficacy and lends itself to be used as a potential biomarker to stratify patients for MSC infusions. Further investigations are warranted to characterize in patients the phenotype of the cytotoxic mediating MSC apoptosis.

MSC recognition by cytotoxic cells is not antigen-specific as neither requires HLA engagement nor results from an alloreaction rejection, thus supporting the current practice of using third-party MSC. MSC must be in physical contact with the activated cytotoxic cells to undergo apoptosis, although immunological synapse is not required. This supports a bystander role for the cytotoxic granules released by the activated cytotoxic cell. Such a mechanism has been described in the context of T-cell target recognition (19), HIV infection (33) and atherosclerosis (34). In these studies, bystander cells are not of mesenchymal origin, thus raising the interesting question of whether nonspecific induction of apoptosis and subsequent immunosuppression is selective for MSC.

Our data suggest an approach to MSC therapeutics that highlights the key role of MSC recipient to orchestrate and determine MSC effector functions. Not only are cytotoxic cells in the recipient required to initiate apoptosis in infused MSC, but also phagocytes which, by engulfing apoptotic MSC and producing IDO, ultimately deliver MSC immunosuppressive activity. Similar mechanisms have been described to explain how apoptotic cells of different lineages, generated *in vitro*, induce immune modulation in GvHD (35–37) and macrophage mediated IDO-dependent immunosuppression in other systemic autoimmune diseases (26). This is also consistent with the described ability of MSC to stimulate recipient immune tolerance networks, like regulatory T cells (38, 39) and macrophages (40, 41).

The depletion of recipient macrophages or the inhibition of IDO activity impairs also the therapeutic activity of live MSC, thereby linking *in vivo* MSC apoptosis with immunosuppression. It is unlikely that any particular phagocyte population



(macrophages or DC) is selectively involved in engulfing apoMSC because they similarly display such an activity *in vivo*. Accordingly, TYRO3, AXL and MERTK (TAM) family members (42, 43) and the T cell immunoglobulin mucin (TIM) proteins 1 and 4 (TIM1/4) (44) play a crucial role in the recognition and phagocytosis of apoptotic cells by both types of phagocytic cells. However, since clodronate exhibit a preferential depleting activity on macrophages compared to dendritic cells, our data suggest macrophages to play a more important role.

One of the impacts of our study is that, although MSC remain the necessary starting point for therapeutic immunosuppression, patient-derived cells play a crucial role in delivering such an immunosuppression. Therefore, the efforts aimed at identifying the most clinically effective MSC subpopulation as well as the potency assays to validate such a selection may prove futile. A further proof supporting this concept is that the administration of *ex-vivo* generated apoMSC can circumvent the requirement for cytotoxic cells in a Th2 inflammatory model (Fig. 6D) and that apoMSC can be effective at suppressing the expansion/infiltration of the GvHD effector cells. Interestingly, apoMSC were mostly effective in the GvHD model only when administered i.p. (Fig. 7, A, B, C, and D). Despite being phagocytosed, apoMSC injected i.v. did not induce IDO production (Fig. 7, J and K), thus suggesting that the site at which MSC apoptosis occur may influence the immunosuppressive function, perhaps by engaging with a subpopulation of phagocytes.

A final question is whether a cytokine-dependent 'licensing' (9, 15, 45) co-exists with the generation of apoptotic MSC. Although our data indicate that cytokine-licensing is not required for the therapeutic activity, we cannot exclude that, before undergoing apoptosis, MSC directly inhibit inflammatory reactions through the conventional

pathways. Furthermore, caspase activation in MSC may trigger cell death independent signals that stimulate the synthesis of immunomodulatory molecules independently of the generation of signals for phagocytosis (46). Consistent with this, it has been shown that MSC activate caspase-dependent IL-1 signalling that enhances secretion of immunomodulatory molecules (47).

Our study constitutes a paradigm shift in MSC therapeutics whereby their apoptotic demise is a key step in the effector mechanism of immunosuppression exerted by MSC. A further impact of our discovery is that the principle underpinning this mechanism can be used as a biomarker to predict clinical responses to MSC and therefore stratify GvHD patients for MSC treatment. We therefore believe that the next generation of clinical trials should swing from choosing the best MSC population to choosing the patients most likely to respond. Furthermore, the intriguing possibility that apoMSC may be effective in patients refractory to MSC, paves the way to new avenues in the manufacturing of MSC.

## **Materials and Methods**

### *Study Design*

This study aimed to verify whether MSC undergo apoptosis after infusion and to test the role played by MSC apoptosis in the initiation of recipient-derived tolerogenic immune response.

A mouse model of GvHD, in which the disease is mediated by the expansion and activation of Mh CD8<sup>+</sup> cells in the recipient, was chosen for three important reasons: it recapitulates a minor mismatch between donor and recipient; T cells effecting GvHD

can be precisely enumerated; there is proof-of-principle that MSC are effective in treating GvHD. Furthermore, human MSC were used in order to avoid the confounding effects of recipient cytokines on MSC immune-modulating function (9, 45). In this system, murine inflammatory cytokines will not cross-react with the corresponding human receptors and will not activate immunosuppressive molecules in human MSC (14–16), whilst retaining the ability to expand murine effector cells mediating GvHD. Depletion of phagocytes (macrophages and dendritic cells) and inhibition of IDO production were conceived as loss of function experiments to assess the requirement of these factors in the delivery of MSC apoptosis-dependent immunosuppression.

A mouse model of ovalbumin (OVA)-induced allergic airway inflammation was used to assess whether the causative relationship between cytotoxic cells and MSC apoptosis in the delivery of MSC immunosuppression is valid in a disease associated with Th2-type inflammation.

All animal procedures were carried out in compliance with the UK Home Office Animals (Scientific Procedures) Act of 1986.

No randomization method was used. In all experiments animals were randomly allocated to control or experimental groups. No blinding approach was adopted. No statistical methods were used to predetermine sample size, which was estimated only on previous experience with assay sensitivity and the different animal models. Unless specified otherwise, three independent experimental replicates were performed.

To demonstrate that the presence of cytotoxic cells against MSC in GvHD patients could be predictive of MSC therapeutic activity, samples from GvHD patients were

collected and tested for their ability to induce MSC apoptosis in a cytotoxic assay within 24 hours before MSC infusion. At the time of performing the assay and cytofluorimetric analysis the operator was blind to patients' clinical details. All patients were affected by steroid-resistant GvHD and received MSC for compassionate use. PBMC from healthy donors were used as controls. All samples were collected after informed consent was obtained in accordance with the local ethics committee requirements.

#### *Mice and disease models*

Acute GvHD was induced as previously described (13). Briefly, after lethal irradiation (11 Gy), recipient C57BL/6 male mice were transplanted with  $1 \times 10^6$  purified CD8<sup>+</sup> cells from female Mh mice,  $5 \times 10^6$  unfractionated BM and  $2 \times 10^6$  purified CD4<sup>+</sup> cells from female C57BL/6 wild-type donors. The control group received BM and purified CD4<sup>+</sup> cells only.

For the depletion of all macrophages and dendritic cells mice received 1 mg liposome clodronate (ClodronateLiposome.com, Amsterdam, The Netherlands) i.v. 72 hours before MSC infusion (23). Recipient IDO activity was inhibited by using 1-DMT treatment (Sigma-Aldrich Company Ltd, Dorset, UK) (2mg/ml) in the drinking water starting from 6 days prior to MSC injection until animals were sacrificed (26).

C57BL/6-Prf1<sup>tm1Sdz</sup>/J (Perforin<sup>-/-</sup>) mice were purchased from J Jackson labs, bred with Matahari Rag2<sup>-/-</sup> mice and the resulting offspring intercrossed for 2 generations to obtain Mh Rag2<sup>-/-</sup>.Perf KO F3 mice.

OVA-induced airway inflammation was induced as previously described (20). Briefly, female Balb/C mice (Harlan Laboratories, Bicester, UK) were injected intraperitoneally with 30 µg of chicken egg albumin (OVA type V) on day 0 and 7. Controls received vehicle (aluminum hydroxide) only. On day 14, 15 and 16 animals were challenged with an aerosolized solution of OVA (3%) for 25 minutes. MSC or ApoMSC were injected 1 hour after the last challenge. After additional 18 hours, mice were terminally anaesthetized, a cannula inserted into the exposed trachea and three aliquots of sterile saline were injected into the lungs. The total number of cells in the lavage fluid was counted.

#### *MSC preparations*

Clinical grade BM-derived human MSC were generated from BM aspirates collected from the iliac crest of healthy donors. The cells were plated at a density of 10-25 million/636 cm<sup>2</sup>. After 3 days at 37 °C and 5% CO<sub>2</sub> non-adherent cells were discarded. When cell confluence of 90-100% was achieved, cells were detached with Trypsin-EDTA and reseeded at a density of 5000 cells/cm<sup>2</sup>. MSC were used at passage 2 for all *in vivo* experiments, whilst they were used by passage 8 for the *in vitro* experiments. In the latter case we did not observe any difference in terms of apoptosis susceptibility between different passages.

#### *Patients*

Between November 2012 and July 2016, 16 patients affected by steroid-resistant GvHD were treated with MSC according to Regulation (EC) No 1394/2007. All patients received GvHD prophylaxis. Of the 16 patients included in the study, 13 developed GVHD following hematopoietic stem cell transplantation, and the remaining 3 after

DLI. 12 patients were affected by acute GvHD, 3 by late onset acute GvHD and 1 by chronic GvHD. The diagnosis of GvHD was made on histological criteria and GvHD staged according to standard criteria (48, 49). Patient characteristics are summarized in Table S1. Samples were collected within 24 hours before MSC injection.

### *Statistics*

Results were expressed as mean $\pm$ SD. The unpaired Student *t* test was performed to compare 2 mean values. One-way ANOVA and Tukey's Multiple Comparison test was used to compare 3 or more mean values. Probability of null hypothesis less than 5% ( $p > .05$ , two-sided) was considered statistically significant.

Please see Supplementary Materials for full experimental procedures.

### **List of Supplementary Materials**

#### Supplemental Experimental Procedures

Figure S1. MSC can be traced in the lungs of mice after infusion

Figure S2. Human MSC immunosuppression is not 'licensed' by murine cytokines

Figure S3. MSC apoptosis is activated by cytotoxic cells in a non-antigen-specific manner

Figure S4. Cytotoxicity against MSC varies amongst PBMC donor but is independent on the percentage of CD8<sup>+</sup> or CD56<sup>+</sup> in GvHD patients

Figure S5. MSC killing is mediated by caspase 3 and effected by GrB and perforin

Figure S6. Infused MSC can be imaged in the lungs of mice with Th2-type lung inflammation

Table S1. Clinical features of GvHD patients

Video S1. Living cell imaging of FRET-MSC plated alone

Video S2. Living cell imaging of FRET-MSC plated with PHA-aPBMC

Video S3. Living cell imaging of FRET-MSC plated with resting PBMC

Video S4. Living cell imaging of FRET-MSC plated with PHA-aPBMC in the presence of the pan-caspase inhibitor Z-VAD-FMK

Video S5. Living cell imaging of FRET-MSC plated with PHA-aPBMC in the presence of the GrB inhibitor Z-AAD-CMK

Video S6. Living cell imaging of FRET-MSC plated with PHA-aPBMC in the presence of the perforin inhibitor EGTA

## References

**Acknowledgements:** We would like to thank Dr Domenico Spina (Institute of Pharmaceutical Science, King's College London) for his kind help and support for statistical analysis.

**Funding:** Bloodwise specialist programme 14019, Bloodwise specialist programme 12006. AG is a recipient of a Bloodwise Clinical Research Training Fellowship 15029.


**Author Contributions:** Conceptualization: A.G, H.W and F.D. Methodology: A.G., F.D, C.B, Y.R.V and R.C. Formal. Analysis: A.G and L.D. Investigation: A.G., Y.R.V., C.T., C.L., L.B., S.W., K.H., T.S.C. and L.W. Resources: F.D., K.O., D.I. M., J.A., M. V. B. and M. B. Visualization: A.G. and L.D. Writing - Original Draft: A.G. and F.D.

Writing – Review & Editing: A.G., C.B., G.L., F. M. W., H.W. and F.D. Funding acquisition: F.D. Supervision: F.D.

**Competing financial interests:** The authors declare no competing financial interests.

**Data and materials availability:** Reasonable requests for additional data or materials will be fulfilled under appropriate agreements.


## Figures






**Figure 1: MSC undergo *in vivo* apoptosis after infusion without affecting delivery of immunosuppression. A:** luc-MSCs were injected i.v. into naïve, BM and GvHD mice 3 days after transplantation. All animals were then injected intraperitoneally with DEVD-aminoluciferin and imaged 1 hour later. *N*: 6 mice per group, grouped from 3 independent experiments. White lines separate multiple photographs assembled in the final image. **B:** TLS was measured from the images of mice in Fig. 1A and shown as mean±SD. **C, D:** Infiltration of GvHD effector cells (CD8<sup>+</sup>Vβ8.3<sup>+</sup>) in the spleen (**C**) and lungs (**D**) of GvHD mice (black circles) and GvHD mice treated with MSC (black squares), 4 days after MSC injection. *N*: 15 (GvHD) and 13 (GvHD+MSC) mice, grouped from 4 independent experiments; mean±SD are shown. Statistics in B: one-way ANOVA, with Tukey's Multiple Comparison Test. \*\*: p<.01, \*\*\*: p<.001, ns: not significant. In C and D: unpaired t-test. \*\*: p<.01.





**Figure 2. MSC apoptosis is indispensable for immunosuppression and requires functionally activated cytotoxic cells in the recipient.** **A:** The percentage of CD8<sup>+</sup>Vβ8.3<sup>+</sup> cells in lung cell suspensions from naïve C57BL/6 male, BM or GvHD mice was analyzed in the lymphocyte population; mean±SD are shown. *N*: 12 (GvHD), 3 (BM) and 3 (naïve) mice, grouped from 3 independent experiments. **B:** CD8<sup>+</sup> cells were sorted from the lungs and spleens of naïve female

Mh (grey bars) or GvHD mice (white bars) 7 days after transplant and tested for their ability to induce MSC apoptosis *in vitro*. The results show annexin V<sup>+</sup>/7-AAD<sup>-</sup> MSC (mean±SD) in 3 independent experiments (N=10 per group), black bar represents the level of apoptosis in MSC cultured alone used as control (N: 3) **C**: luc-MSCs were infused in three independent experiments in GvHD (N=7) and GvHDPerf<sup>-/-</sup> (N=7) mice 3 days after transplantation. 1 hour later mice were injected with DEVD-aminoluciferin and imaged. White lines separate multiple photographs assembled in the final image. **D**: TLS was obtained from Fig. 2C and expressed as mean±SD. **E**, **F**: infiltration of effector GvHD cells (CD8<sup>+</sup>Vβ8.3<sup>+</sup>) in the spleen (**E**) and lungs (**F**) of untreated GvHDPerf<sup>-/-</sup> (N=16) and GvHDPerf<sup>-/-</sup> (N=17) mice treated with MSC (mean±SD of 4 independent experiments). Statistics in A and B: one-way ANOVA, with Tukey's Multiple Comparison Test. \*: p<.05; \*\*\*: p<.001. In D, E and F: unpaired t-test. \*\*\*: p<.001. ns: not significant.




**Figure 3. Cytotoxic activity against MSC predicts clinical responses to MSC in GvHD patients. A, B:** PBMC obtained from healthy controls (HC) or patients with GvHD receiving MSC in the following 24 hours were incubated in 24-well plates with MSC at a 20/1 PBMC/MSR ratio for 4 hours. The level of apoptosis was measured in MSC assessing the level of annexin-V/7-AAD by flow-cytometry. **(A)** Representative plots for HC, clinical responders (R) and non-responders (NR). **(B)** The level of apoptosis was compared among HC (circles,  $N=5$ ), R (triangles;  $N=5$ ) and NR


(squares;  $N=12$ ). Statistics: one-way ANOVA and Tukey's Multiple Comparison test.

\*\*\*:  $p < .0001$ . ns: not significant.


A




B




C




D




E




F




G



H




I




**Figure 4. MSC apoptosis is mediated by activated CD8<sup>+</sup> and CD56<sup>+</sup> cytotoxic cells and is the result of a bystander effect.** **A:** PBMC from healthy donors were activated using phytohemagglutinin (PHA) (PHA-aPBMC) or Mixed Lymphocyte Reaction (MLR) (MLR-aPBMC). Resting (light grey bars), PHA-aPBMC (black bars) or MLR-aPBMC (dark grey bars) were incubated with MSC at the indicated ratios for 4 hours. ND: Not done. **B, E, F, I:** MLR-aPBMC or PHA-aPBMC were cultivated with MSC in the presence or absence of the pan-caspase inhibitor Z-VAD-FMK (10  $\mu$ M) (**B**), GrB inhibitor Z-AAD-CMK (300  $\mu$ M), perforin inhibitor ethylene glycol-bis(2-aminoethylether)-N,N,N',N'-tetraacetic acid (EGTA) (4 mM) (**E**), neutralizing concentrations of FAS-L mAb anti-CD178 (**F**), or escalating doses (10 to 75  $\mu$ M) of PKC $\zeta$ -PS (**I**). **H:** MLR-aPBMC were cultivated with MSC in direct contact or separated by a transwell<sup>®</sup>. **C, D:** MLR-aPBMC were used unfractionated, positively selected for CD11b<sup>+</sup>, CD4<sup>+</sup>, CD8<sup>+</sup> or CD56<sup>+</sup> cells (**C**) or depleted of CD56<sup>+</sup>, CD8<sup>+</sup> or both (**D**). **G:** apoptosis in MSC after culture with autologous (black bars) or allogeneic (grey bars) PHA-aPBMC in the presence or absence of neutralizing doses of anti-HLA-A-B-C or anti-HLA-DR antibodies. In **B- I:** the PBMC/MSK ratio was 20/1. Results represent the mean  $\pm$  SD of 3 or 6 (H) independent experiments. Statistics: one-way ANOVA, with Tukey's Multiple Comparison Test. \*: p<0.5. \*\*: p<.01. \*\*\*: p<.001. ns: not significant.






**Figure 5. MSC apoptosis does not interfere with the antigen-specific cytotoxic cell recognition of the cognate target.** **A:** apoptosis in T2-cell after culture with 4D8 cells at a 20/1 4D8:T2 ratio. Where indicated increasing concentrations of MSC (used as *cold* target) were added. Apoptotic T2 cells were identified as annexin V<sup>+</sup>/7-AAD<sup>+</sup> cells. **B:** apoptosis in K562 cultured with NK cells (20/1 NK:K562 ratio). Where indicated increasing concentrations of MSC (used a *cold* target) were added. **C:** apoptosis in MSC cultured with 4D8 cells (20/1 4D8:MSC ratio). Where indicated increasing concentrations of T2 cells (used as *cold* target) were added. **D:** apoptosis in MSC cultured with NK cells at a 20/1 NK:MSC ratio. Where indicated increasing dilutions of K562 (used as *cold* target) were added. In all experiments the level of MSC, T2 or K562 cell apoptosis was assessed after 4 hours of co-culture by flow

cytometry. Results represent the mean  $\pm$ SD of 3 independent experiments. Statistics in **A**, **B**, **C** and **D**: unpaired T-test. \*:  $p < .05$ . ns: not significant.




**Figure 6. Apoptotic MSC exert *in vivo* immunosuppressive in a Th2-type inflammation model in the absence of cytotoxic cells.** **A:** luc-MSCs were injected into naïve ( $N=3$ ) and OVA+MSC ( $N=6$ ) mice one hour after the last challenge. One hour later, mice received DEVD-aminoluciferin and were imaged in 3 independent experiments. White lines separate multiple photographs assembled in the final image. **B:** TLS was measured from Fig. 6A (mean $\pm$ SD). **C:** Eighteen hours after MSC infusion, eosinophil infiltration was assessed in the BAL of naïve ( $N=3$ ), naïve infused with MSC ( $N=3$ ), OVA ( $N=6$ ) and OVA+MSC ( $N=6$ ) mice in two independent experiments and mean $\pm$ SD are shown. **D:** eosinophil infiltration (mean $\pm$ SD) in BAL of OVA-sensitized mice treated with ApoMSC. Groups were: OVA without ApoMSC ( $N=6$ ), OVA treated with  $1 \times 10^6$  ApoMSC ( $N=7$ ); and naïve mice receiving  $1 \times 10^6$  ( $N=2$ ) ApoMSC. Results represent the mean $\pm$ SD of 3 independent experiments.

Statistics in **B**: unpaired t-test. ns: not significant. Statistics in **C and D**: one-way ANOVA and Tukey's Multiple comparison test. \*:  $p < .05$ . ns: not significant.



**Figure 7. ApoMSC exert immunosuppressive activity in GvHD and are engulfed by recipient phagocytic cells in which they elicit IDO production. A-**

**D:** Infiltration of GvHD effector cells was assessed in spleen (A, C) and lungs (B, D) of GvHD mice (black circles) and GvHD mice treated with ApoMSC (black squares). ApoMSC were infused i.p. (GvHD mice  $N=10$ , GvHD+ApoMSC mice  $N=8$ ) (A, B), or i.v. (GvHD mice  $N=9$ , GvHD+ApoMSC mice  $N=7$ ) (C, D). Results represent the mean $\pm$ SD of 3 independent experiments. Statistics: unpaired t-test. \*:  $p<.05$ ; \*\*:  $p<.01$ . ns: not significant. **E-K:** MSC were labelled using CellTrace™ Violet and subjected to apoptosis induction using GrB/FAS-L (5  $\mu$ g/ml and 10  $\mu$ g/ml, respectively). ApoMSC were injected i.p. (E, F and J) or i.v. (G, H, I and K) into GvHD mice 3 days after the transplant. After 2 hours, animals were sacrificed and mesenteric lymph nodes (E, F and J) or lungs (G, H, I and K) were harvested. Cells engulfing ApoMSC were identified as Violet<sup>+</sup> cells within the CD11b<sup>+</sup> (E), CD11c<sup>+</sup> (F), CD11b<sup>high</sup>CD11c<sup>int</sup> (G), CD11c<sup>+</sup>CD11b<sup>-</sup> (H) and CD11b<sup>high</sup>CD11c<sup>-</sup> (I) subpopulations. The corresponding subpopulations were gated in GvHD mice which had not received violet-labelled ApoMSC. **J** and **K:** IDO expression was assessed in CD11c<sup>+</sup> and CD11b<sup>+</sup> (J) or CD11b<sup>high</sup>CD11c<sup>int</sup>, CD11c<sup>+</sup>CD11b<sup>-</sup> and CD11b<sup>high</sup>CD11c<sup>-</sup> (K) cells positive for CellTrace Violet (engulfing apoMSC) and compared with the corresponding populations in GvHD mice that had not received ApoMSC. Data are representative of similar results obtained from three mice in 2 independent experiments.



**Figure 8.** Recipient phagocytes and IDO production are required for MSC immunosuppressive activity in GvHD. **A, B:** GvHD mice were treated with liposomal clodronate 10 minutes after the transplant. Where indicated, MSC were infused 3 days later. The infiltration of GvHD effector cells ( $CD8^+V\beta 8.3^+$ ) in spleen (A) or lungs (B) was quantitated in spleen and lungs after 4 additional days. Mean $\pm$ SD obtained grouping three independent experiments with  $N$ : 12 (GvHD) and 10 (GvHD+MSC) mice per group. **C, D:** GvHD effector cell infiltration was studied in spleen (C) and lungs (D) of GvHD mice treated with the IDO-inhibitor 1-DMT. In the treated mice, MSC were infused 3 days after the transplant ( $N=11$ ). Controls consisted of GvHD mice which did not receive MSC ( $N=9$ ). Results refers to the mean $\pm$ SD of 3 independent experiments. Statistics: unpaired t-test. \*:  $p<.05$ ; \*\*:  $p<.01$ . ns: not significant.

## Supplementary Materials

### **Apoptosis in mesenchymal stromal cells is required to initiate *in vivo* recipient-mediated immunomodulation**

Antonio Galleu, Yanira Riffo-Vasquez, Cristina Trento, Cara Lomas, Luigi Dolcetti, Tik Shing Cheung, Malte von Bonin, Laura Barbieri, Krishma Halai, Sophie Ward, Ling Weng, Ronjon Chakraverty, Giovanna Lombardi, Fiona M. Watt, Kim Orchard, David I. Marks, Jane Apperley, Martin Bornhauser, Henning Walczak, Clare Bennet, and Francesco Dazzi

#### **List of Supplementary Materials**

Supplemental Experimental Procedures

Figure S1. MSC can be traced in the lungs of mice after infusion

Figure S2. Human MSC immunosuppression is not 'licensed' by murine cytokines

Figure S3. MSC apoptosis is activated by cytotoxic cells in a non-antigen-specific manner

Figure S4. Cytotoxicity against MSC varies amongst PBMC donor but is independent on the percentage of CD8<sup>+</sup> or CD56<sup>+</sup> in GvHD patients

Figure S5. MSC killing is mediated by caspase 3 and effected by GrB and perforin

Figure S6. Infused MSC can be imaged in the lungs of mice with Th2-type lung inflammation

Video S1. Living cell imaging of FRET-MSK plated alone

Video S2. Living cell imaging of FRET-MSK plated with PHA-aPBMC

Video S3. Living cell imaging of FRET-MSK plated with resting PBMC



Video S4. Living cell imaging of FRET-MSC plated with PHA-aPBMC in the presence of the pan-caspase inhibitor Z-VAD-FMK

Video S5. Living cell imaging of FRET-MSC plated with PHA-aPBMC in the presence of the GrB inhibitor Z-AAD-CMK

Video S6. Living cell imaging of FRET-MSC plated with PHA-aPBMC in the presence of the perforin inhibitor EGTA

Table S1. Clinical features of GvHD patients

## Supplemental Experimental Procedures

### *Mice and disease models*

C57BL/6 (H2b) and Balb/C (H2d) mice were purchased from Harlan Laboratories (Bicester, UK). Mh (C57BL/6 background, CD8<sup>+</sup>Tg, H-2b, CD45.2<sup>+</sup>, H-2Db-restricted) (50) mice are transgenic for a T-cell receptor specific for the male antigen UTY presented in the context of H-Db. Mice were bred in-house and maintained at the Biological Service Unit of the Royal Free and University College London Medical School (London, UK). All mice were used between 6 and 12 weeks of age.

Acute GvHD was induced as previously described (13). Briefly, after lethal irradiation (11 Gy), recipient C57BL/6 male mice were transplanted with 1x10<sup>6</sup> purified CD8<sup>+</sup> cells from female Mh mice, 5x10<sup>6</sup> unfractionated BM and 2x10<sup>6</sup> purified CD4<sup>+</sup> cells from female C57BL/6 wild-type donors. The control group received BM and purified CD4<sup>+</sup> cells only. CD4<sup>+</sup> and CD8<sup>+</sup> T cells were obtained by positive selection using magnetic beads (Miltenyi Biotec Ltd, Biscley, UK). Live MSC (1x10<sup>6</sup>) were injected i.v. at day +3, whilst apoMSC (1x10<sup>6</sup>) were administered i.v. or i.p. at day +1, +3 and +6 from the transplant. In all cases, animals were euthanized for analysis at day +7. The infiltration of GvHD effector cells was assessed by flow-cytometry and the percentage was expressed as proportion of cells in the lymphocyte gate, based on the physical characteristics of the cells.

OVA-induced airway inflammation was induced as previously described (20). Briefly, female Balb/C mice (Harlan Laboratories, Bicester, UK) were injected i.p. with 30 µg of chicken egg albumin (OVA type V) (Sigma-Aldrich Company Ltd, Dorset, UK) on day 0 and 7. Controls received vehicle (aluminum hydroxide) only. On day 14, 15 and

16 animals were challenged with an aerosolized solution of OVA (3%) for 25 minutes. MSC or ApoMSC were injected 1 hour after the last challenge. After additional 18 hours, mice were terminally anaesthetized with urethane (2 g/kg i.p.), (Sigma-Aldrich Company Ltd, Dorset, UK), a cannula inserted into the exposed trachea and three 0.5 mL aliquots of sterile saline were injected into the lungs. The total number of cells in the lavage fluid was counted. For differential cell counts, cytospin preparations were stained with Diff Quick (DADE Behring, Germany) and cells counted using standard morphological criteria.

No randomization method was used. In all experiments animals were randomly allocated to control or experimental groups. No blinding approach was adopted.

Animal procedures were carried out in compliance with the UK Home Office Animals (Scientific Procedures) Act of 1986.

#### *Cell preparations and media*

Cultures were carried out in complete RPMI 1640 medium containing GlutaMAX™, HEPES (25mM), Penicillin 5000 U/ml and Streptomycin 5000 µg/ml (ThermoFisher Scientific, Paisley, UK), fetal bovine serum 10% (Labtech.com, Uckfield, UK).

Human peripheral blood samples from healthy donors were procured by the National Blood Service (Colindale, UK) as leukocyte cones. Samples from GvHD patients were collected within 24 hours before MSC injection. Informed consent was obtained in accordance with the local ethics committee requirements. PBMC were isolated by density gradient separation on Histopaque-1077 (SigmaAldrich Company Ltd, Dorset, UK).

mSpl were isolated through a cell strainer (BD Falcon, Oxford, UK), whilst lungs were cut into small pieces and incubated with Collagenase type IV (250 U/ml) (Lorne Laboratories, Reading, UK), DNase I from bovine pancreas (250 U/ml) (Merk Millipore, Watford, UK) and fetal bovine serum 6.25% at 37<sup>o</sup> C for 1 hour.

### *MSC preparations*

Clinical grade BM-derived human MSC were generated from BM aspirates collected from the iliac crest of healthy donors. Briefly, 2 ml of BM aspirate was collected in a tube with 100 µl preservative-free heparin. The cells were plated within 24 hours at a density of 10-25 million/636 cm<sup>2</sup> by using alpha modified Eagle's medium (ThermoFisher Scientific, Paisley, UK), conservative-free heparin (1 UI/ml) (Wockhardt UK Limited, Wrexham, UK) and 5% platelet lysate and then incubated for 3 days at 37 °C and 5% CO<sub>2</sub> ambience. Non-adherent cells were discarded using phosphate buffered saline (ThermoFisher Scientific, Paisley, UK). When cell confluence of 90-100% was achieved cells were detached with Trypsin-EDTA (0.05% trypsin, 0.5γ mM EDTA•4Na) (ThermoFisher Scientific, Paisley, UK) and reseeded at a density of 5000 cells/cm<sup>2</sup>. MSC were used at passage 2 for all in vivo experiments, whilst they were used by passage 8 for the in vitro experiments. In the latter case we did not observe any difference in terms of apoptosis susceptibility between different passages. Released criteria were based on positivity (> 80%) for CD105, CD90, CD73, negativity (<2%) for CD3, CD14, CD19, CD31, CD45.

ApoMSC were obtained by plating 5x10<sup>5</sup> cells per well in a 96 round-bottom well plate in the presence of synthetic human GrB (5 µg/ml) (Enzo Life Sciences, Exeter, UK) and anti-FAS human (activating, clone CH11) (10 µg/ml) (Merk Millipore, Watford, UK)

for 24 hours in complete RPMI. The concentration of GrB and FasL was chosen to produce at least 80% of MSC apoptosis. Where indicated (Z-Apo), the pan-caspase inhibitor Z-VAD-FMK (10 µg/ml) was added for the entire duration of the treatment.

### *Patients*

Between November 2012 and July 2016, 16 patients affected by steroid-resistant GvHD were treated with MSC in the Department of Haematology at Imperial College London, Southampton University Hospital, Bristol Haematology and Oncology Centre and the University Hospital Carl Gustav Carus, Dresden. MSC were administered for compassionate use (according to Regulation (EC) No 1394/2007). Patients had received a myeloablative or reduced-intensity conditioning prior to hematopoietic stem cell transplantation. All patients received GvHD prophylaxis with 3 or 4 doses of methotrexate combined with cyclosporin. T-cell depletion with alemtuzumab or ATG was performed in all adult patients transplanted in the UK centers. Of the 16 patients included in the study, 13 developed GVHD following hematopoietic stem cell transplantation, and the remaining 3 after DLI. 12 patients were affected by acute GvHD, 3 by late onset acute GvHD and 1 by chronic GvHD. The diagnosis of GvHD was made on histological criteria and GvHD staged according to standard criteria (48, 49).

Patients were considered to be steroid-refractory if: (a) those with aGVHD failed to respond to high-dose methylprednisolone after 6 days; (b) the one with cGVHD failed to respond to high-dose steroids after 2-4 weeks, with the addition of MMF and cyclosporin at 1 and 4 weeks respectively. Clinical responses to MSC were assessed

1 week after MSC infusion and defined as an improvement of at least 50% in at least one organ affected by GvHD. Patient characteristics are summarized in Table S1.

### *Imaging of MSC*

Luc-MSCs were transfected with the pGL3-Control vector containing the SV40 promoter for the expression of luc<sup>+</sup> (Promega, Southampton, UK) or with pECFP-DEVDR-Venus (Addgene, Teddington, UK) using electroporation (Gene Pulser Xcell, BioRad, Kidlington, UK). Cells were suspended in a total volume of 250 µl of buffer and electroporated in 0.4 cm gap cuvettes using 10 µg of DNA at 250 volts and 950 F. When pECFP-DEVDR-Venus was used, the donor fluorophore pECFP and the acceptor Venus-YFP were linked through the flexible linker DEVDR which is recognized and cleaved by the active form of caspase 3. In this system caspase 3 activity can be monitored through the analysis of the FRET between pECFP and Venus-YFP. When caspase 3 is not active, the flexible linker DEVDR remains intact and energy transfer from pECFP is allowed with emission of YFP signal. Conversely, in the presence of caspase 3 activation DEVDR is cleaved, thus energy transfer is lost and the pECFP signal increases.

For confocal imaging, pECFP-DEVDR-Venus transfected MSCs were plated in complete RPMI at a concentration of  $1 \times 10^5$  cells in a 30mmx10 mm dish (Corning, Flintshire, UK) and let adhere overnight. The following day PHA-aPBMCs were added at a PBMC:MSC ratio of 40/1. Where indicated, pancaspase inhibitor Z-VAD-FMK (50 µM), EGTA (4 mM), GrB inhibitor Z-AAD-CMK (300 µM) were used. Living cell imaging was acquired every 3 minutes for 180 minutes using a Leica TCS-SP5 II Confocal

Microscope, with 488 nm and 407 nm lasers. The images were processed and analyzed by using the software “R” and EBIImage package.

In vivo imaging was performed injecting i.v.  $1 \times 10^6$  luc-MSCs into naïve C57BL/6, BM or GvHD mice 3 days after the transplant in the GvHD model. In the airway inflammation model, luc-MSCs were infused i.v. in naïve Balb/C or OVA-treated mice 1 hour after the last OVA challenge. After one additional hour, mice were anesthetized with isoflurane (1.5% isoflurane, 98.5% Oxygen), injected i.p. with 3 mg of VivoGlo™ Casp 3/7 Substrate Z-DEVD Aminoluciferine (Promega, Southampton, UK) and imaged using IVIS® Lumina III (PerkinElmer, Waltham, USA) system for a total time of 5 minutes. Images were analyzed by using the software “R” and EBIImage package to obtain mean TLS. Confirmation of the presence of transfected MSC was obtained injecting mice with VivoGlo™ Luciferin (Promega, Southampton, UK).

#### *Detection of efferocytosis*

MSCs were first labelled using CellTrace™ Violet labelling (ThermoFisher Scientific, Paisley, UK) at a final concentration of 5  $\mu$ M. Then ApoMSCs were obtained as described above, using synthetic human GrB (5  $\mu$ g/ml) and anti-FAS human (10  $\mu$ g/ml) for 24 hours.  $10 \times 10^6$  labelled apoMSCs were then injected i.p. or i.v. and mice sacrificed after 2 hours post-injection. Spleen, lungs, peritracheal, paratracheal, pericardial, mesenteric, periportal and celiac lymph nodes were collected and analysed by flow-cytometry. Positivity of CellTrace Violet was assessed as a measure of ApoMSC engulfment in CD11b<sup>+</sup> and CD11c<sup>+</sup> gated subpopulations of phagocytic cells. Cells positive for the CellTrace Violet were then assessed for their expression of IDO.

### *Pre-activation of human PBMC and murine CD8<sup>+</sup> cells*

PHA-aPBMC were obtained plating  $5 \times 10^6$  human PBMC in 24-well plate in the presence of PHA (5  $\mu\text{g/ml}$ ) (Sigma-Aldrich Company Ltd, Dorset, UK) in a final volume of 2 ml of complete RPMI for 72 hours. MLR-aPBMC were obtained using one-way MLR in which PBMC from one donor (stimulators) were irradiated (30 Gy) and co-cultured with the PBMC of an unrelated donor (responder) at a stimulator:responder ratio of 1/1 in complete medium at a density of  $0.75 \times 10^6$  cells/cm<sup>2</sup>. Cells were then incubated at 37<sup>0</sup> C, 5% CO<sub>2</sub> for 5 days.

NK cells were purified by positively selecting CD56<sup>+</sup> cells from healthy donor PBMC (Miltenyi Biotec Ltd, Bisley, UK) and activated with rh-IL-2 (1000 U/ml).

NY-ESO1-specific CD8<sup>+</sup> T cell clone (Clone 4D8) was kindly supplied by Prof. Vincenzo Cerundolo (Institute of Molecular Medicine, Oxford university, UK). The clone was expanded in complete RPMI 1640 with Sodium Pyruvate (1 mM), 2-Mercaptoethanol (0.05 mM) (ThermoFisher Scientific, Paisley, UK), recombinant human-IL-2 (rh-IL-2) (400 U/ml) (Peprotec EC Ltd, London, UK) and PHA (5  $\mu\text{g/ml}$ ) (Sigma-Aldrich Company Ltd, Dorset, UK) (51).

Mh CD8<sup>+</sup> were stimulated using the following protocol:  $5 \times 10^6$  purified CD8<sup>+</sup> Mh cells were plated in 24-well plates in the presence of CD3/CD28-coated beads (Dynabeads®) (ThermoFisher Scientific, Paisley, UK) in a final volume of 2 ml of complete RPMI and incubated for 72 hours.

### *Immunosuppressive assay*

Serial dilutions of human MSC were plated in a flat bottom 96-well plate and let adhere overnight in 100  $\mu\text{l}$  of complete RPMI. Where indicated, MSC cultures were exposed



to hIFN- $\gamma$  and hTNF- $\alpha$ , mIFN- $\gamma$  and mTNF- $\alpha$  (20 ng/ml each) (all cytokines were from Peptotec EC Ltd, London, UK), supernatant from PHA-aPBMC or from ConA-aSpl. The following day,  $5 \times 10^5$  Balb/C mSpl were labelled with Carboxyfluorescein Diacetate Succinimidyl Ester (ThermoFisher Scientific, Paisley, UK) dye and plated with MSC at escalating MSC/mSpl ratios. The culture controls consisted of mSpl plated without MSC in the presence (positive control) or in the absence of ConA (negative control). Proliferation of mSpl was then assessed by flow-cytometry after 72 hours and expressed as the percentage of the proliferation obtained at each MSC/mSpl dilution in comparison with the one obtained in the positive control culture. Results were expressed as percentage of inhibition.

#### *Cytotoxic Assay*

$1 \times 10^5$  MSC were plated overnight in a total volume of 500  $\mu$ l. The day after pre-activated immune cells were plated at different concentrations (2.5 to 40/1 effector:MSC ratios). MSC apoptosis was then tested at different time points using flow-cytometry or confocal microscopy analysis. Eventually, the assay was performed for 4 hours in the vast majority of the cases. At flow-cytometry MSC were identified as CD45- cells.

Antigen-specific cytotoxic activity of clone 4D8 was tested using T2 cells pulsed with NY-ESO-1 antigen (epitope SLLMWITQC) at a concentration of 0.1  $\mu$ M for 1 hour.

In the competition assay, T2 (from Hans Stauss, University College London) and K562 cells (from Junia Melo, Imperial College London) were discriminated from effector cells by CellTrace<sup>TM</sup> Violet labelling. The tracer concentration was optimized for the T2 (1  $\mu$ M) and K562 (2.5  $\mu$ M) cells. Cell lines were tested for mycoplasma contamination before use.

When flow-cytometry was used, the level of apoptosis was assessed using the PE Annexin V apoptosis detection kit (BD Biosciences, Oxford, UK). Unless specified, apoptotic cells were identified as annexin V<sup>+</sup>/7-AAD<sup>-</sup> cells.

### *Inhibitors*

Where indicated cultures were supplemented with pan-caspase inhibitor Z-VAD-FMK (10 µM in the flow-cytometry experiments or 50 µM in the living cell confocal experiments) (R&D System, Oxon, UK), perforin inhibitor EGTA (4 mM) (Sigma-Aldrich Company Ltd, Dorset, UK), GrB inhibitor Z-AAD-CMK (300 µM) (Merk Millipor, Watford, UK), neutralizing antibodies against HLA-DR (clone L243) (50 µg/ml), human HLA-A,B,C (clone W6/32) (100 µg/ml) (BD Biosciences, Oxford, UK), TNF-α antagonist Etanercept (Enbrel®) (10 µg/ml or 100 µg/ml) (Amgen, Cambridge, UK). Each reagent was incubated with MSC 1 hour before the culture with effector killer cells. In all cases, the concentration of the corresponding inhibitor was kept for the duration of the cytotoxic assay.

The neutralizing anti-CD178 (Clone NOK-1) (10 µg/ml or 100 µg/ml) (BD Biosciences, Oxford, UK), anti-TRAIL (clone 2E2) (10 µg/ml or 100 µg/ml) (Enzo Life Sciences, Exeter, UK) antibodies, MYR Protein Kinase-Cζ Pseudosubstrate (PKCζ-PS) (10 µM, 25 µM or 75 µM) (ThermoFisher Scientific, Paisley, UK) and Etanercept (10 µg/ml or 100 µg/ml) were incubated with effector killer cells for 2 hours before the cultures with MSC. In all cases, the concentration of the corresponding inhibitor was kept for the duration of the cytotoxic assay.

### *Flow-cytometry*

The following antibodies specific for murine molecules were used: anti-CD45 (FITC, Clone 30-F11) (eBiosciences Ltd, Hatfield, UK), anti-V $\beta$ 8.3 (FITC, Clone 1B3.3), anti-CD8 (APC, Clone 53-6.7), antiCD4 (PE, Clone H129.19), anti-CD19 (APC-H7, Clone 1D3), anti-NK1.1 (PerCP-Cy5.5, Clone PK136) (BD Biosciences, Oxford, UK), anti-CD11b (PerCP-Cy5.5, clone M1/70), anti-CD11c (APC-Cy7, clone n418), Ido1 (Alexa Fluo647, clone 2e2) (BioLegend, London, UK). For human specific molecules, the following antibodies were used: antiCD45 (FITC, clone 2D1), anti-CD8 (APC, Clone SK1), anti-CD4 (PE, Clone SK3), anti-CD11b (PerCP-Cy5.5, clone M1/70), anti-CD56 (FITC, clone HCD56) (BD Biosciences, Oxford, UK).

All samples were acquired using BD FACS Canto II using the software FACS Diva and analyzed with Flow-jo software. FRET and CAf were assessed by flow-cytometry as previously described (52).


### *Real Time quantitative PCR*

MSC RNA was obtained from TRIzol® (ThermoFisher Scientific, Paisley, UK) lysates and extracted using RNeasy Mini Kit (Qiagen, Manchester, UK). Real Time quantitative PCR (qRT-PCR) was performed following TaqMan® RNA-to-CT™ 1-Step Kit instructions (ThermoFisher Scientific, Paisley, UK), using 20 ng of RNA template per reaction. Assays were carried out in duplicates on an StepOnePlus RT PCR system thermal cycler (Applied Biosystem, UK) using TaqMan primers (all purchased from ThermoFisher Scientific, Paisley, UK). The human primers used were the following: IDO2 (Hs01589373\_m1), TSG6 (Hs01113602\_m1) and PTSG2 (Hs00153133\_m1) and HPRT1 (Hs02800695\_m1) as housekeeping gene. Data were

then analysed using StepOne™ software version 2.1 and relative quantification obtained with  $\Delta\Delta C_t$  method, considering untreated MSC as reference.






### *Statistics*

Results were expressed as mean $\pm$ SD. The unpaired Student *t* test was performed to compare 2 mean values. One-way ANOVA and Tukey's Multiple Comparison test was used to compare 3 or more mean values. Probability of null hypothesis less than 5% ( $p > .05$ , two-sided) was considered statistically significant. No statistical methods were used to predetermine sample size, which was estimated only on previous experience with assay sensitivity and the different animal models.




**Figure S1. MSC can be traced in the lungs of mice after infusion. A:** lethally irradiated C57BL/6 male mice were transplanted with bone marrow (BM) and CD4<sup>+</sup>-purified cells from female syngeneic donors with or without CD8<sup>+</sup> cells purified from Mh mice (CD8<sup>+</sup>V $\beta$ 8.3<sup>+</sup>) (GvHD and BM groups, respectively). At day +3 post-transplant, luc-MSC were infused and mice imaged one hour later for the analysis of caspase 3 activation after i.p. injection of DEVD-aminoluciferin. At day +7 post-

transplant, mice were sacrificed and the infiltration of GvHD effector cells (CD8<sup>+</sup>V $\beta$ 8.3<sup>+</sup>) in lungs and spleen was analyzed by flow-cytometry. **B**: in order to confirm the presence of luc-MSK in the lungs of all groups of mice infused with MSC, the same mice imaged in Figure 1A were injected with D-Luciferin. White lines separate multiple photographs assembled in the final image. **C**: TLS was measured from the images of mice in Fig. S1B and shown as mean $\pm$ SD. Statistics: 1-way ANOVA, with Tukey's Multiple Comparison Test. ns: not significant.




**Figure S2. Human MSC immunosuppression is not ‘licensed’ by murine cytokines.** **A:** human MSC were plated overnight at serial dilutions alone or in the presence of human interferon- $\gamma$ /human Tumour Necrosis- $\alpha$  (hIFN- $\gamma$ /hTNF- $\alpha$ ) (20 ng/ml each) or murine IFN- $\gamma$ /murine TNF- $\alpha$  (mIFN- $\gamma$ /mTNF- $\alpha$ ) (20 ng/ml each) or supernatant obtained from PBMC activated with PHA (PHA-aPBMC) for 72 hours or mSpl activated with ConA (ConA-aSpl) for 72 hours, as indicated. MSC were then tested for the ability to inhibit the proliferation of ConA-stimulated mSpl labelled with carboxyfluorescein succinimidyl ester. Proliferation was determined after 72 hours by flow-cytometry. The curve was obtained plotting the percentage of inhibition against the corresponding MSC/mSpl ratio. **B, C:** human MSC were plated overnight either untreated or exposed to hIFN- $\gamma$ /hTNF- $\alpha$  (20 ng/ml each) as indicated and then tested for the ability to suppress mSpl proliferation at 1:10 MSC/mSpl ratio. The histogram plot (**B**) is representative of 3 independent experiments, while bars (**C**) represent the mean $\pm$ SD of 3 independent experiments. Statistics: one-way ANOVA and Tukey’s Multiple Comparison test. \*\*\*:  $p < .001$ . ns: not significant. **D:** human MSC were incubated alone or in the presence of hIFN- $\gamma$ /hTNF- $\alpha$  (20 ng/ml each), mIFN- $\gamma$ /mTNF- $\alpha$  (20 ng/ml each), supernatants obtained from PHA-aPBMC or ConA-aSpl. After 24 hours, *IDO*, *TSG6* and *PTSG2* expressions were assessed by real time PCR and calculated as relative expression in comparison to untreated MSC. Representative results of three independent experiments are shown.
















**Figure S3. MSC apoptosis is activated by cytotoxic cells in a non-antigen-specific manner. A:** CD8<sup>+</sup> cells isolated from naïve female Mh mice were stimulated for 3 days with anti-CD3/CD28 beads and cultured with MSC at a 20/1 Mh T-cell:MSC ratio. After 4 hours the level of apoptosis was assessed in MSC by annexin V/7AAD staining. Results represent the mean±SD of 3 independent experiments. Statistics: one way ANOVA, with Tukey's Multiple Comparison Test. \*\*\*: p<.001. **B:** in order to confirm the presence of luc-MSC in the lungs of all groups of mice infused with MSC, the same mice imaged in Figure 2C were injected with D-Luciferin. White lines separate multiple photographs assembled in the final image. **C:** TLS was measured from the images of mice in Fig. S3B and shown as mean±SD. Statistics: unpaired t-test. ns: not significant.



**Figure S4. Cytotoxicity against MSC varies amongst PBMC donor but is independent on the percentage of CD8<sup>+</sup> or CD56<sup>+</sup> in GvHD patients.** **A:** PBMC obtained from 2 different GvHD patients (Patient 1 and Patient 2) were tested for their cytotoxic activity against MSC from two different donors (MSC1 and MSC2). **B:** apoptosis in MSC obtained from different donors (MSC1, MSC2 and MSC3) after incubation with PBMC from four different MLR responder/stimulator combinations (MLR1, MLR2, MLR3, MLR4). In A and B the level of apoptosis was assessed by flow-cytometry after 4 hours of co-culture. **C, D:** PBMC obtained from 11 GvHD patients (R: 3, NR: 8) were analysed for the percentage of CD8<sup>+</sup> (**C**) and CD56<sup>+</sup> (**D**) cells. Statistics: unpaired t-test. ns: not significant.



**Figure S5. MSC killing is mediated by caspase 3 and effected by GrB and perforin.** **A:** PHA-aPBMC were incubated with MSC at escalating PBMC/MSC ratios. MSC apoptosis was assessed by annexin V/7-AAD at different time-points by flow-cytometry. Results represent the mean $\pm$ SD of 3 independent experiments. **B, C:** MSC were transfected with the pECFP-DEVDR-Venus vector (FRET-MSC) and the Förster Resonance Energy Transfer (FRET) between pECFP and Venus-YFP FRET was studied by flow-cytometry and Caspase activity ( $CA_t$ ) calculated. FRET-MSC were cultured alone, with PHA-aPBMC, or PHAaPBMC in the presence of Z-VAD-FMK (50  $\mu$ M) (**B**), GrB inhibitor Z-AAD-CMK (300  $\mu$ M) or the perforin inhibitor EGTA (4 mM) (**C**). Results of 5 (**B**) or 3 (**C**) independent experiments are shown. When PBMC were present, the PBMC:MSC ratio was 40/1. Statistics: one-way ANOVA and Tukey's Multiple Comparison test. \*\*:  $p > .01$ . \*\*\*:  $p > .001$ . ns: not significant. **D:** MLR-aPBMC were cultivated with MSC (20/1 ratio) and apoptosis evaluated by flow-cytometry 4 hours later. Where indicated, the TNF- $\alpha$  inhibitor Etanercept or the mAb antiTRAIL were used at 10  $\mu$ g/ml or 100  $\mu$ g/ml. Results represents the mean $\pm$ SD of 3 independent experiments.



**Figure S6. Infused MSC can be imaged in the lungs of mice with Th2-type lung inflammation.** **A:** Balb/C mice were immunized i.p. with OVA at day 0 and 7 and subsequently challenged with OVA through aerosol at days 14, 15 and 16 (OVA group). Experimental group was treated with MSC one hour after the last challenge (OVA+MSC). When luc-MSCs were used, mice were imaged one hour after infusion for the analysis of caspase 3 activation after i.p. injection of DEVD-aminoluciferin. After 18 hours from treatment, eosinophils infiltration in BAL was evaluated. **B-E:** Percentage (**B, D**) and absolute numbers (**C, E**) of different cellular types in the BAL (**B, C**) and lungs (**D, E**) of naïve (with bars) ( $N=3$ ) and OVA-sensitized (black bars) ( $N=3$ ) mice. Results represent the mean $\pm$ SD of 3 independent experiments. In OVA-sensitized mice, the analysis was performed 1 hour after the last aerosol challenge. **F:** in order to confirm the presence of luc-MSCs in the lungs of all groups of mice infused with MSC, the same mice imaged in Figure 6A were injected with D-Luciferin. White lines separate multiple photographs assembled in the final image. **G:** TLS was measured from the images of mice in Fig. S6F and shown as mean $\pm$ SD. Statistics: unpaired t-test. ns: not significant.

**Supplementary Table.**

**Table S1. Clinical features of GvHD patients**

<b>Diagnosis</b>	<b>Donor type</b>	<b>GvHD</b>	<b>Grade</b>	<b>Organs involved</b>	<b>Concomitant therapy for GvHD</b>	<b>MSC Dose (x10<sup>6</sup>/Kg)</b>	<b>Response</b>
DLBCL	SIB	Late onset	3	skin, liver	Steroid, MMF, Infliximab	1.6	NR
CLL	SIB	Late onset	4	gut	Steroid, MMF, Infliximab, Alemtuzumab	2.8	R
HL	VUD	Acute	3	skin, Gut	Steroid	3.0†	NR†
						7.4‡	R‡
CML	VUD	Late onset*	3	Skin, Liver	MMF	3	NR
AML	VUD	Chronic*	N/A	Skin	Steroid, CSA	2.7	NR
AML	SIB	Acute	3	Gut	Steroid, CSA	2.1	R
CML	VUD	Acute*	4	Gut	Steroid, CSA	2.9	R
AML	VUD	Acute	4	Skin, gut	Steroid, CSA	3.1	NR
FL	SIB	Acute	4	Skin, gut, Liver	Steroid, CSA, MMF	1.6	R
MM	SIB	Acute	4	Gut	Steroid, Infliximab	2.1	NR
AML	VUD	Acute	4	Gut	Steroids, Budenofalk, CsA	1.28	NR
pre-B ALL	UUD	Acute	4	Skin	Steroids, Topic glucocorticoids	1.03	NR
MDS/RAEB-2	VUD	Acute	4	Gut	Steroids, Tacrolimus,MMF, Etanercept, Ruxolitinib, MTX, Alemtuzumab, CsA	1.55	NR
Mixed AML/T-ALL	VUD	Acute	4	Gut	Steroids, CSA	1.33	NR
MM	VUD	Acute	3	Gut	Steroids, CSA	1.01	NR
B-ALL, BCRABL <sup>+</sup>	SIB	Acute	3	Skin, Liver	Steroids, ECP, CsA	1.11	NR

†: first dose

‡: second dose

\*: GvHD post-Donor Lymphocyte Infusion

AML: Acute Myeloid Leukemia; CML: Chronic Myeloid Leukemia; CLL: Chronic Lymphocytic Leukemia; CSA: Cyclosporine; DLBCL: Diffuse Large B-Cell Lymphoma; FL: Follicular Lymphoma; HL: Hodgkin Lymphoma; NR: no response; MM: Multiple Myeloma; MMF: Mycophenolate; R: response; SIB: HLA-identical sibling; VUD: Volunteer Unrelated Donor.



## Supplementary Video Legends

**Video S1. Living cell imaging of FRET-MSC plated alone.** MSC were transfected with the pECFP-DEVDR-Venus vector (FRET-MSC) and caspase 3 activation studied through the analysis of the FRET between pECFP and Venus-YFP. Living cell imaging was acquired every 3 minutes for 180 minutes using a Leica TCS-SP5 II Confocal Microscope, with 488 nm and 407 nm lasers. The images were processed and analyzed by using the software “R” and EBIImage package. Red and blue colors correspond to high (high caspase 3 activity) or low (low caspase 3 activity) ECP/FRET ratios, respectively.

**Video S2. Living cell imaging of FRET-MSC plated with PHA-aPBMC.** As in Video S1 but with FRET-MSC plated with PHA-aPBMC at a PBMC:MSC ratio 40/1.

**Video S3. Living cell imaging of FRET-MSC plated with resting PBMC.** As in Video S1 but with FRET-MSC plated with resting PBMC at a PBMC:MSC ratio of 40/1.

**Video S4. Living cell imaging of FRET-MSC plated with PHA-aPBMC in the presence of the pan-caspase inhibitor Z-VAD-FMK.** As in Video S2 but in the presence of the pan-caspase inhibitor Z-VADFMK (50  $\mu$ M).

**Video S5. Living cell imaging of FRET-MSC plated with PHA-aPBMC in the presence of the GrB inhibitor Z-AAD-CMK.** As in Video S2 but in the presence of the GrB inhibitor Z-AAD-CMK (300  $\mu$ M).

**Video S6. Living cell imaging of FRET-MSC plated with PHA-aPBMC in the presence of the perforin inhibitor EGTA.** As in Video S2 but in the presence of the Perforin inhibitor EGTA (4 mM).