



King's Research Portal

Document Version
Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Cho, I., Lui, P., & Ali, N. (2020). Treg regulation of epithelial stem cell lineage. *Journal of Immunology and Regenerative Medicine*.

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Treg Regulation of the Epithelial Stem Cell Lineage

Inchul Cho¹, Prudence Lui¹, Niwa Ali^{1,2}

¹Centre for Stem Cells and Regenerative Medicine, School of Basic and Biomedical Sciences,
King's College London, London, UK

²The Francis Crick Institute, London, UK

Corresponding author: Niwa Ali

Niwa.Ali@kcl.ac.uk,

Phone number: +44 207 188 7188,

Summary

Tissue repair and maintenance in adult organisms is dependent on the interactions between stem cells (SCs) and constituent cells of their microenvironment, or niche. Recent and accumulating evidence suggests that immune cells, specifically Foxp3⁺ CD4⁺ Regulatory T cells (Tregs), play an important role as a component of the SC niche. Undisputedly, Tregs are the major immunosuppressive lineage of the CD4⁺ T cell compartment, and reside within numerous secondary lymphoid organs, where they exert their functions. These cells are also specialised in facilitating protective functions specific to their tissue of residence. In this review, we discuss the emerging concepts supporting the SC-regulatory functions of tissue-resident Tregs, during both the steady-state and SC-mediated regeneration. We highlight the skin, intestines, and lung as model organs which are subject to recurrent microinjury and exposure to microbiota. An in-depth understanding of the biology of the Treg-SC axis will inform ongoing immunotherapeutic endeavours to target specific subpopulations of tissue-resident Tregs.

Introduction

Successful tissue repair is largely dependent upon the efficient self-renewal and plasticity of stem cell (SC) populations that differentiate towards multiple cellular lineages. Some SC populations, including those present in the hematopoietic system, intestinal epithelium, or epidermis, constitutively regenerate tissue throughout the life-span of an organism. Long-term maintenance of a healthy tissue requires a finely tuned coordination between stem cells and constituent cells within the niche, while exempting exogenous threats, such as pathogens and toxins.

The epithelial barrier constitutes the first line of defence against external physical and chemical injury. The relationship between tissue maintenance and protection from external insults are well-illustrated during wound healing responses upon epithelial barrier breach. Initial injury invokes the recruitment and/or local activation of tissue-resident immune cells (TRICs) to sites of damage. This early immune response serves to protect the tissue against invading microorganisms, and to clear damaged cells or cellular debris. The removal of damaged cells provides the spatial cue(s) necessary to induce epithelial SC proliferation and differentiation, thus replenishing the epithelium. The benefits and negative effects of immune inflammation on SC activation has been demonstrated by several previous studies, which have been reviewed^{1,2}. However, only a limited number of studies demonstrate direct immune cell regulation of epithelial SC activity.

Tissue-specific functions of immune cells, in particular, regulatory T cells (Tregs), have been documented in multiple non-lymphoid tissues, such as muscle and adipose tissue³. Historically, research elucidating the existence of a direct immune cell-SC axis has been largely underexplored. This is despite the notion that TRIC activity is intimately associated with SC function, as observed during regenerative responses⁴. Instead, the mechanisms influencing SC activity have been extensively studied in light of the surrounding epithelial cells and other stromal cells, which were largely driven by popularisation of the SC niche as a regulatory mechanism. The idea of the SC niche posits that SCs are regulated by cells and extracellular matrices directly within their microenvironment. Hence, immune cells, both resident and migratory in nature, have not been fully explored in this context.

Recent evidence suggests that multiple immune cell populations can directly interact with SCs to modulate their behaviour. Of which, Tregs are a prominent immune cell subset that reside in numerous peripheral tissues, where they are heavily implicated in SC regulation. In this review, we highlight recent evidence that supports the role of tissue-resident Tregs, not only as sentinels of the immune response, but as constituents of the epithelial SC niche. This is primarily exemplified in model organs that are subject to recurrent microinjury and exposure to microbiota, such as the intestines, lungs and the hair follicles (HFs) of skin.

The Treg Lineage

Maintenance of healthy tissues requires the immune system to distinguish between self and non-self. In several organs, such as the skin, lungs and the intestines, where microorganisms thrive, it is important to regulate over-active immune responses against self and commensal microorganisms. As such, there are several distinct subsets of TRICs that exert immunosuppressive effects, such as tolerogenic dendritic cells, innate lymphoid cells (ILCs), and Tregs. The conventional immune functions of these cell types have been previously reviewed⁵⁻⁷.

Regulatory T cells constitute a subset of CD4⁺ T cells that express the lineage defining transcription factor forkhead box protein 3 (Foxp3). The majority of thymic Tregs (tTregs) develop in the thymus during thymocyte differentiation into mature T cells; whereas peripheral Tregs (pTregs) develop from naïve T cells in secondary lymphoid organs. The pTregs are then seeded into non-lymphoid organs, where they encounter tissue-specific antigens. Both within peripheral organs and in secondary lymphoid organs, Tregs suppress effector immune responses directed against self-antigens. This process is referred to as ‘immune tolerance’.

In humans, mutations in Foxp3 manifests in an autoimmune disease, termed immunodysregulation poly-endocrinopathy, X linked, or IPEX⁸. The study of this disease, or the function of Foxp3-expressing Tregs have been facilitated by the ‘scurfy’ mouse line, which harbour an insertional mutation at the Foxp3 locus. The resultant premature stop codon leads to the development of spontaneous autoimmunity⁹. Follow up studies in mice further revealed specific expression of Foxp3 in CD4⁺ CD25⁺ peripheral Tregs¹⁰, and that deletion of *Foxp3* phenocopies *Foxp3* mutant scurfy mice. It is now widely accepted that the high-affinity IL-2 receptor alpha chain, CD25 and the transcription factor Foxp3 are critical factors for Treg development, maintenance, and function¹⁰⁻¹².

The delicate balance between immunosuppressive Tregs and effector T cells (Teffs) is crucial for the regulation of inflammation and self-tolerance. An imbalance in this dynamic, for instance, as a result of Treg ablation or dysregulation, underlies the pathology of numerous autoimmune diseases in both mice and humans¹³. Moreover, genome-wide association studies have attributed autoimmune disease susceptibility in humans to impaired Treg function¹⁴. Whilst functional Tregs are important for maintaining tissue homeostasis, the pathways they utilise may be hijacked by cancer cells to modulate the host immune system and dampen anti-tumour responses. As such, inhibition of Tregs and/or downstream pathways have emerged as promising candidates to promote anti-tumour immune responses¹⁵. However, the current understanding of the diversity and molecular mechanisms that dictate tissue-specialised functions of resident Tregs remains incomplete. Establishing a solid foundational knowledge of the TRIC-SC axis will be crucial to fully elucidate the biology of auto-inflammatory and tissue-regenerative diseases.

Tregs and the Skin Stem Cell Niche

The skin constitutes the primary defence system against external insults and is the first organ to encounter pathogenic microbes. To facilitate defence against foreign bodies, the skin is equipped with a diverse range of active TRICs that communicate with local epithelia, ensuring robust immune responses upon the breach of the skin’s barrier function. Amongst the numerous TRIC in the skin, Tregs account for 20-60% of CD4⁺ T cells in the mouse skin^{16,17}. Similarly, Tregs in human skin comprise 20% of total CD4⁺ T cells, as opposed to 5% in the peripheral circulation¹⁸.

It has recently been shown that an abrupt wave of Tregs are seeded into the skin early during postnatal development between postnatal day 6 to day 13, likely originating from the thymus¹⁷. This Treg migratory event is facilitated by the expression of the chemokine CCL20 by the developing epithelial cells of the HF, attracting Tregs that express the chemokine receptor CCR6¹⁹. Intriguingly, mice grown under germ-free conditions do not accumulate Tregs in this manner, suggesting that the process is dependent on the presence of microbiota, and that Tregs are involved in the regulation of immune homeostasis in the skin¹⁹.

Anatomically, the skin, or the epidermis, can be subdivided into the interfollicular epidermis (IFE), which forms the protective barrier, and the HFs, or the pilosebaceous unit, whose epithelium is contiguous with the IFE. The pilosebaceous unit is composed of distinct stem cell compartments, which can be identified by the expression of their respective molecular markers: Blimp1 (sebaceous gland stem cells); Gli1 and Lgr6 (lower isthmus stem cells); and integrin alpha-6 (ITGA6), Lgr5, CD34 and Krt15 (hair follicle stem cells (HFSCs)). It is speculated that these stem cells are restricted to their own lineage boundaries based on fate-mapping experiments^{20–23}. The HFSCs are located in the lower portion of the epithelia, anatomically referred to as the ‘bulge’ region. These cells are responsible for mediating perpetual bouts of growth arrest (telogen) and activation (anagen) during the hair follicle (HF) cycle to form a new hair shaft²⁴. The HFSCs depend on signalling ligands, such as BMP, TGF- β and Wnt inhibitors to maintain quiescence^{25–28}; whereas the transition into a proliferative status requires activation of Wnt signalling^{28,29}. The differentiation trajectories of HFSCs are regulated by microenvironmental signals, which includes Notch signalling³⁰.

Histological studies of human skin reveal that the majority of CD4⁺/Foxp3⁺ Tregs are preferentially associated with HFs, and rarely the interfollicular dermis or within the epidermis itself^{16–19,31}. Flow cytometric analyses revealed higher number of Tregs in the scalp and the face, which contain more HFs than other body sites. This is further supported by intravital multiphoton imaging of healthy mouse flank skin, which showed bulge-associated Tregs displayed more dynamic activity than Tregs localised to other skin niches^{16,31}. Together, they present evidence that both human and murine skin Tregs are concentrated in the vicinity of follicular epithelium where HFSCs reside^{18,32}.

The spatial proximity between HFSCs and Tregs suggest a possibility that the two cell types may engage in regulatory interactions. In line with this, genome-wide association studies have linked single nucleotide polymorphisms in Treg-associated genes, including Foxp3, to alopecia areata, a disorder of immune-mediated disruption of HF regeneration^{33,34}. Importantly, augmenting Tregs clinically has shown efficacy in treating patients with this disease³⁵.

Curiously, the number of Foxp3⁺ Tregs declines during the growth phase of the HF cycle, anagen, and increases during the latter stages of the growth arrest, telogen (Figure 1). Specific ablation of murine Tregs during telogen leads to impairment of the telogen-to-anagen transition, and consequently impacts hair regeneration. The impairment of hair regeneration was attributed to a reduced proliferation and differentiation capacity of CD34⁺ ITGA-6⁺ HFSCs¹⁶. Transcriptomic profiling further revealed a deficient Notch target gene expression profile in HFSCs in the absence of Tregs, and preferential expression of the Notch ligand, Jagged-1 (Jag1), in skin resident Tregs. Conditional deletion of Jag1 in Tregs using Foxp3^{YFP-Cre}/Jag1^{fl/fl} mice largely phenocopied depletion of all skin Tregs, suggesting that the HFSC-regulatory role of Tregs is mediated via Jag1¹⁶. It was further demonstrated using 2-photon intravital microscopy that Tregs reside within 5 μ m of the HF epithelium, suggesting that a subpopulation of Tregs may directly interact with HFSCs.




Figure 1. Major functions of skin-resident Tregs in the regulation of epithelial SCs. Tregs are highly abundant during the quiescent phase of the HF cycle, telogen, relative to the growth phase, anagen. Depletion of Tregs in Foxp3-DTR mice impairs HFSC activation and transition into anagen. A Treg-specific deletion of a notch ligand, Jagged1, phenocopies Treg depletion, suggesting that Tregs regulate HFSCs in a Jagged1-dependent manner. Upon epidermal injury, HFSCs are recruited to sites of wounding, where they contribute to epidermal regeneration. However, under inflammatory conditions, neutrophils inhibit the migration of HFSCs to sites of injury within the epidermis. Tregs facilitate the regenerative potential of HFSCs by inhibiting CXCL5-mediated activation of neutrophils, which would otherwise inhibit HFSC migration. Red arrows indicate inflammatory response. Green arrows indicate regenerative responses.

After breach of the epithelial barrier upon full thickness wounding, Tregs rapidly accumulate in skin, and display a highly activated phenotype, as indicated by upregulation of Treg-associated activation markers - CD25⁺/CTLA-4⁺/ICOS⁺. Specific ablation of these cells in Foxp3-DTR mice via diphtheria toxin (DT) administration early post-wounding attenuates wound closure and re-epithelialization. The impairment of regeneration was attributed to an interferon-gamma dependent accumulation of Ly-6C^{high} pro-inflammatory macrophages (CD45⁺CD11b^{high}F4/80⁺Ly-6C^{high}Ly-6G^{low}CD206^{low}), in Treg-deficient mice³⁶. It was further discovered that infiltrating Tregs express the epidermal growth factor receptor (EGFR). Conditional deletion of EGFR in Tregs, using Foxp3^{YFP-Cre}/EGFR^{fl/fl} mice, delayed wound closure and led to an increased accumulation of pro-inflammatory macrophages³⁷. These results suggest that the activation of Tregs upon injury may be dependent on EGF signalling. Cutaneous Tregs appear to co-opt a highly conserved regenerative pathway, often used by SCs as signals to enter the cell cycle, to mediate this specialised function of tissue repair. While these studies have demonstrated an active role for cutaneous Tregs during wound healing, whether Tregs directly modulate epithelial SC function, post full thickness wounding of skin, remains to be elucidated.

Indeed, a recent study has demonstrated an indirect role for Tregs in epidermal repair using *Foxp3-DTR* and *Lgr5-TdTom* mice. In this model, Tregs can be specifically ablated and *Lgr5*⁺ HFSCs simultaneously lineage-traced³⁸. Under conditions of epidermal injury as a result of repetitive tape stripping, injured epidermis secretes CXCL5, which causes an increase in neutrophilic infiltrate at sites of injury³⁸. Depletion of Tregs in this setting abrogates the regenerative impairment and is associated with neutrophilic infiltration and increased CXCL5. Recapitulating Treg-mediated repression of CXCL5 via neutralisation of CXCL5, partially rescued the impaired epidermal regeneration in Treg-depleted mice. These experiments indicate that Tregs control immune cell recruitment to regulate the inflammatory status of the skin. This, in turn, affects the migratory capacity of HFSCs and their ability to contribute to epidermal regeneration³⁸.

These studies reveal that skin Tregs play a critical role in both epithelial and HFSC biology. Additionally, these findings provide a mechanistic link between tissue-resident Tregs and epithelial SCs that is necessary for adequate tissue function.

Tregs and the Intestinal Stem Cell Niche

In addition to the skin, the gut is under constant insult from chemicals and foreign microbes within the digestive tract. Hence, like the skin, the gut hosts a diverse repertoire of TRICs to ensure a robust response to injury. However, recurrent inflammation exposes the gut to an increased frequency of reactions directed against self-antigens. As such, the gut also hosts a large number of Tregs, perhaps in order to mitigate the anti-self response. In line with this, Treg numbers and functionality are significantly decreased in mouse models of inflammatory bowel disease^{39,40}. In human patients suffering from inflammatory bowel disease, circulating Treg numbers are decreased, whereas those that infiltrate the inflamed tissue increases. Moreover, reduction in circulating Treg numbers correlates with disease severity⁴¹⁻⁴³.

Amongst murine intestinal resident CD4⁺ T cells, Tregs account for 25-35% in the colon⁴⁴. The number of colonic Tregs are significantly reduced in germ-free mice, suggesting their recruitment is dependent on chemokines that are secreted in the presence of microbiota⁴⁵. Numerous subsets of Tregs exist in the intestines that express transcription factors associated with T cell lineage differentiation, that in turn dictates the distinct nature of the immune response to be elicited.

Gut-resident Tregs can be subdivided into 1) GATA3⁺ Helios⁺ 2) ROR γ t⁺ and 3) ROR γ t⁻ Helios⁻ fractions (Figure 2). The transcription factor GATA3 drives T helper 2 differentiation, but may also associate with the *Foxp3* genomic locus, thus reinforcing *Foxp3* expression to direct the Treg fate⁴⁶. Supporting this notion, Treg-specific deletion of GATA3 leads to destabilisation of the Treg phenotype under inflammatory conditions⁴⁶. Helios expression in Tregs was initially believed to definitively indicate thymic origin^{47,48}. However, this is challenged by a finding that the number of Helios⁺ Tregs increase in peripheral organs *in vivo* after stimulation with an antigenic peptide⁴⁹. Lastly, retinoic acid orphan receptor γ t (ROR γ t) is the master regulator of Th17 cell differentiation⁵⁰. Overexpression of the RORC gene promotes expression of the main effector cytokine IL-17⁵¹. Recent and emerging evidence suggests that Tregs acquire the lineage markers of T effector subsets that they regulate. For example, Tregs expressing the type 2 lineage marker, *Gata3*, preferentially regulate type 2 immune responses⁵².




Figure 2. Major functions of intestinal-resident Tregs in the regulation of epithelial SCs. Several subpopulations of Tregs have been reported in the intestines with distinct roles in mediating tolerance towards dietary antigens and microbiota. Namely, ROR γ t⁻ Tregs appear to be responsible for mediating dietary tolerance; ROR γ t⁺ Tregs mediate microbial tolerance; whereas Gata3⁺ Tregs respond to the epithelial-derived interleukin-33, suggesting a possibility that they mediate epithelial-immune crosstalk. Indeed, IL-33 induces the production of anti-inflammatory cytokines TGF β and IL-10 in Tregs. Finally, naïve T cells differentiated towards a Treg-like phenotype (iTreg, induced Tregs) by administration of TGF β , support ISC growth *in vitro* by secretion of IL-10. Brown arrows indicate findings from *in vitro* studies.

The three subtypes of Tregs play distinct roles in the tissue micro-environment. GATA3⁺ Tregs have been shown to respond to interleukin-33 (IL-33) that is constitutively expressed by epithelial cells. Upon injury, IL-33 production is increased, which activates GATA3⁺ Tregs to further produce IL-10 and tumour growth factor-beta (TGF- β), both of which are anti-inflammatory⁵³. Whether these GATA3⁺ Tregs also express Helios, however, has not been studied. ROR γ t⁺ Tregs are responsible for tolerance to microbiota within the lumen⁵⁴.

Tolerance is induced by the action of the transcription factor cMAF, which acts to dampen Th17 differentiation in colonic T cells⁵⁵. The last sub-type, ROR γ ^T Helios⁻ Tregs are less well defined functionally. Antibiotic treatment of mice does not affect their population size, whilst mice on antigen-free diets have reduced numbers of ROR γ ^T Helios⁻ Tregs, indicating a possible function in mediating tolerance against dietary antigens⁵⁶.

Stem cell regulation mediated by Tregs has also been demonstrated in the intestines by a recent finding by Biton *et al.*⁵⁷. The authors analysed intestinal epithelial cells via single cell RNA-sequencing (scRNA-seq) and identified three clusters of ISCs (ISC-I/II/III) that all express the SC marker, Lgr5. Amongst these clusters, ISC-I cluster consisted of quiescent ISCs that expressed low levels of MHC-II, as opposed to the proliferative ISC-II/III clusters. The differences in MHC-II expression amongst these clusters suggest possible differences in their interactions with local immune cells. As such, the authors assessed possible interactions between ISCs and distinct T cell types using an *in vitro* organoid co-culture. In this setting, they observed an enhanced intestinal organoid renewal when co-cultured with induced-Tregs (T cells cultured *in vitro* in the presence of TGF- β) or IL-10, the main effector cytokine of Tregs. However, depletion of Tregs in Foxp3-DTR mice resulted in increased ISC proliferation and differentiation into mature intestinal cells, such as tuft cells and goblet cells. The discrepancy between findings from *in vitro* and *in vivo* experiments may arise from several factors. Namely, the Tregs used for co-culture were derived from naïve T cells differentiated towards the Treg lineage via *in vitro* stimulation with TGF- β . In order to reconcile the different phenotypes, it will be necessary to fully characterise the cultured Tregs and determine which *in vivo* subtypes they most closely resemble. Moreover, there are several drawbacks of the method used in this study to quantify ISCs. The number of ISCs were indirectly assessed by quantification of Lgr5 transcripts, rather than total cell numbers. Taking this into account, the *in situ* immunofluorescence analyses did not reveal significant changes in the population size, or the number, of ISCs. Hence, a definitive functional role for Tregs in the direct modulation of ISC numbers remains an open question.

Other than Treg-mediated regulation of SC activity, Biton *et al.* demonstrated that local TRICS, other than Tregs, also possess the capacity to facilitate ISC differentiation. Specifically, antimicrobial Th1 responses direct ISC towards Paneth cells, which produce antimicrobial peptides. In contrast, the Th2 cells, which directs anti-parasitic responses, induces differentiation into tuft cells. These specialised cells possess the potential to differentiate into goblet cells, producing mucus directed to parasite eradication⁵⁷. Overall, these findings suggest an intriguing possibility that immune cells are able to utilise SC differentiation to enhance host defence. This was suggested by a previous study⁵⁸, which utilised ROR- α ^{sg/sg} mice in which ROR α is rendered non-functional via the introduction of a premature stop codon⁵⁹. This study reported that type 2 innate lymphoid cells (ILC2) fail to develop under stimulatory conditions in the presence of IL-25, a cytokine that expands and sustains ILC2s⁶⁰. Using this experimental approach, the authors assessed the importance of functional anti-parasitic ROR α ⁺ ILC2s and their impact on mucus secreting goblet cells. The lack of functional ILC2s was associated with a marked decrease in the absolute number of goblet cells⁵⁸. These findings are consistent with previous studies that report a reduction in the level of type 2 cytokines in ROR α ^{sg/sg} mice⁶¹. Importantly, these findings shed light on the pathology of pathological conditions such as asthma, which has been associated with ROR α mutations in a GWAS study⁶². A recent *in vivo* finding demonstrated that upon infection, tuft cells constitutively express IL-25 to sustain and activate ILC2s, which subsequently secrete IL-13 to promote goblet cell differentiation⁶⁰. Given Foxp3 expression and Treg suppressive functions are enhanced by IL-25, it would be of interest to investigate whether Tregs are implicated in this immune-mediated differentiation⁶³.

Overall, a direct or even indirect role, of Tregs in modulating ISCs is yet to be fully elucidated. An interesting avenue for further research would be to study how Tregs impact the classical signalling pathways that are known for the maintenance of the ISC niche. For instance, the activation of Wnt signalling pathway is sufficient to induce ISC proliferation^{64,65}, whereas genetic deletion of Wnt effector genes, such as β -catenin, results in loss the intestinal crypt⁶⁶. Intriguingly, it has recently been shown that Tregs express the membrane-bound Wnt antagonist, Dkk1⁶⁷. Whether tissue-resident Tregs also express Dkk1 is yet to be determined, however. Nonetheless, it raises an intriguing possibility that Dkk1-expressing Tregs, either resident or circulatory, may impact the Wnt signalling pathway in ISC clusters during homeostasis or injury-induced differentiation⁶⁸

Tregs and the Lung Stem Cell Niche

As an integral part of the circulatory system, the most critical functions of the lungs are to expel respiratory waste products and to take in oxygen from the atmosphere. Consequently, the luminal surface of the lungs is constantly exposed to air-borne foreign particles. These include the influenza virus and particulate matters, which invoke immune cell responses. Therefore, the lungs are also home to a diverse repertoire of TRICs^{69,70}.

The lungs can be largely divided into three anatomically distinct segments; large airway, small airway, and the alveolus. Distinct SC lineages are known to exist along the epithelium, with the most multipotent basal cells populating the large airway, club cells lining the small airway, and unipotent type 2 alveolar epithelial cells (AEC2) in the alveolus. The presence of bronchoalveolar stem cells have been demonstrated in the mouse bronchoalveolar duct junction, but not in humans⁷¹.

Perhaps owing to the presence of diverse SCs dedicated to maintaining their respective anatomical compartments, the lungs have remarkable regenerative capacity. Upon removal of one lobe of the lungs, the remaining lobes undergo compensatory growth to maintain the same level of respiratory volume⁷². However, in end-stage lung diseases, such as chronic obstructive pulmonary disease and idiopathic pulmonary fibrosis (IPF), regenerative capacity is impaired. For instance, in IPF, AEC2s form bronchiolar-like structures rather than differentiating into type 1 alveolar epithelial cells (AEC1s) that line the alveolar epithelium⁷³.

The pathological mechanisms of IPF are unclear. The current consensus posits that repeated insults overwhelm the regenerative potential of the AEC2s. Subsequent mobilisation of tissue-resident cells such as fibroblasts to aid restoration of barrier function leads to irreversible remodelling of the extracellular matrices. Such changes within the microenvironment is likely to impact the behaviour of resident cells, including epithelial SCs, fibroblasts and immune cells, which remain incompletely understood.

The role of immune cells in the repair of lung epithelium upon injury has been previously demonstrated. For instance, depletion of tissue-resident ILC2s impaired regeneration post-influenza infection⁷⁴. Gene expression profiling of ILC2s revealed an abundant expression of amphiregulin, which alone was sufficient to restore barrier function in ILC2-depleted mice⁷⁴. Similarly, Tregs in the lungs have been shown to express amphiregulin, which was necessary for the re-establishment of barrier function in the lungs post-influenza infection⁷⁵. However, in both cases, whether barrier function was restored through extracellular matrix remodelling, or direct modulation of lung epithelial SCs has not been assessed. Nonetheless, these findings from mouse studies may be relevant to humans as amphiregulin-expressing Tregs have also

been found in peripheral organs, such as the lungs, where they also exert pro-regenerative functions ⁷⁶.

It is currently unclear whether Tregs exert their regenerative functions in the lungs by acting on the stem cells, or the fibroblasts, to restore barrier functions in the lungs (Figure 3). Additionally, the role of Tregs during degenerative processes, such as pulmonary fibrosis, remains controversial^{77,78}. Given that recurrent activation of fibroblasts underlie the pathology of idiopathic pulmonary fibrosis, it is of critical importance to fully characterise the target cells of Tregs, for an effective regenerative therapy that is without harmful side-effects, such as fibrosis.




Figure 3. Major functions of lung-resident Tregs in the regulation of epithelial SCs. In the lungs, the alveolar epithelium is lined by AEC1s, which differentiate from AEC2s. The maintenance of AEC2s is partially regulated by Notch signalling activities. As such, it presents a possibility that Jagged1-expressing Tregs regulate AEC2 differentiation. A Treg-secreted molecule, amphiregulin, has been associated with restoration of barrier function in the lungs. However, whether Tregs are the primary source of amphiregulins in the lungs, and whether the functional recovery of the lungs were achieved as a result of epithelial stem cell proliferation, or deposition of extracellular matrix proteins, is incompletely addressed. Studies thus far suggest a dual role for amphiregulin. Amphiregulin has been shown to promote the differentiation of fibroblasts into myofibroblasts, which are specialised in remodelling the extracellular matrix. Additionally, recombinant amphiregulin protects AEC2s from apoptosis. Brown arrows indicate findings from *in vitro* studies.

Studies reported thus far indicate an essential role for immune cells during pulmonary repair, although the precise mechanisms by which the repair is mediated has not taken a direct TRIC-SC perspective. The role of amphiregulin has been discussed in this regard, which is secreted by Tregs and is critical for the recovery of barrier function. However, to our knowledge, a direct role for Tregs, or amphiregulin, in the induction of AEC proliferation is not established. There is also a lack of consensus on the role of amphiregulin in different modes of pulmonary injury.

Outlook for the Treg-stem cell Axis in Tissues

The widely held view that the dominant and sole function of the tissue immune system is protection against external insults is beginning to change. In particular, tissue-resident Foxp3⁺ Tregs possess numerous tissue-specific functions, outside of their conventional immunoregulatory role. In this review, we have discussed the SC-regulatory role of Tregs, evidence for which is beginning to accumulate. For instance, in the muscles, Tregs secrete the growth factor, amphiregulin to promote muscle satellite cell proliferation upon injury⁷⁹. Genetic ablation of Tregs has also been reported to result in impairment of pulmonary regeneration in multiple models of injury⁸⁰. We have also discussed SC-regulatory functions of Tregs in the skin, which is supported by conclusive evidence from Treg-specific deletion of a single factor in a model of SC-mediated regeneration. Jagged1-expressing Tregs facilitate HFSC proliferation and differentiation during the telogen-to-anagen transition of the HF cycle. The close proximity between skin Tregs and HFSCs would suggest either a direct interaction between Tregs and HFSCs or an indirect regulatory mechanism involving another proximal tissue cell type occurs. Indeed, the absence of Gata3⁺ Tregs in the skin leads to aberrant fibroblast activation and ECM remodelling, suggesting that there are additional mechanisms via which Tregs may regulate the HFSC microenvironment⁵².

Despite the presence of a diverse repertoire of TRICs in the lung, little is known of their influence on lung regeneration. Additionally, the lungs also harbour haematopoietic progenitors⁸¹, inciting the notion that *de novo* generation of immune cells may occur to sustain immunological demands via processes that are regulated by TRICs, including Tregs. Crucially, we do not yet understand whether there are changes in processes such as haematopoietic activity and the interactions between TRICs and SCs upon injury, or in diseased states.

In light of pulmonary regeneration, it will be critical to understand the interactions between TRIC and SCs and the signalling mechanisms they employ. For example, resident alveolar macrophages (rAMs), a yolk-sac derived, self-renewing population of macrophages interacts with the airway epithelium. Cell-to-cell communication during the steady state and rAM-derived soluble mediators (e.g. CD200/R; IL-10, TGF- β , GM-CSF) maintain homeostasis and quiescence⁸². Loss of rAMs during injury may release these suppressive signals on AEC proliferation. In addition to these TRICs, others like CD103⁺ dendritic cells, intra-epithelial CD103⁺ memory T cells, interstitial macrophages (IMs), mucosa-associated invariant T cells (MAIT), invariant natural killer T cells (iNKT) cells and $\gamma\delta$ -T cells also reside either within the airway epithelium or in the mesenchyme surrounding the AEC. However, their contribution to AEC regeneration is unknown. Given that AEC2s rely on Notch signalling for differentiation into AEC1s⁸³, it is interesting to speculate that Jag1-expressing Tregs may be involved in alveolar differentiation. Currently, a role for alveolar Tregs in AEC2 maintenance or proliferation post-injury has not been identified. Given the recent demonstration that a distinct subset of cutaneous Tregs are poised for the regulation of fibroblasts, and therefore, fibrosis⁵², it may be functionally relevant to fully characterise alveolar resident Tregs. It is possible a subset of Tregs in this location can influence fibroblast behaviour, and/or epithelial SC homeostasis, in a manner similar to that of skin. Such identifications will allow for subsequent therapies that ameliorate fibrosis.

To date, regenerative functions of tissue-resident Tregs have largely been explored in light of amphiregulin secretion and its downstream targets^{75,79,84,85}. However, tissue-specific molecular signatures of Tregs, other than amphiregulin, has been suggested to play a critical role in mediating regeneration of the spinal cord, heart and retina of zebrafish⁸⁶. Indeed, recent reports

form studies of mouse skin suggest that Tregs can regulate regeneration via additional mechanisms, such as Notch signalling pathways, and modulating the activation of other infiltrating immune cells^{16,38}. As such, an interesting avenue for further research would be to fully characterise the diversity of Tregs and their tissue-specific molecular signatures that are vital for the homeostasis of epithelial SCs within individual organs.

Why are immunosuppressive Tregs so intimately associated with the regulation of SC function? One line of relevant evidence is the recent demonstration that quiescent SCs, such as those that reside in HFs, downregulate the expression of MHC-I to evade the immune system. In contrast, fast-cycling SCs begin to express MHC-I and are thereby recognised and targeted for immune-mediated clearance⁸⁷. This appears paradoxical as an immune-mediated attack on SCs would compromise the ability of SCs to re-populate the damaged epithelium. One possibility is that Tregs suppress over-activated immune cell subsets that secrete or promote factors that negatively perturb SC as they enter the cycle. Intriguingly, Treg suppressive phenotypes are reinforced by receiving EGF signalling, such as amphiregulin⁸⁵. This suggests a requirement for a local immunosuppressive environment during the post-injury tissue regenerative phase.

The SC-Treg axis sheds an important and new light on cancer immunology. In this context, Tregs are viewed as suppressors of the immune system that nullify host defence mechanisms⁸⁸. Given the many parallels between the mechanisms by which cancer and SCs are regulated, it is possible that tumour initiating, or progenitor-like SCs, may interact with Tregs in manners similar to that reviewed here, which will be a clinically relevant avenue for further research.

The collective view indicates that Tregs mediate the majority of their functions in their respective tissues of residence. It is therefore likely that strategies targeting the augmentation or suppression of specific Treg subsets will result in optimal therapeutic outcomes. Ultimately, a fundamental understanding of how Tregs impact tissue regeneration, via modulation of SC activity, will be of critical importance in the maintenance and restoration of tissue homeostasis.

Acknowledgements

The authors acknowledge the support of the Wellcome Trust and the Medical Research Council.

References

1. Michael S, Achilleos C, Panayiotou T, Strati K. Inflammation Shapes Stem Cells and Stemness during Infection and Beyond. *Front Cell Dev Biol.* 2016. doi:10.3389/fcell.2016.00118
2. Kizil C, Kyritsis N, Brand M. Effects of inflammation on stem cells: together they strive? *EMBO Rep.* 2015. doi:10.15252/embr.201439702
3. Panduro M, Benoist C, Mathis D. Tissue Tregs. *Annu Rev Immunol.* 2016. doi:10.1146/annurev-immunol-032712-095948
4. Wynn TA, Vannella KM. Macrophages in Tissue Repair, Regeneration, and Fibrosis. *Immunity.* 2016. doi:10.1016/j.immuni.2016.02.015
5. Steinman RM. Decisions about dendritic cells: past, present, and future. *Annu Rev Immunol.* 2012. doi:10.1146/annurev-immunol-100311-102839
6. Mauri C, Menon M. Human regulatory B cells in health and disease: Therapeutic potential. *J Clin Invest.* 2017. doi:10.1172/JCI85113

7. White AM, Wraith DC. Tr1-like T cells - An enigmatic regulatory t cell lineage. *Front Immunol.* 2016. doi:10.3389/fimmu.2016.00355
8. Wildin RS, Ramsdell F, Peake J, et al. X-linked neonatal diabetes mellitus, enteropathy and endocrinopathy syndrome is the human equivalent of mouse scurfy. *Nat Genet.* 2001. doi:10.1038/83707
9. Brunkow ME, Jeffery EW, Hjerrild KA, et al. Disruption of a new forkhead/winged-helix protein, scurfy, results in the fatal lymphoproliferative disorder of the scurfy mouse. *Nat Genet.* 2001. doi:10.1038/83784
10. Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol.* 2003;4(4):330-336. doi:10.1038/ni904
11. Sakaguchi S, Sakaguchi N, Asano M, Itoh M, Toda M. Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J Immunol.* 1995.
12. Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *J Immunol.* 2017. doi:10.1126/science.1079490
13. Bennett CL, Christie J, Ramsdell F, et al. The immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) is caused by mutations of FOXP3. *Nat Genet.* 2001. doi:10.1038/83713
14. Rai E, Wakeland EK. Genetic predisposition to autoimmunity - What have we learned? *Semin Immunol.* 2011. doi:10.1016/j.smim.2011.01.015
15. Tanaka A, Sakaguchi S. Regulatory T cells in cancer immunotherapy. *Cell Res.* 2017. doi:10.1038/cr.2016.151
16. Ali N, Zirak B, Rodriguez RS, et al. Regulatory T Cells in Skin Facilitate Epithelial Stem Cell Differentiation. *Cell.* 2017. doi:10.1016/j.cell.2017.05.002
17. Scharschmidt TC, Vasquez KS, Truong HA, et al. A Wave of Regulatory T Cells into Neonatal Skin Mediates Tolerance to Commensal Microbes. *Immunity.* 2015. doi:10.1016/j.immuni.2015.10.016
18. Rodriguez RS, Pauli ML, Neuhaus IM, et al. Memory regulatory T cells reside in human skin. *J Clin Invest.* 2014. doi:10.1172/JCI72932
19. Scharschmidt TC, Vasquez KS, Pauli ML, et al. Commensal Microbes and Hair Follicle Morphogenesis Coordinately Drive Treg Migration into Neonatal Skin. *Cell Host Microbe.* 2017. doi:10.1016/j.chom.2017.03.001
20. Morris RJ, Liu Y, Marles L, et al. Capturing and profiling adult hair follicle stem cells. *Nat Biotechnol.* 2004. doi:10.1038/nbt950
21. Jaks V, Barker N, Kasper M, et al. Lgr5 marks cycling, yet long-lived, hair follicle stem cells. *Nat Genet.* 2008. doi:10.1038/ng.239
22. Youssef KK, Van Keymeulen A, Lapouge G, et al. Identification of the cell lineage at the origin of basal cell carcinoma. *Nat Cell Biol.* 2010. doi:10.1038/ncb2031
23. Page ME, Lombard P, Ng F, Göttgens B, Jensen KB. The epidermis comprises autonomous compartments maintained by distinct stem cell populations. *Cell Stem Cell.* 2013. doi:10.1016/j.stem.2013.07.010
24. Müller-Röver S, Handjiski B, Van Der Veen C, et al. A comprehensive guide for the accurate classification of murine hair follicles in distinct hair cycle stages. *J Invest Dermatol.* 2001. doi:10.1046/j.0022-202X.2001.01377.x
25. Plikus M V., Baker RE, Chen CC, et al. Self-organizing and stochastic behaviors during the regeneration of hair stem cells. *Science (80-).* 2011. doi:10.1126/science.1201647
26. Hsu YC, Pasolli HA, Fuchs E. Dynamics between stem cells, niche, and progeny in the

- hair follicle. *Cell*. 2011. doi:10.1016/j.cell.2010.11.049
27. Oshimori N, Fuchs E. Paracrine TGF- β signaling counterbalances BMP-mediated repression in hair follicle stem cell activation. *Cell Stem Cell*. 2012. doi:10.1016/j.stem.2011.11.005
 28. Plikus M V., Mayer JA, De La Cruz D, et al. Cyclic dermal BMP signalling regulates stem cell activation during hair regeneration. *Nature*. 2008. doi:10.1038/nature06457
 29. Greco V, Chen T, Rendl M, et al. A Two-Step Mechanism for Stem Cell Activation during Hair Regeneration. *Cell Stem Cell*. 2009. doi:10.1016/j.stem.2008.12.009
 30. Yang H, Adam RC, Ge Y, Hua ZL, Fuchs E. Epithelial-Mesenchymal Micro-niches Govern Stem Cell Lineage Choices. *Cell*. 2017. doi:10.1016/j.cell.2017.03.038
 31. Chow Z, Mueller SN, Deane JA, Hickey MJ. Dermal Regulatory T Cells Display Distinct Migratory Behavior That Is Modulated during Adaptive and Innate Inflammation. *J Immunol*. 2013. doi:10.4049/jimmunol.1203205
 32. Gratz IK, Truong H-A, Yang SH-Y, et al. Cutting Edge: Memory Regulatory T Cells Require IL-7 and Not IL-2 for Their Maintenance in Peripheral Tissues. *J Immunol*. 2013. doi:10.4049/jimmunol.1300212
 33. Petukhova L, Duvic M, Hordinsky M, et al. Genome-wide association study in alopecia areata implicates both innate and adaptive immunity. *Nature*. 2010. doi:10.1038/nature09114
 34. Conteduca G, Rossi A, Megiorni F, et al. Single nucleotide polymorphisms in the promoter regions of Foxp3 and ICOSLG genes are associated with Alopecia Areata. *Clin Exp Med*. 2014. doi:10.1007/s10238-012-0224-3
 35. Castela E, Le Duff F, Butori C, et al. Effects of low-dose recombinant interleukin 2 to promote T-regulatory cells in alopecia areata. *JAMA Dermatology*. 2014. doi:10.1001/jamadermatol.2014.504
 36. Nosbaum A, Ettinger M, Truong H, et al. Regulatory T cells facilitate cutaneous wound healing. *J Invest Dermatol*. 2015. doi:10.4049/jimmunol.1502139
 37. Nosbaum A, Prevel N, Truong H-A, et al. Cutting Edge: Regulatory T Cells Facilitate Cutaneous Wound Healing. *J Immunol*. 2016. doi:10.4049/jimmunol.1502139
 38. Mathur AN, Zirak B, Boothby IC, et al. Treg-Cell Control of a CXCL5-IL-17 Inflammatory Axis Promotes Hair-Follicle-Stem-Cell Differentiation During Skin-Barrier Repair. *Immunity*. 2019. doi:10.1016/j.immuni.2019.02.013
 39. Boehm F, Martin M, Kesselring R, et al. Deletion of Foxp3⁺ regulatory T cells in genetically targeted mice supports development of intestinal inflammation. *BMC Gastroenterol*. 2012. doi:10.1186/1471-230X-12-97
 40. Ishikawa D, Okazawa A, Corridoni D, et al. Tregs are dysfunctional in vivo in a spontaneous murine model of Crohn's disease. *Mucosal Immunol*. 2013. doi:10.1038/mi.2012.67
 41. Holmén N, Lundgren A, Lundin S, et al. Functional CD4⁺CD25^{high} regulatory T cells are enriched in the colonic mucosa of patients with active ulcerative colitis and increase with disease activity. *Inflamm Bowel Dis*. 2006. doi:10.1097/00054725-200606000-00003
 42. Makita S, Kanai T, Oshima S, et al. CD4 + CD25 bright T Cells in Human Intestinal Lamina Propria as Regulatory Cells. *J Immunol*. 2004. doi:10.4049/jimmunol.173.5.3119
 43. Maul J, Loddenkemper C, Mundt P, et al. Peripheral and intestinal regulatory CD4⁺CD25^{high} T cells in inflammatory bowel disease. *Gastroenterology*. 2005. doi:10.1053/j.gastro.2005.03.043
 44. Sharma A, Rudra D. Emerging functions of regulatory T cells in tissue homeostasis. *Front Immunol*. 2018. doi:10.3389/fimmu.2018.00883

45. Atarashi K, Tanoue T, Shima T, et al. Induction of colonic regulatory T cells by indigenous *Clostridium* species. *Science (80-)*. 2011. doi:10.1126/science.1198469
46. Wohlfert EA, Grainger JR, Bouladoux N, et al. GATA3 controls Foxp3 + regulatory T cell fate during inflammation in mice. *J Clin Invest*. 2011. doi:10.1172/JCI57456
47. Weiss JM, Bilate AM, Gobert M, et al. Neuropilin 1 is expressed on thymus-derived natural regulatory T cells, but not mucosa-generated induced Foxp3 + T reg cells . *J Exp Med*. 2012. doi:10.1084/jem.20120914
48. Yadav M, Louvet C, Davini D, et al. Neuropilin-1 distinguishes natural and inducible regulatory T cells among regulatory T cell subsets in vivo. *J Exp Med*. 2012. doi:10.1084/jem.20120822
49. Gottschalk RA, Corse E, Allison JP. Expression of Helios in Peripherally Induced Foxp3 + Regulatory T Cells. *J Immunol*. 2012;188(3):976-980. doi:10.4049/jimmunol.1102964
50. Ivanov II, McKenzie BS, Zhou L, et al. The Orphan Nuclear Receptor ROR γ t Directs the Differentiation Program of Proinflammatory IL-17+ T Helper Cells. *Cell*. 2006. doi:10.1016/j.cell.2006.07.035
51. Manel N, Unutmaz D, Littman DR. The differentiation of human TH-17 cells requires transforming growth factor- β and induction of the nuclear receptor ROR γ t. *Nat Immunol*. 2008;9(6):641-649. doi:10.1038/ni.1610
52. Kalekar LA, Cohen JN, Prevel N, et al. Regulatory T cells in skin are uniquely poised to suppress profibrotic immune responses. *Sci Immunol*. 2019. doi:10.1126/sciimmunol.aaw2910
53. Schiering C, Krausgruber T, Chomka A, et al. The alarmin IL-33 promotes regulatory T-cell function in the intestine. *Nature*. 2014. doi:10.1038/nature13577
54. Xu M, Pokrovskii M, Ding Y, et al. C-MAF-dependent regulatory T cells mediate immunological tolerance to a gut pathobiont. *Nature*. 2018. doi:10.1038/nature25500
55. Xu J, Yang Y, Qiu G, et al. c-Maf Regulates IL-10 Expression during Th17 Polarization. *J Immunol*. 2009. doi:10.4049/jimmunol.0900123
56. Kim KS, Hong SW, Han D, et al. Dietary antigens limit mucosal immunity by inducing regulatory T cells in the small intestine. *Science (80-)*. 2016. doi:10.1126/science.aac5560
57. Biton M, Haber AL, Rogel N, et al. T Helper Cell Cytokines Modulate Intestinal Stem Cell Renewal and Differentiation. *Cell*. 2018. doi:10.1016/j.cell.2018.10.008
58. Wong SH, Walker JA, Jolin HE, et al. Transcription factor ROR α is critical for nuocyte development. *Nat Immunol*. 2012. doi:10.1038/ni.2208
59. Dussault I, Fawcett D, Matthyssen A, Bader JA, Giguère V. Orphan nuclear receptor ROR α -deficient mice display the cerebellar defects of staggerer. *Mech Dev*. 1998. doi:10.1016/S0925-4773(97)00187-1
60. Von Moltke J, Ji M, Liang HE, Locksley RM. Tuft-cell-derived IL-25 regulates an intestinal ILC2-epithelial response circuit. *Nature*. 2016. doi:10.1038/nature16161
61. Jaradat M, Stapleton C, Tilley SL, et al. Modulatory role for retinoid-related orphan receptor α in allergen-induced lung inflammation. *Am J Respir Crit Care Med*. 2006. doi:10.1164/rccm.200510-1672OC
62. Moffatt MF, Gut IG, Demenais F, et al. A large-scale, consortium-based genomewide association study of asthma. *N Engl J Med*. 2010. doi:10.1056/NEJMoa0906312
63. Tang J, Zhou X, Liu J, et al. IL-25 promotes the function of CD4+CD25 + T regulatory cells and prolongs skin-graft survival in murine models. *Int Immunopharmacol*. 2015. doi:10.1016/j.intimp.2015.03.036
64. Barker N, Van Es JH, Kuipers J, et al. Identification of stem cells in small intestine and colon by marker gene *Lgr5*. *Nature*. 2007. doi:10.1038/nature06196

65. Barker N, Ridgway RA, Van Es JH, et al. Crypt stem cells as the cells-of-origin of intestinal cancer. *Nature*. 2009. doi:10.1038/nature07602
66. Pinto D, Gregorieff A, Begthel H, Clevers H. Canonical Wnt signals are essential for homeostasis of the intestinal epithelium. *Genes Dev*. 2003. doi:10.1101/gad.267103
67. Chae WJ, Park JH, Henegariu O, Yilmaz S, Hao L, Bothwell ALM. Membrane-bound Dickkopf-1 in Foxp3+ regulatory T cells suppresses T-cell-mediated autoimmune colitis. *Immunology*. 2017. doi:10.1111/imm.12766
68. Yan KS, Janda CY, Chang J, et al. Non-equivalence of Wnt and R-spondin ligands during Lgr5 + intestinal stem-cell self-renewal. *Nature*. 2017. doi:10.1038/nature22313
69. Kopf M, Schneider C, Nobs SP. The development and function of lung-resident macrophages and dendritic cells. *Nat Immunol*. 2015. doi:10.1038/ni.3052
70. Whitsett JA, Alenghat T. Respiratory epithelial cells orchestrate pulmonary innate immunity. *Nat Immunol*. 2015. doi:10.1038/ni.3045
71. Kim CF, Jackson EL, Woolfenden AE, et al. Identification of bronchioalveolar stem cells in normal lung and lung cancer. *Cell*. 2005. doi:10.1016/j.cell.2005.03.032
72. Fehrenbach H, Voswinckel R, Michl V, et al. Neoalveolarisation contributes to compensatory lung growth following pneumonectomy in mice. *Eur Respir J*. 2008. doi:10.1183/09031936.00109407
73. Plantier L, Crestani B, Wert SE, et al. Ectopic respiratory epithelial cell differentiation in bronchiolised distal airspaces in idiopathic pulmonary fibrosis. *Thorax*. 2011. doi:10.1136/thx.2010.151555
74. Monticelli LA, Sonnenberg GF, Abt MC, et al. Innate lymphoid cells promote lung-tissue homeostasis after infection with influenza virus. *Nat Immunol*. 2011. doi:10.1038/ni.2131
75. Arpaia N, Green JA, Moltedo B, et al. A Distinct Function of Regulatory T Cells in Tissue Protection. *Cell*. 2015. doi:10.1016/j.cell.2015.08.021
76. Lam AJ, MacDonald KN, Pesenacker AM, et al. Innate Control of Tissue-Reparative Human Regulatory T Cells. *J Immunol*. 2019;202(8):2195-2209. doi:10.4049/jimmunol.1801330
77. Chakraborty K, Bhattacharyya A. Treg depletion reduces bleomycin-induced pulmonary fibrosis by inducing Th17 and other CD4+T cell subset responses in lung. In: ; 2016. doi:10.1183/13993003.congress-2016.pa2095
78. Kotsianidis I, Nakou E, Bouchliou I, et al. Global impairment of CD4+CD25+FOXP3+ regulatory T cells in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med*. 2009. doi:10.1164/rccm.200812-1936OC
79. Burzyn D, Kuswanto W, Kolodin D, et al. A Special Population of regulatory T Cells Potentiates muscle repair. *Cell*. 2013. doi:10.1016/j.cell.2013.10.054
80. Mock JR, Garibaldi BT, Aggarwal NR, et al. Foxp3 + regulatory T cells promote lung epithelial proliferation. *Mucosal Immunol*. 2014. doi:10.1038/mi.2014.33
81. Lefrançais E, Ortiz-Muñoz G, Caudrillier A, et al. The lung is a site of platelet biogenesis and a reservoir for haematopoietic progenitors. *Nature*. 2017. doi:10.1038/nature21706
82. Hussell T, Bell TJ. Alveolar macrophages: Plasticity in a tissue-specific context. *Nat Rev Immunol*. 2014. doi:10.1038/nri3600
83. Tsao P-N, Matsuoka C, Wei S-C, et al. Epithelial Notch signaling regulates lung alveolar morphogenesis and airway epithelial integrity. *Proc Natl Acad Sci*. 2016. doi:10.1073/pnas.1511236113
84. Minutti CM, Modak R V., Macdonald F, et al. A Macrophage-Pericyte Axis Directs Tissue Restoration via Amphiregulin-Induced Transforming Growth Factor Beta

- Activation. *Immunity*. 2019. doi:10.1016/j.immuni.2019.01.008
85. Wang S, Zhang Y, Wang Y, et al. Amphiregulin confers regulatory T cell suppressive function and tumor invasion via the EGFR/GSK-3 β /Foxp3 axis. *J Biol Chem*. 2016. doi:10.1074/jbc.M116.717892
 86. Hui SP, Sheng DZ, Sugimoto K, et al. Zebrafish Regulatory T Cells Mediate Organ-Specific Regenerative Programs. *Dev Cell*. 2017. doi:10.1016/j.devcel.2017.11.010
 87. Agudo J, Park ES, Rose SA, et al. Quiescent Tissue Stem Cells Evade Immune Surveillance. *Immunity*. 2018. doi:10.1016/j.immuni.2018.02.001
 88. Takeuchi Y, Nishikawa H. Roles of regulatory T cells in cancer immunity. *Int Immunol*. 2016. doi:10.1093/intimm/dxw025