



King's Research Portal

DOI: 10.1038/s41415-020-2170-x

Document Version Peer reviewed version

Link to publication record in King's Research Portal

Citation for published version (APA): Bomfim, D., Rahim, N., & Austin, R. (2020). Biomechanical planning for minimally invasive indirect restorations. *British Dental Journal*, 229(7), 425-429. https://doi.org/10.1038/s41415-020-2170-x

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Biomechanical planning for minimally invasive indirect restorations

Bomfim, DI; Rahim NM and Austin, RS.

Contents

Introduction
Planning Indirect Restorations in vital teeth, non-vital teeth and worn dentitions
What does the evidence say on direct vs. indirect restorations for vital teeth?
Full vs. Partial coverage restorations long term outcome data in general dental practice
Minimally Invasive Restoration of Vital Teeth – why shift towards partial coverage indirect restorations
for vital teeth?
Minimally Invasive Restoration of the Endodontically treated teeth9
Minimally Invasive Restoration of Erosive Tooth Wear10
Conclusion11
Figures and tables
References

Introduction

Increasingly, a worldwide consensus exists towards minimally invasive approaches for managing dental caries (1); which for non-cavitated carious lesions often involves non-invasive or micro-invasive management (2). In contrast, no such consensus yet exists for the prescription, design and preparation of indirect restorations. The majority of the half-a-billion dental restorations placed worldwide every year are direct composite-resin restorations (which are more conservative than amalgam restorations), whereas, in contrast, the majority of indirect restorations placed are still full coverage crowns, which are less conservative than biomimetic partial coverage indirect restorations (3).

This suggests that biomechanical preparations which allow for a more conservative biomimetic approach to planning restorations are still not popular when it comes to indirect restorations. Indeed, in the United States 95% of indirect restorations are still full coverage crowns rather than partial coverage indirect restorations (4). Moreover, surveys of UK General Dental Practice reveal that the most commonly used methods for planning and designing tooth preparation are the dimensions of the preparation burs and the form of the opposing/adjacent teeth (5), as opposed to minimally invasive partial coverage designs. This is a missed opportunity, given the technological advances for planning restorative treatments, including the ability for digital planning of minimally invasive endodontic access cavities and digital diagnostic wax ups having potential to be more precise than conventional wax ups.

This paper will review the principles behind biomechanical planning for minimally invasive indirect restorations such as the case shown in Figure 3, and consider the with many different variables influencing their outcome as shown in Figure 1 (6-8). This paper will also explore the planning and executing of indirect partial coverage restorations and will outline practical recommendations for maximizing the outcomes for minimally invasive approaches to indirect restorations, with a special focus on vital teeth, endodontically treated teeth and worn dentitions. Throughout the paper, the supporting

evidence for each rationale for partial coverage restorations will be considered, as well as the risks and benefits of adopting a minimally invasive approach to indirect restorations.

Planning Indirect Restorations in vital teeth, non-vital teeth and worn dentitions.

What does the evidence say on direct vs. indirect restorations for vital teeth?

The most important consideration regarding minimally invasive indirect dentistry is to consider whether to provide a direct restoration as opposed to an indirect restoration. Each clinician reading this article will have developed a personal threshold for provision of indirect restorations. This is based on the clinician's own expertise, personal protocols and practice setting, balanced by the available scientific literature, on one hand, and patient values and expectations, on the other (9). Provision of indirect restorations sacrifices more healthy tooth tissue than direct restorations and moreover full coverage crown restorations require more tooth tissue removal than partial coverage restorations (10).

Online evidence-based syntheses algorithms are emerging to help resolve this dilemma. To date more than 10 systemic reviews with meta-analyses (11-20), provide data from prospective clinical outcome studies looking into indirect restorations which are summarized by online clinical-decision supporting applications, e.g. *'Crown-or-fill^{®'}* www.Crownorfill.com (21). As shown in Figure 2, the *Crown-or-fill* app summarizes the evidence regarding indirect or direct restoration of teeth, depending on specific preexisting factors (i.e. location of tooth in dentition, pulpal status, amount of coronal tooth tissue remaining and quality of pre-existing root canal treatment). The algorithm includes the fixed prosthodontic replacement options in terms of bridges and implant supported restorations, which is particularly helpful for discussions of all possible options with patients.

Once the clinical features are entered, prospective outcome data is presented (5-year % failure rate) for each restorative option, in terms of either direct (amalgam / composite resin) or indirect (crown) restoration (with or without a post for non-vital teeth), as well as the fixed prosthodontic replacement options (implant crown /bridge). There is no doubt that this type of evidence synthesis will increasingly guide the decision-making process, when considering whether to directly restore, crown, root fill or extract and replace. This will help consent and communication with patients and potentially save clinicians time and money whilst improving patient outcomes. The main key benefits for dentists using the algorithm are listed in

Decision making Topic		Options appraisal		Strengths	Limitations
Direct or	indirect	Crown vs.	direct	Especially helpful given	Doesn't explicitly
restoration?		restoration?		number of dentine walls	mention the ferrule
		If direct restor	ation;	available and location of	effect
		amalgam vs. compo	site?	tooth in dental arch.	Doesn't take into
				Can be combined with	consideration of
				measurements taken	periodontal health
				from a restorability	
				assessment or intra-oral	
				scan of tooth with	
				restorations removed	
Core with or	without a	Core plus post?		Very useful to help	Doesn't consider

Table 1 Key benefits to dentists for using the Crown-of-Fill Evidence Based Algorithm

post?			justify added time and	indirect vs/ direct
			expense of post	posts or other
			placement	factors (eg cement,
				which root etc.)
Extraction	and	Whether to endo treat or	Ensures that patients	Doesn't' factor in
replacement	VS.	extract?	are aware of annual	costs and cost-
restoration		Whether to replace with	failure rates especially	effectiveness, would
		a bridge or dental	when a tooth is heavily	be good to be able to
		Implant?	broken down and	input financial
			requires endo plus	aspects of care.
			crown.	Doesn't factor in risk
			Helps consider whether	factors for
			an implant/bridge	development of peri-
			maybe more cost-	implantitis.
			effective than a	
			endo/crown	

However, one of the key drawbacks of this application for general dental practice is that although this provides highly standardized data, there are many modifying dentist and patient factors (e.g. case selection, bruxism and caries risk) have a major influence on restoration survival. In addition, the above evidence doesn't discriminate between choice of material or restoration design (partial or full coverage). Factors such as practice setting, patient cohort, renumeration arrangements and clinicians' experience, also make generalisation about indirect restoration outcomes very difficult and therefore we must consider practice-based data which is more applicable to the general dental practitioner (22, 23).

Full vs. Partial coverage restorations long term outcome data in general dental practice

Many readers of this series on minimally invasive (MI) dentistry may already have a preference towards more MI adhesive strategies for teeth that were previously deemed as requiring full coverage crowns. However, full coverage crowns remain a popular choice amongst dentists, with long term practice-based outcome data supporting their use.

Outcome data from single operators, caring for highly motivated cohorts of patients, have 5 decades of follow-up data of indirect restorations following traditional principles. These data reveal that, in patients with regular recall intervals, including strict control of oral hygiene and meticulous occlusal management, the mean survival of metal-ceramic crowns can be up to 47.53 years in a general practice cohort (24) and 25-year survival of 85.40% for a cohort of tooth wear cases in a specialist prosthodontic practice (25). Whilst not generalizable to all UK General Dental Practices, it does reinforce that experienced operators and case selection play an important role in survival of restorations.

Minimally Invasive Restoration of Vital Teeth – why shift towards partial

coverage indirect restorations for vital teeth?

In the last few decades, 3 major drivers are changing clinician's attitudes towards design of indirect restorations, away from full coverage conventional mechanical preparations and towards provision of indirect tooth colored restorations for vital teeth.

Firstly, indirect restorative dentistry is benefitting from the adhesive bonding protocols pioneered in direct adhesive dentistry, whilst also benefitting from more aesthetic translucent etchable/sandblasted/silanated indirect materials. This is allowing a profound shift away from mechanical preparation designs (26) towards adhesive approaches focusing on preservation of enamel and dentine (27).

The second major shift is adoption of biomimetic additive approaches, which refers to the use of digital or analogue wax-ups. The use of digital wax ups have evolved from simple additive approaches based on digital smile design and intra-oral try-in to more extensive full-arch restorative cases as shown in , where digital superimposition of a subtractive wax up is used to guide tooth structure removal. This allows for a more precise plan of different preparations for each specific surface of each individual tooth based on the final proposed contour rather than a standardized tooth removal for all teeth.

Finally, the third profound transformation driving partial coverage indirect restorations is the use of digital dentistry and both additive and subtractive in-surgery computer aided manufacturing, as shown in Figure 3.

However, whilst these technological advances have brought benefits, the introduction of novel hybrid ceramic-polymer materials also has brought some challenges. Early practice-based data shows clinically concerning outcomes in certain situations, namely premature de-bonding or terminal fracture of the material when used as a full coverage crown in load bearing situations. Indeed some practice-based data is finding almost a third of novel ceramic-polymer hybrid restorations experienced debonding after 1 year and a quarter novel non-crystallized lithium disilicate restorations experienced terminal fracture after 1 year, indeed the same data showed almost 60% of Zirconia crowns causing greater than expected wear of the opposing dentition after 1 year (28). This clinically concerning data led to some manufactures withdrawing indications for their novel ceramic-polymer hybrid materials as a full coverage crown (29). The poor performance of these materials as full coverage restorations is thought to be due to the hoop stress concentration at the occlusal/axial transition when used as a conventional full coverage crown, as highlighted by Finite Element Analysis which resulted in the fracture at the transition from occlusal to axial (30). Therefore, these practice-based data of novel indirect materials show clinically concerning results –So, where does this leave us? Should we be changing our preparation

design away from full coverage restorations and towards partial coverage tooth-coloured materials and if so how to maximize the success of these restorations?

Preservation of tooth structure has always been the first key principle of all indirect restorations and there have been several recent major changes which are increasing the predictability of minimally invasive indirect restorations. When the survival of partial coverage ceramic restorations is considered, the largest practice based research data is the Ceramic Survival Analysis (CSA) project (mostly Germany and US). Analysis of 5791 ceramic inlay or onlays in 5523 patients shows that inlays and onlays constructed from many glass ceramic materials (mostly lithium disilicate) have good outcomes for a large number of dentists (167) working in different settings over two decades (22). The CSA reported annual failure rates (AFR) of 1.6 % at 10 years regardless of type of glass ceramic and or whether CAD/CAM or pressed techniques. However, perhaps predictably, restorations with a deep marginal extension into dentine showed an increased failure rate of 78 % compared to restorations in enamel. Unsurprisingly, use of glass-ionomer cement as a core material and use of simplified adhesive protocols (single or dual step dentine bonding agents) also presented a 142 % increased risk of failure as opposed to gold standard 3-step etch/prime/bond protocols.

Therefore any clinician considering a change to their clinical practice, such as adopting a novel CAD/CAM material with a minimally invasive approach, is encouraged to enroll onto the open-source CSA (<u>www.csa-online.net</u>) and enter as a minimum, data on their first 50 consecutive cases using the new material, with full details of their entire protocol. On subsequent review of every case, clinicians determine as soon as possible they are experiencing any issues with their cases, which may require modification of case selection, preparation design or bonding protocol for example.

Minimally Invasive Restoration of the Endodontically treated teeth

Root canal treatment is needed when pulpal health is irreversibly damaged by caries, cracks, restorations and trauma. Timely restoration back to form, function and aesthetics is as important as the quality of the root canal treatment, in order to protect endodontically treated teeth from future fracture or loss of coronal seal and re-infection (ETT). (15, 31). Many clinical studies are now available to support cuspal coverage of ETT in order to improve outcomes for ETT.

Even prior to the restoration, a key aim during endodontic treatment is the preservation of tooth structure with multiple prospective clinical studies correlating increased tooth structure with improved outcomes (32). Provision of cuspal coverage for ETT has specific benefits for premolars and molars, with a non-functional cusp sparing approach advocated to preserve tooth tissue, as opposed to full coverage crowns (4). However, this approach, whilst showing promise in finite element analyses (33), has not been widely adopted. It is also perhaps surprising that more clinical research has not been carried out towards direct cuspal coverage restoration of ETT, even when extensively broken down (34).

In terms of which cusps to cover, as shown in Figure 4, a key parameter to consider is the thickness of the remaining dentine walls. Krifka et al (2009) investigated the influence of remaining cusp wall thickness on the marginal integrity and enamel crack formation and concluded that if non fuctional cusps are to be left uncovered they should be of at least 2 mm thickness (35). As shown in Figure 4, this measurement maybe carried out simply using Iwanson calipers or alternatively by scanning the remaining tooth structure once all the old restorations and caries has been removed and using cross sectional measurement tools provided within chairside CAD/CAM software.

Minimally Invasive Restoration of Erosive Tooth Wear

Clearly, when managing a disease process which involves gradual loss of hard tooth tissue, a minimally invasive approach aimed at preservation of tooth tissue is required. In the first instance, a thorough diagnostic phase is required to ascertain whether indeed a non-operative strategy can be employed, in order to avoid restorations (36). Once it has been decided to provide restorations, initial prescription of direct adhesive restorations, in addition with other minimally invasive methods such as tooth whitening is the recommended course of action, especially for mild erosive tooth wear cases where most of the enamel has been retained.

However, the long-term outcome of minimally invasive direct restorations for patients with moderate to severe tooth wear with extensive dentine exposure coupled with reduced surface area for bonding and parafunctional habits is uncertain and reported annual failure rates vary widely (37), from 0.4-26.3% for direct resin composite, 0-14.9% for indirect resin composite and 2.7% for indirect ceramic restorations.

This leads to the conclusion that choice of a direct or indirect approach is largely operator and patient dependent, therefore it seems sensible to try to avoid indirect restorations until direct restorations have been tried (38). Recent prospective clinical trials of direct restorations for tooth wear have found that a key factor influencing success of anterior composite restorations for tooth wear cases is ensuring that the anterior restorations are provided at a single visit, thus maximizing control of both the adhesive bonding to a fresh substrate and also provision of an appropriate occlusal scheme (39).

For cases whose tooth wear is primarily of an erosive component, such as the case highlighted in Figure 5 who presented with amelogenesis imperfecta, hypodontia and erosive tooth wear primarily related to acid erosion, partial coverage ceramic or resin restorations are likely to have a good prognosis and be the first choice of restorative treatment. However for cases with parafunctional bruxism, whose tooth wear is so severe that all the enamel has been lost for adhesive bonding, full coverage conventional crowns provided with traditional retention and resistance form are the more appropriate treatment choice.

Conclusion

This paper has considered the benefits of adopting a minimally invasive approach when considering the design and execution of indirect restorations. Although we are now in an era whereby prescription of high-gold content partial coverage restorations are declining, new technologies using to preserve tooth structure are arising due to increased availability of bondable aesthetic indirect materials and modern imaging and CAD/CAM for a more bespoke approach to cuspal coverage restorations.

However, more research is needed for long term prospective clinical trials of these novel materials to ensure that clinicians are aware of the correct case selection, design and bonding protocols so that patient outcomes are optimized.

Figures and tables



Figure 1 Variables which influence the outcome of partial coverage indirect restorations



Figure 2 Algorithm for case selection – reproduced from <u>www.crownorfill.com</u> (21)











Figure 3 Case showing the benefits of a minimally invasive approach to indirect restorations.



Figure 4 Step-by-step process for decision making regarding which cusps to cover, according to thickness of residual cusps. 'Courtesy of Dr. Nathaniel Lawson'







Dentinaltube

non-functional cusps









bonding agent applied to all freshly cut enamel & dentin

Dentinaltube

flowable composite placed to blockout all undercuts

Dentinaltube









Figure 5 A case of amelogenesis imperfecta, hypodontia and erosive tooth wear of primarily erosive nature, treated with anterior direct composite resin restorations, posterior partial coverage lithium disilicate restorations and single unit implant supported crowns.

References

1. Schwendicke F, Frencken JE, Bjørndal L, Maltz M, Manton DJ, Ricketts D, et al. Managing carious lesions: consensus recommendations on carious tissue removal. Advances in dental research. 2016;28(2):58-67.

2. Schwendicke F, Splieth C, Breschi L, Banerjee A, Fontana M, Paris S, et al. When to intervene in the caries process? An expert Delphi consensus statement. Clinical oral investigations. 2019;23(10):3691-703.

3. Zöchbauer H. Number of dental restorations worldwide. Market Research Ivoclar Vivadent. 2011.

4. Christensen GJ. The case for onlays versus tooth-colored crowns. The Journal of the American Dental Association. 2012;143(10):1141-4.

5. Jum'ah A, Creanor S, Wilson N, Burke F, Brunton P. Dental practice in the UK in 2015/2016. Part 3: aspects of indirect restorations and fixed prosthodontics. British dental journal. 2019;226(3):192.

6. Austin RS, Chan M. Digital Restoration Design based on Biologic Principles. In: Rekow ED, editor. Digital dentistry: a comprehensive reference and preview of the future: Quintessence Publishing Company Limited; 2018. p. 14.

7. Kelleher M, Bomfim D, Austin R. Biologically Based Restorative Management of Tooth Wear. International Journal of Dentistry. 2012;2012.

8. Magne P. Pascal Magne:'It should not be about aesthetics but tooth-conserving dentistry'. Interview by Ruth Doherty. British dental journal. 2012;213(4):189.

9. Richards D. Evidence-based dentistry: Quintessence Publishing; 2019.

10. Edelhoff D, Sorensen JA. Tooth structure removal associated with various preparation designs for anterior teeth. The Journal of prosthetic dentistry. 2002;87(5):503-9.

11. Afrashtehfar KI, Emami E, Ahmadi M, Eilayyan O, Abi-Nader S, Tamimi F. Failure rate of singleunit restorations on posterior vital teeth: a systematic review. The Journal of prosthetic dentistry. 2017;117(3):345-53. e8.

12. Afrashtehfar KI, Ahmadi M, Emami E, Abi-Nader S, Tamimi F. Failure of single-unit restorations on root filled posterior teeth: A systematic review. Int Endod J. 2017;50(10):951-66.

13. Pjetursson BE, Sailer I, Makarov NA, Zwahlen M, Thoma DS. All-ceramic or metal-ceramic toothsupported fixed dental prostheses (FDPs)? A systematic review of the survival and complication rates. Part II: Multiple-unit FDPs. Dent Mater. 2015;31(6):624-39.

14. E. Jung R, Zembic A, Pjetursson BE, Zwahlen M, S. Thoma D. Systematic review of the survival rate and the incidence of biological, technical, and aesthetic complications of single crowns on implants reported in longitudinal studies with a mean follow-up of 5 years. Clinical oral implants research. 2012;23:2-21.

15. Ng YL, Mann V, Gulabivala K. Tooth survival following non-surgical root canal treatment: a systematic review of the literature. Int Endod J. 2010;43(3):171-89.

16. Nixdorf DR, Moana-Filho EJ, Law AS, McGuire LA, Hodges JS, John MT. Frequency of persistent tooth pain after root canal therapy: a systematic review and meta-analysis. Journal of endodontics. 2010;36(2):224-30.

17. Torabinejad M, Corr R, Handysides R, Shabahang S. Outcomes of nonsurgical retreatment and endodontic surgery: a systematic review. Journal of endodontics. 2009;35(7):930-7.

18. Teughels W, Merheb J, Quirynen M. Critical horizontal dimensions of interproximal and buccal bone around implants for optimal aesthetic outcomes: a systematic review. Clinical oral implants research. 2009;20:134-45.

19. Ng YL, Mann V, Rahbaran S, Lewsey J, Gulabivala K. Outcome of primary root canal treatment: systematic review of the literature -- Part 2. Influence of clinical factors. Int Endod J. 2008;41(1):6-31.

20. Ng YL, Mann V, Rahbaran S, Lewsey J, Gulabivala K. Outcome of primary root canal treatment: systematic review of the literature–part 1. Effects of study characteristics on probability of success. Int Endod J. 2007;40(12):921-39.

21. Afrashtehfar KI, Tamimi F. An online tool that provides access to evidence-based literature on dental restorations: www. crownorfill. com. J Prosthet Dent. 2017;118(6):696-7.

22. Collares K, Corrêa MB, Laske M, Kramer E, Reiss B, Moraes RR, et al. A practice-based research network on the survival of ceramic inlay/onlay restorations. Dent Mater. 2016;32(5):687-94.

23. Schwass DR, Lyons KM, Purton DG. How long will it last? The expected longevity of prosthodontic and restorative treatment. NZ Dent J. 2013;109(3):98-105.

24. Olley RC, Andiappan M, Frost PM. An up to 50-year follow-up of crown and veneer survival in a dental practice. The Journal of prosthetic dentistry. 2018;119(6):935-41.

25. Walton TR. The up to 25-year survival and clinical performance of 2,340 high gold-based metalceramic single crowns. International Journal of Prosthodontics. 2013;26(2).

26. Shillingburg HT, Hobo S, Whitsett LD, Jacobi R, Brackett S. Fundamentals of fixed prosthodontics: Quintessence Publishing Company; 1997.

27. Edelhoff D, Stimmelmayr M, Schweiger J, Ahlers MO, Güth J-F. Advances in materials and concepts in fixed prosthodontics: a selection of possible treatment modalities. British dental journal. 2019;226(10):739-48.

28. Christensen GJ. Gordon J. Christensen's Clinicians Report Volume 11 Issue 11, November 2018, Pages 1–3 3707 N. Canyon Rd, Bldg 7, Provo UT 84604: CR Foundation[®]; 2018 [Available from: https://www.ivoclarvivadent.us/mam/celum/celum_assets/9516343197726_emaxCRreport.pdf?0.

29. Notice: Change in Indication — Lava[™] Ultimate Restorative [press release]. St. Paul, MN: 3M12 Jun 2015

30. Shembish FA, Tong H, Kaizer M, Janal MN, Thompson VP, Opdam NJ, et al. Fatigue resistance of CAD/CAM resin composite molar crowns. Dent Mater. 2016;32(4):499-509.

31. Ray H, Trope M. Periapical status of endodontically treated teeth in relation to the technical quality of the root filling and the coronal restoration. Int Endod J. 1995;28(1):12-8.

32. Al-Nuaimi N, Patel S, Austin RS, Mannocci F. A prospective study assessing the effect of coronal tooth structure loss on the outcome of root canal retreatment. Int Endod J. 2017:n/a-n/a.

33. Lin C-L, Chang Y-H, Liu P-R. Multi-factorial analysis of a cusp-replacing adhesive premolar restoration: a finite element study. Journal of dentistry. 2008;36(3):194-203.

34. ŞİŞMANOĞLU S. Restoration of Endodontically Treated Teeth: A Review of Direct Restorative Approach. Aurum Journal of Health Sciences. 2019;2(1):21-40.

35. Krifka S, Anthofer T, Fritzsch M, Hiller K, Schmalz G, Federlin M. Ceramic inlays and partial ceramic crowns: influence of remaining cusp wall thickness on the marginal integrity and enamel crack formation in vitro. Operative dentistry. 2009;34(1):32-42.

36. Loomans B, Opdam N, Attin T, Bartlett D, Edelhoff D, Frankenberger R, et al. Severe Tooth Wear: European Consensus Statement on Management Guidelines. The journal of adhesive dentistry. 2017;19:111-9.

37. Mesko ME, Sarkis-Onofre R, Cenci MS, Opdam NJ, Loomans B, Pereira-Cenci T. Rehabilitation of severely worn teeth: A systematic review. Journal of dentistry. 2016;48:9-15.

38. Varma S, Preiskel A, Bartlett D. The management of tooth wear with crowns and indirect restorations. Bdj. 2018;224:343.

39. Loomans B, Kreulen C, Huijs-Visser H, Sterenborg B, Bronkhorst E, Huysmans M, et al. Clinical performance of full rehabilitations with direct composite in severe tooth wear patients: 3.5 Years results. Journal of dentistry. 2018;70:97-103.