



King's Research Portal

DOI:

[10.1038/s41591-018-0115-6](https://doi.org/10.1038/s41591-018-0115-6)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Quek, L., David, M. D., Kennedy, A., Metzner, M., Amatangelo, M., Shih, A., Stoilova, B., Quivoron, C., Heiblig, M., Willekens, C., Saada, V., Alsafadi, S., Vijayabaskar, M. S., Peniket, A., Bernard, O. A., Agresta, S., Yen, K., MacBeth, K., Stein, E., ... Vyas, P. (2018). Clonal heterogeneity of acute myeloid leukemia treated with the IDH2 inhibitor enasidenib. *Nature Medicine*, 24(8), 1167-1177. <https://doi.org/10.1038/s41591-018-0115-6>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1 **Clonal heterogeneity of acute myeloid leukemia treated with the IDH2 inhibitor**

2 **Enasidenib**

3
4
5 **Lynn Quek^{1,2,3*}, Muriel D. David^{4*}, Alison Kennedy^{1,2}, Marlen Metzner^{1,2}, Michael**
6 **Amatangelo⁵, Alan Shih⁶, Bilyana Stoilova^{1,2}, Cyril Quivoron⁴, Maël Heiblig⁴,**
7 **Christophe Willekens^{4,7}, Véronique Saada^{4,7}, Samar Alsafadi¹⁵, MS Vijayabaskar¹¹,**
8 **Andy Peniket³, Oliver A Bernard⁴, Sam Agresta⁸, Katharine Yen⁸, Kyle MacBeth⁵, Eytan**
9 **Stein⁶, George S. Vassiliou^{11,12,13,14}, Ross Levine^{6,9,10}, Stephane De Botton^{4,7*}, Anjan**
10 **Thakurta^{5*}, Virginie Penard-Lacronique^{4*} and Paresch Vyas^{1,2,3*}**

11 ¹MRC Molecular Hematology Unit, WIMM, University of Oxford, UK; ²Haematology Theme
12 Oxford Biomedical Research Centre and ³Department of Hematology, Oxford University
13 Hospitals NHS Foundation Trust, UK; ⁴INSERM U1170, Gustave Roussy, Université Paris-
14 Saclay, Equipe labellisée Ligue Nationale Contre le Cancer Villejuif, France, ⁵Celgene
15 Corporation, Summit, New Jersey, USA; ⁶Department of Medicine, Leukemia Service,
16 Memorial Sloan Kettering Cancer Center, New York, USA; ⁷Département d'Hématologie,
17 Gustave Roussy, Université Paris-Saclay, Villejuif, France, ⁸Agios Pharmaceuticals, Inc.,
18 Cambridge, Massachusetts, USA; ⁹Center for Hematologic Malignancies, Memorial Sloan
19 Kettering Cancer Center, New York, USA; ¹⁰Human Oncology and Pathogenesis Program,
20 Memorial Sloan Kettering Cancer Center, New York; USA. ¹¹Haematological Cancer
21 Genetics, Wellcome Trust Sanger Institute, Hinxton, Cambridge, CB10 1SA, UK;
22 ¹²Department of Haematology, University of Cambridge, Cambridge CB2 0PT, UK;
23 ¹³Department of Haematology, Cambridge University Hospitals NHS Trust, Cambridge CB2
24 0QQ, UK; ¹⁴Cancer Research UK Cambridge Institute, University of Cambridge, Robinson
25 Way, Cambridge, CB2 0RE, UK; ¹⁵Département de Recherche Translationnelle/ Inserm
26 U830, Institut Curie, Université Paris Sciences et Lettres, Paris, France.

27 *These authors contributed equally to this work

28 Corresponding authors: Paresh Vyas (paresh.vyas@imm.ox.ac.uk), Lynn Quek
29 (lynn.quek@imm.ox.ac.uk), Virginie Penard-Lacronique (virginie.penard-
30 lacronique@inserm.fr), Stephane de Botton (stephane.debotton@gustaveroussy.fr) and
31 Anjan Thakurta (athakurta@celgene.com)

32 Total words (main text) : 4473

33 Abstract 177 words

34 **Running Title: Clonal Responses in IDH2 mutant AML**

35 **Mutations in the gene encoding isocitrate dehydrogenase 2 (*IDH2*) occur in several**
36 **types of cancer, including acute myeloid leukemia (AML). In model systems, mutant**
37 **IDH2 causes hematopoietic differentiation arrest. Enasidenib, a selective small-**
38 **molecule inhibitor of mutant IDH2, produces a clinical response in 40% of treated**
39 **relapsed/ refractory AML patients by promoting leukemic cell differentiation. Here, we**
40 **studied the clonal basis of response and acquired resistance to enasidenib treatment.**
41 **Using sequential patient samples, we determined the clonal structure of**
42 **hematopoietic cell populations at different stages of differentiation. Pre-therapy IDH2**
43 **mutant clones showed variable differentiation arrest. Enasidenib treatment promoted**
44 **hematopoietic differentiation from either terminal or ancestral mutant clones; less**
45 **frequently, treatment promoted differentiation of non-mutant cells. Analysis of paired**
46 **diagnosis/relapse samples did not identify second site mutations in *IDH2* at relapse.**
47 **Instead, relapse arose by clonal evolution, or selection, of terminal or ancestral**
48 **clones, highlighting multiple bypass pathways that could potentially be targeted to**
49 **restore differentiation arrest. Mapping clonal structure in cell populations at different**
50 **stages of differentiation during therapy illustrates how different clones respond and**
51 **evolve during relapse.**

52 Differentiation arrest is a common feature of many cancer cells. Though intratumoral clonal
53 heterogeneity is well documented, we lack a detailed understanding of how the complement
54 of driver mutations within a clone, contribute to differentiation arrest. Furthermore, though it
55 is clear that genetic and functional, intratumoral heterogeneity helps determine clinical
56 outcome to cancer therapy^{1,2}, few studies have investigated the relationship between clonal
57 structure and therapy response, particularly for therapies targeting oncogenic epigenetic
58 processes. With an average of 13 somatic mutations per patient, arranged in an limited
59 number of clones³, AML is simpler than most cancers from a genomic context, providing a
60 potential paradigm to answer these questions.

61 Somatic mutations in conserved arginine residues R140 and R172 in *ISOCITRATE*
62 *DEHYDROGENASE 2 (IDH2)* occur in 15-25% of AML patients⁴⁻⁶. The mutant proteins have
63 neomorphic activity producing R-2-hydroxyglutarate (2-HG) that competitively inhibits α -
64 ketoglutarate-dependent enzymes including the TET family of 5-methylcytosine (5mC)
65 hydroxylases and Jumonji-C domain histone demethylases^{7,8}. This leads to DNA
66 hypermethylation⁹, increased repressive histone methylation⁸ and impaired hematopoietic
67 differentiation that is reversed by mutant IDH inhibition in model systems^{8,10-12}.

68 We recently showed that enasidenib (AG-221/CC-90007), a first-in-class, allosteric inhibitor
69 of mutant IDH2 (mIDH2)¹³, reduces serum 2-HG, reverses DNA hypermethylation and
70 promotes hematopoietic differentiation in preclinical models^{13,14}. In a phase 1/2 clinical trial,
71 enasidenib monotherapy produced a response rate of 40.3% in relapsed/refractory AML
72 patients¹⁵. In most responding patients terminally mature blood cells were *IDH2* mutant,
73 consistent with response due to enasidenib-induced differentiation of *IDH2* mutant cells¹⁶.
74 However, in 9/71 (12.6%) responding patients, *IDH2* mutant cells were eliminated from
75 peripheral blood cells¹⁶. Failure to respond to enasidenib was associated with a higher co-
76 mutational burden and *NRAS* mutations¹⁶. Finally, most patients who initially responded
77 eventually relapsed. These initial studies did not assess which clone(s) differentiated in
78 response to enasidenib and the clonal mechanism of acquired enasidenib resistance.

79 We have addressed these two questions by studying sequential samples from a subset of
80 trial patients. We showed marked variation in the degree of differentiation arrest of mutant
81 *IDH2* clones. Restoration of differentiation by inhibition of mutant *IDH2* was also clone
82 dependent, varying between patients, arising from either ancestral or terminal clones. In a
83 minority of patients differentiation occurred from wild type progenitor cells, consistent with
84 molecular remission in a subset of patient. Acquired resistance to enasidenib leading to
85 differentiation arrest and relapse did not occur by second site mutations in *IDH2*. Instead,
86 differentiation arrest was restored by multiple mechanisms through clonal evolution or clonal
87 selection.

88 **Results**

89 **Patient cohort studied**

90 The trial enrolled 176 relapsed/refractory *IDH2* mutant AML patients¹⁵. Here, we studied a
91 cytogenetically and genetically representative subset of 37 patients enriched for enasidenib
92 responders (30/37 responders, **Supplementary Fig. 1a-c**). An extended mutational profile
93 was determined in 33/37 patients by either whole exome sequencing (WES) (16/36 patients
94 at read depth of 19-843x, average 121x at loci where variants were called), or targeted
95 resequencing (17/36 patients) (**Supplementary Tables 1 and 2**). Compared to the entire
96 trial cohort, the patient cohort studied here had similar serum baseline 2-HG levels and the
97 mean level of 2-HG suppression was similar in the two cohorts (i.e. on-target response to
98 enasidenib; **Supplementary Fig. 1d-e**).

99 **Enasidenib rebalanced progenitor and precursor compartment sizes and restored** 100 **progenitor function**

101 In AML there are two orthogonal potential hierarchies (**Supplementary Fig. 2a**). There is
102 clonal hierarchy with an initiating clone, transitional and terminal clones. We refer to all non-
103 terminal clones as ancestral clones. These mutant clones exist in a second hierarchy, a
104 hematopoietic cell hierarchy. AML initiating mutations occur in stem, or long-lived progenitor,
105 cells but initiating clones are not usually arrested in differentiation¹⁷⁻¹⁹. However, with

106 acquisition of additional, transforming mutations and epigenetic alterations, clones fail to
107 complete maturation. In the fully transformed state, haemopoiesis in human AML is
108 dominated by expansion either of progenitor-like cells, presumably because of a
109 differentiation block between progenitor and downstream precursor cells²⁰, or less
110 commonly, by precursor-like cells, presumably because of a differentiation block between
111 precursor and mature cells²¹. In both cases, expanded leukemic progenitor-like, or
112 precursor-like, populations have functional leukemic stem cell activity^{20,21}. Thus, we set out
113 to address three questions: (i) where are individual clones arrested in the hematopoietic
114 hierarchy; (ii) which clone(s) responded to mutant IDH2 inhibition by differentiating; (iii) which
115 clone(s) were responsible for loss of response to enasidenib, after an initial response, and
116 by what mechanism?

117 The experimental approach is set out in **Supplementary Fig. 2b**. We first performed flow
118 cytometric quantitation of hematopoietic stem/progenitor (Lin⁻CD34⁺CD117⁺), precursor (Lin⁻
119 CD34⁻CD117⁺) and mature myeloid cell (Lin⁻CD34⁻CD117⁻) populations²¹ at trial entry in 15
120 patients (**Fig. 1a-b and Supplementary Fig. 3a-b**). Sizes of individual stem/progenitor
121 populations within Lin⁻CD34⁺ cells were also quantitated. 11/15 patients had abnormally
122 expanded progenitor-like compartments (mainly LMPP-like and GMP-like; termed progenitor
123 AML) and 4/15 abnormally large myeloid precursor-like populations (termed precursor AML).
124 The ratio of progenitor: precursor AML is consistent with previous studies^{20,21}.

125 Next, we analysed the bone marrow (BM) stem/progenitor/precursor populations in 5
126 patients who achieved complete remission (CR) with enasidenib (**Fig. 1b-c**). In all 5 there
127 was near normalization of the sizes of the stem/progenitor compartments. Pre-treatment, 2
128 patients had pathologically expanded LMPP- and GMP-like progenitor populations (#201-
129 023, #201-011) and 2 patients had expanded myeloid precursor compartments (#201-010
130 and #203-002). Functionally, Lin⁻CD34⁺ progenitor cells from these 5 patients in CR formed
131 myeloid/ erythroid colonies nearly as efficiently as normal cells, in contrast to cells from
132 patients who did not achieve CR (**Fig. 1d**)(**Supplementary Fig. 3c-d**). Thus, enasidenib

133 therapy rebalanced the sizes of hematopoietic stem/progenitor/precursor/mature populations
134 at CR with reacquisition of normal myeloid progenitor function.

135 **Wild type hemopoiesis occurs occasionally with enasidenib therapy**

136 Next, we investigated the clonal basis of differentiation. In principle, enasidenib could have,
137 directly or indirectly, restored differentiation from either wild type cells, ancestral clones or
138 terminal clones in a clonal hierarchy (**Fig. 2a**). We established the clonal basis of response
139 in 6 patients using samples taken at multiple time points before and through treatment,
140 including relapse (**Fig. 2b**). We used WES and karyotype of bone marrow mononuclear cells
141 (BMMNCs) to determine chromosomal copy number and mutational changes
142 (**Supplementary Tables 2-3**). Next, we used WES data to design patient-specific mutation
143 panels to test variant allele frequencies (VAFs) of mutations in unsorted BMMNCs and flow-
144 cytometric sorted hematopoietic stem/progenitor/precursor/mature cell populations and
145 genotype flow-cytometric sorted single cells and hematopoietic colonies derived from single
146 cells (**Supplementary Table 3**). A combination of all these data was used to establish clonal
147 structures. Details on setting false positive and negative thresholds in single cell genotyping
148 (SCG) are presented in Methods.

149 In our initial study, 9 out of 29 patients who achieved CR, for whom samples were available,
150 had loss of mutant *IDH2* peripheral blood cells (complete molecular remission in peripheral
151 blood¹⁶). However, it was unclear if these differentiated blood cells were truly wild type, or
152 from a genetically mutant clone(s) that just lacked mutant *IDH2*? We studied this question in
153 a patient with mutations in *IDH2* (*I*), *PEX26* (*P*), *FEZ2 S208T* (*F*), *DNMT3B* (*D*), *ZCCHC1*
154 (*Z*), *NPM1* (*N*), and *ELMO3* (*E*) in AML blasts pre-enasidenib (#201-022, **Fig. 2c**) who
155 achieved mutant *IDH2* molecular remission. Pre-therapy, imputation from variant allele
156 frequency (VAF) suggested presence of wildtype cells and three possible mutant clones: a
157 clone with *IDH2* mutation alone (*I*), a clone with genotype *IPFDZN* and a minor clone that
158 was either *IPFDZNE* or *IE* (**Supplementary Fig. 4a-c**), though the exact clonal structure
159 could not established unambiguously (**Supplementary Fig. 4d-h**). At CR, the VAF of all

160 mutations was <1.6% (**Fig. 2c, Supplementary Table 4 for depth of sequencing**).
161 Concordantly, at CR the majority (94.6%) of 111 individually genotyped hematopoietic
162 colonies did not contain any mutations present pre-therapy (**Fig. 2d**). Functionally, the
163 colonies produced a normal ratio of myeloid to erythroid colonies consistent with a wild-type
164 genotype (**Fig. 2e**). Thus, in a minority of patients, enasidenib therapy resulted in restoration
165 of wild type terminal blood cell production and progenitor function from wild-type cells.

166 **Enasidenib restored differentiation from ancestral and terminal clones in a clone** 167 **dependent manner**

168 In one patient enasidenib promoted differentiation from an ancestral clone (#201-023, **Fig.**
169 **3**). WES and targeted resequencing of BMMNCs pre-enasidenib detected mutations in
170 *SRSF2* (S), *IDH2* (I), *ASXL1* (A), *GATA2* (G) and two mutations in *RUNX1* (R, r)
171 (**Supplementary Tables 2-3**). Pre-therapy imputation of clonal structure based on VAF in
172 unsorted BMMNCs by exome and targeted resequencing suggested initial acquisition of
173 *SRSF2* mutation (clone S) followed by an *IDH2* mutation (clone SI) followed by acquisition of
174 *ASXL1* (A), the two *RUNX1* and *GATA2* mutations (**Supplementary Fig. 5a and**
175 **Supplementary Tables 2-3**). At CR, only mutations *SRSF2* (S), *IDH2* (I), *ASXL1* (A) were
176 detected in mature cells supporting the existence of an SIA clone that preferentially
177 completes terminal maturation in presence of enasidenib (**Supplementary Fig. 5a-c**).

178 To clarify clonal structure, and position clones within the haemopoietic hierarchy, we
179 performed targeted resequencing for driver variants in 63 single cells and cell populations,
180 from flow-cytometric sorted progenitor and mature myeloid cell compartments, from both
181 pre-therapy and CR samples (**Fig. 3a-b and Supplementary Table 3**). This confirmed a
182 linear clonal evolution pattern: clone SI preceded clone SIA, followed by clones that
183 sequential acquired *RUNX1* mutations (clones SIAR and SIARr) and finally the terminal
184 clone acquired a *GATA2* mutation (SIARrG). The mutational profile in 4 single cells did not fit
185 into this clonal evolution pathway. We detected a single cell with an *ASXL1* mutation (clone
186 A) and three single cells with both *ASXL1* and *SRSF2* mutations (clone AS). In three cells

187 with genotype AS, we detected allele dropout (ADO) of the *IDH2* allele in two out of three
188 cells. Thus, we are unable to determine the mutational state of *IDH2* in those two cells. In
189 contrast, in twelve cells of the SI clone, ADO of the *ASXL1* allele was detected in only three
190 out of twelve cells (**Supplemental Fig. 12a**). Thus, though our results do not exclude a rarer,
191 parallel clonal evolution pathway where clones A and AS exist and failed to acquire
192 mutations in the order shown in the main pathway, they also are consistent with these cells
193 being part of the main clonal evolution pathway.

194 Pre-therapy, 90% of BMMNCs were progenitors (LMPP and GMP) (**Fig. 1c and 3d**) and 90-
195 100% of these leukemic progenitors were the SIARrG clone (**Fig 3c**). Thus, the SIARrG
196 clone is arrested in differentiation at the progenitor stage and expands to dominate the
197 marrow. Less than 10% of BMMNCs were mature myeloid cells (**Fig 1c and Fig 3d**). We
198 were only able to genotype 8 mature myeloid cells pre-therapy (**Fig. 3c**) and they are
199 composed of a mixture of wild type cells and of cells with genotypes A, SI, AS and SIA. At
200 CR, the mature myeloid cell compartment comprised 60% of BMMNCs (**Fig 1c and Fig 3d**)
201 and 85% of mature cells have the *SIA* genotype (**Fig. 3d and Supplementary Fig. 5b,**
202 **Supplementary Table 3**).

203 The progenitor compartment was only 20% of BMMNCs at CR (**Fig. 1c and Fig 3d**) and
204 composed of mixed ancestral clones SI, SIA, SIAR but not the terminal clone SIARrG.
205 Concordantly, the majority of colonies generated by progenitors at CR were “SIA”, with a
206 minority of the SI and SIA genotype (**Fig. 3e**). The ratio of myeloid:erythroid colonies was
207 within normal limits (**Fig. 3f**) Taken together, this data demonstrated a complex, clone-
208 dependent pattern of enasidenib-induced differentiation with mature myeloid cell production
209 sustained principally by a self-renewing ancestral *SIA* clone.

210 In four patients, differentiation of mature cells was principally seen from terminal clones (**Fig.**
211 **2b**). In patient #201-011, WES (**Supplementary Table 2**) and targeted resequencing of
212 BMMNC revealed two mutations in *DNMT3A* and mutations in *IDH2*, *ASXL1* and *XPO1* pre-
213 therapy that persisted at different VAFs at CR (**Supplementary Table 3, Supplementary**

214 **Fig. 6a**). However, it was not possible to impute the clonal structure from the VAF
215 (**Supplementary Fig. 6b**). We genotyped 110 single cells, pre-therapy and at CR (**Fig. 4a**).
216 This revealed an initiating *DNMT3A* clone (clone D) that acquired an *IDH2* mutation (clone
217 DI) (**Fig. 4b**). Subsequently, there is a branching clonal structure with two terminal clones;
218 one acquired an *XPO1* mutation (clone DIX), whereas the other acquired two mutations, a
219 second *DNMT3A* mutation and an *ASXL1* mutation (clone DIdA). SCG suggested the DIdA
220 clone may have arisen by convergent evolution through intermediate DId and DIA clones.
221 There is a caveat with this interpretation as ADO was detected in 6 out of 7 DIA clone cells
222 (**Supplemental Fig. 12d**). In the DId cells, though there were no heterozygous germline
223 single nucleotide polymorphisms in the *ASXL1* gene, the estimated ADO frequency of the
224 *ASXL1* allele was 12.1% (**Supplemental Fig. 12e**). Thus, it is also possible that the DIdA
225 clone may also have arisen through just one mutational pathway.

226 Pre-enasidenib, 89% of BMMNCs were leukemic progenitors (LMPP and GMP) virtually
227 exclusively composed of the DIX clone (**Fig. 4c-d**). A small mature myeloid population is
228 present pre-enasidenib composed of the DIdA clone (**Supplementary Fig. 6c**). These
229 observations suggest that the *IDH2* mutation in the context of the DIdA clone is not fully
230 effective at imposing a complete differentiation block, whereas the same *IDH2* mutation in
231 the context of the DIX clone fully arrests at a progenitor stage.

232 At CR, 82% of BMMNCs were composed of mature myeloid cells, 85% of which were a mix
233 of two terminal branching clones DIdA (54%) and DIX (31%) suggesting enasidenib
234 promoted differentiation from both terminal clones (**Fig. 4c-d, Supplementary Table 3**). To
235 address which progenitors contribute to mature cell output we genotyped single flow
236 cytometric sorted progenitors. The DIdA clone dominated mature GMP, CMP and MEP
237 progenitor compartments (**Fig. 4c-d**). In contrast, clone DIX was detected only in the more
238 immature LMPP progenitor compartment. GMP, CMP and MEP are more clonogenic than
239 the LMPP^{20,22} and concordantly, ~95% of colonies had the DIdA genotype (**Fig. 4e**) that
240 were myeloid-biased (**Fig. 4f**). Interestingly, there is a substantial decrease in the size of the

241 LMPP compartment at CR compared to pre-therapy (**Fig. 4d**) (1240-fold decrease within the
242 Lin⁻ compartment and 81-fold within the CD34⁺ compartment).

243 In three additional patients the terminal clone contributed to mature myeloid cells at CR
244 based on imputed clonal structures, genetic analysis of mature myeloid cells at CR (patient
245 #201-010, **Supplementary Fig. 6d-e**) and genotyping of myeloid colonies at CR (patient
246 #201-027, **Supplementary Fig. 7a-d**; patient #201-006 **Supplementary Fig. 7e-h**). Depth of
247 coverage for each of the mutations in all three patients is in **Supplementary Table 4**.

248 In summary, enasidenib therapy provides relief of differentiation arrest at a progenitor-like or
249 precursor-like stage, normalizing the sizes of these abnormally expanded compartments.
250 The ability of mutant *IDH2* to impose differentiation block is dependent on the context of co-
251 associated mutations within a clone. Consequently, efficacy of enasidenib-induced
252 differentiation is also likely to dependent mutational landscape within a clone.

253 **Relapse of *IDH2* mutant patients on enasidenib occurs by clonal selection/evolution**
254 **and not second site mutations in *IDH2* gene**

255 Although responding patients have a median survival of 18-21 months, many patients
256 relapse¹⁵. To study mechanisms leading to relapse we measured 2-HG levels in 16 patients
257 at diagnosis and relapse, mutational profiles in 12 patients (by WES in 11 cases and
258 targeted sequencing in the other patient) and performed karyotype analysis in 11 subjects
259 (**Fig. 5a-b** and **Supplementary Table 5**). We did not detect second site *IDH2* mutations at
260 relapse in any patient but instead documented 7 patterns of clonal evolution/selection with
261 acquisition of recurrent AML-associated genetic changes (**Fig. 5a**). These are recurrent
262 missense mutations in myeloid malignancy, or nonsense and frameshift mutations in
263 cancers, as documented in the COSMIC database (<http://cancer.sanger.ac.uk/cosmic>). For
264 patient #201-007, mutations were detected prior to relapse, but increased in frequency at
265 relapse (Supplementary Table 3).

266 In 14/16 patients 2-HG levels remained suppressed between best response (CR or PR) and
267 relapse suggesting drug was on target in suppressing neomorphic enzyme function (**Fig.**
268 **5b**). However, in 2 patients (#201-014 and #201-022) rising 2-HG levels and BM leukemic
269 cells (blasts) were seen (**Fig. 5c-d**). Exome sequencing revealed *IDH1* R132C/H mutations
270 albeit accompanied by other genetic abnormalities, some of which are recurrent in AML
271 (point mutations in *RUNX1*, *NPM1* and *t(3:12)*) (**Fig. 5e-f**). These *IDH1* mutations were
272 previously undetectable by high depth NGS (10000x) pre-enasidenib therapy. Surprisingly,
273 in both cases, the VAF indicates that *IDH1* mutations were present in *IDH2* mutant clones.

274 Relapse also associated with increasing VAF of oncogenic gain of function mutations in
275 genes encoding cytokine receptors *CSF3R* (patient #104-021, **Supplementary Fig. 8a-b**)
276 and *FLT3* (patients #201-013, #201-004 **Supplementary Fig. 8c-d** and #201-007
277 **Supplementary Fig. 9f-h**) and predicted loss of function mutation in the negative regulator
278 of cytokine signalling *CBL* (#201-004 **Supplementary Fig. 8d**). For patient #104-021 we
279 could not resolve the clonal structure from targeted resequencing of BMMNC
280 (**Supplementary Fig. 8b, Supplementary Table 2**) but genotyping of 214 single cells pre-
281 enasidenib and at relapse (**Fig. 6a**) demonstrated an initiating *DNMT3A/ IDH2* mutant clone
282 that spawned the major clone with a recurrent *U2AF1* mutation (DIU clone). A minor DIUF
283 clone with a D200E variant in the *FLT3* (not previously described in AML) was also present
284 pre-enasidenib. At relapse, the major clone in the expanded arrested LMPP and GMP
285 compartments had acquired an oncogenic T618I mutation in the cytokine receptor *CSF3R*
286 that is well described in myeloid leukemias²³ (*DIUC clone*). DIUC further evolved, acquiring a
287 variant in *NFKB1* that has not been described previously in AML. Mutations in both *CSF3R*
288 and *NFKB1* were detectable at threshold of sensitivity pre-enasidenib (**Supplementary Fig.**
289 **8a**).

290 Relapse, and re-imposition of differentiation block, was also associated with previously
291 described mutations in hematopoietic transcription factors in myeloid cancers. These
292 included frame shift mutations in *RUNX1* (**Fig. 5e and Supplementary Fig. 8e** one of which

293 has been previously described in AML (*RUNX1* F416fs)²⁴) and *BCORL1*²⁵; non-synonymous
294 variants, in the DNA- and protein-partner binding N-terminal zinc finger of *GATA2*
295 (**Supplementary Fig. 8d**)^{26,27} and in one of the zinc fingers of *BCL11A*²⁸ (**Supplementary**
296 **Fig. 8c**). In all patients these mutations were not detected pre-enasidenib.

297 Deletion of all (monosomy 7), or part (del 7q) of chromosome 7 is common in myeloid
298 malignancy²⁹. Chromosomal abnormalities were present in 18% of the enasidenib cohort¹⁵
299 and 20% in the cohort studied here (**Supplementary Fig 1**). Del 7q was detected in 4 out of
300 12 patients who relapsed (#201-010, #201-007, #201-019 and #201-003) (**Fig. 5a**,
301 **Supplementary Table 5**) but are not enriched in relapsed patients (Amatangelo, M.,
302 Thakurta, A., unpublished data). In all four patients it was detected either cytogenetically, or
303 by WES, pre-enasidenib therapy (**Supplementary Tables 2 and 5**). In three out of the four
304 cases the del 7q clone was selected at relapse (**Supplementary Fig. 9a-e, k and l**). In one
305 case where it was not, it was the dominant clone pre-therapy and at relapse
306 (**Supplementary Fig. 9g-j**).

307 Clonal evolution at relapse also highlighted variants in genes less well studied in AML
308 including *NFKB1* M216I (**Fig. 6a-d**), *DDX1* G699A (**Supplementary Fig. 8c**), *MTUS1*
309 Q781H (**Supplementary Fig. 9f-i**), *DHX15* R222G and *DEAF1* N372K (**Fig. 6e-h**). Of these,
310 acquisition of latter two variants by patient #201-011 at relapse is worthy of comment. Pre-
311 enasidenib the patient had expanded LMPP and GMP populations composed of an arrested
312 DIX clone (**Fig. 4b**). At CR the DIX clone was able to differentiate but only persisted within
313 the LMPP compartment (**Fig. 4d**). At relapse, we detected 30 cells with DIX mutations that
314 acquired a missense mutation in the DExD/H-box helicase *DHX15* R222G (mutation H), 7
315 cells with the DIX mutations that had acquired a variant in the transcription factor *DEAF1*
316 N372L (variant F) and 12 cells with the DIXHF variants (**Fig. 6e-f**). However in 6 out of 7
317 DIXF cells there was ADO for the *DHX15* allele with the *DHX15* R222G mutation and
318 therefore we are unable to determine whether the *DHX15* R222G is present in those cells
319 (**Supplementary Fig. 12c**).

320 Acquisition of the additional *DHX15* and *DEAF1* mutations was associated with
321 differentiation arrest, and a re-expansion of LMPP and GMP progenitor compartments,
322 comprising 89% of MNCs. 83% of LMPP cells were composed of the DIXH and DIXHF
323 clones whereas in the GMP-like compartment there was a more even contribution by the
324 *DIXHF*, *DIXH* and *DIXF* clones (**Fig. 6g-h**).

325 Finally, both *DHX15* and *DDX1* regulate RNA splicing. Human *DHX15* is structurally closely
326 related to its yeast homologue, Prp43^{30,31}. Both proteins have been shown to contribute to
327 disassembly of spliceosomes, efficient debranching and turnover of excised introns^{32,33}.
328 *DHX15* R222G mutations have been previously described in AML^{34,35}. To determine if
329 acquisition of *DHX15* R222G results in altered splicing we performed RNA-Seq of AML
330 blasts at relapse from patient #201-011 and compared splicing to AML cells from the same
331 patient that were wild type for *DHX15* at trial entry (**Supplementary Fig. 8f**). In cells
332 expressing *DHX15* R222G, there were alterations in exon skipping and intron retention
333 compared to cells wild type for *DHX15*. *DDX1* is a DEAD-box RNA helicase with 5' single
334 stranded RNA exonuclease activity postulated to have multiple roles in RNA metabolism³⁶.
335 *DDX1* G699A has not been previously described as a cancer-associated mutation nor has
336 its impact on RNA splicing been studied. Our data shows an increase in intron retention and
337 use of alternative 5' and 3' splice sites, and a decrease in spliced exons in mutant compared
338 to wild type AML cells from the same patient (**Supplementary Fig. 8g**).

339 **Discussion**

340 This study of clonal response and acquired resistance in sequential paired samples from
341 AML patients treated with an IDH2 inhibitor extends prior preclinical studies^{13,14}. Pre-
342 enasidenib there were complex patient- and clone-specific patterns of differentiation arrest.
343 At CR, wildtype dominated cellular reconstitution was less common but does occur. More
344 commonly, enasidenib causes a clone-specific differentiation response, either from ancestral
345 or terminal clones, leading to near normalization of the sizes and functionality of progenitor
346 and precursor hematopoietic compartments with altered clonal mix. Acquired resistance was

347 never due to a second site mutation in the same *IDH2* allele but instead due to either clonal
348 evolution or clonal selection. At least 7 different mutational mechanisms led to re-imposition
349 of differentiation arrest (**Supplementary Fig.10**).

350 In most patients enasidenib was unable to promote terminal differentiation and eradication of
351 *IDH2* mutant clones; ancestral and/or terminal clones remained at CR. In patients with
352 restitution of wild type hemopoiesis, we infer that enasidenib most likely promoted terminal
353 differentiation of arrested self-renewing *IDH2* mutant cells, allowing normal cells to dominate
354 hemopoiesis. Longer term clinical follow up of molecular CR patients will determine if
355 molecular CR patients have a better clinical outcome and if they relapse, the clonal origin of
356 relapse. More generally, understanding the molecular mechanisms of relief from
357 differentiation arrest by *IDH2* inhibitors will require in depth study of changing patterns of
358 epigenetic marks and transcriptional programmes within highly purified, clone-specific,
359 hematopoietic stem, progenitor and precursor populations, before and after drug exposure,
360 as transcriptional and epigenetic profiles are so highly plastic through differentiation.

361 Drug resistance to targeted cancer therapy arises by multiple mechanisms. Resistance to
362 kinase inhibitors in AML^{37,38}, chronic myeloid leukemia³⁹, chronic lymphoid leukemia⁴⁰ and
363 lung cancer^{41,42} often involves second site mutations in the mutant allele modulating drug or
364 substrate binding or copy number changes of the mutant kinase. We did not observe this in
365 enasidenib treated patients.

366 Acquired resistance led to *IDH2* could arise by either epigenetic or genetic mechanisms or a
367 combination of the two. In most patients 2HG remained suppressed at relapse suggesting
368 that enasidenib remained on target and relapsed clones were not dependent on mutant
369 *IDH2*. In hematologic malignancies, genome-wide epigenetic variation (DNA methylation for
370 example) can be several orders more variable than genetic change⁴³⁻⁴⁵, is somatically
371 heritable, and subject to selection. Locus-specific DNA methylation (epiallele) variation
372 shows dynamic change in AML between diagnosis and relapse and can occur with distinct
373 kinetics, such that some patients have a high epiallele diversity and low somatic mutation

374 burden and vice versa⁴⁵. In melanoma, resistance to a B-RAF inhibitor arose in rare cells
375 through stochastic, transient variation in gene expression that was selected for by therapy⁴⁶.
376 This is consistent with prior work on chromatin mediated drug resistance in cancer cell
377 lines⁴⁷.

378 Clones acquiring gene mutations, or grosser genetic changes, have previously been
379 reported in therapy resistant chronic lymphocytic leukemia⁴⁸ and medulloblastoma⁴⁹.
380 Acquisition of an *IDH1* mutation in two patients is an example of how this may occur. Here,
381 these AML propagation is likely to be highly dependent to high 2HG. 2HG addiction may be
382 AML cell autonomous or alternatively, the 2HG requirement may be in BM niche supporting
383 cells or other non-AML cell populations. AML clone-specific and non-AML cell specific
384 analysis of the impact of 2HG on epigenetic and transcriptional programs and metabolism of
385 cells⁵⁰ is needed to understand this dependency. Other examples genetic changes leading
386 to enasidenib resistance include gain of function mutations in proliferative cytokine signalling
387 pathways and loss of or altered function in transcriptional regulators of hemopoiesis.
388 However, mutations in the RAS pathway that are correlate with failure of initial response¹⁶
389 were not associated with acquired resistance.

390 We also detected variants at relapse not previously well studied in AML. An example of this
391 is the *DHX15* R222G mutation, recently described in *RUNX1-RUNX1T1* AML³⁴. The yeast
392 homologue of *DHX15*, Prp43, regulates RNA splicing and ribosome biogenesis. Loss of wild
393 type *DHX15* and overexpression of mutant *DHX15* increases alternative splicing. In contrast,
394 a role for *DEAF1* has not been previously published in normal or malignant hemopoiesis.
395 Curiously *DEAF1* is a paralog of the transcription factor, *RUNX1T1*. *DEAF1* is expressed
396 throughout hemopoiesis but particularly in GMP and AML blasts²¹. In non-hemopoietic
397 tissues it binds to *LMO4*, a member of the *LMO* transcriptional adapter protein family. In
398 blood cells, *LMO2*, a closely related *LMO* family member, partners transcription factors
399 including *GATA*, *E-box* proteins and *LDB1* to form regulatory and oncogenic protein
400 complexes. Thus, the role of *DEAF1* in hemopoiesis and its interaction with *DHX15* merits

401 further study. More broadly, a deeper mechanistic understanding of how wild-type IDH2
402 promote hematopoietic differentiation, which is currently poorly understood, will also
403 increase our understanding of how bypass pathways could re-impose differentiation block,
404 for example by altering transcriptional programmes, as seen in BET inhibitor therapy
405 resistance⁵¹.

406 Finally, this study demonstrates how any cancer therapy alters clonal structure across a fully
407 transformed and pre-malignant cellular hierarchy. By defining clonal structures and mapping
408 where clones were arrested across differentiation, in purified hematopoietic compartments,
409 we obtained a previously unavailable view of where different clones were arrested. This
410 provides the necessary information to study why clones are arrested at different stages of
411 differentiation. Furthermore, analysis of sequential samples through therapy shows how
412 clones differentially responded to therapy. This provides the basis to study clone specific
413 relief of IDH2 inhibition. More generally, our approach could be applied to any cancer
414 therapy, where single cell suspensions and purification of cells at different stages of
415 differentiation is possible. This would then provide a clone specific understanding of how
416 therapy alters clonal structure through a cellular hierarchy. Our approach also provides
417 insights towards a rational basis for combination therapies to reduce drug resistance.

418 **Acknowledgments:** We thank patients and clinical staff for samples studied. L.Q. was
419 supported by an Oxford-Celgene Fellowship; P.V. acknowledges funding from the MRC
420 Disease Team Awards (G1000729/94931 and MR/L008963/1) MRC Molecular Haematology
421 Unit and the Oxford Partnership Comprehensive Biomedical Research Centre (NIHR BRC
422 Funding scheme. oxfbrc-2012-1). V.P.-L. and S.dB. acknowledge funding by the French
423 National Institute of Health (INSERM-AVIESAN), the National Cancer Institute (INCa-DGOS-
424 Inserm_6043 and INCa 2012-1-RT-09), SIRIC-SOCRATE 2.0 and the Fondation Association
425 pour la Recherche sur le Cancer (ARC, Programme). M.D.D. is funded by a fellowship from
426 the Institut National du Cancer (INCa-DGOS_5733). M.H. is a fellow of the Fondation
427 Philanthropia, Ecole des Sciences du Cancer, Gustave Roussy, Villejuif, France. We

428 acknowledge the Core Flow Cytometry and Next Generation Sequencing Facilities at the
429 WIMM; the Imaging, Cytometry and Integrated Biology platforms at Gustave Roussy
430 (Philippe Rameau, Yann Lecluse, Nathalie Droin, M'boyba Khadija Diop, UMS AMMICA);
431 clinical departments at Gustave Roussy (Jean-Baptiste Micol for clinical specimens, Nathalie
432 Auger for cytogenetic analyses and Christophe Marzac and Edwidge Leclercq for FLT3
433 genotyping) and Memorial Sloan Kettering. R.L.L. is supported by grants from the NIH,
434 including R35 CA197594-01A1 and the Memorial Sloan Kettering Cancer Center Support
435 Grant (NIH P30 CA008748, including a supplement to R.L.L.). The views expressed are
436 those of the authors and not necessarily those of the NHS, the NIHR or the Department of
437 Health.

438 **Author contributions:**

439 L.Q., M.D.D. designed/ performed experiments, analysed data; A.K., M.M., M.A., B.S., C.Q.,
440 M.H., C.W., V.S., S.AI. performed experiments/ analysed data; M.S.V. and G.S.V. analysed
441 data; M.A., A.S., A.P., K.Y., S.Ag., S.dB., R.L.L., E.S., K.M., A.T. provided
442 reagents/samples/clinical data; O.A.B., S.dB., A.T., R.L.L., V.P.-L. and P.V. designed the
443 experiments/ analysed the data. L.Q. and P.V. wrote the manuscript. All authors edited the
444 manuscript.

445

446 **Competing interests:**

447 PV has received research grant support from Celgene and is on its speaker bureau. LQ has
448 received research grant support from Celgene

449 **Data and materials availability:** Exome sequencing data has been deposited in
450 ArrayExpress under accession E-MTAB-6299. RNA sequencing data has been deposited in
451 ArrayExpress under accession E-MTAB-6660.

452 **References**

453 1. Abbosh, C., *et al.* Phylogenetic ctDNA analysis depicts early-stage lung cancer
454 evolution. *Nature* **545**, 446-451 (2017).

- 455 2. Jamal-Hanjani, M., *et al.* Tracking the Evolution of Non-Small-Cell Lung Cancer. *N Engl J Med* **376**, 2109-2121 (2017).
456
- 457 3. Cancer Genome Atlas Research, N. Genomic and epigenomic landscapes of adult
458 de novo acute myeloid leukemia. *N Engl J Med* **368**, 2059-2074 (2013).
- 459 4. Paschka, P., *et al.* IDH1 and IDH2 mutations are frequent genetic alterations in acute
460 myeloid leukemia and confer adverse prognosis in cytogenetically normal acute
461 myeloid leukemia with NPM1 mutation without FLT3 internal tandem duplication. *J
462 Clin Oncol* **28**, 3636-3643 (2010).
- 463 5. Green, C.L., *et al.* The prognostic significance of IDH2 mutations in AML depends on
464 the location of the mutation. *Blood* **118**, 409-412 (2011).
- 465 6. Craddock, C.F., *et al.* Outcome of Azacitidine Therapy in Acute Myeloid Leukemia Is
466 not Improved by Concurrent Vorinostat Therapy but Is Predicted by a Diagnostic
467 Molecular Signature. *Clin Cancer Res* (2017).
- 468 7. Xu, W., *et al.* Oncometabolite 2-hydroxyglutarate is a competitive inhibitor of alpha-
469 ketoglutarate-dependent dioxygenases. *Cancer Cell* **19**, 17-30 (2011).
- 470 8. Lu, C., *et al.* IDH mutation impairs histone demethylation and results in a block to cell
471 differentiation. *Nature* **483**, 474-478 (2012).
- 472 9. Figueroa, M.E., *et al.* Leukemic IDH1 and IDH2 mutations result in a
473 hypermethylation phenotype, disrupt TET2 function, and impair hematopoietic
474 differentiation. *Cancer Cell* **18**, 553-567 (2010).
- 475 10. Losman, J.A., *et al.* (R)-2-hydroxyglutarate is sufficient to promote leukemogenesis
476 and its effects are reversible. *Science* **339**, 1621-1625 (2013).
- 477 11. Wang, F., *et al.* Targeted inhibition of mutant IDH2 in leukemia cells induces cellular
478 differentiation. *Science* **340**, 622-626 (2013).
- 479 12. Kats, L.M., *et al.* Proto-oncogenic role of mutant IDH2 in leukemia initiation and
480 maintenance. *Cell Stem Cell* **14**, 329-341 (2014).
- 481 13. Yen, K., *et al.* AG-221, a First-in-Class Therapy Targeting Acute Myeloid Leukemia
482 Harboring Oncogenic IDH2 Mutations. *Cancer discovery* **7**, 478-493 (2017).

- 483 14. Shih, A.H., *et al.* Combination Targeted Therapy to Disrupt Aberrant Oncogenic
484 Signaling and Reverse Epigenetic Dysfunction in IDH2- and TET2-Mutant Acute
485 Myeloid Leukemia. *Cancer discovery* **7**, 494-505 (2017).
- 486 15. Stein, E.M., *et al.* Enasidenib in mutant IDH2 relapsed or refractory acute myeloid
487 leukemia. *Blood* **130**, 722-731 (2017).
- 488 16. Amatangelo, M.D., *et al.* Enasidenib induces acute myeloid leukemia cell
489 differentiation to promote clinical response. *Blood* **130**, 732-741 (2017).
- 490 17. Jan, M., *et al.* Clonal evolution of preleukemic hematopoietic stem cells precedes
491 human acute myeloid leukemia. *Sci Transl Med* **4**, 149ra118 (2012).
- 492 18. Corces-Zimmerman, M.R., Hong, W.J., Weissman, I.L., Medeiros, B.C. & Majeti, R.
493 Preleukemic mutations in human acute myeloid leukemia affect epigenetic regulators
494 and persist in remission. *Proc Natl Acad Sci U S A* **111**, 2548-2553 (2014).
- 495 19. Shlush, L.I., *et al.* Identification of pre-leukaemic haematopoietic stem cells in acute
496 leukaemia. *Nature* **506**, 328-333 (2014).
- 497 20. Goardon, N., *et al.* Coexistence of LMPP-like and GMP-like leukemia stem cells in
498 acute myeloid leukemia. *Cancer Cell* **19**, 138-152 (2011).
- 499 21. Quek, L., *et al.* Genetically distinct leukemic stem cells in human CD34- acute
500 myeloid leukemia are arrested at a hemopoietic precursor-like stage. *J Exp Med* **213**,
501 1513-1535 (2016).
- 502 22. Karamitros, D., *et al.* Human lympho-myeloid progenitors are heterogeneous at the
503 single cell level. *Nature Immunology* **19**, 85-97 (2018).
- 504 23. Maxson, J.E., *et al.* Oncogenic CSF3R mutations in chronic neutrophilic leukemia
505 and atypical CML. *N Engl J Med* **368**, 1781-1790 (2013).
- 506 24. Gaidzik, V.I., *et al.* RUNX1 mutations in acute myeloid leukemia: results from a
507 comprehensive genetic and clinical analysis from the AML study group. *J Clin Oncol*
508 **29**, 1364-1372 (2011).
- 509 25. Li, M., *et al.* Somatic mutations in the transcriptional corepressor gene BCORL1 in
510 adult acute myelogenous leukemia. *Blood* **118**, 5914-5917 (2011).

- 511 26. Papaemmanuil, E., *et al.* Genomic Classification and Prognosis in Acute Myeloid
512 Leukemia. *N Engl J Med* **374**, 2209-2221 (2016).
- 513 27. Fasan, A., *et al.* GATA2 mutations are frequent in intermediate-risk karyotype AML
514 with biallelic CEBPA mutations and are associated with favorable prognosis.
515 *Leukemia* **27**, 482-485 (2013).
- 516 28. Mason, C.C., *et al.* Age-related mutations and chronic myelomonocytic leukemia.
517 *Leukemia* **30**, 906-913 (2016).
- 518 29. Grimwade, D., *et al.* Refinement of cytogenetic classification in acute myeloid
519 leukemia: determination of prognostic significance of rare recurring chromosomal
520 abnormalities among 5876 younger adult patients treated in the United Kingdom
521 Medical Research Council trials. *Blood* **116**, 354-365 (2010).
- 522 30. Tauchert, M.J., Fourmann, J.B., Luhrmann, R. & Ficner, R. Structural insights into the
523 mechanism of the DEAH-box RNA helicase Prp43. *eLife* **6**(2017).
- 524 31. Murakami, K., Nakano, K., Shimizu, T. & Ohto, U. The crystal structure of human
525 DEAH-box RNA helicase 15 reveals a domain organization of the mammalian
526 DEAH/RHA family. *Acta Crystallogr F Struct Biol Commun* **73**, 347-355 (2017).
- 527 32. Arenas, J.E. & Abelson, J.N. Prp43: An RNA helicase-like factor involved in
528 spliceosome disassembly. *Proc Natl Acad Sci U S A* **94**, 11798-11802 (1997).
- 529 33. Fourmann, J.B., *et al.* Dissection of the factor requirements for spliceosome
530 disassembly and the elucidation of its dissociation products using a purified splicing
531 system. *Genes Dev* **27**, 413-428 (2013).
- 532 34. Faber, Z.J., *et al.* The genomic landscape of core-binding factor acute myeloid
533 leukemias. *Nat Genet* **48**, 1551-1556 (2016).
- 534 35. Farrar, J.E., *et al.* Genomic Profiling of Pediatric Acute Myeloid Leukemia Reveals a
535 Changing Mutational Landscape from Disease Diagnosis to Relapse. *Cancer Res* **76**,
536 2197-2205 (2016).
- 537 36. Cordin, O., Banroques, J., Tanner, N.K. & Linder, P. The DEAD-box protein family of
538 RNA helicases. *Gene* **367**, 17-37 (2006).

- 539 37. Cools, J., *et al.* Prediction of resistance to small molecule FLT3 inhibitors:
540 implications for molecularly targeted therapy of acute leukemia. *Cancer Res* **64**,
541 6385-6389 (2004).
- 542 38. Heidel, F., *et al.* Clinical resistance to the kinase inhibitor PKC412 in acute myeloid
543 leukemia by mutation of Asn-676 in the FLT3 tyrosine kinase domain. *Blood* **107**,
544 293-300 (2006).
- 545 39. Shah, N.P., *et al.* Multiple BCR-ABL kinase domain mutations confer polyclonal
546 resistance to the tyrosine kinase inhibitor imatinib (STI571) in chronic phase and
547 blast crisis chronic myeloid leukemia. *Cancer Cell* **2**, 117-125 (2002).
- 548 40. Woyach, J.A., *et al.* Resistance mechanisms for the Bruton's tyrosine kinase inhibitor
549 ibrutinib. *N Engl J Med* **370**, 2286-2294 (2014).
- 550 41. Kobayashi, S., *et al.* EGFR mutation and resistance of non-small-cell lung cancer to
551 gefitinib. *N Engl J Med* **352**, 786-792 (2005).
- 552 42. Choi, Y.L., *et al.* EML4-ALK mutations in lung cancer that confer resistance to ALK
553 inhibitors. *N Engl J Med* **363**, 1734-1739 (2010).
- 554 43. Figueroa, M.E., *et al.* DNA methylation signatures identify biologically distinct
555 subtypes in acute myeloid leukemia. *Cancer Cell* **17**, 13-27 (2010).
- 556 44. Kulis, M., *et al.* Epigenomic analysis detects widespread gene-body DNA
557 hypomethylation in chronic lymphocytic leukemia. *Nat Genet* **44**, 1236-1242 (2012).
- 558 45. Li, S., *et al.* Distinct evolution and dynamics of epigenetic and genetic heterogeneity
559 in acute myeloid leukemia. *Nat Med* **22**, 792-799 (2016).
- 560 46. Shaffer, S.M., *et al.* Rare cell variability and drug-induced reprogramming as a mode
561 of cancer drug resistance. *Nature* **546**, 431-435 (2017).
- 562 47. Sharma, S.V., *et al.* A chromatin-mediated reversible drug-tolerant state in cancer
563 cell subpopulations. *Cell* **141**, 69-80 (2010).
- 564 48. Landau, D.A., *et al.* Evolution and impact of subclonal mutations in chronic
565 lymphocytic leukemia. *Cell* **152**, 714-726 (2013).

- 566 49. Morrissy, A.S., *et al.* Divergent clonal selection dominates medulloblastoma at
567 recurrence. *Nature* **529**, 351-357 (2016).
- 568 50. Raffel, S., *et al.* BCAT1 restricts alphaKG levels in AML stem cells leading to
569 IDHmut-like DNA hypermethylation. *Nature* **551**, 384-388 (2017).
- 570 51. Rathert, P., *et al.* Transcriptional plasticity promotes primary and acquired resistance
571 to BET inhibition. *Nature* **525**, 543-547 (2015).

572

573 **Figure Legends**

574 **Figure 1. Enasidenib treatment induces differentiation of AML progenitor and** 575 **precursor cell populations and restores progenitor function.**

576 a) Top, immunophenotyping of hematopoietic stem/progenitor/precursor and mature cell
577 populations in AML bone marrow (BM) samples pre-treatment showing expanded progenitor
578 (n=11 biologically independent samples) or precursor (n=4 biologically independent
579 samples) populations with normal BM (n=8 biologically independent samples). Below,
580 detailed composition of stem/progenitor compartments in AML pre-treatment (n=15
581 biologically independent samples) and normal BM (n=12 biologically independent samples).
582 Error bars in normal BM= 95% confidence interval. HSC: hematopoietic stem cell, MPP:
583 multipotent progenitor, LMPP: lymphoid-primed multipotent progenitor, CMP: common
584 myeloid progenitor, MEP; megakaryocyte-erythroid progenitor; GMP: granulocyte-
585 macrophage progenitor.

586 b) Top, schematic representation of flow cytometric approach and sequential gates used to
587 analyse samples in (a). Lin⁻, lineage negative; BMMC, bone marrow mononuclear cells.
588 Bottom, example of flow plots from a representative sample prior to enasidenib treatment
589 (Pre-ENA) and at complete remission (CR) in patients with expanded progenitor-like
590 populations (#201-023; left, experiment performed once) or expanded myeloid precursor-like
591 population (#201-010; right, experiment performed once). Numbers shown within the gate
592 indicate percentage of the corresponding cell population compared to all cells in the plot.

593 c) Top, immunophenotyping of hematopoietic cell populations in normal BM (left, as in (a))
594 and in samples from 5 patients (#201-023, #201-011, #201-022, #201-010, #203-002) pre-
595 ENA, at intermediate time points during treatment, and at CR. Bottom, sizes of stem and
596 progenitor compartments. Abbreviations and error bars in normal BM are as in (a). C=cycle,
597 D=day.

598 d) Number of mixed erythroid-myeloid colonies (GEM), granulocyte-macrophage (GM),
599 granulocyte (G), macrophage (M) and erythroid (E) colonies produced per 100 plated flow-
600 sorted CD34⁺ cells from normal BM (n=4 biologically independent samples), enasidenib-
601 treated patients in CR (n=5 biologically independent samples) and enasidenib-treated
602 patients not in CR (n=3 biologically independent samples). Patient samples were plated with
603 addition of Enasidenib (1 μ M) to semi-solid media. Error bars= standard error of the mean. P-
604 values determined by 2-sided Student's paired t-test.

605 **Figure 2. Differentiation response arising from wild-type cells in patients treated with**
606 **enasidenib.**

607 a) Schematic representation of varying possible clonal responses to enasidenib. Four
608 mutations (A, B, C and D) are present in four clones that are arranged in a branching
609 structure. A differentiation response to enasidenib treatment could potentially occur from
610 either wild type cells or from ancestral or terminal clones.

611 b) Summary of the type of differentiation response (from either wild type cells, ancestral or
612 terminal clones) in samples from 6 patients.

613 c) Variant allele frequencies (VAF) of the indicated mutations in AML blasts of patient #201-
614 022 prior to enasidenib treatment (Pre-ENA) and in peripheral blood mononuclear cells
615 (PMNC) at CR, as assessed by targeted re-sequencing.

616 d) Clonal contribution to colony output from the CR sample from patient #201-022, as a
617 percentage of all individually picked colonies genotyped. Clones were identified as wild type
618 (WT), carrying the *FEZ2* P118S mutation (F), or carrying the *FEZ2* P118S and *DNMT3B*

619 N738S mutations (FD). Lineage affiliations of the colonies are as in **Fig. 1d**. Numbers next
620 to the bars indicate the number of colonies analyzed.

621 e) Bar graph showing the lineage affiliation of colonies from Lin⁻CD34⁺ normal cord blood
622 (CB) cells (n=5 biologically independent samples) and CD34⁺ BM cells in the CR sample of
623 patient #201-022. Numbers next to the bar indicate the number of colonies produced per 100
624 plated cells. The GM:E (granulocyte-macrophage:erythroid) ratio of colonies and the 95%
625 confidence interval for the GM:E ratio in normal BM are shown.

626 **Figure 3. Enasidenib induces differentiation from an ancestral IDH2 mutant clone.**

627 All data shown refer to samples from patient #201-023.

628 a) Heat map of targeted re-sequencing of mutations (rows) in single cells (columns, n=63
629 cells) from flow-sorted BM populations isolated pre-ENA and at CR which are shown
630 together. Clonal identification of each cell is shown below the heat map and the key to
631 mutations is denoted by letters on the right. Mutation detection key: red=detected; blue=not
632 detected; white= sequencing failed.

633 b) Clonal structure of the AML sample based on single cell genotyping (SCG). Number next
634 to a clone indicates the number of cells identified in that clone (data from a). The most likely
635 clonal structure is shown in solid arrows with alternatives in dotted arrows. (*) indicates
636 genotype "A" or "SIAR", which were each detected in only 1 cell. § indicates genotype "AS"
637 with ADO of an *IDH2* allele in 2/3 cells. Ø "SI" with ADO of the *ASXL1* allele in 3/12 cells.
638 See also Supplementary Fig. 12a.

639 c) Clonal composition in different immunophenotypic compartments pre-ENA and at CR.
640 Number of cells studied are indicated.

641 d) Clonal contribution (vertical bars) to immunophenotypic stem, progenitor, myeloid
642 precursor and terminal mature GM populations in patient samples pre-ENA and at CR
643 (horizontal bars). Data is from SCG except for mature GM population at CR, where the flow-
644 sorted cell population was genotyped (*). Normal BM is shown for comparison of
645 immunophenotypic populations.

646 e) Clonal contribution to colonies grown from CR sample (percentage of genotyped,
647 individually picked colonies). Key to mutations detected are as in (b). Numbers next to the
648 bars indicate the numbers of colonies analyzed. Lineage affiliations are as in **Fig. 1d**.
649 f) Lineage affiliation of colonies from BM CD34⁺ cells purified from CR sample compared
650 with normal CB (as in **Fig. 2e**).

651 **Figure 4. Enasidenib induces differentiation from a terminal IDH2m clone.**

652 All the data here are from patient #201-011.

653 a) Heat map of targeted re-sequencing of mutations (rows) in single cells (columns, n=110
654 cells) from flow-sorted BM populations pre-treatment and CR which are shown together. The
655 key is as in **Fig. 3a**.

656 b) Clonal structure of the AML sample based on SCG. The key to the panel is as in **Fig. 3b**.

657 * marks the genotype "DIA" where ADO was detected in 6/7 cells. Ø denotes genotype "DId".

658 No heterozygous germline SNPs were available in the *ASXL1* gene. Estimated ADO
659 frequency of the *ASXL1* allele was 12.1%. See also Supplementary Fig. 12c.

660 c) Clonal composition in different immunophenotypic compartments pre-ENA and at CR, as
661 in **Fig. 3c**.

662 d) Clonal contribution (vertical bars) to immunophenotypic BM haematopoietic populations in
663 patient samples pre-ENA and at CR (horizontal bars). Data is from SCG except for mature
664 GM population pre-ENA, where the flow-sorted cell population was genotyped (*).

665 e) Clonal contribution to colonies grown from CR sample (percentage of genotyped,
666 individually picked colonies) as in **Fig. 3e**.

667 f) Lineage affiliation of colonies from BM CD34⁺ cells purified from CR sample compared
668 with normal CB (as in **Fig. 2e**).

669 **Figure 5. Mechanisms leading to relapse of enasidenib-treated patients.**

670 a) Summary of mechanisms (rows) leading to relapse in 12 patients (columns). Selected
671 mutations detected at relapse by WES (all patients except #104-021) or by Heme Panel bait
672 capture sequencing (#104-021) are shown.

673 b) Longitudinal analysis of the percentage suppression of serum 2-HG concentrations prior
674 to enasidenib treatment (pre-ENA), at best response (CR or PR) and at relapse in 14
675 patients with an *IDH2* R140 codon mutation.

676 c-d) Serum 2-HG levels and bone marrow blast percentages prior to enasidenib treatment
677 (pre-ENA), at CR or CRp (complete remission without platelet recovery) during the course of
678 treatment (C=cycle, D=day of treatment) and at relapse in patients #201-014 (c) and #201-
679 022 (d).

680 e-f) Serial mutation analyses in flow-sorted blasts prior to enasidenib treatment (pre-ENA)
681 and at relapse in patients #201-014 (e) and #201-022 (f).

682 **Figure 6. Relapse post-enasidenib occurs through clonal evolution/selection.**

683

684 a) Patient #104-021: Heat map of targeted re-sequencing of mutations (rows) in single cells
685 (n=214 cells, columns) from flow-sorted BM populations pre-treatment and at relapse which
686 are shown together. The key is as in **Fig. 3a**.

687 b) Clonal structure of patient #104-021 based on SCG. The key to the panel is as in **Fig. 3b**.
688 Boxes in dotted red lines highlight clones which are only detected at relapse.

689 c) Clonal composition in different immunophenotypic compartments pre-ENA and at relapse,
690 as in **Fig. 3c**.

691 d) Clonal contribution (vertical bars) to immunophenotypic BM haematopoietic populations in
692 patient samples pre-ENA and at relapse (horizontal bars). Data is from SCG.

693 e) Patient #201-011: Heat map of targeted re-sequencing of mutations (rows) in single cells
694 (n=87 cells, columns) from flow-sorted BM populations at relapse. The key is as in **Fig. 3a**.

695 f) Clonal structure of patient #201-011 at relapse. * indicates 6 cells with genotype "DIA"
696 where we detected ADO in 4/5 cells in the *DNMT3A* allele. Ø denotes "DIId". The estimated
697 ADO frequency of the *ASXL1* allele was 12.1%. § indicates 6/7 "DIXF" cells where there was
698 ADO for the *DHX15* R222G mutant allele. See also Supplementary Fig. 12c. Boxes in dotted
699 red lines as in (b).

700 g) Clonal composition in different immunophenotypic compartments at relapse, as in **Fig. 3c**.
701 h) Clonal contribution (vertical bars) to immunophenotypic BM haematopoietic populations in
702 patient samples at relapse (horizontal bars). Data is from SCG.

703 **Online Materials and Methods**

704 Please see also the Life Sciences Reporting Summary

705 **Patient samples**

706 Bone marrow (BM) or blood samples from normal donors and AML patients were obtained
707 with informed consent and collected by research ethics committee-approved Biobanks
708 (MDSBio Study, MREC 06/Q1606/110, Oxford Musculoskeletal Biobank, MREC
709 09/H0606/11: South Central Oxford C REC), Gustave Roussy (Department of Clinical
710 Hematology and Drug Development Department (DITEP), Gustave Roussy, Villejuif) and
711 MSKCC Biobanks. Cytogenetic analyses were performed in clinical laboratories.
712 Mononuclear cells (MNCs) were isolated by Ficoll density gradient. MNCs were viably frozen
713 in 90% FCS/10% DMSO in liquid nitrogen.

714 **Measurement and analysis of 2-hydroxyglutarate (2-HG)**

715 Serum samples were collected within 28 days before the first dose of enasidenib
716 ('screening') and/or pre-dose on day 1 of each treatment cycle. 2-HG concentration was
717 determined by liquid chromatography tandem mass spectrometry (Covance, Inc USA
718 according to their validated method). Baseline 2-HG was either the average of the screening
719 sample and pre-dose cycle 1 sample, or either sample if both were not available. Percentage
720 suppression of 2-HG was determined by comparing the lowest level of 2-HG observed on-
721 treatment to baseline.

722 **Hematopoietic cell immunophenotyping**

723 Frozen BMMNCs from normal donors or AML samples were washed with Iscove's Modified
724 Dulbecco's Medium (Thermo Fisher Scientific, UK), 10% fetal bovine serum (Sigma, UK)
725 and 1mg/ml bovine pancreatic DNase I (Sigma, UK). Cells were stained for flow cytometry
726 with antibodies in Supplementary Table 8. Analysis was carried out on either BD LSR

727 Fortessa or BD FACSAria Fusion (Becton Dickinson, Oxford UK). Antibodies in the lineage
728 (Lin) depletion are: anti-CD2,CD3,CD4,CD8a,CD10,CD19,CD20 and CD235a. 7-
729 aminoactinomycin-D (7AAD, Becton Dickinson, UK) was used as a live/dead stain.
730 Hematopoietic stem/progenitor cells were defined as subsets of Lin⁻CD34⁺ and myeloid
731 precursor as Lin⁻CD34⁻CD117⁺ as previously described²¹. Terminally mature myeloid cells
732 were defined as Lin⁻CD34⁻CD117⁻.

733 **Mutational analysis by FoundationOne® Heme Panel**

734 Analysis of samples in **Fig. 1a** by FoundationOne® Heme panel was conducted by
735 Foundation Medicine, Inc. Nucleic acid libraries were prepared from DNA and RNA extracted
736 from fresh patient BM samples and captured using custom bait-sets targeting 405 cancer-
737 related genes by DNA-sequencing (DNA-seq), and 265 frequently rearranged genes by
738 RNA-sequencing (RNA-seq). Genes included in this analysis encode known or likely targets
739 of therapies, either approved or in clinical trials, or are otherwise known drivers of
740 oncogenesis published in the literature⁵².

741 **Mutational analysis by Fluidigm Access Array**

742 Selected samples from **Fig. 1a** and **Supplementary Table 2** was performed using highly
743 multiplexed PCR-based targeted re-sequencing with a custom panel of 373 amplicons
744 covering areas of high frequency AML mutations in 35 genes, using the Fluidigm Access
745 Array platform as previously described⁶.

746 **Mutational analysis by whole exome sequencing (WES).**

747 Genomic DNA from flow-sorted AML blasts or CD3-positive cells (germline control) was
748 purified with Allprep DNA/RNA mini or micro Kits (Qiagen, France). After exome capture with
749 SureSelect V5 Mb All Exon kit (Agilent, Les Ulis, France), paired-end 100 bp sequencing
750 was performed on HiSeq2000 (Illumina, Paris France). Read alignment to hg19 reference
751 genome was performed using the BWA algorithm, v0.7.10 with corrections using GATK
752 (v3.3.0) after removal of PCR duplicates. Variant detection was carried out with VARSCAN
753 (v2.3.7). Somatic variants (**Supplementary Table 2**) were selected using the following
754 criteria: minimum depth: 8x, VAF>10% in AML blasts and lower than 10% in germline

755 control, and p-value <0.001. All variants were manually inspected using IGV (v2.3) software.
756 Artfactual variants from DNA oxidation where read pair orientation bias was observed (i.e.
757 predominant F2R1 orientation for C to A variations or F1R2 for G to T variations) were
758 filtered out. Median depth at the positions reported in the **Supplementary Table 2** was 121
759 (range: 19-843).

760 **Selection of mutations for targeted re-sequencing in populations and single cells.**

761 On average 23 somatic mutations were detected per AML sample by WES. We selected
762 mutations for further validation by targeted re-sequencing and SCG based on the following
763 criteria: 1) known recurrent mutations in AML, 2) non-recurrent mutations in genes
764 commonly mutated in AML. In addition, mutations not meeting above criteria, but where the
765 VAF varied by at least 5% between sequential samples in the same patient were of interest
766 as these may be markers of clonal shifts. Where multiple mutations had similar VAFs and
767 showed similar patterns of change in VAFs in sequential samples, a representative mutation
768 was selected. This is illustrated in the example from patient #201-011 (Supplementary
769 Figure 11a). Mutations which were not validated by targeted re-sequencing were excluded
770 in subsequent data analysis. Chromosomal loss of heterozygosity detected by WES (or by
771 karyotyping) was examined using germline single nucleotide polymorphisms (SNP) present
772 in the affected chromosomal region.

773 **Mutational analysis by targeted re-sequencing**

774 Mutations detected by targeted re-sequencing in hematopoietic cell populations are in
775 **Supplementary Table 3**. Average and range of read depths for each mutation is shown in
776 **Supplementary Table 4**. DNA was extracted (DNeasy Blood and Tissue extraction kit,
777 #69506 Qiagen Manchester UK) from bulk and flow-sorted cells from patient samples.
778 Where material was limiting, whole genome amplification (WGA, RepliG, Qiagen, UK)) was
779 performed. Targeted PCR was performed using high Fidelity Phusion Taq polymerase (NEB,
780 UK) or KAPA2G Multiplex DNA Polymerase (KAPA Biosystems, UK) with 10ng of gDNA.
781 Primers used are in **Supplementary Table 7**. A second PCR reaction added Illumina
782 barcodes and sequencing oligonucleotides prior to sample purification, quantitation, pooling

783 and library preparation for sequencing on Illumina MiSeq (Illumina, Saffron Walden, UK).
784 Raw data (average depth ~996x) was aligned using Stampy (v1.0.20)⁵³. A minimum
785 sequencing depth of 100 was set as a threshold for inclusion of data for analysis. >94% of
786 reads had Phred scores of >30. VAF was obtained using the Unix 'grep' (globally search
787 regular expression and print) command line.

788 **Mutant *IDH2* variant allele frequency by quantitative PCR.**

789 *mIDH2* VAF was assessed in gDNA extracted from flow-sorted patient blood CD14⁺
790 monocytes, CD16⁺ neutrophils, or polymorphonuclear neutrophils (Ficoll gradient
791 purification) at various time points during enasidenib treatment. Quantitative SNP assay
792 PCR (12.5ng DNA per test) was performed using TaqMan® Universal PCR Master Mix
793 (Applied Biosystems, France), TaqMan® probes (specific for either *IDH2* wild type (FAM) or
794 mutated R140Q (VIC) alleles (rs121913502, Applied Biosystems, France)). PCR was
795 performed on an ABI 7500 Fast Real-Time PCR analyzer (Applied Biosystems, France) with
796 cycling conditions: initial 1min at 60°C, 10min at 95°C, followed by 40 cycles of 15sec at
797 95°C and 1min at 60°C; and then 1min at 60°C.

798 **Single cell genotyping**

799 Mutations interrogated by SCG are in **Supplementary Table 3**. Single cells were flow-sorted
800 into 96 well plates containing 2µl of phosphate buffered saline. WGA was carried out using
801 Single Cell RepliG kit (Qiagen, Crawley UK). Briefly, following cell lysis, alkali denaturation
802 and neutralisation, a master mix containing Phi29 polymerase, dNTPs and random
803 oligonucleotide primers was added. WGA was carried out at 30⁰C for 8 hours followed by
804 heat inactivation. Diluted (1:20) amplified DNA was used in single or multiplex PCR using
805 primers relevant to the sample and high Fidelity Phusion Taq polymerase (NEB, UK) or
806 KAPA2G Multiplex DNA Polymerase (KAPA Biosystems, UK). Barcoding and sequencing
807 oligonucleotides were added by PCR and sequencing performed on Illumina MiSeq
808 (Illumina, Saffron Walden, UK). ~94% of reads had Phred scores of >30. A threshold of 50
809 reads was set for analysis inclusion. VAF thresholds for determining detection of mutations
810 were determined by genotyping 48 single cells derived from normal bone marrow, and set at

811 the 95% confidence level (mean \pm 1.96 x standard error of mean (SEM); i.e. <5% chance of
812 false positive, **Supplementary Table 6**).

813 **Imputation of clonal structures using bulk VAFs**

814 The most common method used to impute clonal structure is based on the assumption that
815 the most abundant mutation resides in the earliest occurring ancestral clone. This method
816 may be applicable to samples with linear clonal structures but it may not be able to
817 accurately resolve more complex or branching clonal structures. Longitudinal bulk
818 genotyping data may offer additional information, particularly where there is evidence of
819 clonal selection or evolution.

820 Putative clonal structure is first solved for each sample independently using bulk VAF data
821 based on the rules below. In samples where there was no colony or single cell (SC)
822 genotyping, longitudinal sampling (pre-ENA, best response and relapse) can provide
823 additional information on the likely clonal structure of that patient. In the absence of colony
824 or SCG data, bulk VAFs were used to estimate the size of the most likely clones in samples.
825 In samples where there is associated colony and/ or single cell (SC) genotyping, the clonal
826 structure was re-drawn based on these data, which provide a higher confidence structure
827 with some resolution of intermediate clones. Once this clonal structure is solved, bulk VAF
828 was used to estimate the sizes of clones in samples.

829 Sequence of acquisition of mutations can be imputed from bulk genotyping data, using VAF
830 as an estimate of clonal contribution (Supplementary Fig. 11b). Mutations are first ranked
831 according to VAF: in these examples, V_A is highest, V_D is lowest.

832 Factors which may cause data error and bias include limited cell equivalent representation in
833 extracted genomic DNA from small cell numbers, bias present in whole genome amplified
834 material and PCR bias (including sequencing bias). While we were unable to control for the
835 first two factors, we could estimate standard error of our sequencing data. We performed
836 technical replicate genotyping of 19 unsorted BMMC populations and obtained the standard
837 error of mean (SEM) from VAFs from each mutation. In total, 142 standard error values

838 were obtained, and the average SEM was 1.9% (range 0.0-20.2%), with a 95% confidence
839 interval upper limit of 2.47%. This limits our ability to reliably distinguish between clones
840 varying in VAF of ~2.5% (~5% of cells if mutations were heterozygous), and we were not
841 able to impute sequence of acquisition of mutations in population genotyping with less than
842 2.5% difference in VAF between them. Where three or four mutations have VAF within 2.5%
843 of each other, the average VAF of the cluster is taken as the VAF of all mutations in that
844 cluster (Supplementary Fig.11c).

845 **Interpreting VAFs and cellular representation in the context of loss of heterozygosity**
846 **or hemizyosity.**

847 We detected multiple occurrences of loss of heterozygosity (LoH, e.g. copy-loss
848 chromosomal deletions or copy-neutral uniparental disomy) and mutations, which were on
849 the X chromosome in male patients (hemizyosity). There is complexity in interpreting bulk
850 VAFs due potential mix of cells with or without LoH within a bulk population.
851 Notwithstanding this caveat, we used the simple models set out below to help interpret VAFs
852 in different contexts. For somatic heterozygous variants (somatic mutation or germline
853 polymorphism) in autosomal chromosomes the estimated percentage cellular representation
854 is $2 \times \text{VAF}\%$ (Supplementary Fig. 11d). Heterozygous variant and chromosomal deletion
855 resulting in copy-loss LoH in autosomal chromosomes is illustrated in Supplementary Fig.
856 11e. Here, there is a non-linear relationship between VAF and cellular representation
857 (Supplementary Fig. 11f). The formula we used to estimate cellular representation was: %
858 cellular representation = $[-2.777 \times (\text{VAF ratio})^2] + (6.145 \times \text{VAF ratio}) - 2.373$. In
859 heterozygous variants and copy-neutral LoH (e.g. uniparental disomy) in autosomal
860 chromosomes, percentage cellular representation is $100\% - (2 \times \text{VAF}\%)$ (Supplementary Fig.
861 11g). In X-linked variants in male subjects in sex chromosomes: percentage cellular
862 presentation is the same as the VAF% (Supplementary Fig. 11h).

863 **Determination of clonal structures using single cell genotyping**

864 Each single cell was assessed for detection or non-detection of mutations in that patient
865 sample by amplicon sequencing of DNA subject to whole genome amplification. A minimum

866 coverage of 30x across an amplicons was required for a amplicon to be called. We assigned
867 the most likely sequence of acquisition of mutations based on the genotype identified in
868 cells. For example, where mutations A, B, C, D and E were identified in a sample, discrete
869 cells with genotypes A, AB, ABC, ABD, ABCD and AE may be called. In most cases the
870 sequence of acquisition, e.g. A->AB, is clear. However, the sequence of acquisition of
871 mutations during the transition for example, from AB to ABCD may not be clear due to allelic
872 drop out (ADO) i.e. the sequence of acquisition may be AB -> ABC -> ABCD or AB-> ABD->
873 ABCD. In such cases, intermediate genotypes represented by most cells may be more likely
874 to be true. In all cases, models of clonal structures which require the least number of
875 discrete mutational steps required are represented⁵⁴, although alternative structures,
876 including ones where the same mutation is acquired twice, are possible. Once the most
877 likely clonal structure is established, cells where there was failure to amplify a locus that did
878 not alter the assignment of a mutational complement (e.g. an early mutation in the hierarchy)
879 were included in the final analysis.

880 **Allele Drop Out Estimation (ADO)**

881 ADO can be measured for by two methods. First, ADO can be determined by determining
882 the phase of germline SNPs near mutations. For patients #201-023 and #201-011, we
883 genotyped germline SNPs which were either in-phase (i.e. on the same allele as) with a
884 mutation or out of phase (i.e. on the opposite allele).

885 In-phase SNPs were rs6597996 and rs11246258, *DEAF1* N372K mutation, patient #201-
886 011. Out-of-phase SNPs were: (i) rs4911231, *ASXL1* G646fs mutation, patient #201-023; (ii)
887 rs2276598, *DNMT3A* R598X, patient #201-011; (iii) rs7657364, *DHX15* R222G, patient
888 #201-011. The SNPs were situated between 157 bases and 4 kilobases from the mutations.
889 The threshold VAF for ascertaining dropout in these SNPs was <2% (homozygous
890 reference) or >98% (homozygous variant).

891 In patient #201-023 (**Fig. 3a-b**) in 2/3 cells with the “AS” genotype there was ADO of one of
892 the *IDH2* alleles (Supplementary Fig. 12a, left) and in 2/12 cells with the genotype “SI” there
893 was ADO in the *ASXL1* allele (Supplementary Fig. 12a, right).

894 Our analysis also showed that in patient #201-011 *DNMT3A* R598X and R736C mutations
895 are on different alleles and the R736C mutation is in phase with rs2276598 (Supplementary
896 Fig. 12b).

897 By studying both VAFs of the SNP rs2276598 and the R736C mutation in cells with
898 genotype “DIA” in patient #201-011 where ADO may have occurred we determined that in
899 6/7 cells at CR, ADO of the allele that harboured the R598X mutation had occurred
900 (Supplementary Fig. 12c left). Where the SNP did not amplify, the VAF of R736C mutation
901 was informative. Similarly, in the same patient at relapse, 5/6 “DIA” cells at relapse had ADO
902 of the R598X allele (Supplementary Fig. 12c center). We also determined that in 6/7 cells
903 with the “DIXF” genotype at relapse there had been ADO of the allele harbouring the *DHX15*
904 R222G mutation (Supplementary Fig. 12c, right).

905 An alternate method to determine ADO more globally is to study the frequency with which a
906 variant is called homozygous (either reference or alternative) when it should be
907 heterozygote. We analysed 6 SNPs in 5 genes (*ASXL1*, *IDH2*, *DNMT3A*, *DEAF1* and
908 *DHX15*) in 402 single cells from 2 patients (#201-011 and #201-023) known to be
909 heterozygous in the germline (confirmed by genotyping population of flow sorted T cells from
910 the patients). Mean VAF was 49.1% for all 6 SNPs with a near-symmetrical distribution of
911 VAFs across these single cells ranging from 0-100% (Supplementary Fig. 12d). Frequencies
912 of homozygous reference (VAF≤1%) or variant calls in the 402 cells (VAF≥99%, thresholds
913 based on analysis of known homozygous SNPs in 237 cells) were 15.9% (64 cells/402 cells)
914 and 15.2% (61 cells/402 cells) respectively (Supplementary Fig. 12d).

915 Next, if we assume that mutations in patients were heterozygous (i.e. in cases where there is
916 was no data to support uniparental disomy or copy number loss) we asked what was the
917 frequency with which mutations were called homozygous (VAF≥99%) (presumably due to

918 ADO). Across the 23 mutations assessed using this method, the average ADO rate of wild
919 type allele was 10.36% (SD 5.7%, Supplementary Fig. 12e). However, we found variation in
920 ADO rates between different patients, even at the same mutation (e.g. *IDH2* R172K),
921 suggesting that ADO is affected by factors additional to that of the activity of the Phi21
922 polymerase. Where possible we have used gene-specific germline SNPs, or alternatively
923 used sample and mutation specific ADO estimated by the 'homozygous mutant' method for
924 our analyses.

925 **Digital Droplet PCR**

926 We confirmed the VAFs detected by next-generation sequencing (NGS) using digital droplet
927 PCR (ddPCR) in 17 amplicons and 113 sorted multi-cell AML populations from 5 patients
928 using the BIORAD platform as previously described²¹. There was good correlation between
929 VAF values obtained using these two methods ($R^2= 0.974$,). There was one AML variant
930 (*DHX15* R222G) where NGS gave an unexpectedly high VAF in single normal BM MNCs
931 (where the mutation was found to be absent in the bulk normal BM sample). Detection of this
932 variant in normal and AML were therefore carried out using ddPCR which showed presence
933 of the mutation in AML bulk and single cells, but confirmed its absence in normal bulk and
934 single cells.

935 **Whole transcriptome sequencing (RNAseq) and analysis of alternative splicing events**

936 Copy DNA libraries were prepared using extracted RNA from AML blast cells flow-sorted
937 from re- and post-relapse samples from patients #201-011 and #201-013. cDNA libraries
938 were prepared for sequencing with tagmentation and indexing using Illumina Nextera
939 Sample Preparation kit (Illumina, Saffron Walden, UK). RNA-seq data were generated as 75
940 bp paired-end unstranded Illumina reads. Reads were aligned using STAR(v2.4.0.1)⁵⁵ to the
941 human genome (GRCh37) with default parameters. On average alignment was 96.7%
942 (range: 95.5-97.4%) with an average of 139×10^6 (range: 116×10^6 - 169×10^6) mapped reads
943 per sample. Differentially spliced events (DSEs) for the wild type (pre-relapse) and post-
944 relapse with spliceosome gene mutations analysed as paired samples, were identified⁵⁶
945 using Mixture of Isoforms (MISO v0.5.4) using default parameters. An event is termed as

946 differentially spliced if the Bayes Factor (BF) ≥ 10 , $|\Delta PSI| > 0.2$ where PSI is 'percentage
947 spliced in', and the event is supported by ≥ 10 reads. DSEs are classified included or
948 skipped spliced exons (SE), alternative 3'/5' splice sites (A3SS, A5SS), mutually exclusive
949 exons (MXE) or retained introns (RI).

950 **Colony assays**

951 25-250 cells were plated in duplicate in 1.2ml of MethoCult GFH4435 (StemCell
952 Technologies, Manchester, UK). AML patient samples were assayed with added Enasidenib
953 (1 μ M). The rest of the procedure was as previously described²⁰.

954 **Statistical Analysis**

955 Where applicable, statistical analyses were performed using with GraphPad Prism software
956 (v7.02) using statistical methods noted in figure and table legends.

957


958 **References**

- 959 6. Craddock, C.F., *et al.* Outcome of Azacitidine Therapy in Acute Myeloid Leukemia Is
960 not Improved by Concurrent Vorinostat Therapy but Is Predicted by a Diagnostic
961 Molecular Signature. *Clin Cancer Res* (2017).
- 962 20. Goardon, N., *et al.* Coexistence of LMPP-like and GMP-like leukemia stem cells in
963 acute myeloid leukemia. *Cancer Cell* **19**, 138-152 (2011).
- 964 21. Quek, L., *et al.* Genetically distinct leukemic stem cells in human CD34- acute
965 myeloid leukemia are arrested at a hemopoietic precursor-like stage. *J Exp Med* **213**,
966 1513-1535 (2016).
- 967 52. He, J., *et al.* Integrated genomic DNA/RNA profiling of hematologic malignancies in
968 the clinical setting. *Blood* **127**, 3004-3014 (2016).
- 969 53. Lunter, G. & Goodson, M. Stampy: a statistical algorithm for sensitive and fast
970 mapping of Illumina sequence reads. *Genome Res* **21**, 936-939 (2011).
- 971 54. Potter, N.E., *et al.* Single-cell mutational profiling and clonal phylogeny in cancer.
972 *Genome Res* **23**, 2115-2125 (2013).


- 973 55. Dobin, A., *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
974 (2013).
- 975 56. Katz, Y., Wang, E.T., Airoidi, E.M. & Burge, C.B. Analysis and design of RNA
976 sequencing experiments for identifying isoform regulation. *Nat Methods* **7**, 1009-1015
977 (2010).
- 978

Figure 1


a



b



c



d





Figure 2

a Clonal origin of complete remission response to Enasidenib




b

Patient ID	#201-022	#201-023	#201-010	#201-011	#201-006	#201-027
Wild type	■					
Ancestral		■				
Terminal			■	■	■	■

c

#201-022		IDH2 R140Q (I)	PEX26 P118S (P)	FEZ2 S208T (F)	DNMT3B N738S (D)	ZCCHC1 T712M (Z)	NPM1 L135 delins LLSIFKG (N)	ELMO3 R725Q (E)
VAF	Pre- ENA AML blasts	36.00%	25.45%	24.96%	23.20%	22.72%	22.17%	1.83%
	CR PBMNCs	0.24%	1.00%	1.23%	1.61%	0.58%	0.12%	0.31%

d



e




Figure 3


#201-023

a

Single cells →


■ Detected
■ Not detected
□ Failed

■ WT ■ SIA
■ A ■ SIAR
■ AS ■ SIARr
■ SI ■ SIARrG




SRSF2 P95L (S)
IDH2 R140Q (I)
ASXL1 G646fs (A)
RUNX1 W477X (R)
RUNX1 Q268X (r)
GATA2 G320V (G)
Clone (n= 63 cells)

b Clonal structure




d Hemopoietic populations (horizontal bars)
Clones (vertical columns)


■ Stem/ progenitor ■ HSC ■ GMP
■ Precursor ■ MPP ■ CMP
■ Mature ■ LMPP ■ MEP



c Single cell genotyping



e



f






Figure 4
#201-011


a




b Clonal structure




c



d Hemopoietic populations (horizontal bars)
Clones (vertical columns)



e



f





Figure 5


a

Patient ID	#201-014	#201-022	#104-021	#201-013	#201-004	#201-006	#201-010	#201-007	#201-019	#201-003	#201-011	#201-012
IDH1	IDH1 R132C	IDH1 R132H										
cytokine receptor signalling			CSF3R T618I	FLT3 D835A	FLT3 ITD			FLT3 I867M				
haematopoietic transcription factor	RUNX1 F416 Gfs*135			BCL11A N391K	GATA2 N317S/ L321R	RUNX1 T188 Hfs*25						
genes implicated in spliceosome function				DDX1 G699A							DHX15 R222G	
other genes recurrently mutated in haematopoietic cancer		ELMO3 R725Q	NFKB1 M216I	BCOR R1375W	BRCA2 A2643V				SCN3A R28H	AKAP8L D89N	DEAF1 N372K	
				CACNA1G G663R	CBLsplice acceptor				SETD1B A1054 delinsAE	PLCL1 A985V		
				UGT2B10 Y16*								
other genes recurrently mutated in other cancer				SLC18A3 W105*			IL17A P130S	MTUS Q835H		DOM3Z Y92*		
deletion chr 7q												


b




c #201-014



e #201-014



d #201-022



f #201-022






Figure 6


#104-021




b Clonal structure C




c Single cell genotyping




d Hemopoietic populations (horizontal bars) Clones (Vertical columns)




#201-011 (Relapse)



f Clonal structure



g Single cell genotyping Relapse



h Hemopoietic populations (horizontal columns) Clonotype (Vertical bars)

