



King's Research Portal

DOI: 10.3389/fimmu.2020.577815

Document Version Publisher's PDF, also known as Version of record

Link to publication record in King's Research Portal

Citation for published version (APA):

Ortega, P. A., Silva-miranda, M., Torres-larios, A., Campos-chávez, E., Franken, K. C. L. C. M., Ottenhoff, T. H. M., Ivanyi, J., & Espitia, C. (2020). Selection of a Single Domain Antibody, Specific for an HLA-Bound Epitope of the Mycobacterial Ag85B Antigen. *Frontiers in Immunology*, *11*, Article 577815. https://doi.org/10.3389/fimmu.2020.577815

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Selection of a Single Domain Antibody, Specific for an HLA-Bound Epitope of the Mycobacterial Ag85B Antigen

Paola A. Ortega¹, Mayra Silva-Miranda^{1,2}, Alfredo Torres-Larios³, Eduardo Campos-Chávez³, Kees C. L. C. M. Franken⁴, Tom H. M. Ottenhoff⁴, Juraj Ivanyi⁵ and Clara Espitia^{1,5*}

¹ Departamento de Inmunología, Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México, Ciudad de México, México, ² CONACyT-Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México, Ciudad de México, México, ³ Department of Biochemistry and Structural Biology, Instituto de Fisiología Celular, Universidad Nacional Autónoma de México, Ciudad de México, México, ⁴ Department of Infectious Diseases, University Medical Centre Leiden, Leiden, Netherlands, ⁵ Center for Host-Microbiome Interactions, King's College London, London, United Kingdom

OPEN ACCESS

Edited by:

Mario Alberto Flores-Valdez, CONACYT Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco (CIATEJ), Mexico

Reviewed by:

Eddie A. James, Benaroya Research Institute, United States Armando Acosta, Universiti Sains Malaysia Health Campus, Malaysia

***Correspondence:** Clara Espitia espitia@iibiomedicas.unam.mx

Specialty section:

This article was submitted to Microbial Immunology, a section of the journal Frontiers in Immunology

Received: 30 June 2020 Accepted: 14 September 2020 Published: 02 October 2020

Citation:

Ortega PA, Silva-Miranda M, Torres-Larios A, Campos-Chávez E, Franken KCLCM, Ottenhoff THM, Ivanyi J and Espitia C (2020) Selection of a Single Domain Antibody, Specific for an HLA-Bound Epitope of the Mycobacterial Ag85B Antigen. Front. Immunol. 11:577815. doi: 10.3389/fimmu.2020.577815 T cells recognizing epitopes on the surface of mycobacteria-infected macrophages can impart protection, but with associated risk for reactivation to lung pathology. We aimed to identify antibodies specific to such epitopes, which carry potentials for development toward novel therapeutic constructs. Since epitopes presented in the context of major histocompatibility complex alleles are rarely recognized by naturally produced antibodies, we used a phage display library for the identification of monoclonal human single domain antibody producing clones. The selected 2C clone displayed T cell receptor-like recognition of an HLA-A*0201 bound 199KLVANNTRL207 peptide from the Ag85B antigen, which is known to be an immunodominant epitope for human T cells. The specificity of the selected domain antibody was demonstrated by solid phase immunoassay and by immunofluorescent surface staining of peptide loaded cells of the T2 cell line. The antibody affinity binding was determined by biolayer interferometry. Our results validated the used technologies as suitable for the generation of antibodies against epitopes on the surface of Mycobacterium tuberculosis infected cells. The potential approaches forward the development of antibody in immunotherapy of tuberculosis have been outlined in the discussion.

Keywords: tuberculosis, single domain antibodies, peptide-human leukocyte antigen complex, mycobacterial Ag85B, T cell receptor-like antibodies

INTRODUCTION

Current problems of Tuberculosis (TB) control are due to the emergence of drug resistant strains, the failure of the existing BCG vaccination and the persistence of factors associated with povertystricken populations. Consequently, there is a global morbidity of 10 million people with the mortality of 1.2 million in HIV-negative people and 0.25 million in HIV positive subjects (1) New

1

approaches toward the control of TB involve the shortening of the current chemotherapy regimen (2) and prophylaxis for HIVrelated TB without interfering with antiretroviral therapy (3). This pilot study, aiming at the immunotherapy of TB had an initial objective to identify monoclonal antibodies with T cell receptor (TCR)-like specificity, binding to peptide epitope/ human leukocyte antigen (HLA) class I complexes on the surface of *M. tuberculosis* infected cells. This approach has been based in the evidence, that such antibodies can be developed for the specific killing of malignant and virusinfected cells (4–10).

The mycobacterial peptides epitopes complexed with MHC class I molecules on the surface of infected cells are known to be recognized and leading to the activation of CD8+ T cells (11-13). Therefore, antibodies with TCR-like specificity following conjugation with suitable apoptosis-inducing ligands could potentially become mycobactericidal and a suitable adjunct to the chemotherapy of TB. The identification of TCR-like antibodies with specificity against M. tuberculosis Acr1 peptides/HLA.A*0201, HLA.A*011 and HLA.A*24 class I complexes has recently been reported, using peptide/MHC complexes generated via UV-induced peptide exchange. The complexes were panned against human DAb (domain antibody) phage display library (14, 15). This approach can be expanded by testing immunodominant epitopes from other M. tuberculosis antigens recognized in the context of HLA class I alleles (16, 17). Our interest focused on the HLA-A*0201 restricted CD8+T-cell epitopes of Ag85B, a major secreted M. tuberculosis protein. Two Ag85B peptides 143FIYAGSLSA151 and 199KLVANNTRL207 had previously been identified in healthy humans and in immunized HLA-A2 transgenic mice (18), while HLA-A*0201 allele specific recognition of 37YLLDGLRAQ45 and 199KLVANNTRL207, was observed in patients with active TB (19). Since epitopes from Esat-6 and TB10.4, are recognized in the context of various HLA-A alleles, the HLA-A*0201 restricted, 199KLVANNTRL207 epitope from Ag85B appeared to be the most immunodominant (17) and was therefore chosen as our target for the selection and characterization of a TCR-like antibody. Using a human DAb phage display library, we selected a clone producing a single domain antibody (sdAb) against the Ag85Bp199-207/HLA-A*0201 complex (Ag85Bp/HLA-A*0201). Its specificity was determined by ELISA, and by its binding capacity to the Ag85Bp/HLA-A*0201 expressed on cells of the human HLA-A*0201 positive T2 cell line.

MATERIALS AND METHODS

Mycobacterium tuberculosis Peptides/ HLA-A*0201 Complexes

Three nonamer peptides, known to be HLA-A*0201 restricted CD8+T cell epitopes from: Ag85B ($_{199}$ KLVANNTRL $_{207}$), Esat-6 ($_{82}$ AMASTEGNV $_{90}$), or Acr1 ($_{120}$ GILTVSVAV $_{128}$) proteins of *M. tuberculosis* (16) were synthesized by Anaspec, Inc. (USA). The peptides were of > 90% purity, and their homogeneity was confirmed by analytical reverse-phase high-performance liquid

chromatography. The biotinylated recombinant complexes of Ag85Bp/HLA-A*0201, Esat-6_{p82-90}/HLA-A*0201 (Esat-6p/HLA-A*0201), and Acr1_{p120-128}/HLA-A*0201 (Acr1p/HLA-A*0201), were produced using extracellular HLA class I molecules, with C-terminal BirA recognition site, and β 2-microglobulin (20). The insoluble aggregates expressed in *Escherichia coli* in the form of inclusion bodies were solubilized in urea and folded with peptide by dilution. Monomers were biotinylated using the BirA enzyme and purified by gel filtration on a Hiload 16/60 Superdex 75 prep grade.

Evaluation of Refolding of Peptide/HLA-A*0201 Complexes by ELISA

An ELISA was carried out to evaluate the correct conformation of the complexes by using the W6/32 mAb (Invitrogen/USA) which recognizes a conformational epitope on the intact heavy chain/ β 2microglobulin complex (21–23). Briefly, 0.5 µg/well of the biotinylated complexes in Phosphate Buffer Saline (PBS), were immobilized on streptavidin coated (ThermoScientific/USA) and on uncoated high protein-binding (ThermoScientific) plates. Samples were incubated overnight (ON) at 4°C. Next day, plates were washed twice with PBS and then incubated 1 h at room temperature (RT) with W6/32 mAb diluted 1/2,000 in PBS-tween-20 0.05%, Bovine Serum Albumin BSA 2% (PBS-TBSA). After 3 washes with PBS-T, wells were incubated by 1 h with Horseradish peroxidase conjugated goat anti-mouse IgG H+L antibody (antimouse IgG-HRP)[1/2,000 (Invitrogen/USA)]. Then, complexes on both streptavidin coated and uncoated plates, were incubated with streptavidin-HRP (Biosource/China), diluted 1/4,000 1 h at RT. Finally the reaction was revealed with TMB (3,3',5,5'tetramethylbenzidine) (ThermoScientific), and stopped with 100 µl of 0.16M H₂SO₄ Absorbance values were measured at 450 nm using an ELISA plate reader (Multiskan Go, Thermo).

Panning of the Human Single Domain Antibody Phage Library Against the Ag85B_{p199-207}/HLA-A*0201 Complex

A human DAb phage display library, containing approximately 3×10⁹ sdAb clones (Geneservice, Cambridge) was used following a modified protocol (24). A negative panning was carried out for elimination of the background reactivity against streptavidin as follow: 5 x10¹² phage library particles were pre-incubated at 4°C for 1h with 30 μ l of streptavidin magnetic beads M280 (Invitrogen/Norway), then tube was placed in a magnet and phage supernatant was incubated with a 7.5 µg of biotinylated Ag85Bp/HLA-A*0201 in PBS at 4°C for 1h. After that, 200 µl of streptavidin beads were added and sample was incubated for 15 min at 4°C with shaking. Beads were pulled down with the magnet and washed 15 times with PBS-T 0.1%. Finally, sdAb phage complexes were eluted by incubation with glycine-HCl pH 2.2 for 15 min at RT and sample was neutralized with Tris-HCl pH 9.0. For the second and the third round of selection, 2.5 and 1.25 µg of complexes were exposed to streptavidin beads and washed 15 and 25 times with PBS-T 0.1% respectively. After the third final round of panning, the eluted sdAb phages were used to infect 5 ml freshly prepared *E. coli* TG1 culture. Bacteria were plated onto TYE medium supplemented with 4% glucose and carbenicillin 100 µg/ml (TYG_{4%}C₁₀₀). After ON culture, 94 individual clones were picked onto a 96 wells plate, containing 200 µl of $2xTYG_{4\%}C_{100}$. Plates were incubated ON at 37° C, with shaking at 200 rpm. Next day, 5 µl of ON culture from each well was transferred to a new plate with 200 µl of fresh $2xTYG_{4\%}C_{100}$. After 3 h of culture at 37° C, 50 µl 2xTY supplemented with $4x10^{8}$ M13 phage was added to each well and plates were incubated for 1 h at 37° C. After centrifugation to 2,800 rpm during 10 min, pellet was re suspended in 200 µl of $2xTYC_{100}K_{100}$ (Kanamycin 100 µg/ml). Cultures were grown at 26° C with shaking at 250 rpm, during 16-24 h.

ELISA Phage

Phage supernatants from each well were collected and evaluated by ELISA. Biotinylated complexes at 0.5 μ g/well were bound to coated streptavidin plates as described before, and incubated with 100 μ l of phage supernatant for 1 h at RT. After several washes with PBS-T 0.05%, wells were incubated with anti-M13 HRP antibody (1/2,500) for 1 h at RT (GE Healthcare/USA). The reaction was developed with o-Phenylenediamine dihydrochloride OPD (SigmaAldrich/USA), and stopped adding 25 μ l of 3M H₂SO₄. Absorbance values were measured at 492 nm using an ELISA plate reader Multiskan–GO. Phage supernatants from positive clones were also evaluated by phage ELISA against non-target complex; Esat-6p/HLA-A*0201 and Acr1p/HLA-A*0201 as described above.

Domain Antibody Sequencing

Double strand phagemid DNA extraction was performed from 3 selected clones in *E. coli* strain TG1 by using GeneJET Plasmid Miniprep kit (ThermoScientific/Lithuania), The primers used for sequencing were LMB3 (5' CAGGAAACAGCTATGAC 3') and pHEN (5'CTATGCGGCCCCATTCA 3'). Sequencing was carried out in the sequencing facility at Instituto de Investigaciones Biomédicas. For translation BioEdit 7.2 software was used and BLAST and Clustalw tools for sequences analysis and alignment.

Production of Soluble Domain Antibody

Transformation of E. coli HB2151 by phage infection and expression of sdAb were done with minor modifications according to (25). Once bacteria were transformed with phages, the positive clones (2C, 3C, and 7E), 50 μ l of 1:10¹² to 1:10⁶ cell dilutions were sub-cultured in TYEC₁₀₀ plates and incubated overnight at 37°C. Three random unit forming colony (UFC) were picked up from each sample and inoculated in a culture flask containing 2xTYC₁₀₀. The culture was grown with shaking (250 rpm) at 37°C until OD_{600nm}=0.6 (26). Then, Isopropyl-β-D-1-thiogalactoside (IPTG) (Promega/USA) was added to a final concentration of 1 mM, culture was continued, at 26°C with shaking (250 rpm), ON. Cells were harvest by centrifugation at 4,500 rpm, and bacterial sediment was treated with an osmotic buffer (750 mM sucrose, 100 mM Tris pH 7.5) as described by (27). The periplasmic fractions obtained from each clone were subjected to affinity chromatography on Protein-A- agarose (Roche/Germany), in order to purify the sdAbs following the manufacturer protocol. The eluted protein fractions were shuffled and concentrated to 500 μ l in PBS pH 7.4, using amicon-15ml, 10.000 WM (Merck/Ireland). Protein quantification was determined by BCA assay (Pierce/USA).

SDS-PAGE and Western Blot

Ten μ g/well of recombinant periplasmic extracts and 1.4 μ g/well from 2C and 7E clones, were resolved on pre-made SDS-PAGE 4%–20%, (ThermoScientific/USA) and transferred to PVDF membranes. After 1 h blocking with PBS-BSA at RT, membranes were incubated with anti-c-Myc mAb (Sigma/USA) diluted 1/750 in PBS-T-BSA and then after washes with PBS-T, membranes were incubated with anti-mouse IgG HRP diluted (1/2,000)(Invitrogen/USA) for 1 h, washed with PBS-T, and developed with 3 mg/ml of 3,3-diaminobenzidine in PBS and 30% hydrogen peroxide diluted 1:1,000.

Evaluation of Specificity of Single Domain Antibodies by ELISA

The specificity of the purified sdAb 2C and 7E, were evaluated by ELISA using 1µg of target Ag85Bp/HLA-A*0201 and nontargets Esat-6p/HLA-A*0201 and Acr1p/HLA-A*0201 as which were immobilized on streptavidin plates. Complexes were incubated with 5 µg of sdAb followed by incubation with 1/1,000 dilution of anti-c-Myc Ab labelled with HRP (Sigma-Aldrich/Ireland Ltd) by 1h. After several washes with PBS-T, the reaction was developed with 50 µl of TMB. The reaction was stopped with 1M of H_2SO_4 . OD_{450nm} was measured in an Infinitum F50 microplate ELISA reader (Tecan/Switzerland). Three experiments were carried out by duplicated for 2C and 2 experiments for 7E.

Ex Vivo Specificity of Single Domain 2C on the Surface of T2 Cells

To assess the ability to sdAb 2C to recognize the Ag85Bp/HLA-A*0201 an *ex vivo* assay was performed, by using the HLA-A*0201 positive T2 lymphoblastic human cell line (kindly donate by Dr. Patricia Gorocica INER-México). These cells are characterized by export empty HLA class I molecules due to a processing defect by homozygous deletion of the MHC class II region located on chromosome 6 including the TAP1 and TAP2 (transporters associated with antigen processing) genes which encode the transporter proteins (28). Cells were maintained in RPMI-1640 medium supplemented with 20% (vol/vol) fetal bovine serum (FBS) (Gibco/USA), at 37°C, 5% C0₂. T2 cells ($6x10^5$) were placed on flat bottom, 24 well cell culture plates (Costar/USA) in RPMI free serum in absence and presence of peptides from Ag85B and Esat-6.

In order to confirm the presence of HLA-ABC on T2 cells, they were incubated with W6/32 mAb (1µg/million cells), for 30 min on ice, after 3 washes, goat anti-mouse IgG Alexa fluor 488 (Invitrogen/USA) at 1/2,000 dilution was added. Samples were fixating with 0.5% PFA and the slides were mounted with vectashield (Vector Laboratories/USA). Then, cells ($6x10^5$) were incubated with 80 µg of either Ag85Bp(199-207) target peptide and Esat-6p(82-90) as non-target. Twenty µg/ml of β 2m (Sigma) was

added according to (29). Cells were incubated for 8 h at 37°C in 5% CO₂ atmosphere, after that cells were washed twice with PBS and incubated with 10 µg of sdAb 2C ON at 4°C in agitation, and then washed 2 times, with PBS, following by incubation with anti-c-Myc Ab (Santa Cruz-/Europe) (1/100) for 1 h at RT. After three washes, anti-mouse IgG Alexa fluor 488 (Invitrogen/USA) (1/2,000) was added and samples were incubated for 1 h at RT. After three washes, Hoechst 33343 (Life technologies/USA) diluted 1/9,000 was used for 10 min for nucleus staining. sdAb 2C was evaluated on cells in the absence of peptides and as a staining control in one condition, no domain was added. Fluorescence images were acquired with Olympus BX41, (Fluorescence-Microcopy) using the 100x magnifying lens, the digital images were captured with Zen 2.6 blue edition software, and the capture parameters, exposure time and intensity for each staining system, were applied in both control and problem samples.

Kinetic Binding Assays for the Assessment of the Interaction of sdAb 2C With Ag85B/ HLA-A*0201 Complex by Biolayer Interferometry

The binding kinetics and the determination of the dissociation constant (K_D) for the sdAb 2C against the Ag85Bp/HLA-A*0201 complex were performed using Biolayer Interferometry (BLI) at 25°C. Streptavidin biosensors in an Octet RED96 system (FortéBio Inc. San Jose, CA, USA) were used. The assays were performed on black bottom 96-well microplates (Greiner Bio-One 655209) in a total volume of 200 µl with orbital shaking at 1000 rpm. Experiments were controlled with the software Data Acquisition 8.2 (ForteBio, Inc.) For the BLI experiment, a baseline was established using 1x Kinetics buffer (FortéBio Inc. San Jose, CA, USA). Then, the biotinylated Ag85Bp/HLA-A*0201 complex at 25 ng was allowed to bind to streptavidin sensor for 5 min, followed by washing with the same buffer to eliminate nonspecific binding. Next, the purified sdAb was bound to the Ag85Bp/HLA-A*0201 complex in the biosensor and the association rate was measured (k_a) . In the last step, the dissociation rate (k_d) of the antibodycomplex was obtained. The BLI experiment was done with six different concentrations of the sdAb 2C from (2.54 to 81.3 µM), one well with 200 µl without sdAb was used as a negative control. New streptavidin biosensors were used for each experiment. The binding of sdAb 2C at 10 μ M, to non-target Esat-6p/HLA-A*0201 was also tested. The data were processed using the Octet Data Analysis Software version 8.2 (FortéBio Inc. San Jose, CA, USA) according to a 1:1 model.

Statistical Analysis

GraphPad Prism version 6.0c software was used to analyze the results. For the statistical analysis, one-way ANOVA multiple comparisons with Sidak's post Hoc correction was used.

RESULTS

Evaluation of Peptide HLA-A Complexes With W6/32 Antibody

Ag85Bp/HLA-A*0201, Esat-6p/HLA-A*0201, and Acr1p/HLA-A*0201 immobilized on streptavidin plates were recognized by W6/32 mAb (**Figure 1A**), indicating this result that peptides HLA-A*0201 complexes were correctly folded. In contrast, the mAb did not recognize the complexes bound to non-coated streptavidin control plate (**Figure 1B**), showing the results that direct binding of the complexes in the plates could lead to a loss of conformation. The biotinylated complexes bound to streptavidin plates were not recognized for streptavidin-HRP, an indication that biotinylated complexes were correctly oriented by streptavidin on coated plates (**Figure 1A**). In contrast, the positive signal obtained with the complexes bound to the non-coated streptavidin plate (**Figure 1B**), was an indication that exposed biotin in unfolded complexes was being recognized by streptavidin-HRP.

Screening of the Human Single DAb Phage Library Binding to Ag85B_{p199-207} HLA-A*0201/Complexes

The number of phage particles from sdAb library during the three rounds of selection was consistent with the published protocol (24). The final output titers of phage particles showed an enrichment factor of 25 (**Table 1**). From 94 clones evaluated by monoclonal phage-ELISA, only 7 clones showed absorbance 10 fold higher than the negative control (clone not reactive to the





SelS Selection Round	Ag85Bp/ HLA-A*0201	Phages Input	Eluted phages/ml	Eluted/input	Enrichment Factor*	Amplified Phages/ml	Tween 20/ # washes
1	7.5µg	5.0×10 ¹²	6.0×10 ⁶	1.2×10 ⁻⁶	1.0	84×10 ¹²	1.0%/15
2	2.5µg	5.0×10 ¹²	1.4×10 ⁸	2.8×10 ⁻⁵	23	7×10 ¹²	0.1%/15
3	1.25µg	5.0×10 ¹²	1.5 ×10 ⁸	3.0×10 ⁻⁵	25	_	0.1%/25

TABLE 1 | Selective enrichment of phage domain antibody after 3 rounds of biopanning.

*Enrichment factor was determined by dividing the eluted/input ratio in each round by the ratio in the first round. according to Bagheri et al. (30) Zhang et al. (31).

Ag85Bp/HLA-A*0201) (**Figure 2A**). From those, clones 2C, 3C, and 7E did not recognized streptavidin (**Figure 2B**) and all of them showed specific binding by ELISA to the target complex and but none bound to non-target complexes, Esat-6p/HLA-A*0201, and Acr1p/HLA-A*0201 (**Figure 2C**).

Domain Antibody Sequencing

BLAST search analysis of sdAb 2C, 3C, and 7E sequences, showed that all them matched with 122 amino acids length of immunoglobulin heavy chain variable region, partial (>ABM67233.1 Homo sapiens). Sequences corresponded to a human dAb with a length of 160 and 159 amino acids residues for 2C and 7E domains, respectively, the protein sequence of the 3C domain was exactly the same as 2C but shorter in length, 3C had only 151 amino acids due to the presence of a stop codon. For all sequences, the three complementarity determining regions (CDR) and the c-Myc tag sequence were identified. Clones 2C and 7E were selected to continue with de antibody expression phase.

Production of Single Domain Antibodies

sdAb 2C was produced in *E. coli* HB2151. The expression and purification of sdAb 2C is shown in **Figure 3A**. Purified sdAb 2C with the expected molecular mass of \approx 15 kDa was detected by Coomassie blue staining (**Figure 3A**, line 3) and antibody was recognized by anti-c-Myc Ab on Western blot (**Figure 3A**, line 4). The yield of production of 2C was 1418 µg from 50 ml of culture, (Results obtained with 7E are not shown).

Specificity of Single Domain Antibodies by ELISA

The recognition of the Ag85Bp/HLA-A*0201 complex by sdAb 2C was evaluated by ELISA, using Esat-6p/HLA-A*0201 and Acr1p/HLA-A*0201 as non-targets. The results are shown in **Figure 3B**. The binding of sdAb 2C to Ag85Bp/HLA-A*0201 complex was highly specific, showing statistically significant differences with respect to non-targed complexes. On the other hand, sdAb 7E, showed a nonspecific signal absorbance ratios for both target and non-target complexes (Results not shown).



FIGURE 2 | Monoclonal phage-domain Immunoassay. (A) Clones from the third round of panning. Positives clones showed signal recognition to Ag85Bp/HLA-A*0201 of 10 fold higher than negative control (NC), well with unrelated sdAb. Line indicates the threshold used to define a positive result. (B) Positive clones exposed to streptavidin coated plates (without complex). (C) Clones 2C, 3C, and 7E that showed the lowest recognition signal towards streptavidin tested with nontarget peptides.



Ex Vivo Specificity of Soluble Domain 2C on T2 Cells Surface

independent experiments and significant differences are indicated by asterisk (p < 0.05).

Surface expression of HLA-A molecules in T2 cells was demonstrable by binding of the W6/32 mAb in absence and in presence of either Ag85Bp₁₉₉₋₂₀₇ or Esat-66_{p82-90} peptides (**Supplementary Figure S1**). However, sdAbs 2C showed different recognition patterns, whereby the sdAb 2C fluorescence signal was observed only on T2 cells exposed to the Ag85Bp₁₉₉₋₂₀₇ (**Figure 4**). About 6.4% of positive events were observed, but no positive signals were detected without sdAb 2C, or without Ag85Bp₁₉₉₋₂₀₇, or by incubation with the no-target Esat-6 peptide.

sdAb 2C Binding Affinity by Biolayer Interferometry

The association of sdAb 2C to Ag85Bp/HLA-A*0201 was measured by BLI. The affinity constant was calculated in terms of equilibrium dissociation constant (K_D) to be 15 + 0.20 μ M (**Figure 5A**). The sdAb interacts with the target complex in a concentration dependent manner and confirms the dissociation constant value (**Figure 5B**). The interaction of sdAb with Esat-6p/HLA-A*0201 was very low and the binding parameters could not be determined.

DISCUSSION

CD8+ T cells recognizing peptide epitopes bound to MHC/HLA class I molecules have the capacity to lyse the *M. tuberculosis* infected cells, thus contributing to the intracellular killing of the infecting organisms (16). Hence, binding of antibodies with TCR-like recognition specificity seemed desirable for developing antibody constructs, with mycobactericidal potentials. TCR-like Ab recognizing MHC class I bound antigenic peptides on antigen presenting cells, have previously been reported for the treatment of cancer, viral infections and autoimmune diseases (6). The aim toward TB immunotherapy has recently been

initiated by the selection of antibodies against the latency expressed Acr1/HLA class I restricted epitopes (14, 15). We report here on a TCR-like sdAb against an immunodominant HLA-A*0201 binding epitope of the Ag85B, selected by screening and selection from a human DAb phage display library. The secreted, fibronectin binding, mycolyl transferase protein is a strong immunogen in both infected and active TB cases and it has been used in several recombinant vaccine constructs (32–35). We chose the p199KLVANNTRL₂₀₇ peptide as the target epitope in this study, because its known immunodominance for the human CD8+ T cell responses in the context of HLA-A*0201 and it's a conserved sequence in the genome (16, 17, 19). These properties favoured the previous application of TCR-like sdAbs for targeting tumors and cells infected with other pathogens (36, 37).

The procedures used for the selection and evaluation of TCRlike sdAbs ensure the specificity of recognition between for the target and non-target molecules. In this work, all the biotinylated complexes were first evaluated through a comparative ELISA, using plates with or without streptavidin. The results showed that the streptavidin coated plates ensured an adequate arrangement and orientation of p/HLA-A complexes, which is necessary for finding the specific TCR-like sdAbs. Similarly, Ag85Bp/HLA-A*0201 was bound to magnetic pearl cover with streptavidin for the selection of recombinant phages (26). From the third-round of selection, clones, 2C and 3C of identical sequence and clone 7E bound to the target complex. However, after conversion to the soluble form, sdAb 7E lost its specificity for the target complex. Such a change in specificity was previously reported to be due to loss of structural support by the phage scaffold pIII protein for the anti-H1N1 influenza virus antibody's antigen-binding site (38). The production of sdAb 2C and 7E in E. coli HB2151 from periplasmatic extracts was satisfactory, compared with the previously reported production outputs (25, 39).

Although sdAb 2C was highly specific against the target complex, its affinity is low, i.e., in the range of 1–100 μM

	Nucleus	sdAb 2C	Merge
without sdAb 2C			
without peptide			
Esat-6p			
Ag85Bp			

FIGURE 4 | Recognition of *M. tuberculosis* peptide/HLA-A*0201 complex on T2 cells surface by sdAb 2C. Cells were exposed to Ag85B and Esat-6 peptides. As negative control cells without peptide were used. Peptide/HLA-A*0201 complexes were detected with sdAb 2C follow by anti-IgG coupled to Alexa Fluor 488 as secondary Ab. More than 200 fields were examined for each condition using the 100x magnification, by fluorescence microscopy.



FIGURE 5 | Real-time biolayer interferometry sensorgrams for determination of the binding affinity. (A) Sensorgram showing reference substrate binding, on streptavidine biosensors, of Ag85Bp/HLA-A*0201 complex with increased concentrations of sdAb 2C. Calculated affinity constant K_D is shown in the upper right of the sensorgram. Association constant, k_a : 1.93 × 10⁴ + 0.02 × 10⁴ M^{-1 s-1}; dissociation constant, k_a : 0.289 + 0.003 s⁻¹. (B) Steady state analysis of the binding response (nm) as a function of sdAb. Calculated K_D 15.0 + 0.7 μ M.

corresponding to the affinity binding interaction of TCR ligands. The low binding affinity known for TCR-like antibody fragments selected from human libraries can be improved using complementary technologies (40). An increased up to 100 fold of the initial affinity from scFv (single chain variable fragment) directed against the HLA-A2-pWT1126 complex, was achived by mutagenesis combined with yeast display based on one specific scFv-clone (41). The affinity could also be improved by recloning and re-selection of clones or by conversion of sdAbs into multivalent formats with higher avidity (4, 42). Though antibody affinity can be significant for its potential for immunotherapy, most interestingly however, it was found that low, rather than high antibody affinity has been reported to essential for the passive antibody therapy of a drosophila based model of Alzheimer's disease (43, 44). This was interpreted on the grounds that the low affinity anti-tau antibody may loosen up intracellular tau aggregates allowing better access of lysosomal degrading enzymes, while high affinity antibody may make these aggregates more compact and therefore more difficult to degrade. Such interpretation may be relevant also for the desired intracellular mycobactericidal action on M. tuberculosis infected macrophages.

Antibodies generated in animals against MHC-I recombinant tetramers are rarely TCR-like, because many of them recognize the α 3 domain of MHC-I and β 2 microglobulin (β 2m) (5) and also due to the reduced stability of recombinant p/HLA-A complex (45). However, the phage display antibody libraries have the advantage that the fusion proteins are exposed on the surface of phage particles, and recombinant p/HLA-A complex targets can be used through several selection rounds (37). The structure of human sdAbs also known as nanobodies, used in the present work represents a single variable domain based on the VH3-23 germline segment heavy chain with synthetic diversity introduced by PCR mutagenesis into all tree complementary determining regions (24). They are highly stable, easily produced in large quantity by E. coli, they are of low immunogenicity, small size (15 kDa), and can be fused with multiple tags (46-48). Consequently, sdAbs have been used for virus detection (49-51), for imaging in vivo, mainly in cancer due to their bio distribution, high tumor penetrance and fast clearance from the blood circulation (52). The immunotherapeutic potentials include therapeutic targets in cancer (53), antagonism of angiogenesis (54) and acting as metastasis inhibitors.

CONCLUSIONS AND PERSPECTIVES

The TCR-like identity of the sdAb 2C has been validated by its specific recognition of Ag85Bp/HLA-A*0201 on the surface of human T2 cell line. It will be of further interest to test, if the sdAb 2C can detect the expression of the HLA-A*0201-bound Ag85B peptide on human macrophages, which contain either replicating or dormant *M. tuberculosis* infection. A positive result would justify further engineering of the sdAb 2C to become an

immunotoxin, by conjugation with suitable apoptosis inducing ligands, i.e., *Pseudomonas* exotoxin A, Granzyme B or BH3 peptide (55, 56), in endeavour to develop a mycobactericidal immunotherapeutic agent.

Further, development of sdAb 2C will need evaluation in both HLA-A*0201 transgenic mice and in humans with active TB disease. The obtained results are showing the feasibility of selecting TCR-like sdAbs to other immunodominant epitopes of M. tuberculosis and their future development as potential immunotherapeutic adjuncts to the chemotherapy of TB.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

AUTHOR CONTRIBUTIONS

CE and JI conceived and supervised the study. CE, PO, MS-M, and KF performed experiments. TO, KF, MS-M, and PO designed experiments, analyzed data, and provided new tools and reagents. CE, JI, and PO wrote the manuscript. All authors contributed to the article and approved the submitted versión.

ACKNOWLEDGMENTS

We thank Cristina Parada for technical assistance, the University program PASPA-UNAM for a scholarships to CE to take out a Sabbatical stay in the Center for Host-Microbiome Interactions at the Guy's Campus of King's College London. This project was funded by a Research Institutional program "NUATEI" from Instituto de Investigaciones Biomédicas UNAM, México and by the EC/H2020-PHC-2014/EMI-TB grant entitled: Eliciting Mucosal Immunity to Tuberculosis. PO is a PhD student from Doctorado en Ciencias Bioquímicas at Universidad Nacional Autónoma de México (UNAM), and has a scholarship from the Consejo Nacional de Ciencia y 417 Tecnología (CONACYT), (CVU/707238).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: https://www.frontiersin.org/articles/10.3389/fimmu.2020. 577815/full#supplementary-material

SUPPLEMENTARY FIGURE 1 | Recognition of HLA-A molecules on T2 cell surface. Cells were loaded with Ag85B and Esat-6 peptides and cells without peptide as negative control. HLA-A molecules were detected with W6/32 as primary Ab, by anti c-Myc Ab and anti IgG coupled to Alexa Fluor 488 as secondary Abs. More than 200 fields were observed for each condition evaluated using the 100x magnification, by fluorescence microscopy.

REFERENCES

- World Health Organization. *Global tuberculosis report* (2018). Available at: https://www.who.int/es/news-room/fact-sheets/detail/tuberculosis (Accessed Jun 16, 2018).
- Hoagland DT, Liu J, Lee RB, Lee RE. New agents for the treatment of drugresistant *Mycobacterium tuberculosis*. *Adv Drug Deliv Rev* (2016) 102:55–72. doi: 10.1016/j.addr.2016.04.02
- Pawlowski A, Jansson M, Sköld M, Rottenberg ME, Källenius G. Tuberculosis and HIV co-infection. *PLoS Pathog* (2012) 8(2):e1002464. doi: 10.1371/ journal.ppat.1002464
- Chames P, Hufton SE, Coulie PG, Uchanska-Ziegler B, Hoogenboom HR. Direct selection of a human antibody fragment directed against the tumor Tcell epitope HLA-A1-MAGE-A1 from a nonimmunized phage-Fab library. *Proc Natl Acad Sci* (2000) 97(14):7969–74. doi: 10.1073/pnas.97.14.796
- Wittman VP, David W, Tiffany N, Francisca AN, Stephen W and Jon AW. Antibody targeting to a class I MHC-peptide epitope promotes tumor cell death. J Immunol (2006) 177:4187–95. doi: 10.4049/jimmunol.177.6.4187
- Dahan R, Reiter Y. T-cell-receptor-like antibodies generation, function and applications. Expert Rev Mol Med (2012) 14:e6. doi: 10.1017/erm.2012.2
- Saeed M, van BM, Zalba S, Schooten E, Rens JA, Koning GA, et al. Targeting melanoma with immunoliposomes coupled to anti-MAGE A1 TCR-like single-chain antibody. *Int J Nanomed* (2016) 11:955–75. doi: 10.2147/ IJN.S96123
- Porgador A, Yewdell JW, Deng Y, Bennink JR, Germain RN. Localization, quantitation, and in situ detection of specific peptide-MHC class I complexes using a monoclonal antibody. *Immunity* (1997) 6:715–26. doi: 10.1016/s1074-7613(00)80447-1
- Cohen CJ, Sarig O, Yamano Y, Tomaru U, Jacobson S, Reiter Y. Direct phenotypic analysis of human MHC class I antigen presentation: visualization, quantitation, and in situ detection of human viral epitopes using peptide-specific, MHC-restricted human recombinant antibodies. *J Immunol* (2003) 170(8):4349–61. doi: 10.4049/jimmunol.170.8.4349
- Denkberg G, Cohen CJ, Lev A, Chames P, Hoogenboom HR, Reiter Y. Direct visualization of distinct T cell epitopes derived from a melanoma tumorassociated antigen by using human recombinant antibodies with MHCrestricted T cell receptor-like specificity. *Proc Natl Acad Sci* (2002) 99 (14):9421–6. doi: 10.1073/pnas.132285699
- Shams H, Klucar P, Weis SE, Lalvani A, Moonan PK, Safi H, et al. Characterization of a *Mycobacterium tuberculosis* peptide that is recognized by human CD4+ and CD8+ T cells in the context of multiple HLA alleles. *J Immunol* (2004) 173 (3):1966–77. doi: 10.4049/jimmunol.173.3.1966
- Klein MR, Smith SM, Hammond AS, Ogg GS, King AS, Vekemans J, et al. HLA-B*35-restricted CD8 T cell epitopes in the antigen 85 complex of Mycobacterium tuberculosis. J Infect Dis (2001) 183(6):928–34. doi: 10.1086/ 319267
- Lewinsohn DA, Winata E, Swarbrick GM, Tanner KE, Cook MS, Null MD, et al. Immunodominant tuberculosis CD8 antigens preferentially restricted by HLA-B. *PLoS Pathog* (2007) 13(9):1240–9. doi: 10.1371/journal.ppat.0030127
- Dass SA, Norazmi MN, Dominguez AA, Miguel MESGS, Tye GJ. Generation of a T cell receptor (TCR)-like single domain antibody (sDAb) against a *Mycobacterium Tuberculosis* (Mtb) heat shock protein (HSP) 16kDa antigen presented by Human Leukocyte Antigen (HLA)-A*02. *Mol Immunol* (2018) 101:189–96. doi: 10.1016/j.molimm.2018.07.001
- Dass SA, Norazmi MN, Acosta A, Sarmiento ME, Tye GJ. TCR-like domain antibody against *Mycobacterium tuberculosis* (Mtb) heat shock protein antigen presented by HLA-A*11 and HLA-A*24. *Int J Biol Macromol* (2020) 155:305–14. doi: 10.1016/j.ijbiomac.2020.03.229
- Comas I, Chakravartti J, Small PM, Galagan J, Niemann S, Kremer K, et al. Human T cell epitopes of *Mycobacterium tuberculosis* are evolutionarily hyperconserved. *Nat Genet* (2010) 6:498–503. doi: 10.1038/ng.590
- Axelsson –R, Loxton AG, Walzl G, Ehlers MM, Kock MM, Zumla A, et al. Broad profile of co-dominant epitopes shapes the peripheral *Mycobacterium tuberculosis* specific CD8+ T-cell immune response in South African patients with active tuberculosis. *PLoS One* (2013) 3:e58309. doi: 10.1371/ journal.pone.0058309
- Geluk A, van Meijgaarden KE, Franken KL, Drijfhout JW, D'Souza S, Necker A, et al. Identification of major epitopes of *Mycobacterium tuberculosis* AG85B that

are recognized by HLA-A*0201-restricted CD8+ T cells in HLA-transgenic mice and humans. *J Immunol* (2000) 165(11):6463–71. doi: 10.4049/jimmunol.165.11.6463

- Weichold FF, Mueller S, Kortsik C, Hitzler WE, Wulf MJ, Hone DM, et al. Impact of MHC class I alleles on the *M. tuberculosis* antigen-specific CD8+ Tcell response in patients with pulmonary tuberculosis. *Genes Immun* (2007) 8 (4):334–43. doi: 10.1038/sj.gene.6364392
- Prezzemolo T, van Meijgaarden KE, Franken KLMC, Caccamo N, Dieli F, Ottenhoff, et al. Detailed characterization of human Mycobacterium tuberculosis specific HLA-E restricted CD8+ T cells. *Eur J Immunol* (2018) 48(2):293–305. doi: 10.1002/eji.201747184
- Denkberg G, Cohen CJ, Segal D, Kirkin AF, Reiter Y. Recombinant human single-chain MHC-peptide complexes made from *E. coli* By in vitro refolding: functional single-chain MHC-peptide complexes and tetramers with tumor associated antigens. *Eur J Immunol* (2000) 30(12):3522–32. doi: 10.1002/1521-4141(2000012)30:12
- Parham P, Barnstable CJ, Bodmer WF. Use of a monoclonal antibody (W6/32) in structural studies of HLA-A,B,C, antigens. J Immunol (1979) 23(1):342–9.
- 23. Mora GM de L, Duenas GA, Hernández MJ, De la Cruz HE, Pérez CE, Weiss SB, et al. Up-regulation of HLA class-I antigen expression and antigen-specific CTL response in cervical cancer cells by the demethylating agent hydralazine and the histone deacetylase inhibitor valproic acid. *J Transl Med* (2006) 4:55. doi: 10.1186/1479-5876-4-55
- Lee CM, Iorno N, Sierro F, Christ D. Selection of human antibody fragments by phage display. *Nat Protoc* (2007) 2(11):3001-8. doi: 10.1038/ nprot.2007.448
- Abou El-Magd RM, Vozza NF, Tuszynski JA, Wishart DS. Isolation of soluble scFv antibody fragments specific for small biomarker molecule, L-Carnitine, using phage display. J Immunol Methods (2016) 428:9–19. doi: 10.1016/ j.jim.2015.11.006
- Santich BH, Liu H, Liu C, Cheung NK. "Generation of TCR-Like Antibodies Using Phage Display". In: G Houen, editor. *Methods in molecular biology*, vol. 1348. New York: NY human Press (2015). p. 191–204.
- Liu JL, Goldman ER, Zabetakis D, Walper SA, Turner KB, Shriver-Lake LC, et al. Enhanced production of a single domain antibody with an engineered stabilizing extra disulfide bond. *Microb Cell Fact* (2015) 14:158. doi: 10.1186/ s12934-015-0340-3
- DeMars R, Spies T. New genes in the MHC that encode proteins for antigen processing. Trends Cell Biol (1992) 3:81–6. doi: 10.1016/0962-8924(92)90077-z
- Atzin-Méndez JA, López-González JS, Báez R, Arenas-Del Angel MC, Montaño LF, Silva-Adaya, et al. Expansion of quiescent lung adenocarcinoma CD8+ T cells by MUC1-8-mer peptide-T2 cell-β2 microglobulin complexes. Oncol Rep (2016) 1:33–42. doi: 10.3892/or.2015.4328
- Bagheri S, Yousefi M, Safaie Qamsari E, Riazi-Rad F, Abolhassani M, Younesi V, et al. Selection of single chain antibody fragments binding to the extracellular domain of 4-1BB receptor by phage display technology. *Tumour Biol* (2017) 39 (3):1010428317695924. doi: 10.1177/1010428317695924
- 31. Zhang F, Chen Y, Ke Y, Zhang L, Zhang B, Yang L, et al. Single Chain Fragment Variable (scFv) Antibodies Targeting the Spike Protein of Porcine Epidemic Diarrhea Virus Provide Protection against Viral Infection in Piglets. *Viruses* (2019) 11(1):58. doi: 10.3390/v11010058
- Kadir NA, Sarmiento ME, Acosta A, Norazmi MN. Cellular and humoral immunogenicity of recombinant *Mycobacterium smegmatis* expressing Ag85B epitopes in mice. *Int J Mycobacteriol* (2016) 5(1):7–13. doi: 10.1016/ j.ijmyco.2015.09.006
- 33. Babaki M, Soleimanpour S, Rezaee SA. Antigen 85 complex as a powerful Mycobacterium tuberculosis immunogene: Biology, immune-pathogenicity, applications in diagnosis, and vaccine design. Microb Pathog (2017) 112:20–9. doi: 10.1016/j.micpath.2017.08.040
- 34. Horwitz MA, Harth G, Dillon BJ, Maslesa-Galic' S. Recombinant bacillus calmette-guerin (BCG) vaccines expressing the *Mycobacterium tuberculosis* 30-kDa major secretory protein induce greater protective immunity against tuberculosis than conventional BCG vaccines in a highly susceptible animal model. *Proc Natl Acad Sci U S A* (2000) 25:13853–8. doi: 10.1073/ pnas.250480397
- Olsen WA, van PLA, Meng OL, Birk RP, Andersen P. Protection of mice with a tuberculosis subunit vaccine based on a fusion protein of antigen 85b and esat-6. *Infect Immun* (2001) 5:2773–8. doi: 10.1128/IAI.69.5.2773-2778.2001

- 36. De Groeve K, Deschacht N, De Koninck C, Caveliers V, Lahoutte T, Devoogdt N, Muyldermans S, De Baetselier P, Raes G. Nanobodies as tools for in vivo imaging of specific immune cell types. J Nucl Med (2010) 51:782–9. doi: 10.2967/jnumed.109.070078
- Ingram JR, Schmidt FI, Ploegh HL. Exploiting Nanobodies' Singular Traits. *Annu Rev Immunol* (2018) 36:695–715. doi: 10.1146/annurev-immunol-042617-053327
- 38. Kaku Y, Noguchi A, Okutani A, Inoue S, Tanabayashi K, Yamamoto Y, et al. Altered specificity of single-chain antibody fragments bound to pandemic H1N1-2009 influenza virus after conversion of the phage-bound to the soluble form. *BMC Res Notes* (2012) 5:483. doi: 10.1186/1756-0500-5-483
- Ruano-Gallego D, Fraile GC, Fernández LÁ. Screening and purification of nanobodies from *E. coli* culture supernatants using the hemolysin secretion system. *Microb Cell Fact* (2019) 18(1):47. doi: 10.1186/s12934-019-1094-0
- Høydahl L, Frick R, Sandlie I, Løset GÅ. Targeting the MHC Ligandome by Use of TCR-Like Antibodies. *Antibodies Basel* (2019) 8(2):32. doi: 10.3390/ antib8020032
- Zhao Q, Ahmed M, Tassev DV, Hasan A, Kuo TY, Guo HF, et al. Affinity maturation of T-cell receptor-like antibodies for Wilms tumor 1 peptide greatly enhances therapeutic potential. *Leukemia* (2015) 29(11):2238–47. doi: 10.1038/leu.2015.125
- Zhu X, Wang L, Liu R, Flutter B, Li S, Ding J, et al. COMBODY: one-domain antibody multimer with improved avidity. *Immunol Cell Biol* (2010) 88 (6):667–75. doi: 10.1038/icb.2010.21
- Krishnaswamy S, Huang HW, Marchal IS, Ryoo HD, Sigurdsson EM. Neuronally expressed anti-tau scFv prevents tauopathy-induced phenotypes in Drosophila models. *Neurobiol Dis* (2020) 137:104770. doi: 10.1016/ j.nbd.2020.104770
- 44. Congdon E, Chukwu J, Shamir D, Deng J, Ujla D, Sait H, et al. Tau antibody chimerization alters its charge and binding, thereby reducing its cellular uptake and efficacy. *EBioMedicine* (2019) 42:157–73. doi: 10.1016/ j.ebiom.2019.03.033
- Cohen M, Reiter Y. T-Cell Receptor-Like Antibodies: Targeting the Intracellular Proteome Therapeutic Potential and Clinical Applications. *Antibodies* (2013) 2(3):517–34. doi: 10.3390/antib2030517
- Salvador JP, Vilaplana L, Marco MP. Nanobody: outstanding features for diagnostic and therapeutic applications. *Anal Bioanal Chem* (2019) 411 (9):1703–13. doi: 10.1007/s00216-019-01633-4
- Wang P, Li G, Yan J, Hu Y, Zhang C, Liu X, et al. Bactrian camel nanobodybased immunoassay for specific and sensitive detection of Cry1Fa toxin. *Toxicon* (2014) 92:186–92. doi: 10.1016/j.toxicon.2014.10.024

- Mass DR, Sepulveda J, Pernthaner A, Shoemaker CB. Alpaca (Lama pacos) as a convenient source of recombinant camelid heavy chain antibodies (VHHs). J Immunol Methods (2007) 324(1-2):13–25. doi: 10.1016/j.jim.2007.04.008
- 49. Ma Z, Tianyu W, Zhiwei L, Xuyang G, Yangsheng T, Li Y, et al. A novel biotinylated nanobody-based blocking ELISA for the rapid and sensitive clinical detection of porcine epidemic diarrhea virus. *J Nanobiotechnol* (2019) 17(1):96. doi: 10.1186/s12951-019-0531-x
- 50. Gelkop S, Sobarzo A, Brangel P, Vincke C, Romão E, Fedida-Metula S, et al. The Development and Validation of a Novel Nanobody-Based Competitive ELISA for the Detection of Foot and Mouth Disease 3ABC Antibodies in Cattle. Front Vet Sci (2018) 5:250. doi: 10.3389/fvets.2018.00250
- Zhu M, Gong X, Hu Y, Ou W, Wan Y. Streptavidin-biotin-based directional double Nanobody sandwich ELISA for clinical rapid and sensitive detection of influenza H5N1. J Transl Med (2014) 12:352. doi: 10.1186/s12967-014-0352-5
- Oliveira S, Heukers R, Sornkom J, Kok RJ, van Bergen En Henegouwen PM. Targeting tumors with nanobodies for cancer imaging and therapy. J Control Release (2013) 172(3):607–17. doi: 10.1016/j.jconrel.2013.08.298
- 53. Ji L, Dong C, Fan R, Qi S. A high affinity nanobody against endothelin receptor type B: a new approach to the treatment of melanoma. *Mol Biol Rep* (2020) 47(3):2137–47. doi: 10.1007/s11033-020-05313-w
- Ebrahimizadeh W, Mousavi SL, Javidan Z, Rajabibazl M. Production of Novel VHH Nanobody Inhibiting Angiogenesis by Targeting Binding Site of VEGF. Appl Biochem Biotechnol (2015) 176(7):1985–95. doi: 10.1007/s12010-015-1695-y
- Rosenblum M. Immunotoxins and toxin constructs in the treatment of leukemia and lymphoma. Adv Pharmacol (2004) 51:209–28. doi: 10.1016/ S1054-3589(04)51009-8
- Walensky LD, Kung AL, Escher I, Malia TJ, Barbuto S, Wright RD, et al. Activation of apoptosis in vivo by a hydrocarbon-stapled BH3 helix. *Science* (2004) 305(5689):1466–70. doi: 10.1126/science.1099191

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Ortega, Silva-Miranda, Torres-Larios, Campos-Chávez, Franken, Ottenhoff, Ivanyi and Espitia. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.