



## King's Research Portal

DOI:

[10.1007/s00208-020-02123-0](https://doi.org/10.1007/s00208-020-02123-0)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Lester, S., & Radziwill, M. (2021). Signs of Fourier coefficients of half-integral weight modular forms. *Mathematische Annalen*, 379(3-4), 1553-1604. <https://doi.org/10.1007/s00208-020-02123-0>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# SIGNS OF FOURIER COEFFICIENTS OF HALF-INTEGRAL WEIGHT MODULAR FORMS

STEPHEN LESTER AND MAKSYM RADZIWIŁŁ

ABSTRACT. Let  $g$  be a Hecke cusp form of half-integral weight, level 4 and belonging to Kohnen's plus subspace. Let  $c(n)$  denote the  $n$ th Fourier coefficient of  $g$ , normalized so that  $c(n)$  is real for all  $n \geq 1$ . A theorem of Waldspurger determines the magnitude of  $c(n)$  at fundamental discriminants  $n$  by establishing that the square of  $c(n)$  is proportional to the central value of a certain  $L$ -function. The signs of the sequence  $c(n)$  however remain mysterious.

Conditionally on the Generalized Riemann Hypothesis, we show that  $c(n) < 0$  and respectively  $c(n) > 0$  holds for a positive proportion of fundamental discriminants  $n$ . Moreover we show that the sequence  $\{c(n)\}$  where  $n$  ranges over fundamental discriminants changes sign a positive proportion of the time.

Unconditionally, it is not known that a positive proportion of these coefficients are non-zero and we prove results about the sign of  $c(n)$  which are of the same quality as the best known non-vanishing results. Finally we discuss extensions of our result to general half-integral weight forms  $g$  of level  $4N$  with  $N$  odd, square-free.

## 1. INTRODUCTION

Let  $k \geq 2$  be an integer and  $g$  be a weight  $k + \frac{1}{2}$  cusp form for  $\Gamma_0(4)$ . Every such  $g$  has a Fourier expansion

$$g(z) = \sum_{n \geq 1} c(n) n^{\frac{(k-1/2)}{2}} e(nz).$$

The Fourier coefficients  $c(n)$  encode arithmetic information. For instance under certain hypotheses, Waldspurger's Theorem shows that for fundamental discriminants  $d$ , the value  $|c(|d|)|^2$  is proportional to the central value of an  $L$ -function, so that the magnitude of the  $L$ -function essentially determines the size of the coefficient  $c(n)$ . However for  $g$  with real Fourier coefficients, their signs remain mysterious. In this article we contribute towards understanding the sign of such coefficients at fundamental discriminants through examining the number of coefficients which are positive (respectively negative) as well as the number of sign changes in-between.

---

*Date:* August 21, 2020.

Signs of Fourier coefficients of half-integral weight forms have been studied by many authors following the works of Knopp-Kohnen-Pribitkin [13] and Bruinier-Kohnen [3], the former of which showed that such forms have infinitely many sign changes. Subsequent works [9, 17, 20] showed that the sequence  $\{c(n)\}$  exhibits many sign changes under suitable conditions (such as the form having real Fourier coefficients). Notably, Jiang-Lau-Lü-Royer-Wu [11] showed for suitable  $g$  that for every  $\varepsilon > 0$  there are more than  $\gg X^{2/9-\varepsilon}$  sign changes along square-free integers  $n \in [1, X]$ . They also showed this result can be improved assuming the Generalized Lindelöf Hypothesis<sup>1</sup> with an exponent of  $1/4$  in place of  $2/9$ .

For an integral weight Hecke cusp form  $f$  the second named author and Matomäki [22] proved a stronger result, establishing a positive proportion of sign changes along the positive integers. This uses the multiplicativity of the Fourier coefficients of  $f$  in a fundamental way. Fourier coefficients of half-integral weight Hecke cusp forms lack this property, except at squares. So one may wonder whether the Fourier coefficients of half-integral weight Hecke cusp forms also exhibit a positive proportion of sign changes, along the sequence of fundamental discriminants.

In this article we answer this question in the affirmative. We show that under the assumption of the Generalized Riemann Hypothesis (GRH) there exists a positive proportion of fundamental discriminants at which the Fourier coefficients of a suitable half-integral weight form are positive as well as a positive proportion at which the coefficients are negative. Moreover, we show under GRH that the coefficients exhibit a positive proportion of sign changes along the sequence of fundamental discriminants.

For simplicity, our results are stated for the Kohnen space  $S_{k+1/2}^+$ , which consists of all weight  $k+1/2$  modular forms on  $\Gamma_0(4)$  whose  $n$ th Fourier coefficient equals zero whenever  $(-1)^k n \equiv 2, 3 \pmod{4}$ . In this space Shimura's correspondence between half-integral weight forms and integral weight forms is well understood. Kohnen proved [14] there exists a Hecke algebra isomorphism between  $S_{k+1/2}^+$  and the space of level 1 cusp forms of weight  $2k$ . Also, every Hecke<sup>2</sup> cusp form  $g \in S_{k+1/2}^+$  can be normalized so that it has real coefficients<sup>3</sup> and from here on we assume that  $g$  has been normalized in this way.

---

<sup>1</sup>Here the Generalized Lindelöf Hypothesis is assumed for  $L$ -functions attached to quadratic twists of level 1 Hecke eigenforms.

<sup>2</sup>We call a weight  $k+1/2$  cusp form on  $\Gamma_0(4)$  a Hecke cusp form if it is an eigenfunction of the Hecke operator  $T_{p^2}$  (see [27]) for each  $p > 2$ .

<sup>3</sup>The numbers  $c(n)n^{(k-1/2)/2}$  lie in the field generated over  $\mathbb{Q}$  by the Fourier coefficients of its Shimura lift, which is a level 1 Hecke eigenform of weight  $2k$ , so these numbers are real and algebraic (see Proposition 4.2 of [19] and also the remarks before Theorem 1 of [18]).

Let  $\mathbb{N}^{\flat}$  denote the set of fundamental discriminants of the form  $8n$  with  $n > 0$  odd, square-free. Also, let  $\mathbb{N}_g^{\flat}(X) = \{n \in \mathbb{N}^{\flat} \cap [1, X] : c(n) \neq 0\}$ .

**Theorem 1.** *Assume the Generalized Riemann Hypothesis. Let  $k \geq 2$  be an integer and  $g \in S_{k+1/2}^+$  be a Hecke cusp form. Then for all  $X$  sufficiently large the number of sign changes of the sequence  $\{c(n)\}_{n \in \mathbb{N}_g^{\flat}(X)}$  is  $\gg X$ .*

*In particular, for all  $X$  sufficiently large we can find  $\gg X$  integers  $d \in \mathbb{N}^{\flat} \cap [1, X]$  such that  $c(d) < 0$ , and we can find  $\gg X$  integers  $d \in \mathbb{N}^{\flat} \cap [1, X]$  such that  $c(d) > 0$ .*

The proof of Theorem 1 uses the explicit form of Waldspurger’s Theorem due to Kohnen and Zagier [18]. Given a Hecke cusp form  $g \in S_{k+1/2}^+$  they show that for each fundamental discriminant  $d$  with  $(-1)^k d > 0$  that

$$(1.1) \quad |c(|d|)|^2 = L(\tfrac{1}{2}, f \otimes \chi_d) \cdot \frac{(k-1)!}{\pi^k} \cdot \frac{\langle g, g \rangle}{\langle f, f \rangle}.$$

Here  $f$  is a weight  $2k$  Hecke cusp form of level 1 which corresponds to  $g$  as described above,  $L(s, f \otimes \chi_d)$  is the  $L$ -function

$$L(s, f \otimes \chi_d) = \sum_{n \geq 1} \frac{\lambda_f(n) \chi_d(n)}{n^s}, \quad \text{Re}(s) > 1$$

where  $\lambda_f(n)$  are the Hecke eigenvalues of  $f$ , normalized so that  $\lambda_f(1) = 1$  and  $|\lambda_f(n)| \leq d(n)$  for all  $n \geq 1$ , and  $\chi_d(\cdot)$  is the Kronecker symbol. Also,

$$\langle f, f \rangle = \int_{\text{SL}_2(\mathbb{Z}) \backslash \mathbb{H}} y^{2k} |f(z)|^2 \frac{dx dy}{y^2}, \quad \langle g, g \rangle = \frac{1}{6} \int_{\Gamma_0(4) \backslash \mathbb{H}} y^{k+1/2} |g(z)|^2 \frac{dx dy}{y^2}.$$

In Theorem 1 we assume GRH for  $L(s, f \otimes \chi_d)$  for every fundamental discriminant  $d$  with  $(-1)^k d > 0$ .

The restriction to  $d \in \mathbb{N}^{\flat}$  is important. As the proof of Theorem 3 below will show, it is easy to produce many positive (resp. negative) coefficients along *the integers*, assuming a suitable non-vanishing result.

**1.1. Unconditional results.** We also are able to prove a quantitatively weaker yet unconditional result. These results are significantly easier to obtain than our conditional result, Theorem 1

**Theorem 2.** *Let  $k \geq 2$  be an integer and  $g \in S_{k+1/2}^+$  be a Hecke cusp form. Then for any  $\varepsilon > 0$  and all  $X$  sufficiently large the sequence  $\{c(n)\}_{n \in \mathbb{N}_g^{\flat}(X)}$  has  $\gg X^{1-\varepsilon}$  sign changes.*

Theorems 1 and 2 quantitatively match the best known non-vanishing results for Fourier coefficients of half-integral weight forms that are proved using analytic techniques, conditionally under GRH [8] and unconditionally [25] (resp.). In particular,

Theorem 1 gives a different proof that a positive proportion of these coefficients are non-zero, for Hecke forms in the Kohnen space. It should be noted that Ono-Skinner [24] have shown that there exist  $\gg X/\log X$  fundamental discriminants at which these coefficients are non-zero for such forms.<sup>4</sup> However, their result does not give a quantitative lower bound on the size of the coefficients, whereas the analytic estimates do provide such information, which is crucial for our argument.

On the other hand, using the result of Ono-Skinner it is not difficult to produce both many positive Fourier coefficients and many negative ones, at *integers*.

**Theorem 3.** *Let  $k \geq 2$  be an integer and  $g \in S_{k+1/2}^+$  be a Hecke cusp form. Then, for all sufficiently large  $X$*

$$\#\{n \leq X : c(n) > 0\} \gg \frac{X}{\log X} \quad \text{and} \quad \#\{n \leq X : c(n) < 0\} \gg \frac{X}{\log X}.$$

**1.2. Extensions beyond the Kohnen plus space.** Since by now Shimura's correspondence is fairly well-understood (see [2]) we show in the Appendix that the conclusion of Theorems 1 and 2 holds for every half-integral weight Hecke cusp form on  $\Gamma_0(4)$ <sup>5</sup>. Additionally, we can prove analogs of Theorems 1 and 2 which hold for weight  $k + \frac{1}{2}$  ( $k \geq 2$ ) cusp forms  $g$  of level  $4N$  with  $N$  odd and square-free provided that  $g$  corresponds (through Shimura's correspondence) to an integral weight newform. The necessary modifications to our argument and precise statements of the results are in the Appendix.

**1.3. Numerical examples.** To illustrate our results above with a concrete example consider the following weight  $\frac{13}{2}$  Hecke cusp form

$$\delta(z) = \frac{60}{2\pi i} (2G_4(4z)\theta'(z) - G_4'(4z)\theta(z)) = \sum_{\substack{n \geq 1 \\ n \equiv 0,1 \pmod{4}}} \alpha_\delta(n)e(nz),$$

where

$$G_k(z) = \frac{1}{2} \cdot \zeta(1-k) + \sum_{n \geq 1} \sigma_{k-1}(n)e(nz) \quad \text{and} \quad \theta(z) = \sum_{n \in \mathbb{Z}} e(n^2 z).$$

The modular form  $\delta(z)$  corresponds to the modular discriminant  $\Delta(z)$  under the Shimura lift. Assuming GRH for  $L(\frac{1}{2}, \Delta \otimes \chi_d)$  for every fundamental discriminant  $d$ , Theorem 1 implies that there is a positive proportion of sign changes of  $\alpha_\delta(n)$  along the sequence of fundamental discriminants of the form  $8d$ . In fact numerical evidence

<sup>4</sup>As discussed above for forms in the Kohnen space the Fourier coefficients are algebraic integers which lie in a number field [19, Proposition 4.2] so that the Fundamental Lemma of [24] applies.

<sup>5</sup>For  $g \notin S_{k+1/2}^+$  is possible that  $c(8n)\mu^2(2n)$  is zero for each  $n \in \mathbb{N}$ , in this case we detect sign changes of  $\{c(n)\}$  where  $n$  ranges over  $\{n \leq X : n \text{ is even and } \mu^2(n)c(n) \neq 0\}$ .

below suggests that the Fourier coefficients of  $\delta(z)$  change sign approximately one half of the time. Given a subset  $\mathcal{L}$ , let  $\mathcal{S}_{\delta, \mathcal{L}}(X)$  denote the number of sign changes of  $\alpha_\delta(n)$  along  $n \in \mathcal{L} \cap [1, X]$ , and denote by  $\mathcal{N}_{\delta, \mathcal{L}}(X)$  the cardinality of  $\mathcal{L} \cap [1, X]$ . We then have the following numerical data.

$X$	$2 \times 10^5$	$2 \times 10^6$	$2 \times 10^7$	$2 \times 10^8$	$2 \times 10^9$
$\mathcal{S}_{\delta, \mathbb{N}}(X)$	50291	501163	5000867	50000368	500027782
$\frac{1}{\mathcal{N}_{\delta, \mathbb{N}}(X)} \mathcal{S}_{\delta, \mathbb{N}}(X)$	0.502915...	0.501163...	0.500086...	0.500003...	0.500027...

If we restrict to  $\mathbb{N}^{\flat} = \{8d : d \text{ odd, square-free}\}$  then we find the following data.

$X$	$2 \times 10^5$	$2 \times 10^6$	$2 \times 10^7$	$2 \times 10^8$	$2 \times 10^9$
$\mathcal{S}_{\delta, \mathbb{N}^{\flat}}(X)$	5049	50734	506589	5065686	50663938
$\frac{1}{\mathcal{N}_{\delta, \mathbb{N}^{\flat}}(X)} \mathcal{S}_{\delta, \mathbb{N}^{\flat}}(X)$	0.498223...	0.500740...	0.499980...	0.499963...	0.500033...

**1.4. Main Estimates.** The main results follow from the following three propositions. The first two of the propositions allow us to control the size of  $c(\cdot)$  by introducing a mollifier  $M(\cdot; \cdot)$  which is defined in Section 4.2. In our application the specific shape of the mollifier is crucial to the success of the method. We have constructed this mollifier to counteract the large values of

$$\exp \left( \sum_{p \leq |d|^\epsilon} \frac{\lambda_f(p)}{\sqrt{p}} \chi_d(p) \right)$$

that contribute to the bulk of the moments of  $L(\frac{1}{2}, f \otimes \chi_d)$ . Essentially, we are mollifying the  $L$ -function through an Euler product, as opposed to traditional methods which use a Dirichlet series. This approach was sparked by innovations in understanding of the moments of  $L$ -functions, such as the works of Soundararajan [30], Harper [6], and Radziwiłł-Soundararajan [26].

**Proposition 1.1.** *Assume GRH. Let  $g \in S_{k+1/2}^+$  be a Hecke cusp form. Also, let  $M(\cdot; \cdot)$  be as defined in Section 4.2. Then*

$$\sum_{\substack{n \leq X \\ 2n \text{ is } \square\text{-free}}} |c(8n)|^4 M((-1)^k 8n; \frac{1}{2})^4 \ll X.$$

**Proposition 1.2.** *Let  $g \in S_{k+1/2}^+$  be a Hecke cusp form. Also, let  $M(\cdot; \cdot)$  be as defined in Section 4.2. Then*

$$\sum_{\substack{n \leq X \\ 2n \text{ is } \square\text{-free}}} |c(8n)|^2 M((-1)^k 8n; \frac{1}{2})^2 \asymp X.$$

The other key ingredient of our results is an estimate for sums of Fourier coefficients summed against a short Dirichlet polynomial over short intervals. This is proved through estimates for shifted convolution sums of half-integral weight forms.

**Proposition 1.3.** *Let  $g$  be a cusp form of weight  $k + \frac{1}{2}$  on  $\Gamma_0(4)$ . Let  $(\beta(n))_{n \geq 1}$  be complex coefficients such that  $|\beta(n)| \ll_\varepsilon n^{\frac{1}{2} + \varepsilon}$  for all  $\varepsilon > 0$ . Let*

$$M((-1)^k n) := \sum_{m \leq M} \frac{\beta(m) \chi_{(-1)^k n}(m)}{\sqrt{m}}$$

where  $\chi_{(-1)^k n}(m)$  denotes the Kronecker symbol. Uniformly for  $1 \leq y, M \leq X^{\frac{1}{1092}}$

$$\begin{aligned} & \sum_{X \leq x \leq 2X} \left| \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} c(8n) M((-1)^k 8n) \right| \\ & \ll_k X \sqrt{y} \cdot \left( \frac{1}{X^{k+\frac{1}{2}}} \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} |c(8n) M((-1)^k 8n)|^2 \cdot n^{k-\frac{1}{2}} e^{-\frac{4\pi n}{X}} \right)^{1/2} + X^{1-\frac{1}{2184}+\varepsilon}. \end{aligned}$$

In particular, Proposition 1.3 holds for  $M(\cdot) = M(\cdot; \frac{1}{2})$  as defined in Section 4.2.

## 2. THE PROOF OF THEOREM 1

The basic method of proof follows a straightforward approach. Observe that since  $M((-1)^k n; \frac{1}{2}) > 0$  (see the discussion before and after (4.18)), if

$$\left| \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} c(8n) M((-1)^k 8n; \frac{1}{2}) \right| < \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} |c(8n)| M((-1)^k 8n; \frac{1}{2})$$

then the sequence  $\{\mu^2(2n)c(8n)\}$  must have at least one sign change in the interval  $[x, x+y]$ . To analyze the sums above we use a direct approach, which was developed in [22], where sign changes of integral weight forms was studied. The main inputs are Propositions 1.1, 1.2 and 1.3.

**Lemma 2.1.** *Let  $\Lambda \geq 1$ . There exists  $\delta > 0$  such that for  $2 \leq y \leq X^\delta$  we have for all but at most  $\ll X/\Lambda$  integers  $X \leq x \leq 2X$  that*

$$\left| \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} c(8n) M((-1)^k 8n; \frac{1}{2}) \right| \leq \Lambda \sqrt{y}.$$

*Proof.* This follows from Markov's inequality combined with Proposition 1.2 and Proposition 1.3, which holds for  $M(\cdot) = M(\cdot; \frac{1}{2})$ .  $\square$

**Lemma 2.2.** *Assume GRH. Let  $2 \leq y \leq X/2$ . Then for all sufficiently small  $\varepsilon > 0$  there exists a subset of integers  $X \leq x \leq 2X$  which contains  $\gg \varepsilon X$  integers such that*

$$\sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} |c(8n)| M((-1)^k 8n; \frac{1}{2}) > \varepsilon y.$$

*Proof.* For sake of brevity write  $C(n) = |c(8n)| M((-1)^k 8n; \frac{1}{2})$ . By Hölder's inequality

$$(2.1) \quad \sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free}}} C(n)^2 \leq \left( \sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free}}} C(n) \right)^{2/3} \left( \sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free}}} C(n)^4 \right)^{1/3}.$$

Applying Proposition 1.2 it follows that the LHS is  $\gg X$ . Applying Proposition 1.1 the second sum on the RHS is  $\ll X$ . Hence, we conclude that

$$\sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free}}} C(n) \gg X.$$

Also, by Proposition 1.2 we have

$$\sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free} \\ C(n) > 1/\varepsilon}} C(n) < \varepsilon \sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free}}} C(n)^2 \ll \varepsilon X.$$

So that for  $\varepsilon$  sufficiently small

$$(2.2) \quad \sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free} \\ C(n) \leq 1/\varepsilon}} C(n) \gg X.$$

Let  $U$  denote the subset of integers  $X \leq x \leq 2X$  such that

$$\sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free} \\ C(n) \leq 1/\varepsilon}} C(n) \leq \varepsilon y.$$

Using (2.2) we get that

$$(2.3) \quad X \ll \sum_{\substack{X+y \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free} \\ C(n) \leq 1/\varepsilon}} C(n) \leq \frac{1}{y} \sum_{X \leq x \leq 2X} \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free} \\ C(n) \leq 1/\varepsilon}} C(n)$$



where to get the last inequality note that each  $C(n)$  which appears in the sum on the LHS is counted  $\lfloor y \rfloor + 1$  times in the sums on the RHS. The RHS equals

$$\begin{aligned} &= \frac{1}{y} \left( \sum_{x \in U} + \sum_{\substack{X \leq x \leq 2X \\ x \notin U}} \right) \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free} \\ C(n) \leq 1/\varepsilon}} C(n) \\ &\ll \frac{1}{y} \cdot \left( (\varepsilon y)X + \frac{y}{\varepsilon} \cdot \sum_{\substack{X \leq x \leq 2X \\ x \notin U}} 1 \right). \end{aligned}$$

Since the first term above is of size  $\varepsilon X$  and using (2.3) we must have that the second term is  $\gg X$  since  $\varepsilon$  is sufficiently small. So  $\#\{X \leq x \leq 2X : x \notin U\} \gg \varepsilon X$ .  $\square$

*Proof of Theorem 1.* In Lemma 2.1 take  $\Lambda = \frac{1}{\varepsilon^2}$ ,  $y = \frac{1}{\varepsilon^6}$  so for except at most  $\ll \varepsilon^2 X$  integers  $X \leq x \leq 2X$  we have that

$$(2.4) \quad \left| \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} c(8n)M((-1)^k 8n; \frac{1}{2}) \right| \leq \Lambda \sqrt{y} = \frac{1}{\varepsilon^5}.$$

In Lemma 2.2 take  $y = \frac{1}{\varepsilon^6}$  so there are  $\gg \varepsilon X$  integers  $X \leq x \leq 2X$  such that

$$(2.5) \quad \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} |c(8n)|M((-1)^k 8n; \frac{1}{2}) > \varepsilon y = \frac{1}{\varepsilon^5}.$$

Combining (2.4) and (2.5) we get that there are  $\gg \varepsilon X$  integers  $X \leq x \leq 2X$  such that

$$\sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} |c(8n)|M((-1)^k 8n; \frac{1}{2}) > \left| \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} c(8n)M((-1)^k 8n; \frac{1}{2}) \right|.$$

Since  $M > 0$  (see the discussion after (4.18)) this implies there exists at least  $\gg \varepsilon X$  integers  $X \leq x \leq 2X$  such that  $[x, x+y]$  contains a sign change of the sequence  $\{\mu^2(2n)c(8n)\}$ . Since every sign change of  $\{\mu^2(2n)c(8n)\}$  on  $[X, 2X]$  yields at most  $\lfloor y \rfloor + 1$  intervals  $[x, x+y]$  which contain a sign change it follows that there are at least  $\gg \varepsilon \frac{X}{y} = \varepsilon^7 X$  sign changes in  $[X, 2X]$ , which completes the proof of Theorem 1.  $\square$

### 3. THE PROOFS OF THEOREM 2 AND THEOREM 3

3.1. **The proof of Theorem 2.** Throughout we will need the following estimate.

**Lemma 3.1.** *We have*

$$\sum_{\substack{X \leq n \leq 2X \\ 2n \text{ is } \square\text{-free}}} |c(8n)|^2 \asymp X.$$

*Proof.* This follows immediately by applying (1.1) along with Proposition 5.1 below with  $u = 1$ .  $\square$

We are now ready to start the preparations for the proof of Theorem 2, which as it turns out is considerably easier to prove than Theorem 1.

**Lemma 3.2.** *Let  $\Lambda \geq 1$ . There exists  $\delta > 0$  such that for all  $2 \leq y \leq X^\delta$  we have for all but at most  $\ll X/\Lambda$  integers  $X \leq x \leq 2X$  that*

$$\left| \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} c(8n) \right| \leq \Lambda \sqrt{y}.$$

*Proof.* This follows from Markov's inequality combined with Proposition 1.3 (with the choice  $\beta(1) = 1$  and  $\beta(m) = 0$  for all  $m \geq 2$ ) and Lemma 3.1.  $\square$

**Lemma 3.3.** *Let  $\varepsilon > 0$  and  $2 \leq y \leq X/2$ . Then there exist  $\gg X^{1-\frac{3}{2}\varepsilon}$  integers  $X \leq x \leq 2X$  such that*

$$\sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} |c(8n)| > \frac{y}{X^\varepsilon}.$$

*Proof.* Note that using (1.1) and Heath-Brown's result [7, Theorem 2] we get that

$$\sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free}}} |c(8n)|^4 \ll X^{1+\varepsilon}.$$

Hence, using the above estimate along with Lemma 3.1 we can apply Hölder's inequality as in (2.1) to get

$$(3.1) \quad \sum_{\substack{X \leq n \leq 2X \\ 2n \text{ is } \square\text{-free}}} |c(8n)| \gg X^{1-\varepsilon/2}.$$

Also, using Lemma 3.1 we have

$$\sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free} \\ |c(8n)| > X^\varepsilon}} |c(8n)| < X^{-\varepsilon} \sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free}}} |c(8n)|^2 \ll X^{1-\varepsilon}.$$

So combining this along with (3.1) we get

$$(3.2) \quad \sum_{\substack{X \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free} \\ |c(8n)| \leq X^\varepsilon}} |c(8n)| \gg X^{1-\varepsilon/2}.$$

Let  $U$  denote the subset of integers  $X \leq x \leq 2X$  such that

$$\sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free} \\ |c(8n)| \leq X^\varepsilon}} |c(8n)| \leq \frac{y}{X^\varepsilon}.$$

Using (3.2) and arguing as in (2.3) we get that

$$\begin{aligned} X^{1-\varepsilon/2} &\ll \sum_{\substack{X+y \leq 8n \leq 2X \\ 2n \text{ is } \square\text{-free} \\ |c(8n)| \leq X^\varepsilon}} |c(8n)| \leq \frac{1}{y} \sum_{X \leq x \leq 2X} \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free} \\ |c(8n)| \leq X^\varepsilon}} |c(8n)| \\ &= \frac{1}{y} \left( \sum_{x \in U} + \sum_{\substack{X \leq x \leq 2X \\ x \notin U}} \right) \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free} \\ |c(8n)| \leq X^\varepsilon}} |c(8n)| \\ &\ll \frac{1}{y} \cdot \left( \frac{y}{X^\varepsilon} X + yX^\varepsilon \cdot \sum_{\substack{X \leq x \leq 2X \\ x \notin U}} 1 \right). \end{aligned}$$

Since the first term is of size  $X^{1-\varepsilon}$  we must have that the second term is  $\gg X^{1-\varepsilon/2}$  so that  $\#\{X \leq x \leq 2X : x \notin U\} \gg X^{1-\frac{3}{2}\varepsilon}$ .  $\square$

We are now ready to prove Theorem 2.

*Proof of Theorem 2.* Let  $y = X^{6\varepsilon}$  and  $\Lambda = \frac{y^{1/2}}{X^\varepsilon}$ . By Lemma 3.2 we have

$$\left| \sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} c(8n) \right| \leq \frac{y}{X^\varepsilon}$$

for all  $X \leq x \leq 2X$  with at most  $X^{1+\varepsilon}y^{-1/2} = X^{1-2\varepsilon}$  exceptions. On the other hand, by Lemma 3.3 we have

$$\sum_{\substack{x \leq 8n \leq x+y \\ 2n \text{ is } \square\text{-free}}} |c(8n)| > \frac{y}{X^\varepsilon}$$

on a subset of cardinality at least  $X^{1-3\varepsilon/2}$ . Therefore the two subsets intersect on at least  $\gg X^{1-3\varepsilon/2}$  values of  $x$ , and therefore give rise to at least  $X^{1-15\varepsilon/2}$  sign changes in  $[X, 2X]$ .  $\square$

**3.2. The proof of Theorem 3.** The proof of Theorem 3 is completely elementary and does not depend on any of the other techniques developed here.

*Proof of Theorem 3.* Let  $g \in S_{k+1/2}^+$  be a Hecke cusp form and  $f$  denote the weight  $2k$  Hecke cusp form of level 1 that corresponds to  $g$ . For  $d$  a fundamental discriminant with  $(-1)^k d > 0$

$$(3.3) \quad c(n^2|d|) = c(|d|) \sum_{r|n} \frac{\mu(r)\chi_d(r)}{\sqrt{r}} \lambda_f\left(\frac{n}{r}\right),$$

where  $\lambda_f(n)$  denotes the  $n$ th Hecke eigenvalue of  $f$  (see [18, Eq'n (2)]). In particular if  $p$  is prime this becomes

$$c(|d|p^2) = c(|d|) \left( \lambda_f(p) - \frac{\chi_d(p)}{\sqrt{p}} \right).$$

Since there exists  $p$  which depends at most on  $k$  such that<sup>6</sup>  $\lambda_f(p) < -2/\sqrt{p}$  it follows that if  $c(|d|) \neq 0$  then  $c(|d|)$  and  $c(|d|p^2)$  have opposite signs. By Ono and Skinner's result there are  $\gg X/(p^2 \log(X/p^2)) \gg X/\log X$  fundamental discriminants  $|d| \leq X/p^2$  such that  $c(|d|) \neq 0$ . So by considering the signs of the Fourier coefficients  $c(n)$  at  $n = |d| \leq X/p^2$  with  $c(|d|) \neq 0$  and at  $m = |d|p^2 \leq X$  we arrive at the claimed result.  $\square$

#### 4. UPPER BOUNDS FOR MOLLIFIED MOMENTS

Let  $f$  be a level 1, Hecke cusp form of weight  $2k$ . The aim of this section is to compute an upper bound for mollified moments of  $L(\frac{1}{2}, f \otimes \chi_d)$ , conditionally under GRH. This gives an upper bound for the mean square of the mollified Fourier coefficients of  $g \in S_{k+1/2}^+$ .

The second moment  $L(\frac{1}{2}, f \otimes \chi_d)$  has been asymptotically computed assuming GRH by Soundararajan and Young [31]. However, a direct adaptation of their method cannot handle the introduction of a mollifier of length  $\gg |d|^\epsilon$ . For this reason we use a different approach, which uses the refinement of Soundararajan's [30] method for upper bounds on moments due to Harper [6] as well as the construction of an appropriate mollifier, which is based on an iterative construction and roughly has

<sup>6</sup>To see this, use the zero free regions for  $L(s, f)$ ,  $L(s, \text{sym}^2 f)$  and the Hecke relation  $\lambda_f(p)^2 = 1 + \lambda_f(p^2)$  to get that for  $y$  sufficiently large in terms of  $k$  that

$$\left| \sum_{\substack{p \leq y \\ |\lambda_f(p)| \geq 2p^{-1/2}}} \lambda_f(p) \right| < \sum_{\substack{p \leq y \\ |\lambda_f(p)| \geq 2p^{-1/2}}} |\lambda_f(p)|,$$

so the claim follows.

the structure of an Euler product. A similar mollifier and iterative approach was developed by Soundararajan and Radziwiłł [26].

The main result is

**Proposition 4.1.** *Assume GRH. Let  $l, \kappa > 0$  such that  $l \cdot \kappa \in \mathbb{N}$  and  $\kappa \cdot l \leq C$ . Also, let  $M(\cdot; \frac{1}{\kappa}) = M(\cdot; \frac{1}{\kappa}, x, C)$  be as in Section 4.2. Then*

$$\sum_{|d| \leq x}^b L(\frac{1}{2}, f \otimes \chi_d)^l M(d; \frac{1}{\kappa})^{l \cdot \kappa} \ll x,$$

where the sum is over fundamental discriminants and the implied constant depends on  $f, l, \kappa$ .

Note that by Waldspurger's formula (see (1.1)), we know  $L(\frac{1}{2}, f \otimes \chi_d) \geq 0$ , so that  $L(\frac{1}{2}, f \otimes \chi_d)^l$  is unambiguous.

Using the proposition above we can easily deduce Proposition 1.1.

*Proof of Proposition 1.1.* Using (1.1) it follows that

$$\sum_{\substack{1 \leq n \leq x \\ 2n \text{ is } \square\text{-free}}} |c(8n)|^4 M((-1)^k 8n; \frac{1}{2})^4 \ll \sum_{|d| \leq 8x}^b L(\frac{1}{2}, f \otimes \chi_d)^2 M(d; \frac{1}{2})^4 \ll x$$

where the sum is over all fundamental discriminants. □

**4.1. Preliminary results.** In this section we introduce our mollifier for the Fourier coefficients of  $g$  at fundamental discriminants. The shape of the mollifier is motivated by Harper's refinement [6] of Soundararajan's [30] bounds for moments.

**Notation.** Let  $\lambda(n) = (-1)^{\Omega(n)}$  denote the Liouville function (which should not be confused with  $\lambda_f(n)$ ). Denote by  $\nu(n)$  the multiplicative function with  $\nu(p^a) = \frac{1}{a!}$  and write  $\nu_j(n) = (\nu * \dots * \nu)(n)$  for the  $j$ -fold convolution of  $\nu$ . A useful observation is that

$$(4.1) \quad \nu_j(p^a) = \frac{j^a}{a!},$$

which may be proved by induction or otherwise. Also for an interval  $I$ ,  $m \in \mathbb{Z}$  and  $a(\cdot)$  a real-valued completely multiplicative function let

$$P_I(m; a(\cdot)) = \sum_{p \in I} \frac{a(p)}{\sqrt{p}} \left( \frac{m}{p} \right).$$

Also, given a statement  $S$  we let  $1_S$  equal one if  $S$  is true and zero otherwise.

**An  $L$ -function inequality.** We now prove an inequality in the spirit of Harper's work on sharp upper bounds for moments of  $L$ -functions, in the context of quadratic twists of  $L$ -functions attached to Hecke cusp forms. The upshot of the inequality is it essentially allows us to almost always bound the  $L$ -function by a very short Dirichlet polynomial.

Let us now introduce the following notation, which will be used throughout this section. Let  $l, \kappa > 0$  be such that  $l \cdot \kappa \in \mathbb{N}$  and suppose that  $1 \leq l \cdot \kappa \leq C$ . Also, let  $\eta_1, \eta_2 > 0$  be sufficiently small in terms of  $C$ . For  $j = 0, \dots, J$  let

$$(4.2) \quad \theta_j = \eta_1 \frac{e^j}{(\log \log x)^5} \quad \text{and} \quad \ell_j = 2 \lfloor \theta_j^{-3/4} \rfloor,$$

where  $J$  is chosen so that  $\eta_2 \leq \theta_J \leq e\eta_2$  (so  $J \asymp \log \log \log x$ ). For  $j = 1, \dots, J$  let  $I_j = (x^{\theta_{j-1}}, x^{\theta_j}]$  and set  $I_0 = (c, x^{\theta_0}]$ , where  $c \geq 2$  is sufficiently large in terms of  $\kappa$  and  $C$ . Note that the choice of parameters  $\theta_j, \ell_j$  depends on  $C$  and is independent of  $l, \kappa$  satisfying  $1 \leq l \cdot \kappa \leq C$ .

For  $j = 0, \dots, J$  and  $t > 0$  let

$$w(t; j) = \frac{1}{t^{1/(\theta_j \log x)}} \left( 1 - \frac{\log t}{\theta_j \log x} \right)$$

and define the completely multiplicative function  $a(\cdot; j)$  by

$$(4.3) \quad a(p; j) = \lambda_f(p) w(p; j),$$

and note  $|a(p; j)| \leq 2$ . The smooth weights  $w(\cdot; j)$  appear for technical reasons and their effect is mild.

Let  $\ell$  be an positive even integer. For  $t \in \mathbb{R}$ , let

$$E_\ell(t) = \sum_{s \leq \ell} \frac{t^s}{s!}.$$

Note  $E_\ell(t) \geq 1$  if  $t \geq 0$  and  $E_\ell(t) > 0$  for any  $t \in \mathbb{R}$  since  $\ell$  is even; the latter inequality may be seen using the Taylor expansion for  $e^t$ . Moreover, using the Taylor expansion it follows for  $t \leq \ell/e^2$  that

$$(4.4) \quad e^t \leq (1 + e^{-\ell}) E_\ell(t)$$

(see [26, Lemma 1]).

For each  $j = 0, \dots, J$  let

$$(4.5) \quad D_j(m; l) = \prod_{r=0}^j (1 + e^{-\ell_j}) E_{\ell_r}(l P_{I_r}(m; a(\cdot; j)))$$

and note  $D_j(m; l) > 0$ , since each term in the product is  $> 0$ . A useful observation is that for a positive integer  $s$

$$\begin{aligned}
P_I(m; a(\cdot))^s &= \sum_{p_1, \dots, p_s \in I} \frac{a(p_1 \cdots p_s)}{\sqrt{p_1 \cdots p_s}} \left( \frac{m}{p_1 \cdots p_s} \right) \\
(4.6) \quad &= \sum_{\substack{p|n \Rightarrow p \in I \\ \Omega(n)=s}} \frac{a(n)}{\sqrt{n}} \binom{m}{n} \sum_{p_1 \cdots p_s = n} 1 \\
&= s! \sum_{\substack{p|n \Rightarrow p \in I \\ \Omega(n)=s}} \frac{a(n)}{\sqrt{n}} \binom{m}{n} \nu(n).
\end{aligned}$$

Additionally, for any real number  $l \neq 0$

$$\begin{aligned}
(4.7) \quad E_\ell(lP_I(m; a(\cdot))) &= \sum_{s \leq \ell} \frac{l^s}{s!} P_I(m)^s \\
&= \sum_{\substack{p|n \Rightarrow p \in I \\ \Omega(n) \leq \ell}} \frac{l^{\Omega(n)} a(n) \nu(n)}{\sqrt{n}} \binom{m}{n}.
\end{aligned}$$

We are now ready to state our main inequality for  $L(\frac{1}{2}, f \otimes \chi_d)$ .

**Proposition 4.2.** *Assume GRH. Suppose that  $1 \leq |d| \leq x$  is a fundamental discriminant and  $l > 0$  is a real number. Also let*

$$A(d) = A(d; f) = \prod_{\substack{p|d \\ p > 3}} \left( 1 + \frac{\lambda_f(p^2) - 1}{p} \right).$$

Let  $\ell_j, \theta_j, I_j$  and  $a(\cdot, j)$  be as in (4.2) and (4.3) (resp.). Then there exists a sufficiently large real number  $C_1 = C_1(l) > 0$  such that for each  $|d| \leq x$  either:

$$(4.8) \quad |P_{I_0}(d; a(\cdot, j))| \geq \frac{\ell_0}{le^2}$$

for some  $0 \leq j \leq J$  or

$$(4.9) \quad (A(d) \log x)^{l/2} L(\frac{1}{2}, f \otimes \chi_d)^l \ll D_J(d; l) + \sum_{\substack{0 \leq j \leq J-1 \\ j+1 \leq u \leq J}} \left( \frac{1}{\theta_j} \right)^{C_1} \exp\left(\frac{3l}{\theta_j}\right) D_j(d; l) \left( \frac{e^2 l P_{I_{j+1}}(d; a(\cdot, u))}{\ell_{j+1}} \right)^{s_{j+1}},$$

for any even integers  $s_1, \dots, s_J$ , where the implied constant depends on  $f$  and  $l$ .

*Proof.* Applying Theorem 2.1 of Chandee [4] gives, for any  $X \geq 10$ , that

$$(4.10) \quad L\left(\frac{1}{2}, f \otimes \chi_d\right) \leq \exp\left(\sum_{p^n \leq X} \frac{\chi_d(p^n)(\alpha_p^n + \beta_p^n) \log X/p^n}{np^{n(\frac{1}{2} + \frac{1}{\log X})}} \frac{\log X}{\log X} + 3 \cdot \frac{\log |d|}{\log X} + O(1)\right),$$

where  $\alpha_p, \beta_p$  are the Satake parameters (note  $\alpha_p^n + \beta_p^n$  is real-valued see [21, Remark 2]). In the sum over primes, the contribution from the prime powers with  $n \geq 3$  is bounded so that it may be included in the  $O(1)$  term. Also, noting that  $\alpha_p^2 + \beta_p^2 = \lambda_f(p^2) - 1$  the squares of primes contribute

$$(4.11) \quad \frac{1}{2} \cdot \sum_{\substack{p \leq X \\ p \nmid d}} \frac{(\lambda_f(p^2) - 1) \log X/p^2}{p^{1 + \frac{2}{\log X}} \log X} = -\frac{1}{2} \log \log X - \sum_{\substack{p \leq X \\ p \nmid d}} \frac{(\lambda_f(p^2) - 1)}{2p} + O(1)$$

where the implied constant depends on  $f$ .

For  $r = 0, \dots, J$ , let  $\mathcal{E}_r$  be the set of fundamental discriminants  $d$  such that  $\max_{r \leq u \leq J} |P_{I_r}(d; a(\cdot; u))| < \frac{\ell_r}{le^2}$ . For each  $d$  we must have one of the following: (i)  $d \notin \mathcal{E}_0$ ; (ii)  $d \in \mathcal{E}_r$  for each  $0 \leq r \leq J$ ; (iii) there exists  $0 \leq j \leq J - 1$  such that  $d \in \mathcal{E}_r$  for  $0 \leq r \leq j$  and  $d \notin \mathcal{E}_{j+1}$ . Hence, we get for each fundamental discriminant with  $|d| \leq x$  either:

$$(4.12) \quad \max_{0 \leq u \leq J} |P_{I_0}(d; a(\cdot; u))| \geq \frac{\ell_0}{le^2};$$

$$(4.13) \quad \max_{j \leq u \leq J} |P_{I_j}(d; a(\cdot; u))| < \frac{\ell_j}{le^2} \quad \text{for each } j = 0, \dots, J;$$

or for some  $j = 0, \dots, J - 1$

$$(4.14) \quad \max_{r \leq u \leq J} |P_{I_r}(d; a(\cdot; u))| < \frac{\ell_r}{le^2} \quad \text{for each } r \leq j$$

and

$$(4.15) \quad \max_{j+1 \leq u \leq J} |P_{I_{j+1}}(d; a(\cdot; u))| \geq \frac{\ell_{j+1}}{le^2}.$$

Here (4.12), (4.13) correspond to possibilities (i) and (ii) above (resp.), while (4.14) and (4.15) corresponds to (iii).

If (4.12) holds for  $d$  then we are done. If (4.13) holds, we apply (4.10) with  $X = x^{\theta_j}$  and so the term  $3 \cdot \frac{\log |d|}{\log X}$  is  $\ll 1$ . Also, the contribution from (4.11) is

$$-\frac{1}{2} \log \log x - \sum_{\substack{p \leq X \\ p \nmid d}} \frac{(\lambda_f(p^2) - 1)}{2p} + O(1).$$



Hence, applying these estimates along with (4.4) we get that

$$\begin{aligned}
(A(d) \log x)^{l/2} L(\tfrac{1}{2}, f \otimes \chi_d)^l &\ll \exp \left( l \sum_{p \leq x^{\theta_j}} \frac{a(p; J)}{\sqrt{p}} \chi_d(p) \right) \\
(4.16) \qquad &\ll \prod_{j=0}^J \exp(lP_{I_j}(d; a(\cdot; J))) \\
&\leq \prod_{j=0}^J (1 + e^{-\ell_j}) E_{\ell_j}(lP_{I_j}(m; a(\cdot; J))) = D_J(d; l).
\end{aligned}$$

Finally, if (4.14) and (4.15) hold we apply (4.10) with  $X = x^{\theta_j}$  and for each  $0 \leq r \leq j$  we argue as above and bound the term  $\exp(lP_{I_r}(d; a(\cdot; j)))$  by  $(1 + e^{-\ell_r})E_{\ell_r}(lP_{I_r}(d; a(\cdot, j)))$ . In (4.10) we use the inequality  $3 \cdot \frac{\log |d|}{\log X} \leq \frac{3}{\theta_j}$  and note that the contribution from the squares of primes (4.11) is

$$-\frac{1}{2} \log \log x - \sum_{p|d} \frac{(\lambda_f(p^2) - 1)}{2p} + O\left(\log \frac{1}{\theta_j}\right).$$

So that applying these estimates, we get that for any non-negative, even integer  $s_{j+1}$

$$\begin{aligned}
(A(d) \log x)^{l/2} L(\tfrac{1}{2}, f \otimes \chi_d)^l \\
(4.17) \qquad &\ll \left(\frac{1}{\theta_j}\right)^{C_1} \exp(3l/\theta_j) D_j(d; l) \max_{j+1 \leq u \leq J} \left( \frac{e^{2lP_{I_{j+1}}}(d; a(\cdot, u))}{\ell_{j+1}} \right)^{s_{j+1}},
\end{aligned}$$

where we have included the extra factor

$$\max_{j+1 \leq u \leq J} \left( \frac{e^{2lP_{I_{j+1}}}(d; a(\cdot, u))}{\ell_{j+1}} \right)^{s_{j+1}} \geq 1.$$

Combining (4.16) and (4.17) and using the inequality  $\max(a, b) \leq a + b$  for  $a, b \geq 0$  gives (4.9).  $\square$

**4.2. The definition of the mollifier.** With Proposition 4.2 in mind we now choose our mollifier so that it will counteract the large values of  $D_J(d; l)$ . For  $j = 0, \dots, J$  let

$$M_j(m; \tfrac{1}{\kappa}) = \sum_{\substack{p|n \Rightarrow p \in I_j \\ \Omega(n) \leq \ell_j}} \frac{\kappa^{-\Omega(n)} a(n; J) \lambda(n)}{\sqrt{n}} \nu(n) \left( \frac{m}{n} \right),$$

where  $\ell_j, \theta_j$  and  $I_j$  are as in (4.2). Let

$$(4.18) \qquad M(m; \tfrac{1}{\kappa}) = (\log x)^{1/(2\kappa)} \prod_{0 \leq j \leq J} M_j(m; \tfrac{1}{\kappa}).$$

A useful observation is that  $M(m; \frac{1}{\kappa}) > 0$ , which can be seen by using (4.7) along with the fact that  $E_\ell(t) > 0$  for even  $\ell$  and  $t \in \mathbb{R}$ . Also, let  $\delta_0 = \sum_{j \leq J} \ell_j \theta_j$ . Observe that by construction  $\delta_0 \asymp 1$  is sufficiently small and the length of the Dirichlet polynomial  $M(m; \frac{1}{\kappa})$  is  $x^{\delta_0}$ . Also,

$$(4.19) \quad M(m; \frac{1}{\kappa})^{l \cdot \kappa} = (\log x)^{l/2} \sum_{n \leq x^{l \cdot \kappa \delta_0}} \frac{h(n) a(n; J) \lambda(n)}{\kappa^{\Omega(n)} \sqrt{n}} \left( \frac{m}{n} \right)$$

where

$$(4.20) \quad h(n) = \sum_{\substack{n_0 \cdots n_J = n \\ p|n_0 \Rightarrow p \in I_0, \dots, p|n_J \Rightarrow p \in I_J \\ \Omega(n_0) \leq l \cdot \kappa \ell_0, \dots, \Omega(n_J) \leq l \cdot \kappa \ell_J}} \nu_{l \cdot \kappa}(n_0; \ell_0) \cdots \nu_{l \cdot \kappa}(n_J; \ell_J).$$

and for  $n, r, \ell \in \mathbb{N}$

$$(4.21) \quad \nu_r(n; \ell) = \sum_{\substack{n_1 \cdots n_r = n \\ \Omega(n_1) \leq \ell, \dots, \Omega(n_r) \leq \ell}} \nu(n_1) \cdots \nu(n_r).$$

Note that  $\nu_r(n; \ell) \leq \nu_r(n)$  and if  $\Omega(n) \leq \ell$  then  $\nu_r(n; \ell) = \nu_r(n)$ . Also,

$$(4.22) \quad M(m; \frac{1}{\kappa})^{l \cdot \kappa} \ll x^{(l \cdot \kappa) \delta_0} (\log x)^{l/2},$$

where the implied constant depends on  $\kappa$  (note that  $\sum_{n \leq y} \kappa^{-\Omega(n)} \ll y(\log y)^{\frac{1}{\kappa}-1}$  where the implied constant depends on  $\kappa$ , for any  $y \geq 3$ .)

**4.3. The Proof of Proposition 4.1.** We first require the following fairly standard lemma, which follows from Poisson summation.

**Lemma 4.1.** *Let  $F$  be a Schwartz function such that its Fourier transform  $\widehat{F}$  has compact support in  $(-A, A)$ , for some fixed  $A > 0$ . Also, let  $I_0, \dots, I_J \subset [1, x]$  be disjoint intervals of the form  $I_0 = (c, x^{\theta_0}]$ ,  $I_j = (x^{\theta_{j-1}}, x^{\theta_j}]$  where  $\theta_0, \dots, \theta_J \in \mathbb{R}$  and  $c \geq 1$  is fixed. Suppose  $\ell_0, \dots, \ell_J \geq 1$  are real numbers such that  $\sum_{j=0}^J \ell_j \theta_j < 1/2$ . Then for any arithmetic function  $g$  we have*

$$(4.23) \quad \sum_{m \in \mathbb{Z}} \prod_{0 \leq j \leq J} \sum_{\substack{p|n \Rightarrow p \in I_j \\ \Omega(n) \leq \ell_j}} \frac{g(n) \left( \frac{m}{n} \right)}{\sqrt{n}} F\left( \frac{m}{x} \right) = \widehat{F}(0) x \prod_{0 \leq j \leq J} \sum_{\substack{p|n \Rightarrow p \in I_j \\ \Omega(n^2) \leq \ell_j}} \frac{g(n^2) \varphi(n^2)}{n^2}.$$

*Proof.* The LHS of (4.23) equals

$$(4.24) \quad \sum_{\substack{n_0, \dots, n_J \in \mathbb{N} \\ p|n_0 \Rightarrow p \in I_0, \dots, p|n_J \Rightarrow p \in I_J \\ \Omega(n_0) \leq \ell_0, \dots, \Omega(n_J) \leq \ell_J}} \frac{g(n_0) \cdots g(n_J)}{\sqrt{n_0 \cdots n_J}} \sum_m \left( \frac{m}{n_0 \cdots n_J} \right) F\left( \frac{m}{x} \right).$$

Using Poisson summation, see Remark 1 of [21], we get that for  $n \leq X^{1/2}$  if  $n = \square$  that

$$\sum_m \left(\frac{m}{n}\right) F\left(\frac{m}{x}\right) = \widehat{F}(0)x \frac{\varphi(n)}{n}$$

and for  $n \neq \square$  the above sum equals zero. By assumption  $n_0 \cdots n_J \leq x^{\ell_0 \theta_0} \cdots x^{\ell_J \theta_J} < x^{1/2}$ . Hence, the inner sum in (4.24) equals zero unless  $n_0 \cdots n_J = \square$  and since  $(n_j, n_i) = 1$  for  $i \neq j$  in this case  $n_0, \dots, n_J = \square$ . We conclude that the inner sum in (4.24) equals

$$\widehat{F}(0)x \frac{\varphi(n_0 \cdots n_J)}{n_0 \cdots n_J} = \widehat{F}(0)x \frac{\varphi(n_0) \cdots \varphi(n_J)}{n_0 \cdots n_J}$$

when  $n_0, \dots, n_J = \square$  and is zero otherwise, thereby giving the claim.  $\square$

To prove Proposition 4.1 we will use Proposition 4.2 and then apply Lemma 4.1. This leaves us with the problem of bounding the resulting sum. This task will be accomplished in the next two lemmas.

**Lemma 4.2.** *Let  $a(\cdot), b(\cdot)$  be real-valued, completely multiplicative functions with  $|a(p)|, |b(p)| \leq 2$ . Then for each  $j = 0, \dots, J$*

(4.25)

$$\begin{aligned} & \sum_{\substack{mn=\square \\ p|mn \Rightarrow p \in I_j \\ \Omega(m) \leq (l \cdot \kappa) \ell_j, \Omega(n) \leq \ell_j}} \frac{l^{\Omega(n)} a(m) b(n) \lambda(m)}{\kappa^{\Omega(m)} \sqrt{mn}} \nu_{l \cdot \kappa}(m; \ell_j) \nu(n) \frac{\varphi(mn)}{mn} \\ &= \left(1 + O\left(\frac{1_{j=0} \cdot (\log x)^{O(1)} + 1}{2^{\ell_j}}\right)\right) \prod_{p \in I_j} \left(1 + l^2 \frac{\alpha(p)}{2p} + O\left(\frac{1}{p^2}\right)\right), \end{aligned}$$

where

$$(4.26) \quad \alpha(p) = \alpha(p; a(\cdot), b(\cdot)) = (a(p) - b(p))^2,$$

and the implied constants depend on  $\kappa$  and  $l$ .

**Remark 1.** *Later we will take  $a(\cdot) = a(\cdot; J)$  and  $b = a(\cdot; j)$  with  $j < J$  (see (4.3)). Observe that*

$$(4.27) \quad \sum_{p \leq x^{\theta_j}} \frac{\alpha(p)}{p} = \sum_{p \leq x^{\theta_j}} \frac{\lambda_f(p)^2}{p} (w(p; J) - w(p; j))^2 \ll \sum_{p \leq x^{\theta_j}} \frac{1}{p} \cdot \frac{\log p}{\theta_j \log x} \ll 1.$$

*Proof.* In the sum over  $m, n$  on the LHS of (4.25) write  $mn = r^2$  and let

$$f_1(m) = \frac{a(m) \lambda(m) \nu_{l \cdot \kappa}(m)}{\kappa^{\Omega(m)}} \quad \text{and} \quad f_2(n) = l^{\Omega(n)} b(n) \nu(n).$$

We will proceed by estimating the sum on the LHS of (4.25) in terms of an Euler product. To this end, observe that if  $\max\{\Omega(n), \Omega(m)\} \geq \ell_j$  then  $2^{\Omega(m)+\Omega(n)}/2^{\ell_j} \geq 1$ ; so using this along with the remarks following (4.21) the sum on the LHS of (4.25) equals

$$(4.28) \quad \sum_{p|r \Rightarrow p \in I_j} \frac{1}{r} (f_1 * f_2)(r^2) \frac{\varphi(r^2)}{r^2} + O\left(\frac{1}{2^{\ell_j}} \sum_{p|r \Rightarrow p \in I_j} \frac{(|f_1| * |f_2|)(r^2)}{r}\right).$$

The error term in (4.28) is

$$(4.29) \quad \ll \frac{1}{2^{\ell_j}} \prod_{p \in I_j} \left(1 + O\left(\frac{1}{p}\right)\right) \asymp \begin{cases} \frac{1}{2^{\ell_j}} \left(\frac{\theta_j}{\theta_{j-1}}\right)^{O(1)} \asymp \frac{1}{2^{\ell_j}} & \text{if } j \neq 0, \\ \frac{(\log x)^{O(1)}}{2^{\ell_0}} & \text{if } j = 0. \end{cases}$$

Recalling (4.1), the main term in (4.28) is

$$= \prod_{p \in I_j} \left(1 - \frac{l^2 a(p)b(p)}{p} + \frac{l^2 a(p)^2}{2p} + \frac{l^2 b(p)^2}{2p} + O\left(\frac{1}{p^2}\right)\right).$$

□

**Lemma 4.3.** *Let  $a(\cdot), b(\cdot)$  be real-valued, completely multiplicative functions with  $a(p), b(p) \ll 1$ . Then for each  $j = 0, \dots, J$  and any even integer  $s \geq 4(l \cdot \kappa)\ell_j$*

$$(4.30) \quad \left| \sum_{\substack{mn = \square \\ p|mn \Rightarrow p \in I_j \\ \Omega(m) \leq (l \cdot \kappa)\ell_j, \Omega(n) = s}} \frac{a(m)b(n)\lambda(m)}{\kappa^{\Omega(m)}\sqrt{mn}} \nu_{\kappa \cdot l}(m; \ell_j) \nu(n) \frac{\varphi(mn)}{mn} \right| \\ \ll \frac{1_{j=0} \cdot (\log x)^{O(1)} + 1}{2^{s/2} \lfloor \frac{3}{8}s \rfloor!} \left(1 + \sum_{p \in I_j} \frac{b(p)^2}{p}\right)^{s/2},$$

where the implied constant depends on  $\kappa$  and  $l$ .

*Proof.* In the sum on the LHS of (4.30) write  $m = gm_1$ ,  $n = gn_1$  where  $g = (m, n)$ , so that  $mn = \square$  implies that  $m_1 = \square$  and  $n_1 = \square$ . Also recall that  $\nu_{\kappa \cdot l}(m; \ell_j) \leq \nu_{\kappa \cdot l}(m)$ . Hence, this sum is

$$(4.31) \quad \ll \sum_{\substack{p|gmn \Rightarrow p \in I_j \\ \Omega(g) \leq (l \cdot \kappa)\ell_j, \Omega(gn^2) = s}} \frac{|a(g)b(g)|a(m)^2b(n)^2}{\kappa^{2\Omega(m)+\Omega(g)}gmn} \nu_{l \cdot \kappa}(g) \nu_{l \cdot \kappa}(m^2) \nu(n^2),$$

where we have also used the estimates  $\nu_r(ab) \leq \nu_r(a)\nu_r(b)$  and  $\nu(a) \leq 1$ , for  $a, b, r \in \mathbb{N}$ , which follow from (4.1). The sum over  $m$  is

$$(4.32) \quad \ll \sum_{p|m \Rightarrow p \in I_j} \frac{a(m)^2 \nu_{l \cdot \kappa}(m^2)}{\kappa^{2\Omega(m)} m} \ll \prod_{p \in I_j} \left( 1 + \frac{(l \cdot \kappa)^2}{2} \cdot \frac{a(p)^2}{\kappa^2 p} \right) \ll \begin{cases} \left( \frac{\theta_j}{\theta_{j-1}} \right)^{O(1)} \asymp 1 & \text{if } j \neq 0, \\ (\log x)^{O(1)} & \text{if } j = 0. \end{cases}$$

Hence, using (4.32) along with the inequality  $\nu(n^2) \leq \nu(n)2^{-\Omega(n)}$  it follows that (4.31) is

$$(4.33) \quad \begin{aligned} & \ll (1_{j=0}(\log x)^{O(1)} + 1) \sum_{\substack{p|g \Rightarrow p \in I_j \\ \Omega(g) \leq (l \cdot \kappa)\ell_j}} \frac{|a(g)b(g)|}{\kappa^{\Omega(g)} g} \nu_{l \cdot \kappa}(g) \sum_{\substack{p|n \Rightarrow p \in I_j \\ \Omega(n) = (s - \Omega(g))/2}} \frac{b(n)^2}{2^{\Omega(n)} n} \nu(n) \\ & = (1_{j=0}(\log x)^{O(1)} + 1) \sum_{\substack{p|g \Rightarrow p \in I_j \\ \Omega(g) \leq (l \cdot \kappa)\ell_j \\ 2|\Omega(g)}} \frac{|a(g)b(g)|}{\kappa^{\Omega(g)} g} \nu_{l \cdot \kappa}(g) \left[ \frac{1}{((s - \Omega(g))/2)!} \left( \frac{1}{2} \sum_{p \in I_j} \frac{b(p)^2}{p} \right)^{(s - \Omega(g))/2} \right], \end{aligned}$$

by (4.6). Using the assumption that  $s \geq 4(l \cdot \kappa)\ell_j$  gives  $s - \Omega(g) \geq \frac{3}{4}s$  for  $g$  with  $\Omega(g) \leq (l \cdot \kappa)\ell_j$ . This allows us to bound the bracketed term on the RHS of (4.33) by

$$2^{\Omega(g)/2} \cdot \frac{1}{2^{s/2} \lfloor \frac{3}{8}s \rfloor!} \left( 1 + \sum_{p \in I_j} \frac{b(p)^2}{p} \right)^{s/2}.$$

Applying this estimate in (4.33) it follows that the RHS of (4.33) is bounded by

$$\begin{aligned} & \ll \frac{(1_{j=0}(\log x)^{O(1)} + 1)}{2^{s/2} \lfloor \frac{3}{8}s \rfloor!} \left( 1 + \sum_{p \in I_j} \frac{b(p)^2}{p} \right)^{s/2} \sum_{p|g \Rightarrow p \in I_j} \frac{(4l)^{\Omega(g)}}{g} \\ & \ll \frac{(1_{j=0}(\log x)^{O(1)} + 1)}{2^{s/2} \lfloor \frac{3}{8}s \rfloor!} \left( 1 + \sum_{p \in I_j} \frac{b(p)^2}{p} \right)^{s/2}. \end{aligned}$$

□

In the next lemma we estimate averages of our mollifier  $M$  as defined in Section 4.2 against the terms which appear in Proposition 4.2.

**Lemma 4.4.** *For  $j = 0, \dots, J$  let  $s_j$  be an even integer with  $4(l \cdot \kappa)\ell_j \leq s_j \leq \frac{2}{5\theta_j}$ . Then we have the following estimates:*

$$(4.34) \quad \sum_{|d| \leq x}^b D_J(d; l) M(d; \frac{1}{\kappa})^{l \cdot \kappa} \ll x(\log x)^{l/2};$$

$$(4.35) \quad \sum_{|d| \leq X}^b M(d; \frac{1}{\kappa})^{l \cdot \kappa} P_{I_0}(d; a(\cdot; u))^{s_0} \ll x(\log x)^{O(1)} \frac{s_0!}{2^{s_0/2} \lfloor \frac{3}{8}s_0 \rfloor!} \left( \sum_{p \in I_0} \frac{a(p; u)^2}{p} \right)^{s_0/2};$$

$$(4.36) \quad \sum_{|d| \leq X}^b D_j(d; l) P_{I_{j+1}}(d; a(\cdot; u))^{s_{j+1}} M(d; \frac{1}{\kappa})^{l \cdot \kappa} \\ \ll x(\log x)^{l/2} e^{2l^2(J-j)} \frac{s_{j+1}!}{2^{s_{j+1}/2} \lfloor \frac{3}{8}s_{j+1} \rfloor!} \left( \sum_{p \in I_{j+1}} \frac{a(p; u)^2}{p} \right)^{s_{j+1}/2}$$

for each  $j = 0, \dots, J-1$ . The implied constants depend at most on  $f, \kappa, l$  (and not on  $j, u$ ).

*Proof.* We will first prove (4.36), which is the most complicated of the bounds, and at the end of the proof we will indicate how to modify the argument to establish (4.34) and (4.35). Recall that  $D_j$  and  $M$  are positive (see the remarks after (4.5) and (4.18)). So for any Schwartz function  $F$  which majorizes  $1_{[-1,1]}$ , such that  $\widehat{F}$  has compact support in  $(-A, A)$  the LHS of (4.36) is

$$\leq \sum_{m \in \mathbb{Z}} D_j(m; l) P_{I_{j+1}}(m; a(\cdot; u))^{s_{j+1}} M(m; \frac{1}{\kappa})^{l \cdot \kappa} F\left(\frac{m}{x}\right).$$

The Dirichlet Polynomial above is supported on integers  $n \leq x^\theta$  where  $\theta = l \cdot \kappa \sum_{0 \leq r \leq j} \theta_r \ell_r + \sum_{0 \leq r \leq j} \theta_r \ell_r + s_{j+1} \theta_{j+1} < 1/2$ . Hence, using (4.5), (4.6), (4.7), (4.19), (4.20) and applying Lemma 4.1 the above equation is bounded by

$$(4.37) \quad \leq x(\log x)^{l/2} \widehat{F}(0) \prod_{0 \leq r \leq J} \sum_{\substack{mn = \square \\ p|mn \Rightarrow p \in I_r \\ \Omega(m) \leq (l \cdot \kappa) \ell_r}} \frac{a(m; J) g(n; u, r) \lambda(m)}{\kappa^{\Omega(m)} \sqrt{mn}} \nu_{\kappa \cdot l}(m; \ell_r) \nu(n) \frac{\varphi(mn)}{mn}$$

where

$$g(n; u, r) = \begin{cases} 1_{\Omega(n) \leq \ell_r} \cdot l^{\Omega(n)} \cdot a(n; j) (1 + e^{-\ell_r}) & \text{if } 0 \leq r \leq j, \\ 1_{\Omega(n) = s} \cdot s! \cdot a(n; u) & \text{if } r = j + 1, \\ 1_{n=1} & \text{if } j + 1 < r \leq J. \end{cases}$$

Let  $\alpha(\cdot)$  be as in (4.26) and take  $a(\cdot) = a(\cdot; J)$ ,  $b(\cdot) = a(\cdot; j)$ . Applying Lemma 4.2, the contribution to (4.37) from the product over  $0 \leq r \leq j$  is bounded in absolute value by

$$(4.38) \quad \ll \prod_{0 \leq r \leq j} \left( 1 + O \left( \frac{1_{r=0} \cdot (\log x)^{O(1)} + 1}{2^{\ell_r}} \right) \right) \prod_{p \in I_j} \left( 1 + l^2 \frac{\alpha(p)}{2p} + O \left( \frac{1}{p^2} \right) \right) \ll 1,$$

where in the last step we applied (4.27). Applying Lemma 4.3 with  $a(\cdot) = a(\cdot; J)$  and  $b(\cdot) = a(\cdot; u)$  the term in (4.37) with  $r = j + 1$  contributes

$$(4.39) \quad \ll \frac{s_{j+1}!}{2^{s_{j+1}/2} \lfloor \frac{3}{8} s_{j+1} \rfloor!} \left( \sum_{p \in I_{j+1}} \frac{a(p; u)^2}{p} \right)^{s_{j+1}/2}.$$

It remains to handle the factors in the product in (4.37) with  $j + 1 < r \leq J$ . For such factors,  $m$  is a square so these terms are bounded by

$$(4.40) \quad \begin{aligned} &\ll \prod_{j+1 < r \leq J} \sum_{p|m \Rightarrow p \in I_r} \frac{a(m; J)^2}{\kappa^{2\Omega(m)} m} \nu_{l \cdot \kappa}(m^2) \\ &\ll \prod_{x^{\theta_j} < p \leq x^{\theta_J}} \left( 1 + \frac{(l \cdot \kappa)^2}{2} \cdot \frac{\lambda_f(p)^2}{\kappa^2 p} \right) \ll \left( \frac{\theta_J}{\theta_j} \right)^{2l^2} = e^{2l^2(J-j)}, \end{aligned}$$

where we used (4.1) and the bound  $|\lambda_f(p)| \leq 2$ , in the last steps.

Using (4.38), (4.39) and (4.40) in (4.37) completes the proof of (4.36). To establish (4.34) we repeat the same argument, the only differences are that in (4.38) the product is over all  $0 \leq r \leq J$  and the terms estimated in (4.39) and (4.40) do not appear. To establish (4.35) note that the term estimated in (4.38) does not appear and in the bound (4.40) the only difference is that  $j = 0$ , so we bound this term by  $O((\log x)^{O(1)})$ . Finally in place of (4.39) we get by using Lemma 4.3 the bound

$$(\log x)^{O(1)} \frac{s_0!}{2^{s_0/2} \lfloor \frac{3}{8} s_0 \rfloor!} \left( \sum_{p \in I_0} \frac{a(p; u)^2}{p} \right)^{s_0/2}.$$

From these estimates (4.35) follows.  $\square$

*Proof of Proposition 4.1.* Let  $A(d)$  be as in the statement of Proposition 4.2. In particular, using  $|\lambda_f(p^2)| \leq d(p^2) = 3$  and  $\lambda_f(p^2) = \lambda_f(p)^2 - 1 \geq -1$  it follows that

$$\left( \frac{\varphi(d)}{d} \right)^2 \ll A(d) \ll \left( \frac{d}{\varphi(d)} \right)^2.$$

Using this observation we note that it suffices to prove that

$$(4.41) \quad \sum_{|d| \leq x}^b (A(d)^{1/2} L(\tfrac{1}{2}, f \otimes \chi_d))^l M(d; \tfrac{1}{\kappa})^{l \cdot \kappa} \ll x,$$

for  $l > 0$  (note the definition of  $M$  is independent of  $l$ , for  $1 \leq l \cdot \kappa \leq C$ ), since by Cauchy-Schwarz (4.41) implies

$$\begin{aligned} & \sum_{|d| \leq x}^b L(\tfrac{1}{2}, f \otimes \chi_d)^l M(d; \tfrac{1}{\kappa})^{l \cdot \kappa} \\ & \leq \left( \sum_{|d| \leq x}^b \frac{1}{A(d)^l} \right)^{1/2} \left( \sum_{|d| \leq x}^b (A(d)^{1/2} L(\tfrac{1}{2}, f \otimes \chi_d))^{2l} M(d; \tfrac{1}{\kappa})^{2l \cdot \kappa} \right)^{1/2} \ll x. \end{aligned}$$

We will now establish (4.41). For  $j = 0, \dots, J$ , let  $s_j = 2 \lfloor \frac{1}{5\theta_j} \rfloor$ . We first divide the sum over  $|d| \leq x$  into two sums depending on whether

$$(4.42) \quad |P_{I_0}(d; a(\cdot; u))| \geq \ell_0 / (le^2)$$

for some  $0 \leq u \leq J$ . To bound the contribution of the terms to the LHS of (4.41) with  $|d| \leq x$  for which (4.42) holds, we use the bound  $A(d) \ll (\log \log x)^2$ , Chebyshev's inequality and then apply Cauchy-Schwarz to see that

$$\begin{aligned} & \sum_{\substack{|d| \leq x \\ |P_{I_0}(d; a(\cdot; u))| \geq \ell_0 / (le^2)}}^b (A(d)^{1/2} L(\tfrac{1}{2}, f \otimes \chi_d))^l M(d; \tfrac{1}{\kappa})^{l \cdot \kappa} \\ & \ll (\log \log x)^l \sum_{|d| \leq x}^b L(\tfrac{1}{2}, f \otimes \chi_d)^l M(d; \tfrac{1}{\kappa})^{l \cdot \kappa} \left( \frac{le^2 P_{I_0}(d; a(\cdot; u))}{\ell_0} \right)^{s_0} \\ & \leq (\log \log x)^l \left( \sum_{|d| \leq x}^b L(\tfrac{1}{2}, f \otimes \chi_d)^{2l} \right)^{1/2} \left( \frac{(le^2)^{2s_0}}{\ell_0^{2s_0}} \sum_{|d| \leq x}^b M(d; \tfrac{1}{\kappa})^{2l \cdot \kappa} P_{I_0}(d; a(\cdot; u))^{2s_0} \right)^{1/2}. \end{aligned}$$

On the RHS, the first sum is  $\ll x(\log x)^{O(1)}$  by Soundararajan's [30] method for upper bounds for moments (see the examples in [30, Section 4]). Using Lemma 4.4, (4.35), applying Stirling's formula and recalling the definition of  $\ell_0, \theta_0$  (see (4.2)) the second sum on the RHS is

$$\begin{aligned} & \ll x(\log x)^{O(1)} \left( \frac{100l^2 s_0^{5/4}}{\ell_0^2} \right)^{s_0} \left( \sum_{p \leq x} \frac{\lambda_f(p)^2}{p} \right)^{s_0} \\ & \ll x(\log x)^{O(1)} \left( 100l^2 \theta_0^{1/4} \log \log x \right)^{s_0} \ll x(\log x)^{O(1)} \exp(-(\log \log x)^5) \ll \frac{x}{(\log x)^A} \end{aligned}$$



for any  $A \geq 1$ . Hence, the contribution to (4.41) from  $|d| \leq x$  satisfying (4.42) for some  $u \leq J$  is  $\ll J \frac{x}{(\log x)^A} \ll \frac{x}{(\log x)^A} \log \log \log x$  for any  $A \geq 1$ .

For the remaining fundamental discriminants  $|d| \leq x$  we apply (4.9) to see that their contribution to (4.41) is bounded by

$$(4.43) \quad \ll \frac{1}{(\log x)^{l/2}} \left( \sum_{|d| \leq x}^b D_J(d; l) M(d; \frac{1}{\kappa})^{l \cdot \kappa} + \sum_{\substack{0 \leq j \leq J-1 \\ j+1 \leq u \leq J}} \left( \frac{1}{\theta_j} \right)^{C_1} \exp\left(\frac{3l}{\theta_j}\right) \left( \frac{e^2 l}{\ell_{j+1}} \right)^{s_{j+1}} \sum_{|d| \leq x}^b P_{I_{j+1}}(d; a(\cdot, u))^{s_{j+1}} D_j(d; l) M(d; \frac{1}{\kappa})^{l \cdot \kappa} \right).$$

To complete the proof it suffices to show that the expression above is  $\ll x$ . Applying (4.34) the first term in (4.43) is

$$(4.44) \quad \frac{1}{(\log x)^{l/2}} \sum_{|d| \leq x}^b D_J(d; l) M(d; \frac{1}{\kappa})^{l \cdot \kappa} \ll x.$$

Using (4.36) the second term in (4.43) is bounded by

$$\ll x \sum_{\substack{0 \leq j \leq J-1 \\ j+1 \leq u \leq J}} \left( \frac{1}{\theta_j} \right)^{C_1} \exp\left(\frac{3l}{\theta_j}\right) \left( \frac{e^2 l}{\ell_{j+1}} \right)^{s_{j+1}} \cdot e^{2l^2(J-j)} \frac{(s_{j+1})!}{2^{s_{j+1}/2} \lfloor \frac{3}{8} s_{j+1} \rfloor!} \left( \sum_{p \in I_{j+1}} \frac{a(p; u)^2}{p} \right)^{s_{j+1}/2}.$$

Applying Stirling's formula and estimating the inner sum over primes trivially as  $\leq 5 \log\left(\frac{\log x^{\theta_{j+1}}}{\log x^{\theta_j}}\right) = 5$ , we see that the above is

$$(4.45) \quad \ll x \sum_{0 \leq j \leq J-1} (J-j) \left( \frac{1}{\theta_j} \right)^{C_1} e^{2l^2(J-j)} \exp\left(\frac{3l}{\theta_j}\right) \left( \frac{e^2 l}{\ell_{j+1}} \cdot \left(\frac{8}{3}\right)^{3/8} \cdot \frac{s_{j+1}^{5/8}}{\sqrt{2} e^{5/8}} \cdot 5 \right)^{s_{j+1}}.$$

By construction  $\frac{s_{j+1}^{5/8}}{\ell_{j+1}} \ll \theta_{j+1}^{1/8}$ . Hence, there exists  $c > 0$  (which may depend on  $l, \kappa$ ) such that (4.45) is

$$(4.46) \quad \ll x \sum_{0 \leq j \leq J-1} (J-j) e^{2l^2(J-j)} \exp\left(-\frac{c}{\theta_j} \log \frac{1}{\theta_j} + O\left(\frac{1}{\theta_j}\right)\right) \ll x \sum_{1 \leq j \leq J} j e^{2l^2 j} \exp\left(-\frac{c}{2} \cdot \frac{j e^j}{\theta_j}\right) \ll x,$$

where we used that  $\theta_J/\theta_j = e^{J-j}$  in the last step. Combining (4.46) with (4.44), gives that (4.43) is  $\ll x$ , which completes the proof.  $\square$

## 5. THE PROOF OF PROPOSITION 1.2

The main input into the proof of Proposition 1.2 is a twisted first moment of  $L(\frac{1}{2}, f \otimes \chi_d)$ . Similar moment estimates were established in [29], [31] and [26] and our proof closely follows the methods developed in those papers. We will include a proof of the following result for completeness.

**Proposition 5.1.** *Let  $\phi$  be a Schwartz function with compact support in the positive reals. Also, let  $u = u_1 u_2^2$ , be odd where  $u_1$  is square-free. Then*

$$(5.1) \quad \sum_{\substack{m \\ 2m \text{ is } \square\text{-free}}} L(\tfrac{1}{2}, f \otimes \chi_{8m}) \chi_{8m}(u) \phi\left(\frac{(-1)^k 8m}{x}\right) = C \int_0^\infty \phi(\xi) d\xi \cdot \frac{x}{u_1^{1/2}} \cdot \vartheta(u) + O(x^{7/8+\varepsilon} u^{3/8+\varepsilon})$$

where  $C > 0$  depends only on  $f$  and  $\vartheta(\cdot)$  is a multiplicative function with  $\vartheta(p^{2j+1}) = \lambda_f(p) + O(1/p)$  and  $\vartheta(p^{2j}) = 1 + O(1/p)$ .

The constant  $C$  is explicitly given in the proof below, see (5.30). Before proving Proposition 5.1, we will use the result to deduce Proposition 1.2.

**5.1. The Proof of Proposition 1.2.** We will only prove the lower bound

$$\sum_{\substack{n \leq x \\ 2n \text{ is } \square\text{-free}}} |c(8n)|^2 M((-1)^k 8n; \tfrac{1}{2})^2 \gg x$$

since the proof of the upper bound is similar. Using (1.1) it follows that for a Schwartz function  $\phi(\cdot)$  which minorizes  $1_{[1/2, 1]}(\cdot)$  with  $\widehat{\phi}(0) > 0$  the sum above is

$$(5.2) \quad \geq \frac{(k-1)!}{\pi^k} \cdot \frac{\langle g, g \rangle}{\langle f, f \rangle} \sum_{\substack{m \\ 2m \text{ is } \square\text{-free}}} L(\tfrac{1}{2}, f \otimes \chi_{8m}) M(8m; \tfrac{1}{2})^2 \phi\left(\frac{(-1)^k 8m}{x}\right).$$

To proceed we now expand  $M(8m; \frac{1}{2})^2$  and see that it equals

$$(5.3) \quad M(8m; \tfrac{1}{2})^2 = (\log x)^{1/2} \sum_{n \leq x^{2\delta_0}} \frac{h(n) a(n; J) \lambda(n)}{2^{\Omega(n)} \sqrt{n}} \left(\frac{8m}{n}\right)$$

where  $h(n)$  is as in (4.20) (note that here  $l \cdot \kappa = 2$ ). Applying Proposition 5.1 gives that the RHS of (5.2) equals

$$(5.4) \quad \begin{aligned} & C' x(\log x)^{1/2} \sum_{\substack{n_1 n_2^2 \leq x^{2\delta_0} \\ n_1 \text{ is } \square\text{-free}}} \frac{h(n_1 n_2^2) a(n_1 n_2^2; J) \lambda(n_1) \vartheta(n_1 n_2^2)}{2^{\Omega(n_1) + 2\Omega(n_2)} n_2 n_1} + O(x^{9/10}) \\ &= C' x(\log x)^{1/2} \prod_{0 \leq j \leq J} \sum_{\substack{p | mn^2 \Rightarrow p \in I_j \\ \Omega(mn^2) \leq 2\ell_j}} \frac{a(mn^2; J) \lambda(m) \vartheta(mn^2) \mu(m)^2}{2^{\Omega(m) + 2\Omega(n)} mn} \nu_2(mn^2; \ell_j) + O(x^{9/10}), \end{aligned}$$

where  $C' = C'(f, g, k, \phi) > 0$  and in the last step we used that  $\vartheta$  is a multiplicative function.

We will now estimate the inner sum on the RHS of (5.4). First note that  $\vartheta(mn^2) \ll d(m) \left(\frac{mn^2}{\varphi(m)\varphi(n^2)}\right)^{O(1)}$  and recall the remarks after (4.21), so we get that the sum equals

$$(5.5) \quad \begin{aligned} & \sum_{p | mn^2 \Rightarrow p \in I_j} \frac{a(mn^2; J) \lambda(m) \vartheta(mn^2) \mu(m)^2}{2^{\Omega(m) + 2\Omega(n)} mn} \nu_2(mn^2) \\ &+ O\left(\frac{1}{4^{\ell_j}} \sum_{p | nm \Rightarrow p \in I_j} \frac{a(m; J) a(n; J)^2 d(m) \nu_2(m) \nu_2(n^2)}{mn} \cdot \left(\frac{mn^2}{\varphi(m)\varphi(n^2)}\right)^{O(1)}\right). \end{aligned}$$

The error term is

$$(5.6) \quad \ll \frac{1}{4^{\ell_j}} \prod_{p \in I_j} \left(1 + O\left(\frac{1}{p}\right)\right) \ll \frac{1_{j=0} \cdot (\log x)^{O(1)} + 1}{4^{\ell_j}}.$$

To estimate the main term in (5.5) consider

$$s(p; a) = \sum_{t \geq 0} \frac{a(p; J)^{2t}}{4^t p^t} \vartheta(p^{2t+a}) \nu_2(p^{2t+a}).$$

Recall  $\vartheta(p) = \lambda_f(p) + O(1/p)$  and  $\vartheta(p^2) = 1 + O(1/p)$ . Evaluating the sum over  $n$  in the main term of (5.5) we see that it equals

$$(5.7) \quad \begin{aligned} & \left(\prod_{p \in I_j} s(p; 0)\right) \sum_{p | m \Rightarrow p \in I_j} \frac{a(m; J) \mu(m)}{2^{\Omega(m)} m} \prod_{p^a || m} \frac{s(p; a)}{s(p; 0)} \\ &= \prod_{p \in I_j} \left(1 + \frac{a(p; J)^2}{2p} - \frac{\lambda_f(p) a(p; J)}{p} + O\left(\frac{1}{p^2}\right)\right). \end{aligned}$$

Using (5.6), (5.7) along with the estimates  $a(p; J) = \lambda_f(p) + O\left(\frac{\log p}{\theta_J \log x}\right)$  and

$$\prod_{p \in I_j} \left(1 + \frac{a(p; J)^2}{2p} - \frac{\lambda_f(p)a(p; J)}{p} + O\left(\frac{1}{p^2}\right)\right)^{-1} \ll 1_{j=0} \cdot (\log x)^{O(1)} + 1$$

we get that the RHS of (5.4) equals

$$(5.8) \quad C' x (\log x)^{1/2} \prod_{c < p \leq x^{\theta_J}} \left(1 - \frac{\lambda_f(p)^2}{2p} + O\left(\frac{1}{\theta_J \log x} \cdot \frac{\log p}{p} + \frac{1}{p^2}\right)\right) \\ \times \prod_{0 \leq j \leq J} \left(1 + O\left(\frac{1_{j=0} \cdot (\log x)^{O(1)} + 1}{4^{\ell_j}}\right)\right).$$

To estimate the second product above note that

$$(5.9) \quad \prod_{0 \leq j \leq J} \left(1 + O\left(\frac{1_{j=0} \cdot (\log x)^{O(1)} + 1}{4^{\ell_j}}\right)\right) = 1 + O\left(\frac{1}{4^{\ell_J}}\right) = 1 + O(\eta_2) \geq \frac{1}{2},$$

since  $\eta_2$  is sufficiently small. Also, there exists some constant  $C'' > 0$  such that the Euler product over  $c < p \leq x^{\theta_J}$  in (5.8) is

$$\geq \prod_{c < p \leq x^{\theta_J}} \left(1 - \frac{\lambda_f(p)^2}{2p} - C'' \left(\frac{1}{\theta_J \log x} \cdot \frac{\log p}{p} + \frac{1}{p^2}\right)\right) \gg (\log x)^{-1/2},$$

since  $c$  is sufficiently large (so each term in the Euler product is positive). Hence, using this and (5.9) we get that (5.8) is  $\gg x (\log x)^{1/2} \cdot (\log x)^{-1/2} = x$ , which completes the proof.

**5.2. The proof of Proposition 5.1.** Let  $f$  be a weight  $2k$ , level 1, Hecke cusp form. For a fundamental discriminant  $d$ , let

$$\Lambda(s, f \otimes \chi_d) = \left(\frac{|d|}{2\pi}\right)^s \Gamma\left(s + \frac{2k-1}{2}\right) L(s, f \otimes \chi_d).$$

The functional equation for  $\Lambda(s, f \otimes \chi_d)$  is given by

$$\Lambda(s, f \otimes \chi_d) = (-1)^k \operatorname{sgn}(d) \Lambda(1-s, f \otimes \chi_d).$$

Note that the central value vanishes when  $(-1)^k d < 0$ .

For  $c > 0$  and  $x > 0$  let

$$W_z(x) = \frac{1}{2\pi i} \int_{(c)} \frac{\Gamma(z+s+k-\frac{1}{2})}{\Gamma(z+k-\frac{1}{2})} (2\pi x)^{-s} \frac{ds}{s}.$$

Our starting point in the proof of Proposition 5.1 is the following approximate functional equation for  $L(s, f \otimes \chi_d)$ . Also, define  $W = W_{1/2}$ .

**Lemma 5.1.** *Let  $f$  be a level 1, Hecke cusp form with weight  $2k$ . For  $s \in \mathbb{C}$  with  $0 \leq \operatorname{Re}(s) \leq 1$  and  $d$  a fundamental discriminant*

$$\begin{aligned} L(s, f \otimes \chi_d) &= \sum_{n \geq 1} \frac{\lambda_f(n) \chi_d(n)}{n^s} W_s \left( \frac{n}{|d|} \right) \\ &\quad + (-1)^k \operatorname{sgn}(d) \left( \frac{|d|}{2\pi} \right)^{1-2s} \frac{\Gamma(1-s)}{\Gamma(s)} \sum_{n \geq 1} \frac{\lambda_f(n) \chi_d(n)}{n^{1-s}} W_{1-s} \left( \frac{n}{|d|} \right). \end{aligned}$$

Moreover, the function  $W = W_{1/2}$  satisfies  $\xi^j W^{(j)}(\xi) \ll \xi^{-A}$  for any  $A \geq 1$  and  $W(\xi) = 1 + O(\xi^{k-\varepsilon})$  as  $\xi \rightarrow 0$ , for any  $\varepsilon > 0$ .

*Proof.* See Lemma 5 of [26] and Lemma 2.1 of [31]. □

Since we will sum over fundamental discriminants we introduce a new parameter  $Y$  with  $1 \leq Y \leq x$  to be chosen later. Also, write  $F(\xi; x, y) = \phi\left(\frac{\xi}{x}\right) W\left(\frac{y}{\xi}\right)$ . Applying Lemma 5.1 the LHS of (5.1) equals

$$\begin{aligned} (5.10) \quad & 2 \sum_{(m,2)=1} \sum_{a^2|m} \mu(a) \sum_{n \geq 1} \frac{\lambda_f(n)}{\sqrt{n}} \left( \frac{8m}{nu} \right) F((-1)^k 8m; x, n) \\ &= 2 \left( \sum_{\substack{a \leq Y \\ (a,2u)=1}} + \sum_{\substack{a > Y \\ (a,2u)=1}} \right) \mu(a) \sum_{\substack{(m,2)=1 \\ a^2|m}} \sum_{n \geq 1} \frac{\lambda_f(n)}{\sqrt{n}} \left( \frac{8m}{nu} \right) F((-1)^k 8m; x, n). \end{aligned}$$

**The terms with  $a > Y$ .** Write  $m = r^2 e$  where  $e$  is square-free, and note since in the sum in (5.10)  $a^2|m$  it follows that  $a|r$ . So the second sum in (5.10) equals

$$\begin{aligned} (5.11) \quad & 2 \sum_{(r,2u)=1} \sum_{\substack{a > Y \\ a|r}} \mu(a) \sum_{\substack{e \text{ is } \square\text{-free} \\ (e,2)=1}} \sum_{(n,r)=1} \frac{\lambda_f(n)}{\sqrt{n}} \left( \frac{8e}{nu} \right) F((-1)^k 8r^2 e; x, n) \\ & \ll \sum_{|r| \leq x^{1/2+\varepsilon}} \sum_{\substack{a > Y \\ a|r, (a,2)=1}} \sum_{\substack{1 \leq (-1)^k e \leq x^{1+\varepsilon}/r^2 \\ e \text{ is } \square\text{-free} \\ (e,2)=1}} \left| \sum_{(n,r)=1} \frac{\lambda_f(n)}{\sqrt{n}} \left( \frac{8e}{n} \right) W\left(\frac{n}{8r^2|e|}\right) \right| + \frac{1}{x}, \end{aligned}$$

where in the second line we have used the rapid decay of  $W$ . Using the definition of  $W$  and for  $\operatorname{Re}(s) > 1$ , writing  $L(s, f \otimes \chi_{8e}) = \prod_p L_p(s)$  where  $L_p(s) =$

$\left(1 - \frac{\lambda_f(p)(\frac{8e}{p})}{p^s} + \frac{(\frac{8e}{p})^2}{p^{2s}}\right)^{-1}$  we get for  $c > 1/2$  that

$$(5.12) \quad \sum_{(n,r)=1} \frac{\lambda_f(n)}{\sqrt{n}} \left(\frac{8e}{n}\right) W\left(\frac{n}{8r^2|e|}\right) = \frac{1}{2\pi i} \int_{(c)} \frac{\Gamma(s+k)}{\Gamma(k)} \left(\frac{8r^2e}{2\pi}\right)^s \frac{L(s+\frac{1}{2}, f \otimes \chi_{8e})}{\prod_{p|r} L_p(s+\frac{1}{2})} \frac{ds}{s}.$$

The integrand is holomorphic for  $1/\log x \leq \operatorname{Re}(s) \leq 2$  and in this region bounded by (note  $r^2|e| \leq x^2$ )

$$\begin{aligned} &\ll \prod_{p|r} \left(1 + \frac{O(1)}{\sqrt{p}}\right) (r^2|e|)^{\operatorname{Re}(s)} |\Gamma(s+k)| |L(s+\frac{1}{2}, f \otimes \chi_{8e})| \cdot \frac{1}{|s|} \\ &\ll x^\varepsilon (r^2|e|)^{\operatorname{Re}(s)} \exp(-|s|) |L(s+\frac{1}{2}, f \otimes \chi_{8e})|. \end{aligned}$$

Hence, shifting contours on the RHS of (5.12) to  $\varepsilon$  and applying the above estimate we get that the LHS of (5.12) is bounded by

$$\ll x^\varepsilon \int_{(\varepsilon)} \exp(-|s|) |L(s+\frac{1}{2}, f \otimes \chi_{8e})| |ds|.$$

Also note, that by Cauchy-Schwarz and Corollary 2.5 of [31] (which follows from Heath-Brown's result [7])

$$\sum_{\substack{1 \leq (-1)^k e \leq x^{1+\varepsilon}/r^2 \\ e \text{ is } \square\text{-free} \\ (e,2)=1}} |L(s+\frac{1}{2}, f \otimes \chi_{8e})| \ll \frac{x^{1+\varepsilon}}{r^2} (1 + |\operatorname{Im}(s)|)^{1/2+\varepsilon},$$

for  $\operatorname{Re}(s) \geq 0$ . Applying this estimate in (5.11) it follows that the second sum in (5.10) is bounded by

$$(5.13) \quad \ll x^{1+\varepsilon} \sum_{|r| \leq x^{1/2+\varepsilon}} \sum_{\substack{a > Y \\ a|r}} \frac{1}{r^2} \ll \frac{x^{1+\varepsilon}}{Y}.$$

**The terms with  $a < Y$ : preliminary lemmas.** It remains to estimate the first sum on the LHS of (5.10). This will be done by applying Poisson summation to the character sum, as developed in [29].

Let  $\ell \in \mathbb{Z}, n \in \mathbb{N}$ . Define

$$G_\ell(n) = \left(\frac{1-i}{2} + \left(\frac{-1}{n}\right) \frac{1+i}{2}\right) \sum_{a \pmod{n}} \left(\frac{a}{n}\right) e\left(\frac{a\ell}{n}\right).$$

In Lemma 2.3 of [29] it is shown that  $G_\ell$  is a multiplicative function. Moreover,  $G_0(n) = \varphi(n)$  if  $n$  is a square and is identically zero otherwise. Also, for  $p^\alpha || \ell$ ,  $\ell \neq 0$

$$(5.14) \quad G_\ell(p^\beta) = \begin{cases} 0 & \text{if } \beta \leq \alpha \text{ is odd,} \\ \varphi(p^\beta) & \text{if } \beta \leq \alpha \text{ is even,} \\ -p^\alpha & \text{if } \beta = \alpha + 1 \text{ is even,} \\ \left(\frac{\ell p^{-\alpha}}{p}\right) p^\alpha \sqrt{p} & \text{if } \beta = \alpha + 1 \text{ is odd,} \\ 0 & \text{if } \beta \geq \alpha + 2. \end{cases}$$

**Lemma 5.2.** *Let  $F$  be a Schwartz function. Then for any odd integer  $n$*

$$\sum_{(d,2)=1} \left(\frac{d}{n}\right) F(d) = \frac{1}{2n} \left(\frac{2}{n}\right) \sum_{\ell} (-1)^\ell G_\ell(n) \tilde{F}\left(\frac{\ell}{2n}\right),$$

where

$$\tilde{F}(\lambda) = \int_{\mathbb{R}} (\cos(2\pi\lambda\xi) + \sin(2\pi\lambda\xi)) F(\xi) d\xi.$$

*Proof.* This is established in the proof of Lemma 2.6 of [29], in particular see the last equation of the proof.  $\square$

Using Lemma 5.1 it follows that  $F(\cdot; x, n)$  is a Schwartz function, since  $\phi$  and its derivatives decay rapidly. Hence, applying Lemma 5.2 the first sum on the RHS of (5.10) equals (note  $u$  is odd)

$$(5.15) \quad \begin{aligned} & 2 \sum_{\substack{a \leq Y \\ (a,2u)=1}} \mu(a) \sum_{(n,a)=1} \frac{\lambda_f(n)}{\sqrt{n}} \left(\frac{8}{nu}\right) \sum_{(m,2)=1} \left(\frac{m}{nu}\right) F\left(m; \frac{(-1)^k x}{8a^2}, \frac{(-1)^k n}{8a^2}\right) \\ &= 2 \sum_{\substack{a \leq Y \\ (a,2u)=1}} \mu(a) \sum_{(n,a)=1} \frac{\lambda_f(n)}{\sqrt{n}} \cdot \frac{1}{2nu} \left(\frac{16}{nu}\right) \sum_{\ell \in \mathbb{Z}} (-1)^\ell G_\ell(nu) \tilde{F}\left(\frac{\ell}{2nu}; \frac{x}{8a^2}, \frac{n}{8a^2}\right), \end{aligned}$$

where

$$\tilde{F}(\lambda; x, y) = \int_{\mathbb{R}} (\cos(2\pi\lambda\xi) + (-1)^k \sin(2\pi\lambda\xi)) F(\xi; x, y) d\xi.$$

Note that for odd  $k$  after applying Lemma 5.2 in (5.15) we also made the change of variables  $\xi \rightarrow -\xi$ .

Before proceeding we require several estimates for  $\tilde{F}$ . The first such result is a basic estimate on the rate of decay of  $\tilde{F}$ .

**Lemma 5.3.** *Let  $x, y > 0$  and  $\lambda \in \mathbb{R}$ . We have for any  $\theta \geq \varepsilon > 0$  that*

$$\tilde{F}(\lambda; x, y) \ll x \left( \frac{x}{y} \right)^\theta$$

where the implied constant depends on  $\varepsilon$ . Additionally, we have for any integer  $A \geq 1$  that

$$\tilde{F}(\lambda; x, y) \ll x \cdot \frac{1}{|y\lambda|^A}$$

where the implied constant depends on  $A$ .

*Proof.* Using the definition of  $W$  and shifting contours of integration to  $\operatorname{Re}(s) = \theta \geq \varepsilon > 0$  we have that

$$W\left(\frac{y}{x\xi}\right) \ll \left(\frac{x\xi}{y}\right)^\theta$$

so that

$$\tilde{F}(\lambda; x, y) \ll x \left(\frac{x}{y}\right)^\theta \int_{\mathbb{R}} \xi^\theta |\phi(\xi)| d\xi \ll x \left(\frac{x}{y}\right)^\theta.$$

Let  $\mathcal{W}(\xi) = W\left(\frac{y}{x\xi}\right)\phi(\xi)$ . For each integer  $0 \leq B \leq A$ , we get by shifting the contour of integration to  $\operatorname{Re}(s) = A$  and switching the order of differentiation and integration that

$$\frac{d^B}{d\xi^B} W\left(\frac{y}{x\xi}\right) \ll \left(\frac{x}{y}\right)^A \xi^{A-B}.$$

Using this and the fact that  $\phi$  has compact support it follows that

$$\mathcal{W}^{(A)}(\xi) \ll \left(\frac{x}{y}\right)^A \cdot \frac{1}{1 + |\xi|^2}.$$

Hence, integrating by parts and using the above bound it follows that

$$\tilde{F}(\lambda; x, y) \ll x \cdot \frac{1}{|x\lambda|^A} \int_{\mathbb{R}} |\mathcal{W}^{(A)}(\xi)| d\xi \ll x \cdot \frac{1}{|y\lambda|^A}.$$

□

We also require the following information about the Mellin transform of  $\tilde{F}$ . Let

$$\Phi(s) = \int_0^\infty \phi(t) t^{-s} dt.$$

**Lemma 5.4.** *For  $\ell \in \mathbb{Z}$  with  $\ell \neq 0$ , let*

$$\check{F}(s; \ell, u, a^2) = \int_0^\infty \tilde{F}\left(\frac{\ell}{2tu}; \frac{x}{8a^2}, \frac{t}{8a^2}\right) t^{s-1} dt.$$



Then for  $\operatorname{Re}(s) > 0$

(5.16)

$$\check{F}(s; \ell, u, a^2) = \frac{x^{1+s}}{8a^2} \Phi(-s) \int_0^\infty W\left(\frac{1}{y}\right) \left( \cos\left(\pi y \cdot \frac{\ell}{8ua^2}\right) + (-1)^k \sin\left(\pi y \cdot \frac{\ell}{8ua^2}\right) \right) \frac{dy}{y^{s+1}}.$$

Moreover, the function  $\check{F}(s; \ell, u, a^2)$  extends to an entire function in the half-plane  $\operatorname{Re}(s) \geq -1 + \varepsilon$  and in this region

$$\check{F}(s; \ell, u, a^2) \ll \frac{x^{1+\operatorname{Re}(s)}}{a^2(1+|s|)^A} \left( \left( \frac{|\ell|}{ua^2} \right)^{\operatorname{Re}(s)} + 1 \right)$$

for any  $A \geq 1$ .

*Proof.* Changing the order of integration and making a change of variables  $x\xi/t \rightarrow t$  in the integral over  $t$  it follows that  $\check{F}(s; \ell, u, a^2)$  equals

$$\begin{aligned} & \frac{x}{8a^2} \int_0^\infty \int_{\mathbb{R}} \left( \cos\left(\pi \frac{\ell}{8ua^2} \cdot \frac{x\xi}{t}\right) + (-1)^k \sin\left(\pi \frac{\ell}{8ua^2} \cdot \frac{x\xi}{t}\right) \right) W\left(\frac{t}{x\xi}\right) \phi(\xi) t^{s-1} d\xi dt \\ &= \frac{x^{1+s}}{8a^2} \int_{\mathbb{R}} \int_0^\infty \phi(\xi) \xi^s \cdot W\left(\frac{1}{t}\right) \left( \cos\left(\pi t \cdot \frac{\ell}{8ua^2}\right) + (-1)^k \sin\left(\pi t \cdot \frac{\ell}{8ua^2}\right) \right) \frac{dt}{t^{s+1}} d\xi, \end{aligned}$$

which establishes the first claim.

The function

$$w(s) = \int_0^\infty W\left(\frac{1}{y}\right) \left( \cos\left(\pi y \cdot \frac{\ell}{8ua^2}\right) + (-1)^k \sin\left(\pi y \cdot \frac{\ell}{8ua^2}\right) \right) \frac{dy}{y^{s+1}}$$

is holomorphic in the region  $\operatorname{Re}(s) \geq \varepsilon$ . Write  $w(s) = I_1(s) + I_2(s)$  where  $I_1$  is the portion of the integral over  $[0, \frac{8au^2}{|\ell|}]$  and  $I_2$  is the rest. Due to the rapid decay of  $W$ ,  $I_1(s)$  is holomorphic in the region  $\operatorname{Re}(s) \geq -1$  and in this region

$$(5.17) \quad I_1(s) \ll \int_0^{\frac{8au^2}{|\ell|}} |W(1/y)| y^{-\operatorname{Re}(s)-1} dy \ll \left( \frac{|\ell|}{ua^2} \right)^{\operatorname{Re}(s)} + 1.$$

Next, write

$$(5.18) \quad I_2(s) = \left( \frac{|\ell|}{8ua^2} \right)^s \int_1^\infty (\cos(\pi y) + (-1)^k \operatorname{sgn}(\ell) \cdot \sin(\pi y)) \frac{dy}{y^{s+1}} + I_3(s).$$

The integral  $I_3(s)$  is holomorphic in  $\operatorname{Re}(s) \geq -1$  and uniformly in this region we have by Lemma 5.1, that  $W(1/y) = 1 + O(y^{-k+\varepsilon})$  we get

$$I_3(s) \ll \int_{\frac{8au^2}{|\ell|}}^\infty \left| 1 - W\left(\frac{1}{y}\right) \right| \frac{dy}{y^{\operatorname{Re}(s)+1}} \ll \left( \frac{|\ell|}{ua^2} \right)^{\operatorname{Re}(s)} + 1.$$

The first integral on the RHS of (5.18) can be analytically continued to  $\operatorname{Re}(s) > -1$  by integrating by parts. This provides the analytic continuation of  $I_2(s)$  to  $\operatorname{Re}(s) \geq -1 + \varepsilon$ , and shows that in this region

$$I_2(s) \ll (|s| + 1) \left( \left( \frac{|\ell|}{ua^2} \right)^{\operatorname{Re}(s)} + 1 \right).$$

Hence applying this estimate along with (5.17) in (5.16) and noting  $\Phi$  decays rapidly establishes the claim.  $\square$

**The terms with  $a < Y$ : main term analysis i.e.  $\ell = 0$ .** Recall that  $G_0(n) = \varphi(n)$  if  $n = \square$  and is zero otherwise. So using Lemma 5.3 the term with  $\ell = 0$  in (5.15) equals

$$(5.19) \quad \sum_{(a,2u)=1} \mu(a) \sum_{\substack{(n,2a)=1 \\ nu=\square}} \frac{\lambda_f(n)}{\sqrt{n}} \frac{\varphi(nu)}{nu} \tilde{F} \left( 0; \frac{x}{8a^2}, \frac{n}{8a^2} \right) + O \left( \frac{x^{1+\varepsilon}}{Y} \right) \\ = \frac{x}{8} \frac{1}{2\pi i} \int_{(c)} \left( \frac{x}{2\pi} \right)^s \frac{\Gamma(s+k)}{\Gamma(k)} \left( \sum_{\substack{a \\ (a,2u)=1}} \frac{\mu(a)}{a^2} \sum_{\substack{(n,2a)=1 \\ nu=\square}} \frac{\lambda_f(n)}{n^{s+1/2}} \frac{\varphi(nu)}{nu} \right) \Phi(-s) \frac{ds}{s} + O \left( \frac{x^{1+\varepsilon}}{Y} \right).$$

We now evaluate the inner sum on the RHS of (5.19) (note  $\operatorname{Re}(s) = c > 0$ ). Since  $nu = \square$  write  $n = e^2 u_1$ , (recall  $u = u_1 u_2^2$ ) so

$$(5.20) \quad \sum_{(a,2u)=1} \frac{\mu(a)}{a^2} \sum_{\substack{(n,2a)=1 \\ nu=\square}} \frac{\lambda_f(n)}{n^{s+1/2}} \frac{\varphi(nu)}{nu} = \sum_{(e,2)=1} \frac{\lambda_f(u_1 e^2)}{(u_1 e^2)^{s+1/2}} \frac{\varphi(e^2 u_1 u)}{e^2 u_1 u} \sum_{(a,2eu)=1} \frac{\mu(a)}{a^2} \\ = \frac{4}{3\zeta(2)u_1^{s+\frac{1}{2}}} \sum_{(e,2)=1} \frac{\lambda_f(u_1 e^2)}{e^{2s+1}} \prod_{p|eu} \frac{1 - \frac{1}{p}}{1 - \frac{1}{p^2}}.$$

The sum on the RHS can be expressed as an Euler product as follows. Let  $r(n) = \prod_{p|n} \frac{1 - \frac{1}{p}}{1 - \frac{1}{p^2}}$ . Also, let

$$\tilde{\sigma}_p(z; \alpha, \beta) = \sum_{j=0}^{\infty} \frac{\lambda_f(p^{2j+\alpha})}{p^{jz}} r(p^{j+\beta}), \quad \mathcal{G}(z; u) = \prod_{p|u_1} \frac{\tilde{\sigma}_p(z; 1, 1)}{\tilde{\sigma}_p(z; 0, 0)} \prod_{\substack{p|u_2 \\ p \nmid u_1}} \frac{\tilde{\sigma}_p(z; 0, 1)}{\tilde{\sigma}_p(z; 0, 0)},$$

and

$$\mathcal{H}(z) = \prod_p \frac{\tilde{\sigma}_p(z; 0, 0)}{L_p(z, \operatorname{sym}^2 f)},$$

where  $L_p(z, \text{sym}^2 f)^{-1} = (1 - \frac{\alpha_p^2}{p^s})(1 - \frac{1}{p^s})(1 - \frac{\beta_p^2}{p^s})$  and  $\alpha_p, \beta_p$  are the Satake parameters of  $f$  (i.e.  $\alpha_p + \beta_p = \lambda_f(p)$  and  $\alpha_p \beta_p = 1$ ). It follows that

$$(5.21) \quad \sum_{(e,2)=1} \frac{\lambda_f(u_1 e^2)}{e^{2s+1}} \prod_{p|eu} \frac{1 - \frac{1}{p}}{1 - \frac{1}{p^2}} = L(1+2s, \text{sym}^2 f) \frac{\mathcal{G}(1+2s; u)}{\tilde{\sigma}_2(1+2s; 0, 0)} \mathcal{H}(1+2s).$$

Note for each prime  $p$  that  $L_p(1, \text{sym}^2 f) > 0$  and  $0 < r(p) < 1$  so that  $\tilde{\sigma}_p(1; 0, 0) > 0$  for each  $p$ .

Also,  $\frac{\mathcal{G}(1+2s; u)}{\tilde{\sigma}_2(1+2s; 0, 0)} \mathcal{H}(1+2s) \ll u^\varepsilon$  for  $\text{Re}(s) \geq -\frac{1}{4} + \varepsilon$  and is analytic in this region. Hence, applying (5.20) and (5.21) in the RHS of (5.19) then shifting contours  $\text{Re}(s) = -\frac{1}{4} + \varepsilon$ , collecting the residue at  $s = 0$  we see that the RHS of (5.19) equals

$$(5.22) \quad \frac{x}{\pi^2 u_1^{1/2}} \cdot L(1, \text{sym}^2 f) \frac{\mathcal{G}(1; u)}{\tilde{\sigma}_2(1; 0, 0)} \mathcal{H}(1) \cdot \Phi(0) + O(u^\varepsilon x^{3/4+\varepsilon}) + O\left(\frac{x^{1+\varepsilon}}{Y}\right).$$

Finally, note  $\mathcal{G}(1; \cdot)$  is a multiplicative function satisfying  $\mathcal{G}(1; p^{2j+1}) = \lambda_f(p) + O(1/p)$  and  $\mathcal{G}(1; p^{2j}) = 1 + O(1/p)$ . Set  $\vartheta(n) = \mathcal{G}(1; n)$ .

**The terms with  $a \leq Y$ : Off-diagonal analysis i.e.  $\ell \neq 0$ .** In (5.15) it remains to bound

$$(5.23) \quad \sum_{\ell \neq 0} \sum_{\substack{a \leq Y \\ (a, 2u) = 1}} \left| \sum_{(n, 2a) = 1} \frac{\lambda_f(n)}{n^{1/2}} \left(\frac{4}{n}\right) \frac{G_\ell(nu)}{nu} \tilde{F}\left(\frac{\ell}{2nu}; \frac{x}{8a^2}, \frac{n}{8a^2}\right) \right|.$$

First, note that by Lemma 5.3 the contribution from the terms with  $|\ell| \geq Y^2 u x^\varepsilon$  to (5.23) is bounded by

$$(5.24) \quad \ll \sum_{|\ell| \geq Y^2 u x^\varepsilon} \sum_{a \leq Y} \frac{x}{a^2} \left( \sum_{n \leq |\ell|^\varepsilon x} \left(\frac{a^2 u}{|\ell|}\right)^A + \sum_{n > |\ell|^\varepsilon x} \left(\frac{x}{n}\right)^\theta \right) \ll \frac{1}{x}$$

where  $A, \theta$  have been chosen sufficiently large with respect to  $\varepsilon$ .

It remains to estimate the terms in (5.23) with  $|\ell| \leq Y^2 u x^\varepsilon$ . Let  $\psi_{4\ell}(n) = \left(\frac{4\ell}{n}\right)$  and note that  $\psi_{4\ell}$  is a character of modulus at most  $4|\ell|$ . Using that  $G_\ell$  is a multiplicative function, and using (5.14) we can write

$$(5.25) \quad \sum_{(n, 2a) = 1} \frac{\lambda_f(n)}{n^s} \left(\frac{4}{n}\right) \frac{G_\ell(nu)}{u\sqrt{n}} = L(s, f \otimes \psi_{4\ell}) R(s; \ell, u, a), \quad \text{Re}(s) > 1,$$

where

$$R(s; \ell, u, a) = \prod_p R_p(s; \ell, u, a).$$

It follows that for  $(p, 4au\ell) = 1$

$$R_p(s; \ell, u, a) = \left(1 + \frac{\lambda_f(p)}{p^s} \left(\frac{4\ell}{p}\right)\right) \cdot \left(1 - \frac{\lambda_f(p)}{p^s} \left(\frac{4\ell}{p}\right) + \frac{1}{p^{2s}}\right) = 1 + O\left(\frac{1}{p^{2\operatorname{Re}(s)}}\right).$$

For  $p|au\ell$  and  $\operatorname{Re}(s) \geq \frac{1}{2} + \varepsilon$ , writing  $p^\theta || u$ , we have that

$$\begin{aligned} |R_p(s; \ell, u, a)| &= \left| \left(1 + O\left(\frac{1}{p^{1/2}}\right)\right) \cdot \left(\frac{G_\ell(p^\theta)}{p^\theta} + O\left(\sum_{j \geq 1} \frac{p^{j+\theta}}{p^{j(\frac{1}{2}+\varepsilon)} p^{j/2+\theta}}\right)\right) \right| \\ &\leq \left(1 + O\left(\frac{1}{p^\varepsilon}\right)\right). \end{aligned}$$

Also  $R_2(s; \ell, u, a) \ll 1$  for  $\operatorname{Re}(s) \geq \frac{1}{2}$ . Hence, for  $0 < a, u, |\ell| \leq x^2$  and  $\operatorname{Re}(s) \geq \frac{1}{2} + \varepsilon$

$$(5.26) \quad R(s; \ell, u, a) \ll \prod_{p|au\ell} \left(1 + \frac{O(1)}{p^\varepsilon}\right) \ll x^\varepsilon.$$

So  $R(\cdot; \ell, u, a)$  is an absolutely convergent Euler product and thereby defines a holomorphic function in the half-plane  $\operatorname{Re}(s) \geq \frac{1}{2} + \varepsilon$ . Hence, applying Mellin inversion and (5.25) we get for  $c > 0$  that

$$(5.27) \quad \begin{aligned} \sum_{(n, 2a)=1} \frac{\lambda_f(n)}{n} \left(\frac{4}{n}\right) \frac{G_\ell(nu)}{u\sqrt{n}} \tilde{F}\left(\frac{\ell}{2nu}; \frac{x}{8a^2}, \frac{n}{8a^2}\right) \\ = \frac{1}{2\pi i} \int_{(c)} \check{F}(s; \ell, u, a^2) L(1+s, f \otimes \psi_{4\ell}) R(1+s; \ell, u, a) ds. \end{aligned}$$

Using Lemma 5.4 and (5.26) we can shift contours in the integral above to  $\operatorname{Re}(s) = -\frac{1}{2} + \varepsilon$ , since the integrand is holomorphic in this region. Using the convexity bound

$$L\left(\frac{1}{2} + \varepsilon + it, f \otimes \psi_{4\ell}\right) \ll (1 + |t|)^{1/2+\varepsilon} |\ell|^{1/2+\varepsilon}$$

along with Lemma 5.4 and (5.26) it follows that for  $|\ell|, u, a \leq x$

$$(5.28) \quad \frac{1}{2\pi i} \int_{(-\frac{1}{2}+\varepsilon)} \check{F}(s; \ell, u, a^2) L(1+s, f \otimes \psi_{4\ell}) R(1+s; \ell, u, a) ds \ll \frac{x^{\frac{1}{2}+\varepsilon}}{a^2} \cdot \left( \left(\frac{ua^2}{|\ell|}\right)^{1/2} + 1 \right) |\ell|^{1/2}.$$

Applying (5.27) and (5.28) in (5.23) the terms with  $|\ell| \leq Y^2 x^\varepsilon u$  in (5.23) are bounded by

$$(5.29) \quad x^{\frac{1}{2}+\varepsilon} \sum_{1 \leq |\ell| \leq Y^2 x^\varepsilon u} \sum_{a \leq Y} \frac{1}{a^2} \left( \left(\frac{ua^2}{|\ell|}\right)^{1/2} + 1 \right) |\ell|^{1/2} \ll x^{\frac{1}{2}+\varepsilon} Y^3 u^{3/2}.$$

**Completion of the proof of Proposition 5.1.** Applying (5.22), (5.24), (5.29) in (5.15) and then using the resulting formula along with (5.13) in (5.10) it follows that

$$(5.30) \quad \sum_{2m \text{ is } \square\text{-free}}^m L\left(\frac{1}{2}, f \otimes \chi_{8m}\right) \chi_{8m}(u) \phi\left(\frac{(-1)^k 8m}{x}\right) \\ = \frac{x}{\pi^2 u_1^{1/2}} \cdot L(1, \text{sym}^2 f) \frac{\mathcal{G}(1; u)}{\tilde{\sigma}_2(1; 0, 0)} \mathcal{H}(1) \cdot \Phi(0) + O\left(\frac{x^{1+\varepsilon}}{Y} + x^{\frac{1}{2}+\varepsilon} Y^3 u^{3/2}\right).$$

Choosing  $Y = x^{1/8} u^{-3/8}$  and recalling the estimates given for  $\mathcal{G}(1; \cdot)$  after (5.22) completes the proof with  $C = \frac{1}{\pi^2} \cdot L(1, \text{sym}^2 f) \frac{\mathcal{H}(1)}{\tilde{\sigma}_2(1; 0, 0)}$ . By the comment directly after (5.21) it follows that  $C > 0$ .

## 6. THE PROOF OF PROPOSITION 1.3

In order to establish Proposition 1.3 we will need the following variant of the shifted convolution problem for coefficients of half-integral weight forms.

**Proposition 6.1.** *Uniformly in  $1 \leq \Delta \leq X^{\frac{1}{52}}$ ,  $0 < |h| < X^{\frac{1}{2}}$  and  $(v, \Delta) = 1$ , we have for every  $\varepsilon > 0$*

$$(6.1) \quad \frac{1}{X^{k-1/2}} \left| \sum_n c(n) n^{\frac{k-1/2}{2}} e\left(\frac{nv}{\Delta}\right) e^{-\frac{2\pi n}{X}} c(n+h) (n+h)^{\frac{k-1/2}{2}} e^{-\frac{2\pi(n+h)}{X}} \right| \ll X^{1-\frac{1}{104}+\varepsilon}.$$

**6.1. The shifted convolution problem.** We begin with a proof of Proposition 6.1. The proof is based on the by now standard combination of the circle method and modularity. Recall that a weight  $k + \frac{1}{2}$  modular form (with trivial character) transforms under  $\Gamma_0(4)$  in the following way

$$(6.2) \quad g(\gamma z) = \nu(\gamma) j_\gamma(z)^{k+\frac{1}{2}} g(z), \quad \forall \gamma \in \Gamma_0(4),$$

where for  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  we set  $j_\gamma(z) = cz + d$  and  $\nu(\gamma) = \left(\frac{c}{d}\right) \varepsilon_d$ , where  $\left(\frac{\cdot}{\cdot}\right)$  denotes the quadratic residue symbol in the sense of Shimura [27] (see *Notation 3*) and

$$\varepsilon_d := \begin{cases} 1 & \text{if } d \equiv 1 \pmod{4}, \\ i^{2k+1} & \text{if } d \equiv -1 \pmod{4}. \end{cases}$$

Also, we record the following estimate for the Fourier coefficients of  $g$

$$(6.3) \quad \sum_{n \leq X} |c(n)|^2 \ll X$$

(see [10, Theorem 5.1] and use partial summation). This implies that

$$(6.4) \quad |c(n)| \ll n^{1/2+\varepsilon}.$$

We record below a few standard lemmas.

**Lemma 6.1.** *Let  $\eta \in (0, 1]$  and  $Q \geq 1$  be given. Let  $Q^{\eta/2} \geq \Delta \geq 1$  be an integer. Let  $I_{d/q}(\cdot)$  denote the indicator function of the interval  $[d/q - Q^{-2+\eta}; d/q + Q^{-2+\eta}]$ . Let  $\mathcal{Q}$  denote the set of integers  $n \in [Q, 2Q]$  that can be written as  $4\Delta r$  with  $r \equiv 1 \pmod{4}$  prime. For  $\alpha \in \mathbb{R}$ , put*

$$\tilde{I}(\alpha) = \frac{Q^{2-\eta}}{2L} \sum_{q \in \mathcal{Q}} \sum_{d \pmod{q}}^* I_{d/q}(\alpha) \quad \text{and} \quad L = \sum_{q \in \mathcal{Q}} \varphi(q).$$

Finally let  $I(\cdot)$  denote the indicator function of the interval  $[0, 1]$ . Then, for every  $\varepsilon > 0$

$$\int_{\mathbb{R}} (I(\alpha) - \tilde{I}(\alpha))^2 d\alpha \ll_{\varepsilon} Q^{-\eta/2+\varepsilon}.$$

*Proof.* This is a consequence of a result of Jutila (see [12] or [5, Proposition 2]). In the notation of [5, Proposition 2] we specialize to  $\delta = Q^{-2+\eta}$  and notice that  $L \asymp \frac{\varphi(4\Delta)}{\Delta^2} \cdot \frac{Q^2}{\log Q}$ .  $\square$

Crucial to our analysis is the following consequence of the modularity of  $g$ .

**Lemma 6.2.** *Let  $g$  be a cusp form of weight  $k + \frac{1}{2}$  of level 4 where  $k \geq 2$  is an integer. Let  $(d, q) = 1$  and  $(v, \Delta) = 1$  be two pairs of co-prime integers. Suppose that  $q = 4\Delta r$  with  $r > 4\Delta$  a prime congruent to 1  $\pmod{4}$ . Write  $d = rb + 4\Delta\ell$  with  $(b, 4\Delta) = 1$  and  $(\ell, r) = 1$ . Let  $\Delta^* = \Delta/(4v + b, \Delta)$  and  $(4v + b)^* = (4v + b)/(4v + b, \Delta)$ . Then, for any  $\beta \in \mathbb{R}$  and any real  $X \geq 1$*

$$g\left(\frac{d}{q} + \frac{v}{\Delta} + \beta + \frac{i}{X}\right) = \overline{\varepsilon_{(4v+b)^*} \chi_{4\Delta^*}}((4v+b)^* r) \chi_r(4\Delta^* \ell) \\ \times \left(\frac{X}{4\Delta^* r}\right)^{k+\frac{1}{2}} \left(\frac{i}{1-i\beta X}\right)^{k+\frac{1}{2}} g\left(-\frac{r^2(4v+b)^*}{4\Delta^*} - \frac{16(\Delta^*)^2 \ell}{r} + \frac{i}{(4\Delta^* r)^2(\frac{1}{X} - i\beta)}\right),$$

where  $\chi_t(\cdot)$  denotes a real Dirichlet character of modulus  $t$ . Finally whenever we write  $\frac{\bar{a}}{q}$  we denote by  $\bar{a}$  an integer such that  $\bar{a} \equiv 1 \pmod{q}$ .

**Remark 2.** *By inspection of the proof below, the conclusion of the lemma also holds when  $v = 0$  and we will use this later.*

*Proof.* Since  $q = 4\Delta r$  we can write

$$\frac{d}{q} + \frac{v}{\Delta} = \frac{d + 4vr}{4\Delta r}.$$

Notice that  $(d + 4vr, 4\Delta r) = (d + 4vr, \Delta)$  since  $(d, 4r) = 1$  by assumption. Therefore the above is equal to

$$(6.5) \quad \frac{(4v + b)^*r + 4\Delta^*\ell}{4\Delta^*r}.$$

Throughout set  $w := (4v + b)^*r + 4\Delta^*\ell$ . We now consider

$$z = -\frac{\bar{w}}{4\Delta^*r} + \frac{i}{(4\Delta^*r)^2(\frac{1}{X} - i\beta)}$$

and the matrix

$$\gamma = \begin{pmatrix} w & \star \\ 4\Delta^*r & \bar{w} \end{pmatrix} \in \Gamma_0(4)$$

where  $\star$  is uniquely determined by the choice of  $\bar{w} \in \mathbb{Z}$ . A slightly tedious computation using (6.5) reveals that

$$(6.6) \quad \gamma z = \left(\frac{d}{q} + \beta\right) + \frac{v}{\Delta} + \frac{i}{X}.$$

We also find that

$$(6.7) \quad j_\gamma(z) = \frac{i}{\frac{4\Delta^*r}{X} - i(4\Delta^*r)\beta} = \frac{1}{4\Delta^*r} \cdot \frac{iX}{1 - iX\beta}$$

and that

$$(6.8) \quad \nu(\gamma) = \overline{\varepsilon_{(4v+b)^*}} \cdot \left(\frac{4\Delta^*r}{\bar{w}}\right),$$

where  $\left(\frac{\cdot}{\cdot}\right)$  denotes the extended quadratic residue symbol in the sense of Shimura [27, Notation 3]. In particular since  $4\Delta^*r$  is divisible by four, this extended quadratic residue symbol coincides with  $\chi_{4\Delta^*r}(w)$ , a real Dirichlet character of modulus  $4\Delta^*r$ , so that  $\chi_{4\Delta^*r}(\bar{w}) = \chi_{4\Delta^*r}(w)$ . Moreover by multiplicativity of the Jacobi symbol

$$\chi_{4\Delta^*r}(w) = \left(\frac{4\Delta^*r}{w}\right) = \left(\frac{4\Delta^*}{w}\right) \left(\frac{r}{w}\right).$$

Since  $4\Delta^*$  is divisible by 4 and  $r$  is congruent to 1 (mod 4) both expressions are Dirichlet characters of modulus  $4\Delta^*$  and modulus  $r$  respectively. In particular the above expression is equal to

$$\chi_{4\Delta^*}(w)\chi_r(w) = \chi_{4\Delta^*}((4v + b)^*r)\chi_r(\Delta^*\ell).$$

Using the modularity of  $g$ , (6.2), and combining this with (6.6), (6.7), (6.8) we conclude that

$$g\left(\frac{d}{q} + \frac{v}{\Delta} + \beta + \frac{i}{X}\right) = \overline{\varepsilon_{(4v+b)^*}} \chi_{4\Delta^*}((4v+b)^*r) \chi_r(4\Delta^*\ell) \cdot \left(\frac{X}{4\Delta^*r}\right)^{k+\frac{1}{2}} \cdot \left(\frac{i}{1-iX\beta}\right)^{k+\frac{1}{2}} \\ \times g\left(-\frac{\bar{w}}{4\Delta^*r} + \frac{i}{(4\Delta^*r)^2 \cdot (\frac{1}{X} - i\beta)}\right).$$

Write  $(\bar{a})_b$  for the inverse of  $a$  modulo  $b$ . Also, let  $a_1 = \overline{(r \cdot (4v+b)^*)}_{4\Delta^*}$  and  $a_2 = \overline{(4\Delta^* \cdot \ell)}_r$ . Observe

$$(\bar{w})_{4\Delta^*r} \equiv a_1 \pmod{4\Delta^*} \quad \text{and} \quad (\bar{w})_{4\Delta^*r} \equiv a_2 \pmod{r}.$$

So by the Chinese Remainder Theorem

$$(\bar{w})_{4\Delta^*r} \equiv a_1(\bar{r})_{4\Delta^*r} + a_2\overline{(4\Delta^*)}_r 4\Delta^* \pmod{4\Delta^*r}.$$

It follows that

$$\frac{\bar{w}}{4\Delta^*r} \equiv \frac{\overline{r^2(4v+b)^*}}{4\Delta^*} + \frac{16(\Delta^*)^2\ell}{r} \pmod{1}$$

and since  $g(x+iy)$  is 1-periodic in  $x$  the claim follows.  $\square$

We are now ready to prove the main proposition of this section.

*Proof of Proposition 6.1.* Using the circle method we can re-write the LHS of (6.1) as

$$\frac{1}{X^{k-\frac{1}{2}}} \left| \int_0^1 g\left(\alpha + \frac{v}{\Delta} + \frac{i}{X}\right) g\left(-\alpha + \frac{i}{X}\right) e(-\alpha h) d\alpha \right|.$$

Let  $Q = X^{1/2+2\eta}$  and  $\eta \in (0, 1]$  to be determined later (we will choose  $\eta = \frac{1}{13}$ ). Let  $1 \leq \Delta \leq X^{\eta/4} \leq Q^{\eta/2}$ . By Lemma 6.1 the above expression is equal to

$$(6.9) \quad \frac{1}{X^{k-\frac{1}{2}}} \frac{Q^{2-\eta}}{2L} \sum_{q \in \mathcal{Q}} \sum_{d \pmod{q}}^* e\left(-\frac{hd}{q}\right) \int_{-Q^{-2+\eta}}^{Q^{-2+\eta}} g\left(\frac{d}{q} + \frac{v}{\Delta} + \beta + \frac{i}{X}\right) g\left(-\frac{d}{q} - \beta + \frac{i}{X}\right) e(-\beta h) d\beta,$$

plus an error term of size  $\ll XQ^{-\eta/4} \leq X^{1-\eta/8}$ , where in the estimation of the error term we have used the bound  $|g(\alpha + i/X)| \ll X^{\frac{k+\frac{1}{2}}{2}}$  which holds uniformly for all  $\alpha \in \mathbb{R}$ . This follows from the fact that  $y^{(k+\frac{1}{2})/2}|g(z)|$  is bounded on  $\mathbb{H}$  since  $g$  is a cusp form.

We now write  $q = 4\Delta r$  with  $r$  a prime congruent to 1 (mod 4) and just as in Lemma 6.2 we write,  $d = (4\Delta)\ell + rb$  with  $(\ell, r) = 1$  and  $(b, 4\Delta) = 1$ . Then by



Lemma 6.2 and the triangle inequality, (6.9) is less than

$$(6.10) \quad \leq X \cdot \frac{Q^{2-\eta}}{2L} \sum_{b \pmod{4\Delta}}^* \sum_{\substack{Q \leq 4\Delta r \leq 2Q \\ r \equiv 1 \pmod{4} \text{ prime}}} \left( \frac{\sqrt{X}}{4\Delta^* r} \right)^{k+\frac{1}{2}} \cdot \left( \frac{\sqrt{X}}{4\Delta r} \right)^{k+\frac{1}{2}} \cdot S,$$

where

$$(6.11) \quad S = \left| \sum_{\ell \pmod{r}}^* e\left(-\frac{h\ell}{r}\right) \int_{-Q^{-2+\eta}}^{Q^{-2+\eta}} g\left(-\frac{r^2(4v+b)^*}{4\Delta^*} - \frac{16(\Delta^*)^2\ell}{r} + \frac{i}{(4\Delta^*r)^2(\frac{1}{X} - i\beta)}\right) \right. \\ \left. \times g\left(\frac{\overline{r^2b}}{4\Delta} + \frac{16\overline{\Delta^2\ell}}{r} + \frac{i}{(4\Delta r)^2(\frac{1}{X} + i\beta)}\right) \cdot \frac{e(-h\beta)d\beta}{|1 - i\beta X|^{2k+1}} \right|,$$

and where we applied Lemma 6.2 with  $v = 0$  in second appearance of  $g$ . Notice now that for any  $q \geq 1$ ,

$$\frac{i}{q^2(\frac{1}{X} - i\beta)} = \frac{iX}{q^2 \cdot |1 - i\beta X|^2} - \frac{X^2\beta}{q^2|1 - i\beta X|^2}.$$

Moreover since  $|\beta X| \leq Q^{-2+\eta}X \leq X^{-\eta}$  the imaginary part of the above expression is  $(1 + o(1))X/q^2$ . Therefore expanding in Fourier coefficients we can write, for any  $\varepsilon > 0$ ,

$$(6.12) \quad g\left(-\frac{r^2(4v+b)^*}{4\Delta^*} - \frac{16(\Delta^*)^2\ell}{r} + \frac{i}{(4\Delta^*r)^2(\frac{1}{X} - i\beta)}\right) \\ = \sum_{n \leq X^\varepsilon(4\Delta^*r)^2/X} \alpha(n; \beta) n^{\frac{k-\frac{1}{2}}{2}} e\left(-\frac{n\overline{16\Delta^*\ell}}{r}\right) + O(X^{-A})$$

for any  $A \geq 1$ , where the coefficients  $\alpha(n; \beta)$  are independent of  $\ell$  and  $|\alpha(n; \beta)| \ll |c(m)| \ll_\varepsilon n^{1/2+\varepsilon}$  for all  $\varepsilon > 0$  and  $\beta \in \mathbb{R}$  by (6.4). Similarly we have for every  $\varepsilon > 0$

$$(6.13) \quad g\left(\frac{\overline{r^2b}}{4\Delta} + \frac{16\overline{\Delta^2\ell}}{r} + \frac{i}{(4\Delta r)^2(\frac{1}{X} + i\beta)}\right) = \sum_{m \leq X^\varepsilon(4\Delta r)^2/X} \gamma(m; \beta) m^{\frac{k-\frac{1}{2}}{2}} e\left(\frac{m\overline{16\Delta^2\ell}}{r}\right) + O(X^{-A})$$

for any  $A \geq 1$ , where the coefficients  $\gamma(m; \beta)$  are independent of  $\ell$  and  $|\gamma(m; \beta)| \ll |c(n)| \ll_\varepsilon m^{1/2+\varepsilon}$  for all  $\varepsilon > 0$  and  $\beta \in \mathbb{R}$ .

Applying (6.12) and (6.13) in (6.11) it follows that

$$S \ll \int_{-Q^{-2+\eta}}^{Q^{-2+\eta}} \sum_{\substack{m \leq X^\varepsilon(4\Delta r)^2/X \\ n \leq X^\varepsilon(4\Delta^*r)^2/X}} \left| \alpha(n; \beta) \gamma(m; \beta) (mn)^{\frac{k-\frac{1}{2}}{2}} S(\overline{m16\Delta^2} - \overline{n16(\Delta^*)^2}, -h; r) \right| d\beta + X^{-100},$$

where  $S(a, b; c)$  is the standard Kloosterman sum. Since  $0 < |h| < r$  and  $r$  is prime the Weil bound shows that the Kloosterman sum above is always  $\ll \sqrt{r} \ll \sqrt{\frac{Q}{\Delta}}$ . Therefore, applying Cauchy-Schwarz along with (6.3) in the above equation it follows that

$$S \ll X^\varepsilon \cdot Q^{-2+\eta} \cdot \left(\frac{(4\Delta r)^2}{X}\right)^{\frac{k+\frac{3}{2}}{2}} \cdot \left(\frac{(4\Delta r)^2}{X}\right)^{\frac{k+\frac{3}{2}}{2}} \cdot \sqrt{\frac{Q}{\Delta}}.$$

Also, recall that  $Q = X^{\frac{1}{2}+2\eta}$ ,  $\Delta \leq X^{\eta/4}$  and  $L \gg \frac{Q^2}{\Delta X^\varepsilon}$ . We conclude that (6.10) is

$$\ll \frac{X^{1+\varepsilon}}{L} \cdot \Delta \cdot \frac{Q}{\Delta} \cdot \frac{Q^2}{X} \cdot \sqrt{\frac{Q}{\Delta}} \ll X^{\frac{3}{4} + \frac{25\eta}{8} + \varepsilon}.$$

We also recall that when using Lemma 6.1 in (6.9) we introduced an error of size  $X^{1-\eta/8}$ . Picking  $\eta = \frac{1}{13}$  gives a total error of size  $\ll X^{1-\frac{1}{104}+\varepsilon}$  and allows  $\Delta$  to go up to  $X^{\frac{1}{52}}$ .  $\square$

**6.2. Proof of Proposition 1.3.** We are now ready to prove Proposition 1.3. Let  $h$  be the indicator function of the integers that can be written as  $8n$  with  $n$  odd and square-free. We find that

$$h(n) = \sum_{\substack{2^{\alpha+3}d^2|n \\ d \text{ odd}}} \mu(d)\mu(2^\alpha).$$

Let

$$h_{\leq Y}(n) = \sum_{\substack{n=2^{\alpha+3}d^2m \\ d \leq Y, \alpha \leq 1 \\ d \text{ odd}}} \mu(d)\mu(2^\alpha), \quad h_{> Y}(n) = \sum_{\substack{n=2^{\alpha+3}d^2m \\ d > Y, \alpha \leq 1 \\ d \text{ odd}}} \mu(d)\mu(2^\alpha).$$

By the triangle inequality,

$$\sum_{X \leq x \leq 2X} \left| \sum_{x \leq n \leq x+y} c(n)M((-1)^k n)h_{> Y}(n) \right| \leq y \sum_{n \leq 4X} |c(n)M((-1)^k n)h_{> Y}(n)|$$

and by the definition of  $h_{> Y}(n)$  this is

$$(6.14) \quad \ll y \sum_{\substack{d > Y \\ d \text{ odd}}} \sum_{\substack{n \leq 4X \\ d^2 | n}} |c(n)M((-1)^k n)|.$$

A trivial bound gives  $M((-1)^k n) \ll_\varepsilon MX^\varepsilon$ . By Shimura's result (3.3) and Deligne's bound,  $|c(d^2n)| \ll_\varepsilon d^\varepsilon |c(n)|$  for all  $\varepsilon > 0$ . Hence, we conclude after an application of

Cauchy-Schwarz and (6.3) that

$$\sum_{\substack{n \leq X \\ d^2 | n}} |c(n)| \ll_\varepsilon d^\varepsilon \sum_{n \leq X/d^2} |c(n)| \ll_\varepsilon d^\varepsilon \cdot \left\lfloor \frac{X}{d^2} \right\rfloor.$$

Therefore (6.14) is bounded by

$$\ll_\varepsilon y \left( \frac{X}{Y} + 1 \right) M X^\varepsilon.$$

This contributes  $\ll X^{1-\delta+\varepsilon}$  provided that  $Y \geq X^\delta M y$ .

Therefore after an application of Cauchy-Schwarz, it remains to obtain a non-trivial upper bound for

$$\sum_{X \leq x \leq 2X} \left| \sum_{x \leq n \leq x+y} c(n) M((-1)^k n) h_{\leq Y}(n) \right|^2.$$

We introduce an auxiliary smoothing, and bound the above by

$$\ll \sum_{x \geq 1} \left| \sum_{x \leq n \leq x+y} c(n) M((-1)^k n) h_{\leq Y}(n) \right|^2 f\left(\frac{x}{X}\right)^2$$

where

$$f(u) = \exp(-2\pi u) u^{\frac{k-1/2}{2}}.$$

We note that the implicit constant is allowed to depend on the weight  $k + \frac{1}{2}$ . Let  $\alpha(n) = c(n) M((-1)^k n) h_{\leq Y}(n)$ . Expanding the square we re-write the above expression as

$$\sum_{0 \leq h_1, h_2 \leq y} \sum_{n \geq 1} \alpha(n+h_1) \alpha(n+h_2) f\left(\frac{n}{X}\right)^2$$

Grouping terms together and using a Taylor expansion we can re-write the above as

$$\begin{aligned} & \sum_{|h| \leq y} \sum_{n \geq 1} \alpha(n) f\left(\frac{n}{X}\right) \alpha(n+h) f\left(\frac{n+h}{X}\right) \sum_{\substack{0 \leq h_1, h_2 \leq y \\ h_1 - h_2 = h}} \left(1 + O\left(\frac{y}{n} + \frac{y}{X}\right)\right) + O(1) \\ (6.15) \quad & = \sum_{|h| \leq y} (y+1-|h|) \sum_{n \geq 1} \alpha(n) f\left(\frac{n}{X}\right) \alpha(n+h) f\left(\frac{n+h}{X}\right) + O(y^3 M^2 X^\varepsilon) \end{aligned}$$

where we set  $\alpha(n) := 0$  for  $n \leq 0$  and where we used (6.3) in the estimation of the error term. The term  $h = 0$  contributes

$$(6.16) \quad \ll y \sum_{n \geq 1} |c(n) h_{\leq Y}(n) M((-1)^k n)|^2 \cdot f\left(\frac{n}{X}\right)^2$$

Repeating the same argument as before we see that

$$y \sum_{n \geq 1} |c(n)h_{>Y}(n)M((-1)^k n)|^2 \cdot f\left(\frac{n}{X}\right)^2 \ll_{\varepsilon} y \left(\frac{X}{Y} + 1\right) X^{\varepsilon} M^2.$$

This gives a total contribution of  $\ll X^{1-\delta+\varepsilon}$  provided that  $Y > X^{\delta} M^2 y$ . It follows that (6.16) is bounded by

$$\ll y \sum_{n \text{ odd}} \mu^2(n) |c(8n)M((-1)^k 8n)|^2 \cdot f\left(\frac{n}{X}\right)^2 + X^{1-\delta+\varepsilon}$$

as needed.

We now focus on the terms with  $h \neq 0$  in (6.15). Opening  $h_{\leq Y}$  and  $M((-1)^k n)$  we see that the RHS of (6.15) restricted to  $h \neq 0$  is bounded by

$$\begin{aligned} &\ll_{\varepsilon} y^2 M^2 Y^2 X^{\varepsilon} \\ &\times \sup_{\substack{0 < |h| \leq y \\ d_1, d_2 \leq Y \\ \alpha_1, \alpha_2 \in \{0,1\} \\ m_1, m_2 \leq M}} \left| \sum_{\substack{d_1^2 2^{\alpha_1+3} | n \\ d_2^2 2^{\alpha_2+3} | n+h}} c(n) \chi_{(-1)^k n}(m_1) f\left(\frac{n}{X}\right) c(n+h) \chi_{(-1)^k(n+h)}(m_2) f\left(\frac{n+h}{X}\right) \right|. \end{aligned}$$

By the Chinese Remainder Theorem the condition  $2^{\alpha_1+3} d_1^2 | n$  and  $2^{\alpha_2+3} d_2^2 | n+h$  can be re-written as a single congruence condition to a modulus of size  $\leq 4Y^4$ . Moreover  $\chi_{(-1)^k n}(m_1)$  is  $4m_1$ -periodic in  $n$ , where-as  $\chi_{(-1)^k(n+h)}(m_2)$  is  $4m_2$ -periodic in  $n$ . Therefore fixing the congruence class of  $n$  modulo  $4[m_1, m_2]$  fixes the value of  $\chi_{(-1)^k n}(m_1) \chi_{(-1)^k(n+h)}(m_2)$ . Therefore we can bound the above supremum by

$$(6.17) \quad \sup_{\substack{0 < |h| \leq y \\ 1 \leq \Delta \leq 4Y^4 M^2 \\ \gamma \pmod{\Delta}}} \left| \sum_{n \equiv \gamma \pmod{\Delta}} c(n) f\left(\frac{n}{X}\right) c(n+h) f\left(\frac{n+h}{X}\right) \right|.$$

Finally, we can write

$$\mathbf{1}_{n \equiv \gamma \pmod{\Delta}} = \frac{1}{\Delta} \sum_{0 \leq v < \Delta} e\left(\frac{vn}{\Delta}\right) e\left(-\frac{v\gamma}{\Delta}\right).$$

Plugging this into (6.17) we see that it is

$$\leq \sup_{\substack{0 < |h| \leq y \\ 1 \leq \Delta \leq 4Y^4 M^2 \\ v \pmod{\Delta}}} \left| \sum_{n \geq 1} c(n) e\left(\frac{nv}{\Delta}\right) f\left(\frac{n}{X}\right) c(n+h) f\left(\frac{n+h}{X}\right) \right|$$

and without loss of generality we can also assume that  $(v, \Delta) = 1$ . It follows that the RHS of (6.15) restricted to  $h \neq 0$  is bounded by

$$\ll_{\varepsilon} X^{\varepsilon} \cdot y^2 M^2 Y^2 \sup_{\substack{0 < |h| \leq y \\ 1 \leq \Delta \leq 4Y^4 M^2 \\ (v, \Delta) = 1}} \left| \sum_{n \geq 1} c(n) e\left(\frac{nv}{\Delta}\right) f\left(\frac{n}{X}\right) c(n+h) f\left(\frac{n+h}{X}\right) \right|.$$

According to Proposition 6.1 the above expression is  $\ll_{\varepsilon} y^2 M^2 Y^2 X^{1 - \frac{1}{104} + \varepsilon}$  provided that  $4Y^4 M^2 \leq X^{\frac{1}{52}}$ . Moreover we introduced earlier error terms that were  $\ll_{\varepsilon} X^{1 - \delta/2 + \varepsilon}$  as long as  $Y \geq yX^{\delta} M$ . Choosing  $Y = yX^{\delta} M$  we obtain the restriction  $y^4 X^{4\delta} M^6 \leq X^{\frac{1}{52}}$ . We also decide to require that  $y^2 M^2 Y^2 X^{1 - \frac{1}{104}} \leq X^{1 - \delta/2}$  which gives  $y^2 M^2 Y^2 X^{\delta/2} \leq X^{\frac{1}{104}}$  and in particular  $y^4 X^{5\delta/2} M^4 \leq X^{\frac{1}{104}}$ . We posit (somewhat arbitrarily) that we will require  $y, M \leq X^{\delta}$ . In that case the previous two conditions are verified if  $X^{21\delta/2} \leq X^{\frac{1}{104}}$  and if  $X^{14\delta} \leq X^{\frac{1}{52}}$ . This leads to the choice of  $\delta = \frac{1}{1092}$ , implying the restriction  $y, M \leq X^{\frac{1}{1092}}$  and giving an error term of  $\ll X^{1 - \frac{1}{2148}}$ .

## 7. APPENDIX

**7.1. The structure of the space of half-integral weight forms.** Due to recent work of Baruch-Purkait [2], Shimura's correspondence between half-integral weight forms and integral weight forms is better understood. Let  $A_{k+1/2}^+$  denote the *conjugate plus space*, which is defined as  $W_4 S_{k+1/2}^+$ , where  $W_4 : S_{k+1/2} \rightarrow S_{k+1/2}$  is given by

$$(W_4 f)(z) = (-2iz)^{-k-1/2} f\left(\frac{-1}{4z}\right).$$

Baruch-Purkait proved that  $A_{k+1/2}^+ \cap S_{k+1/2}^+ = \{0\}$ . Letting  $S_{k+1/2}^-$  denote the orthogonal complement of  $A_{k+1/2}^+ \oplus S_{k+1/2}^+$  (w.r.t. the Petersson inner product) they proved that Niwa's isomorphism (see the main theorem in [23]) induces a Hecke algebra isomorphism between  $S_{k+1/2}^-$  and  $S_{2k}^{\text{new}}(2)$  (the space of weight  $2k$  newforms on  $\Gamma_0(2)$ ).

Also, let  $U_m$  be the operator, which acts on power series as follows

$$U_m \left( \sum_{n \geq 1} a_n q^n \right) = \sum_{n \geq 1} a_{mn} q^n.$$

Niwa proved that  $U_4 W_4$  is Hermitian on  $S_{k+1/2}$  and that  $(U_4 W_4 - \alpha_1)(U_4 W_4 - \alpha_2) = 0$  where  $\alpha_1 = 2^k \left(\frac{2}{2k+1}\right)$  and  $\alpha_2 = -\frac{1}{2} \cdot \alpha_1$ . The Kohnen plus space  $S_{k+1/2}^+$  is the subspace of cusp forms in  $S_{k+1/2}$  with  $U_4 W_4$ -eigenvalue equal to  $\alpha_1$  (Kohnen [14, Proposition 2]).

**7.2. Results.** Recall that  $\mathbb{N}^{\flat} = \{n \in \mathbb{N} : n = 8m \text{ and } \mu^2(2m) = 1\}$  and  $\mathbb{N}_g^{\flat}(X) = \{n \leq X : n \in \mathbb{N}^{\flat} \text{ and } c(n) \neq 0\}$ . Also, let  $S_{2k}(N)$  denote the space of weight  $2k$  cusp forms on  $\Gamma_0(N)$ .

**Theorem 4.** *Let  $k \geq 2$  be an integer and  $g$  be a Hecke cusp form of weight  $k + \frac{1}{2}$  on  $\Gamma_0(4)$ . Then it is possible to normalize  $g$  so that its  $n$ th Fourier coefficient is real for  $n \in \mathbb{N}^{\flat}$ .*

*In addition, suppose that  $c(n) \neq 0$  for some  $n \in \mathbb{N}^{\flat}$ . Then for every  $\varepsilon > 0$  the sequence  $\{c(n)\}_{n \in \mathbb{N}_g^{\flat}(X)}$  has  $\gg X^{1-\varepsilon}$  sign changes. Assuming GRH the sequence  $\{c(n)\}_{n \in \mathbb{N}_g^{\flat}(X)}$  has  $\gg X$  sign changes.*

**Remark 3.** *We will see that if  $g$  is of the form*

$$(7.1) \quad g(z) = aG(z) + b(W_4G)(z)$$

*with  $G \in S_{k+1/2}^+$ , and  $a, b \in \mathbb{C}$  with  $\frac{a}{b} = -\frac{1}{\alpha_2} \lambda_F(2)$ , where  $F \in S_{2k}(1)$  is the Shimura lift of  $G$ , then  $c(8n)\mu^2(2n) = 0$  for each  $n \in \mathbb{N}$ . Otherwise, for a Hecke cusp form  $g \in S_{k+1/2}$  not as above, the proof below and Lemma 3.1 imply  $c(8n)\mu^2(2n) \neq 0$  for some  $n$  so that the conclusion of Theorem 4 holds for  $g$ .*

*Moreover, for  $g$  as in (7.1) with  $b \neq 0$  the subsequent argument shows that  $c(2n)\mu^2(2n) = \frac{b}{\alpha_2} c_G(8n)\mu^2(2n)$ , where  $c_G(n)$  denotes the  $n$ th Fourier coefficient of  $G$ . Hence in this case we conclude that after suitable normalization, the sequence  $\{c(2n)\mu^2(2n)\}_{n \in \mathbb{N}}$  has  $\gg X$  sign changes as  $n$  ranges over  $[1, X]$  under GRH and  $\gg X^{1-\varepsilon}$  such sign changes unconditionally.*

*Proof.* Niwa (see the main theorem of [23]) proved that there is a Hecke algebra isomorphism between  $S_{k+1/2}$  and  $S_{2k}(2)$ . Denoting Niwa's isomorphism by  $\psi$ , write  $f = \psi(g) \in S_{2k}(2)$ . By Atkin-Lehner [1, Theorem 5] either  $f \in S_{2k}^{\text{new}}(2)$  or  $f$  is an oldform, i.e.  $f(z) = \alpha F(z) + \beta F(2z)$  with  $F \in S_{2k}(1)$ ,  $\alpha, \beta \in \mathbb{C}$ .

We first consider the case  $\psi(g) = f \in S_{2k}^{\text{new}}(2)$ . Here we can apply Shimura's explicit version of Waldspurger's Theorem, which for a fundamental discriminant of the form  $(-1)^k d = 8n > 0$  with  $2n$  square-free gives

$$|c(2n)|^2 = L\left(\frac{1}{2}, f \otimes \chi_d\right) \frac{(k-1)!}{2\pi^k} \cdot \frac{\langle g, g \rangle}{\langle f, f \rangle}$$

(see [28, Theorem 3B.4]). The Fourier coefficients can also be normalized so that they are real (this immediately follows from [28, Corollary 3B.5]). Also, in this case  $c(8n) = \lambda_2 \cdot c(2n)$  where  $\lambda_2 \in \{\pm 1\}$  and is independent of  $n$  (this follows from [27, Main Theorem], Niwa's isomorphism, and Atkin-Lehner [1, Theorem 3]; see the proof of Theorem 5 of [2]). Hence, our argument proceeds just as before and  $\{c(n)\}_{n \in \mathbb{N}_g^{\flat}(X)}$  has  $\gg X^{1-\varepsilon}$  sign changes unconditionally and  $\gg X$  under GRH.

Let us now suppose  $f \in S_{2k}(2)$  is an oldform. Then by Kohnen [14, Theorem 1] and Baruch-Purkait [2, Theorem 5, Corollary 6.1], there exist  $G_1, G_2 \in S_{k+1/2}^+$  and  $a, b \in \mathbb{C}$  such that  $g(z) = a \cdot G_1(z) + b \cdot (W_4 G_2)(z)$ . Since  $g$  is a Hecke cusp form it follows from multiplicity one for  $S_{k+1/2}^+$  (see [14, lemma, p. 256]) that  $G_1$  is a scalar multiple of  $G_2$ . Hence, we can write  $g(z) = a \cdot G(z) + b \cdot (W_4 G)(z)$  for some  $a, b \in \mathbb{C}$  and  $G \in S_{k+1/2}^+$ . If  $W_4 G = 0$  then  $g \in S_{k+1/2}^+$ , so we are done. Suppose  $W_4 G \neq 0$ , since  $S_{k+1/2}^+ \cap A_{k+1/2}^+ = \{0\}$ ,  $W_4 G \notin S_{k+1/2}^+$  so  $U_4 G = U_4 W_4(W_4 G) = \alpha_2 W_4 G$  since  $W_4$  is an involution. Hence,

$$g(z) = \sum_{n \geq 1} \left( a \cdot c_G(n) + \frac{b}{\alpha_2} \cdot c_G(4n) \right) e(nz).$$

For  $n \in \mathbb{N}^b$  write  $n = 8m$  where  $2m$  is square-free then by (3.3)

$$c_g(8m) = a \cdot c_G(8m) + \frac{b}{\alpha_2} \cdot c_G(2^2 \cdot 8m) = c_G(8m) \cdot \left( a + \frac{b}{\alpha_2} \cdot \lambda_F(2) \right),$$

where  $\lambda_F(\cdot)$  denotes the Hecke eigenvalues of the level 1 modular form which corresponds to  $G$  under  $\psi$ . By assumption  $a + \frac{b}{\alpha_2} \cdot \lambda_F(2) \neq 0$ , since otherwise  $c_g(8m)\mu^2(2m) = 0$  for every  $m \in \mathbb{N}$ . Moreover, since  $G \in S_{k+1/2}^+$  we know  $\{c_G(n)\}_{n \in \mathbb{N}_G^b(X)}$  has  $\gg X$  sign changes under GRH and  $\gg X^{1-\varepsilon}$  unconditionally, so the result follows.  $\square$

Additionally, it is possible to extend our result to level  $4N$  with  $N$  square-free and odd if we also restrict to fundamental discriminants that lie in a suitable progression. For each prime divisor of  $p|N$  we require that  $\chi_d(p) = w_p$  where  $w_p$  is the eigenvalue of the Atkin-Lehner operator  $W_p$ . So by the Chinese Remainder Theorem there exists  $\eta \pmod{N}$  such that for  $d \equiv \eta \pmod{N}$  we have  $\chi_d(p) = w_p$ . Let  $\mathbb{N}_N^b = \{n \in \mathbb{N} : n = 8m, \mu^2(2m) = 1, \text{ and } (-1)^k n \equiv \eta \pmod{N}\}$  and  $\mathbb{N}_{N,g}^b(X) = \{n \leq X : n \in \mathbb{N}_N^b \text{ and } c(n) \neq 0\}$ . Note that for  $n \in \mathbb{N}_N^b$  by construction  $(N, n) = 1$ .

Let  $S_{k+1/2}(4N)$  denote the space of weight  $k + 1/2$  cusp forms of level  $4N$ , with  $N$  odd and square-free. Also, let  $S_{k+1/2}^-(4N)$  be as defined by Baruch-Purkait (see [2, Section 6.3]), who showed that this space is isomorphic to  $S_{2k}^{\text{new}}(2N)$ . This complements Kohnen's result [15] that  $S_{k+1/2}^{\text{new}}(4N)$  is isomorphic to  $S_{2k}^{\text{new}}(N)$ .

We can also prove the following result.

**Theorem 5.** *Let  $k \geq 2$  and  $N$  be odd and square-free. Suppose  $g \in S_{k+1/2}^+(4N)$  or  $g \in S_{k+1/2}^-(4N)$ , is a Hecke cusp form. Then it is possible to normalize  $g$  so that its  $n$ th Fourier coefficient is real for  $n \in \mathbb{N}_N^b$ . Moreover, for every  $\varepsilon > 0$  the sequence  $\{c(n)\}_{n \in \mathbb{N}_{N,g}^b(X)}$  has  $\gg X^{1-\varepsilon}$  sign changes. Assuming GRH the sequence  $\{c(n)\}_{n \in \mathbb{N}_{N,g}^b(X)}$  has  $\gg X$  sign changes.*

*Proof.* We will only sketch the argument, since our main propositions need to be modified when  $N \neq 1$ . First, suppose  $g \in S_{k+1/2}^-(4N)$ . Then by [2, Theorem 8]  $g \in S_{k+1/2}^-(4N)$  and multiplicity one holds in  $S_{k+1/2}^-(4N)$  in the whole space  $S_{k+1/2}^-(4N)$ <sup>7</sup>. Hence for  $(-1)^k d \in \mathbb{N}_N^\flat$  we can apply Shimura's result [28, Theorem 3B.4] to get that

$$(7.2) \quad |c(2n)|^2 = 2^{\omega(N)} \frac{(k-1)! \langle g, g \rangle}{2\pi^k \langle f, f \rangle} L\left(\frac{1}{2}, f \otimes \chi_d\right)$$

where  $\omega(N) = \sum_{p|N} 1$ . Here we have used the condition  $d \equiv \eta \pmod{N}$  to estimate the Euler product present in the statement of the theorem along with [1, Theorem 3]. Using [28, Corollary 3B.5] it also follows that the Fourier coefficients of  $g$  can be normalized so that they are real. Also, in this case just as before we have  $c(8n) = \lambda_2 \cdot c(2n)$  with  $\lambda_2 \in \{\pm 1\}$ .

Next, suppose  $g(z) \in S_{k+1/2}^+(4N)$  and without loss of generality assume  $g \in S_{k+1/2}^{+, \text{new}}(4N)$ . Kohlen [15] proved  $S_{k+1/2}^{+, \text{new}}(4N)$  and  $S_{2^k}^{\text{new}}(N)$  are isomorphic as Hecke algebras. In this setting we can apply Proposition 4.2 of Kumar and Purkait [19] and it follows  $g \in S_{k+1/2}^{+, \text{new}}(4N)$  can be normalized so that it has real (and algebraic) Fourier coefficients. Moreover, for discriminants  $(-1)^k d \in \mathbb{N}_N^\flat$  Corollary 1 of [16] implies

$$(7.3) \quad |c(8n)|^2 = 2^{\omega(N)} \frac{\Gamma(k) \langle g, g \rangle}{\pi^k \langle f, f \rangle} L\left(\frac{1}{2}, f \otimes \chi_d\right).$$

Using the formulas (7.2) and (7.3) we have in each case above that

$$\sum_{n \in \mathbb{N}_{g,N}^\flat(X)} |c(8n)|^4 M\left((-1)^k 8n; \frac{1}{2}\right)^4 \ll \sum_{|d| \leq X}^b L\left(\frac{1}{2}, f \otimes \chi_d\right) M\left(d; \frac{1}{2}\right)^4.$$

To bound this mollified moment, the only modification needed in the proof of Proposition 4.1 is to change the definition of  $I_0$  so that  $I_0 = (c, X^{\theta_0}]$  where  $c$  is also sufficiently large in terms of  $N$ . Repeating the argument (with no further modifications) we arrive at

$$\sum_{|d| \leq X}^b L\left(\frac{1}{2}, f \otimes \chi_d\right) M\left(d; \frac{1}{2}\right)^4 \ll X.$$

We also need to prove

$$\sum_{n \in \mathbb{N}_{g,N}^\flat(X)} |c(8n)|^2 M\left((-1)^k 8n; \frac{1}{2}\right)^2 \asymp X.$$

---

<sup>7</sup>Multiplicity one also holds in the space  $S_{k+1/2}^+$ , but fails in the entire space  $S_{k+1/2}$  (consider  $g \in S_{k+1/2}^+$  and  $W_4 g$ ).



This follows in the same way as before once we have established an analog of Proposition 5.1 for  $f \in S_{2k}^{\text{new}}(M)$  with  $M = N$  or  $M = 2N$ , where we average over discriminants  $(-1)^k d \in \mathbb{N}_N^b$ . The necessary modifications for this computation have already been worked out in the paper of Radziwiłł and Soundararajan [26]. Finally, we need to establish the estimate

$$\sum_{X \leq x \leq 2X} \left| \sum_{\substack{x \leq 8n \leq x+y \\ n \text{ odd} \\ (-1)^k 8n \equiv \gamma \pmod{N}}} \mu^2(n) c(8n) M((-1)^k 8n) \right| \ll X \sqrt{y} + X^{1 - \frac{1}{2148} + \epsilon}.$$

To do this we first need to modify the proof of Lemma 6.2 in a straightforward way. From here we arrive at the analog of Proposition 6.1, for any  $g \in S_{k+1/2}(4N)$ . To establish the above bound we repeat the argument used in the proof of Proposition 1.3. The only modification necessary is that the range of  $\Delta$  in (6.17) will now be  $1 \leq \Delta \leq 16NY^4M^2$  to account for the progression  $(-1)^k 8n \equiv \eta \pmod{N}$ .

Combining the three estimates above we argue as in the proof of Theorem 1 thereby finishing the proof.  $\square$

## 8. ACKNOWLEDGEMENTS

We would like to thank Abhishek Saha for many helpful conversations regarding Kohnen's plus space and for pointing out Shimura's paper [28]. We would also like to express our gratitude to Dinakar Ramakrishnan who provided us with comments on an earlier draft of the appendix. Finally, we would like to thank the anonymous referee for a careful reading of the paper.

## REFERENCES

1. A. O. L. Atkin and J. Lehner, *Hecke operators on  $\Gamma_0(m)$* , Math. Ann. **185** (1970), 134–160. MR 0268123
2. E. M. Baruch and S. Purkait, *Newforms of half-integral weight: the minus space counterpart*, Canad. J. Math. **72** (2020), no. 2, 326–372. MR 4081695
3. J-H. Bruinier and W. Kohnen, *Sign changes of coefficients of half integral weight modular forms*, Modular forms on Schiermonnikoog, Cambridge Univ. Press, Cambridge, 2008, pp. 57–65. MR 2512356
4. V. Chandee, *Explicit upper bounds for L-functions on the critical line*, Proc. Amer. Math. Soc. **137** (2009), no. 12, 4049–4063. MR 2538566
5. G. Harcos, *An additive problem in the Fourier coefficients of cusp forms*, Math. Ann. **326** (2003), no. 2, 347–365. MR 1990914
6. A. Harper, *Sharp conditional bounds for moments of the Riemann zeta function*, Available on the arXiv [arXiv:1305.4618](https://arxiv.org/abs/1305.4618) (2013), 20 pages.
7. D. R. Heath-Brown, *A mean value estimate for real character sums*, Acta Arith. **72** (1995), no. 3, 235–275. MR 1347489
8. ———, *The average analytic rank of elliptic curves*, Duke Math. J. **122** (2004), no. 3, 591–623. MR 2057019

9. T. A. Hulse, E. M. Kiral, C. I. Kuan, and L-M. Lim, *The sign of Fourier coefficients of half-integral weight cusp forms*, Int. J. Number Theory **8** (2012), no. 3, 749–762. MR 2904928
10. H. Iwaniec, *Topics in classical automorphic forms*, Graduate Studies in Mathematics, vol. 17, American Mathematical Society, Providence, RI, 1997. MR 1474964
11. Y. J. Jiang, Y. K. Lau, G. S. Lü, E. Royer, and J. Wu, *Sign changes of Fourier coefficients of modular forms of half integral weight, 2*, Available on the arXiv [arXiv:1602.08922](https://arxiv.org/abs/1602.08922) (2016), 19 pages.
12. M. Jutila, *Transformations of exponential sums*, Proceedings of the Amalfi Conference on Analytic Number Theory (Maiori, 1989), Univ. Salerno, Salerno, 1992, pp. 263–270. MR 1220470
13. M. Knopp, W. Kohlen, and W. Pribitkin, *On the signs of Fourier coefficients of cusp forms*, Ramanujan J. **7** (2003), no. 1-3, 269–277, Rankin memorial issues. MR 2035806
14. W. Kohlen, *Modular forms of half-integral weight on  $\Gamma_0(4)$* , Math. Ann. **248** (1980), no. 3, 249–266. MR 575942
15. ———, *Newforms of half-integral weight*, J. Reine Angew. Math. **333** (1982), 32–72. MR 660784
16. ———, *Fourier coefficients of modular forms of half-integral weight*, Math. Ann. **271** (1985), no. 2, 237–268. MR 783554
17. W. Kohlen, Y.-K. Lau, and J. Wu, *Fourier coefficients of cusp forms of half-integral weight*, Mathematische Zeitschrift **273** (2013), no. 1, 29–41.
18. W. Kohlen and D. Zagier, *Values of  $L$ -series of modular forms at the center of the critical strip*, Invent. Math. **64** (1981), no. 2, 175–198. MR 629468
19. N. Kumar and S. Purkait, *A note on the Fourier coefficients of half-integral weight modular forms*, Arch. Math. (Basel) **102** (2014), no. 4, 369–378. MR 3196964
20. Y.-K. Lau, E. Royer, and J. Wu, *Sign of Fourier coefficients of modular forms of half-integral weight*, Mathematika **62** (2016), no. 3, 866–883. MR 3521358
21. Stephen Lester and Maksym Radziwiłł, *Quantum unique ergodicity for half-integral weight automorphic forms*, Duke Math. J. **169** (2020), no. 2, 279–351. MR 4057145
22. K. Matomäki and M. Radziwiłł, *Sign changes of Hecke eigenvalues*, Geom. Funct. Anal. **25** (2015), no. 6, 1937–1955. MR 3432161
23. S. Niwa, *On Shimura’s trace formula*, Nagoya Math. J. **66** (1977), 183–202. MR 0562506
24. K. Ono and C. Skinner, *Non-vanishing of quadratic twists of modular  $L$ -functions*, Invent. Math. **134** (1998), no. 3, 651–660. MR 1660945
25. A. Perelli and J. Pomykała, *Averages of twisted elliptic  $L$ -functions*, Acta Arith. **80** (1997), no. 2, 149–163. MR 1450922
26. M. Radziwiłł and K. Soundararajan, *Moments and distribution of central  $L$ -values of quadratic twists of elliptic curves*, Invent. Math. **202** (2015), no. 3, 1029–1068. MR 3425386
27. G. Shimura, *On modular forms of half integral weight*, Ann. of Math. (2) **97** (1973), 440–481. MR 0332663
28. ———, *On Hilbert modular forms of half-integral weight*, Duke Math. J. **55** (1987), no. 4, 765–838. MR 916119
29. K. Soundararajan, *Nonvanishing of quadratic Dirichlet  $L$ -functions at  $s = \frac{1}{2}$* , Ann. of Math. (2) **152** (2000), no. 2, 447–488. MR 1804529
30. ———, *Moments of the Riemann zeta function*, Ann. of Math. (2) **170** (2009), no. 2, 981–993. MR 2552116
31. K. Soundararajan and M. P. Young, *The second moment of quadratic twists of modular  $L$ -functions*, J. Eur. Math. Soc. (JEMS) **12** (2010), no. 5, 1097–1116. MR 2677611

SCHOOL OF MATHEMATICAL SCIENCES, QUEEN MARY UNIVERSITY OF LONDON, 327 MILE  
END ROAD, LONDON E1 4NS, UK

*Current address:* Department of Mathematics, King's College London, London WC2R 2LS, UK

*E-mail address:* `steve.lester@kcl.ac.uk`

DEPARTMENT OF MATHEMATICS, CALTECH, 1200 E CALIFORNIA BLVD, PASADENA, CA  
91125, USA

*E-mail address:* `maksym.radziwill@gmail.com`