



King's Research Portal

DOI: 10.1017/S0033291721000088

Document Version Peer reviewed version

Link to publication record in King's Research Portal

Citation for published version (APA):

NIMA Consortium, Cowen, P., Cavanagh, J., Harrison, N., Bullmore, E., & Pariante, C. (2021). The influence of comorbid depression and overweight status on peripheral inflammation and cortisol levels. *Psychological Medicine*. https://doi.org/10.1017/S0033291721000088

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1	The influence of comorbid depression and overweight status on peripheral
2	inflammation and cortisol levels
3	Authors: Anna P. McLaughlin ^{1,2} , Naghmeh Nikkheslat ¹ , Caitlin Hastings ¹ , Maria A. Nettis ^{1,2} ,
4	Melisa Kose ¹ , Courtney Worrell ¹ , Zuzanna Zajkowska ¹ , Nicole Mariani ¹ , Daniela Enache ¹ ,
5	Giulia Lombardo ¹ , Linda Pointon ³ , NIMA Consortium ⁴ , Philip Cowen ⁵ , Jonathan Cavanagh ⁶ ,
6	Neil Harrison ^{7,} Edward Bullmore ³ , Carmine M. Pariante ^{1,2} and Valeria Mondelli ^{1,2} .
7	
8	1 King's College London, Department of Psychological Medicine, Institute of Psychiatry,
9	Psychology and Neuroscience, London, UK.
10	2 National Institute for Health Research (NIHR) Maudsley Biomedical Research Centre,
11	South London and Maudsley NHS Foundation Trust, King's College London, London, UK
12	3 Department of Psychiatry, University of Cambridge, UK.
13	4 See supplementary material.
14	5 University Department of Psychiatry, Warneford Hospital, Oxford, UK.
15	6 Mental Health and Wellbeing, Sackler Institute, Neurology block, Queen Elizabeth
16	University hospital, Glasgow, UK.
17	7 Cardiff University Brain Research Imaging Centre (CUBRIC), Division of Psychological
18	Medicine and Clinical Sciences, Cardiff, UK.
19	Corresponding Author: Dr Valeria Mondelli ^a
20	Email: valeria.mondelli@kcl.ac.uk
21	Word Count: 3,713.

^a Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, The Maurice Wohl Clinical Neuroscience Institute, Stress, Psychiatry and Immunology Laboratory, Cutcombe Road, London, SE5 9RT.

- 1 Abstract
- 2

Background: Depression and overweight are each associated with abnormal immune system
activation. We sought to disentangle the extent to which depressive symptoms and overweight
status contributed to increased inflammation and abnormal cortisol levels.

6 **Methods:** Participants were recruited through the Wellcome Trust NIMA Consortium. The 7 sample of 216 participants consisted of 69 overweight patients with depression; 35 overweight 8 controls; 55 normal-weight patients with depression and 57 normal-weight controls. Peripheral 9 inflammation was measured as high-sensitivity C-Reactive Protein (hsCRP) in serum. Salivary 10 cortisol was collected at multiple points throughout the day to measure cortisol awakening 11 response and diurnal cortisol levels.

12 **Results:** Overweight patients with depression had significantly higher hsCRP compared with 13 overweight controls (p=0.042), normal-weight depressed patients (p<0.001) and normal-14 weight controls (p < 0.001), after controlling for age and gender. Multivariable logistic 15 regression showed that comorbid depression and overweight significantly increased the risk of 16 clinically elevated hsCRP levels ≥ 3 mg/L (OR: 2.44, 1.28-3.94). In a separate multivariable 17 logistic regression model, overweight status contributed most to the risk of having hsCRP 18 levels \geq 3mg/L (OR: 1.52, 0.7-2.41), while depression also contributed a significant risk (OR: 19 1.09, 0.27-2). There were no significant differences between groups in cortisol awakening 20 response and diurnal cortisol levels.

21 Conclusion: Comorbid depression and overweight status is associated with increased hsCRP,
22 and the coexistence of these conditions amplified the risk of clinically elevated hsCRP levels.
23 Overweight status contributed most to the risk of clinically elevated hsCRP levels, but
24 depression also contributed to a significant risk. We observed no differences in cortisol levels
25 between groups.

1 Introduction

2 Rates of depression and obesity have risen dramatically in recent years, with each 3 disorder separately posing a major health concern and economic cost to society (Abdelaal, le 4 Roux, & Docherty, 2017; James et al., 2018; Malhi & Mann, 2018). Depression and obesity 5 are highly comorbid disorders, and the coexistence of these conditions significantly increases 6 the risk for developing subsequent disorders, likely due to the chronic inflammatory state that 7 they induce (Ouakinin, Barreira, & Gois, 2018). High levels of inflammatory markers have 8 been widely described in depression (Baumeister, Russell, Pariante, & Mondelli, 2014; Enache, 9 Pariante, & Mondelli, 2019; Haapakoski, Mathieu, Ebmeier, Alenius, & Kivimäki, 2015; 10 Howren, Lamkin, & Suls, 2009; Valkanova, Ebmeier, & Allan, 2013), with a recent meta-11 analysis finding that approximately 25% of patients with depression presented with elevated 12 peripheral inflammation (Osimo, Baxter, Lewis, Jones, & Khandaker, 2019). However, weight 13 gain also increases peripheral inflammation, and studies of patients with depression have found 14 that 20% of patients to be obese and 50% to be overweight (Papakostas et al., 2005). Therefore, 15 it is unclear whether the increased inflammation observed in patients with depression is 16 predominantly due to the high rates of overweight and obesity in this group. As inflammation 17 may be both a causal mechanism and potential treatment target for depressive symptoms, it is 18 crucial to understand to what extent depression and weight gain each contribute to the increased 19 inflammation observed in patients with depression (Ambrosio et al., 2018).

20

Inflammation likely plays a key role in modifying the bidirectional relationship between depression and obesity, but not all patients with depression demonstrate increased inflammation. Patients with depression who have both increased inflammation and metabolic disturbances may have a distinct 'immuno-metabolic' form of depression (Milaneschi, Lamers, Berk, & Penninx, 2020). This type of depression is a significant risk factor for weight gain and

1 subsequent obesity (Hasler, 2004) and was specifically associated with increased peripheral 2 levels of inflammatory markers, such as C-reactive protein (CRP) (Lamers et al., 2013). 3 Increased CRP levels have clinical relevance for depression (Andrew H. Miller & Charles L. 4 Raison, 2016), as CRP \geq 3 mg/L was associated with an increased risk of developing depression 5 later in life (Au, Smith, Gariepy, & Schmitz, 2015), and a lack of response to antidepressant 6 medication (Chamberlain et al., 2019; Zhang et al., 2019). Conversely, reductions in body 7 mass index (BMI) following weight loss interventions are associated with an improvement in 8 depressive symptoms and lower CRP levels (Capuron et al., 2011; Perez-Cornago et al., 2014). 9 These data demonstrate the crucial role of inflammation, as measured by CRP, in the interface 10 between depression and weight gain (Ambrosio et al., 2018).

11

12 Depression with metabolic disturbances has also been associated with dysregulated hypothalamic-pituitary-adrenal (HPA) axis activity (Gold, 2015). The HPA axis produces the 13 14 anti-inflammatory stress hormone cortisol, which is involved in regulating mood, metabolism, 15 and circadian rhythms. Meta-analyses have suggested that patients with this subtype of depression have lower diurnal cortisol levels, meaning HPA axis hypoactivity (Lamers, 16 17 Vogelzangs et al. 2013, Juruena, Bocharova et al. 2018). In contrast, patients who exhibit the 18 more classic symptoms of depression, such as reduced appetite and insomnia, tend to 19 demonstrate HPA axis hyperactivity (Juruena, Bocharova, Agustini, & Young, 2018; Lamers 20 et al., 2013). This association was mechanistically supported by longitudinal studies finding 21 that lower cortisol levels in children are associated with higher BMI at age 18 (Ruttle et al., 22 2014), and that blunted cortisol responses lead to further weight gain and increased 23 inflammation (Champaneri et al., 2013). HPA axis hypoactivity may therefore be indicative of 24 a pathophysiological process, capable of influencing mood and weight gain, which is unique 25 to depressed patients with metabolic disturbances.

1 Research investigating inflammatory markers and their association with depressive 2 symptoms in individuals with obesity is typically complicated by a high incidence of other 3 inflammatory disorders, such as atherosclerosis, diabetes, cardiovascular disease, and 4 hypertension (Upadhyay, Farr, Perakakis, Ghaly, & Mantzoros, 2018). Once patients have 5 developed comorbid depression and obesity, they become more vulnerable to developing 6 subsequent inflammatory conditions. These conditions are associated with an increased risk of 7 recurrent depressive episodes (Nigatu, Bultmann, & Reijneveld, 2015), demonstrating the 8 vicious cycle to which these patients may be liable. Therefore, it is vital to confirm the 9 association between depressive symptoms and inflammation in a sample free of comorbid 10 disorders, as the presence of other disorders may confound this relationship. It is also important 11 to identify which subgroup of patients with depression are most vulnerable so that they can be 12 prioritised for clinical interventions targeted at reducing their risk of developing further 13 comorbidities.

14

15 Therefore, we investigated inflammatory mechanisms, in the form of pro-inflammatory 16 hsCRP and anti-inflammatory cortisol, to determine how these biomarkers are associated with 17 depression and overweight status. We also investigated whether the presence of comorbid 18 depression and overweight increased inflammation to a clinically relevant level. Crucially, our 19 study used a sample of patients with depression who were free of other comorbid disorders, 20 with a comparison group of overweight controls, who were otherwise healthy.

21

This is the first study to explore differences in hsCRP levels and HPA axis activity in overweight individuals with and without depression, as well as normal-weight individuals with and without depression. Our cross-sectional study aimed to investigate: 1) whether overweight patients with depression had increased inflammation relative to all other groups; 2) what extent overweight status and depression status contributed to clinically elevated hsCRP levels
 ≥3mg/L; 3) whether overweight patients with depression had lower diurnal cortisol levels
 relative to all other groups.

4

5 <u>Methods</u>

6 Clinical data, blood samples, and saliva samples were collected in a cross-sectional, 7 observational design, as part of a multi-centre study investigating immune Biomarkers in 8 Depression (BIODEP), through the Wellcome Trust Consortium for Neuroimmunology of 9 Mood Disorders and Alzheimer's disease (NIMA). The study was approved by the Research 10 Ethics Committee (National Research Ethics Service East of England, Cambridge Central, UK; 11 approval number: 15/EE/0092). All procedures contributing to this work comply with the 12 ethical standards of the relevant national and institutional committees on human 13 experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All participants 14 provided written informed consent and underwent eligibility screening prior to taking part in 15 any study procedures.

16

17 Participants

18 Participants were aged 25-50 years inclusive and recruited at five clinical research study 19 centres (King's College London, Oxford, Cambridge, Brighton and Glasgow) from primary 20 and secondary NHS health services and the general population. Participants were excluded if 21 they 1) were pregnant or breastfeeding; 2) were underweight (defined as BMI <18); 3) were 22 taking medication likely to compromise the interpretation of immunological data (including, 23 but not limited to, statins, corticosteroids, antihistamines and anti-inflammatory medications); 24 4) met criteria for alcohol abuse, drug abuse or dependence in the last six months; 5) had 25 participated in a clinical trial of an investigational drug within the last 12 months; 6) had

1 lifetime history of any serious medical disorder likely to compromise the interpretation of 2 immunological data; 7) had a recent infection or illness likely to compromise the interpretation 3 of immunological data. Patients with depression were considered eligible if they met the 4 criteria for Major Depressive Disorder (MDD). Patients with depression were excluded if they 5 had a lifetime history of bipolar disorder or non-affective psychosis. Healthy controls were 6 considered eligible if they had no personal history of MDD or treatment with a monoaminergic 7 antidepressant for depressive symptoms or any other indication, as well as no current or 8 lifetime history of any major psychiatric disorder as defined by Diagnostic and Statistical 9 Manual Version 5 (DSM-5).

10

11 Sample groups

12 The current sample was selected from a larger sample of participants taking part in the 13 BIODEP study, where participants were recruited based on their clinical response to 14 antidepressants as described by our group previously (Chamberlain et al., 2019; Nikkheslat et 15 al., 2019). For the purpose of the current study, participants from the BIODEP pool were 16 grouped according to their BMI status (overweight participants with BMI \geq 25) and then 17 categorised these participants according to the presence of depression (MDD patients versus 18 healthy controls).

19

20 Demographic and clinical measures

Age, gender, smoking status, and medical history were documented by semi-structured clinical interviews. Height and weight were measured for calculation of BMI (kg/m²) and overweight status was defined as BMI \geq 25. History and diagnosis of MDD and other psychiatric disorders were assessed with the Structured Clinical Interview for DSM-5 (Kübler,

2013). The severity of depressive symptoms was assessed using 17-item Hamilton Rating Scale
 for Depression (HAM-D) (Hamilton, 1960).

- 3

4 High sensitivity C-Reactive Protein (hsCRP)

5 Peripheral inflammation was measured as serum levels of high-sensitivity CRP 6 (hsCRP), which has been demonstrated as a reliable biomarker of inflammation associated with 7 MDD. Participants fasted for 8 hours and abstained from strenuous exercise for 72 hours prior 8 to their blood draw, which was carried out between 8:00 and 10:00. Blood samples were 9 collected in clotting tubes, allowed to coagulate at room temperature for 30-60 minutes, then 10 centrifuged at 1600 Relative Centrifugal Force for 15 minutes. The serum samples were 11 separated and transported to a central laboratory (Q2 solutions) where they were analysed on 12 the day of collection. Samples were exposed to anti-CRP-antibodies on latex particles, and the 13 increase in light absorption due to complex formation was used to quantify hsCRP levels, using 14 Turbidimetry on Beckman Coulter AU analysers. Inter and intra-assay coefficient of variations 15 were <10%. The measure of hsCRP was calculated from one blood draw taken at the time of 16 clinical assessment, for each participant. In line with previous studies, clinically elevated 17 hsCRP levels were defined as hsCRP≥3mg/L (A. H. Miller & C. L. Raison, 2016; Pearson et 18 al., 2003).

19

20 Salivary cortisol

Participants were issued with the materials and instructions for collecting the saliva samples at the time of their clinical interview. Our previously published literature describes the collection procedure for saliva samples in more detail (Nikkheslat et al., 2019). Using salivette sampling devices (Sarstedt, Leicester, UK), samples were self-collected by participants at home at six time points throughout the day; at awakening, 15, 30 and 60 minutes after

1 awakening, 12:00 and at 20:00. Individuals who described problems during sample collection 2 in the self-recorded questionnaire, or who did not respect the time-intervals required, were 3 removed from the analysis. The current study included only participants who completed saliva 4 sample collections accurately and who provided an adequate amount of saliva for cortisol 5 measurement. Salivary cortisol levels were measured using a commercially available high-6 sensitivity salivary cortisol enzyme immunoassay kit from Salimetrics. SoftMax Pro 4.8 7 software was used to calculate the cortisol values, following a 4-parameter fit. The analytical 8 sensitivity was set to 0.19 nmol/l. Inter and intra-assay coefficient of variations ranged from 8-9 10% and 6-10%, respectively. To investigate the activity and responsiveness of the HPA axis, 10 we first compared the mean values at the various time points of salivary cortisol collection; 11 and secondly, we calculated the area under the curve with respect to the increase (AUCi) for 12 the cortisol awakening response using the four time points of 0, 15, 30, and 60 minutes after 13 awakening; and the area under the curve with respect to the ground (AUCg) for the diurnal 14 cortisol using the three points: awakening, noon and 20:00. AUCg indicated the total amount 15 of cortisol produced and overall HPA axis activity during the day. AUCi indicated the variation 16 (either positive or negative) in cortisol concentration and thus signified the HPA axis reactivity 17 and response to the stress of awakening. The formulas for the calculations of the AUC were 18 derived from the trapezoidal formula introduced by Pruessner et al. (Pruessner, Kirschbaum, 19 Meinlschmid, & Hellhammer, 2003).

20

21 Statistical analysis

All statistical analyses were performed using R software, version 4.0.0. Data were evaluated for normality and logarithmic transformed if required. To assess group differences in demographic variables, Chi-squared or Kruskal-Wallis tests were used, as appropriate. To address the first hypothesis, group differences in logarithmic-transformed hsCRP levels were

1 assessed using two-way analysis of covariance (ANCOVA), with group and gender as factors 2 and age as a covariate. Pairwise comparisons were carried out using the False Discovery Rate 3 (FDR) correction, with the alpha set at 0.05 (5%). Effect size differences in hsCRP groups 4 relative to the normal-weight controls were calculated using Cohen's d. To address the second 5 hypothesis, two models of multivariable logistic regression were used. The first assessed how 6 much risk comorbid depression and overweight contributed to the hsCRP levels ≥3mg/L 7 relative to other participant groups, with age and gender as covariates. The second assessed 8 how much risk depression and overweight status (BMI ≥25) separately contributed to hsCRP 9 levels ≥ 3 mg/L, with age and gender as covariates. To address the third hypothesis, group 10 differences in cortisol awakening response (AUCi) and logarithmic-transformed diurnal 11 cortisol (AUCg) were assessed using two-way analysis of covariance (ANCOVA), with 12 adjustment for age and gender. The threshold for statistical significance of all tests was defined 13 as two-tailed $p \leq 0.05$.

14

15 <u>Results</u>

16 Sample demographics

17 The demographic and clinical characteristics of the sample are presented in Table 1. 18 Age and gender were significantly different between groups, while ethnicity and smoking 19 status were similar across groups, in both samples. Although the proportion of smokers were 20 not significantly different between groups, we repeated all analyses excluding smokers to 21 determine if smoking status significantly influenced hsCRP and cortisol levels. As the direction 22 of the results did not change for any of the relevant tests after excluding smokers, we have 23 reported results for the full sample, with the data for non-smokers presented in the supplement. 24 A similar number of overweight patients with depression and normal-weight patients with 25 depression were currently taking antidepressant medication (47 patients vs. 39 patients, 1 $x^2=0.019$, p=0.889) and a two-way ANCOVA showed that both groups had a similar age of 2 onset of depressive symptoms (F(1, 109)=1.617, p=0.206, with adjustment for age and gender.

3

4 High sensitivity C-Reactive Protein (hsCRP)

5 A two-way ANCOVA showed a significant effect of group on hsCRP, with adjustment 6 for age and gender F(3, 211)=16.43, p<0.001. As there was no significant interaction between 7 group and gender on hsCRP, F(3, 211)=1.4, p=0.245, pairwise comparisons with the FDR 8 correction were carried out, comparing the main effects of group, but not gender. Overweight patients with depression had significantly higher hsCRP compared with overweight controls 9 10 (p=0.042), normal weight depressed patients (p<0.001) and normal weight controls (p<0.001); 11 see Figure 1). Within the group of overweight patients with depression there were six 12 individuals with extreme BMI values \geq 40 indicating morbid obesity. When excluding these 13 individuals from analysis, a two-way ANCOVA showed a significant effect of group on 14 hsCRP, with adjustment for age and gender F(3, 201)=15.11, p<0.001. As there was no 15 significant interaction between group and gender on hsCRP, F(3, 207)=1.36, p=0.257, pairwise 16 comparisons with the FDR correction were carried out, comparing the main effects of group, 17 but not gender. Overweight patients with depression had higher hsCRP compared with 18 overweight controls at trend-level (*p*=0.068), and significantly higher hsCRP compared with 19 normal weight depressed patients (p < 0.001) and normal weight controls (p < 0.001).

20

Multivariable logistic regression analysis was performed to investigate whether group
status contributed to the risk of having hsCRP ≥3mg/L, after controlling for age and gender.
Overweight patients with depression were at a significantly increased risk of having hsCRP
≥3mg/L (OR: 2.44, 95% confidence interval (CI)=1.28-3.94, *p*<0.001), overweight controls
did not have a significantly increased risk of hsCRP ≥3mg/L (OR: 1.51, 95% CI=-0.33-2.87,

p=0.13), normal-weight patients with depression did not have a significantly increased risk of
hsCRP ≥3mg/L (OR: 1.08, 95% CI=-0.6-2.4, p=0.28), while normal-weight controls were at a
significantly decreased risk of hsCRP ≥3mg/L (OR: -4.39, 95% CI=-7.04 to -1.97, p<0.001).
Age and gender were not significantly associated with hsCRP ≥3mg/L (OR:0.01, 95%
confidence interval (CI) = -0.04-0.06, p=0.621, OR:0.61, 95% CI=-0.19-1.47, p=0.145,
respectively).

7

Multivariable logistic regression analysis was also performed to investigate the
association between depression and overweight status on hsCRP ≥3mg/L, after controlling for
age and gender. Depression status was significantly associated with hsCRP ≥3mg/L (OR: 1.09,
95% CI=0.27-2.01, p=0.013) and overweight status was significantly associated with hsCRP
≥3mg/L (OR: 1.52, 95% CI=0.7-2.41, p<0.001). Age and gender were not significantly
associated with hsCRP ≥3mg/L (OR:0.62, 95% confidence interval (CI)=-0.04-0.06, p=0.647,
OR:0.01, 95% confidence interval (CI)=-0.19-1.47, p=0.137 respectively).

15

16 Cortisol

17 A two-way ANCOVA showed no significant differences between groups in CARi, after 18 adjustment for age and gender, F(3, 162)=0.473, p=0.701. Similarly, a two-way ANCOVA 19 showed no significant differences between groups in logarithmic-transformed AUCg, after 20 adjustment for age and gender, F(3, 167)=0.719, p=0.542.

21 Discussion

To our knowledge, this is the first study to observe significantly higher hsCRP levels in overweight patients with depression relative to overweight controls, as well as depressed and non-depressed normal-weight participants. Of clinical relevance, overweight patients with depression had the highest risk for hsCRP levels \geq 3mg/L, indicating clinically elevated peripheral inflammation. Our results demonstrate that coexisting depression and overweight
 may exacerbate peripheral inflammation and highlights the urgent need to optimise treatment
 strategies for these patients. We observed no differences in HPA axis activity between groups.

4

5 Our findings regarding elevated hsCRP levels in overweight patients with depression 6 are consistent with previous studies in normal-weight patients with depression and obese 7 patients with depression (Haapakoski et al., 2015; Rethorst, Bernstein, & Trivedi, 2014). Of 8 note, our study design separating overweight and normal-weight participants, from those with 9 and without depression, was able to demonstrate the separate and comorbid association 10 between these disorders and hsCRP levels ≥ 3 mg/L. Peripheral inflammation is believed to 11 initiate and perpetuate a range of sickness-related behaviour, such as fatigue, weakness, 12 malaise, sleep, and disturbed appetite (Dantzer, 2006), which may further maintain the vicious 13 cycle between weight gain and depressive symptoms. The increased risk for overweight 14 patients with depression to have clinically elevated hsCRP \geq 3mg/L is concerning, as CRP is a 15 predictor of all-cause mortality (Li et al., 2017). Increased mortality and risk of developing 16 subsequent comorbid disorders in patients with depression places a substantial burden on 17 healthcare infrastructure and the economy (Tremmel, Gerdtham, Nilsson, & Saha, 2017), in 18 addition to reduced quality of life for these individuals (Nigatu, Reijneveld, de Jonge, van 19 Rossum, & Bultmann, 2016). Answering the question of whether depression or weight gain 20 contributed more to inflammation was identified as a key objective by a recent meta-analysis 21 (Ambrosio et al., 2018), so that research for this field can progress into therapeutic 22 interventions designed to break the cyclic pattern between these disorders.

23

Testing CRP levels in overweight patients with depression could be useful for recommending adjunct anti-inflammatory treatments or predicting their likelihood of

1 developing additional comorbidities in the future. Indeed, several clinical trials have observed 2 that higher BMI at baseline predicted a poorer response to antidepressant treatments (Jha et al., 3 2018; Uher et al., 2009), while we have previously shown that immune-metabolic status was 4 associated with a poorer response to antipsychotic medication in individuals with psychosis 5 (Nettis et al., 2019). Anti-inflammatory drugs improve the therapeutic action of antidepressants 6 (Bai et al., 2020; Haroon, Raison, & Miller, 2012), which could be particularly useful in the 7 context of chronically inflamed overweight patients with depression. Overweight patients with 8 hsCRP \geq 3mg/L may prove ideal criterion to test this, as stratifying patients to specific 9 treatments according to inflammatory markers seems to improve treatment response (Cuthbert 10 & Insel, 2013; Kohler, Krogh, Mors, & Benros, 2016).

11

12 The lack of significant differences between groups in HPA axis activity was surprising, 13 as HPA axis dysregulation has been implicated in both obesity and depression (Incollingo 14 Rodriguez et al., 2015), however, the nature of the dysregulation is not always consistent. A 15 systematic review comparing atypical depression with the melancholic (typical) presentation, 16 found that atypical patients demonstrated lower cortisol relative to melancholic patients, but 17 not so low that they were significantly different from controls (Juruena et al., 2018). In contrast, our study observed no differences in cortisol levels, despite hsCRP levels varying dramatically 18 19 between groups. Dysregulation of the HPA axis may change over illness course, with 20 hypotheses suggesting that at first cortisol levels are highly responsive to peripheral 21 inflammation, but over time this responsivity declines (Gold, 2015; Perrin, Horowitz, Roelofs, 22 Zunszain, & Pariante, 2019). Another explanation is that specific symptoms are responsible for 23 changes in HPA axis activity (Iob, Kirschbaum, & Steptoe, 2020), and that antidepressant use additionally modifies HPA axis activity, as our group recently observed (Nikkheslat et al., 24

2019). Ultimately, our sample of patients may have been too heterogeneous in terms of
 depressive symptoms, antidepressant use and illness course to identify group differences.

3

4 Our study design was substantially strengthened by our strict exclusion criteria, encompassing 5 recent illnesses and comorbid disorders, such as atherosclerosis, diabetes and cardiovascular 6 disease, which often confounds research in patients with obesity. We also excluded patients 7 taking medications that could influence immunological data, such as statins, corticosteroids, 8 antihistamines, anti-inflammatory medications Additionally, the direction of our results was 9 maintained after we removed smokers and individuals with extreme BMI values ≥ 40 , 10 suggesting that our findings are robust. Our study was limited by a lack of additional body 11 composition and clinical measures, as these would have enabled us to define an immune-12 metabolic subgroup of patients within our sample. However, using overweight status as defined 13 by BMI has high translational value for clinical settings. Our sample consisted of 14 predominantly white participants, which limits the generalisability of the results. Our study 15 could have been improved by the measurement of lifestyle factors, particularly physical 16 activity, sleep and diet, as these factors may confound the relationship between depression and 17 weight gain (Schmidt et al., 2015). Future studies should be designed to include more diverse 18 participant samples and adjust for lifestyle factors as covariates as they may explain some of 19 the heterogeneity in the association between depressive symptoms and inflammation.

20

21 Chronic inflammation is likely a key modifier in the vicious cycle between depressive 22 symptoms and weight gain. If the cycle can be reversed, meaning that a reduction in depressive 23 symptoms is accompanied by healthy behaviour such as exercise and healthy eating, which 24 indirectly reduces inflammation and leads to excess weight loss, this would result in a better 25 quality of life for patients and a vast reduction in global healthcare burdens and economic cost

I (Jantaratnotai, Mosikanon, Lee, & McIntyre, 2017; Nigatu et al., 2016; Tremmel et al., 2017).
Ultimately, our data suggest that depression and overweight are separately associated with
increased inflammation, but this association is amplified when these conditions coexist.
Patients with depression with BMI ≥25 would likely benefit from further research to determine
if a higher level of clinical support or anti-inflammatory treatment approaches could potentially
improve depressive symptoms and reduce the risk of developing further comorbidities in this
subgroup of patients.

8

9 Acknowledgments

10 This research has been supported by the National Institute for Health Research (NIHR) 11 Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation 12 Trust and King's College London. The views expressed are those of the authors and not 13 necessarily those of the NHS, the NIHR or the Department of Health. The authors would like 14 to thank the research team at Brighton, Cambridge, Glasgow, King's College London and 15 Oxford; and all the study participants without whom this work would not have been possible 16 and completed. The members of NIMA Consortium are thanked and acknowledged (see 17 Annex). VM is supported by MQ: Transforming Mental Health (Grant: MQBF1) and by the 18 Medical Research Foundation (Grant: MRF-160-0005). This work was supported by the 19 NIHR Cambridge Biomedical Research Centre (Mental Health).

20

21 Financial support

This work was funded by a grant from the Wellcome Trust (Grant number: 104025/Z/14/Z) to the NIMA Consortium, which is also funded by Janssen, GlaxoSmithKline, Lundbeck and Pfizer. Recruitment of patients was supported by the National Institute of Health Research (NIHR) Clinical Research Network: Kent, Surrey and Sussex & Eastern. The work

is also supported by the Medical Research Council (UK) MR/J002739/1 and the Commission
of European Communities Seventh Framework Programme (Collaborative Project Grant
Agreement no. 22963, Mood Inflame); and part funded by the NIHR/Wellcome Trust, King's
Clinical Research Facility and the NIHR Biomedical Research Centre [and Dementia Unit] at
South London and Maudsley NHS Foundation Trust and King's College London.

6

7 Conflicts of interest

8 SRC consults for Cambridge Cognition and Shire; and his input in this project was 9 funded by a Wellcome Trust Clinical Fellowship (110049/Z/15/Z). ETB was employed half-10 time by GlaxoSmithKline, and held stock in GSK, until May 2019; he has since worked full 11 time for University of Cambridge; he was holding stock (not currently) in GSK. ETB is an 12 NIHR Senior Investigator. NAH consults for GSK. PdB, DJ and WCD are employees of 13 Janssen Research & Development, LLC., of Johnson & Johnson, and hold stock in Johnson & 14 Johnson. The other authors report no financial disclosures or potential conflicts of interest.

1 <u>References</u>

2	Abdelaal, M., le Roux, C. W., & Docherty, N. G. (2017). Morbidity and mortality associated
3	with obesity. The Annals of Translational Medicine, 5(7), 161.
4	doi:10.21037/atm.2017.03.107
5	Ambrosio, G., Kaufmann, F. N., Manosso, L., Platt, N., Ghisleni, G., Rodrigues, A. L. S.,
6	Kaster, M. P. (2018). Depression and peripheral inflammatory profile of patients with
7	obesity. Psychoneuroendocrinology, 91, 132-141.
8	doi:10.1016/j.psyneuen.2018.03.005
9	Au, B., Smith, K. J., Gariepy, G., & Schmitz, N. (2015). The longitudinal associations
10	between C-reactive protein and depressive symptoms: evidence from the English
11	Longitudinal Study of Ageing (ELSA). International Journal of Geriatric Psychiatry,
12	30(9), 976-984. doi:10.1002/gps.4250
13	Bai, S., Guo, W., Feng, Y., Deng, H., Li, G., Nie, H., Tang, Z. (2020). Efficacy and safety
14	of anti-inflammatory agents for the treatment of major depressive disorder: a
15	systematic review and meta-analysis of randomised controlled trials. Journal of
16	Neurology, Neurosurgery, and Psychiatry, 91(1), 21-32. doi:10.1136/jnnp-2019-
17	320912
18	Baumeister, D., Russell, A., Pariante, C. M., & Mondelli, V. (2014). Inflammatory biomarker
19	profiles of mental disorders and their relation to clinical, social and lifestyle factors.
20	Social Psychiatry and Psychiatric Epidemiology, 49(6), 841-849.
21	doi:10.1007/s00127-014-0887-z
22	Capuron, L., Poitou, C., Machaux-Tholliez, D., Frochot, V., Bouillot, J. L., Basdevant, A.,
23	. Clement, K. (2011). Relationship between adiposity, emotional status and eating
24	behaviour in obese women: role of inflammation. Psychological Medicine, 41(7),
25	1517-1528. doi:10.1017/S0033291710001984

1	Chamberlain, S. R., Cavanagh, J., de Boer, P., Mondelli, V., Jones, D. N. C., Drevets, W. C.,						
2	Bullmore, E. T. (2019). Treatment-resistant depression and peripheral C-reactive						
3	protein. British Journal of Psychiatry, 214(1), 11-19. doi:10.1192/bjp.2018.66						
4	Champaneri, S., Xu, X., Carnethon, M. R., Bertoni, A. G., Seeman, T., DeSantis, A. S.,						
5	Golden, S. H. (2013). Diurnal salivary cortisol is associated with body mass index and						
6	waist circumference: the Multiethnic Study of Atherosclerosis. Obesity (Silver						
7	Spring), 21(1), E56-63. doi:10.1002/oby.20047						
8	Cuthbert, B. N., & Insel, T. R. (2013). Toward the future of psychiatric diagnosis: the seven						
9	pillars of RDoC. BMC Medicine, 11, 126. doi:10.1186/1741-7015-11-126						
10	Dantzer, R. (2006). Cytokine, sickness behavior, and depression. Neurologic Clinics, 24(3),						
11	441-460. doi:10.1016/j.ncl.2006.03.003						
12	Enache, D., Pariante, C. M., & Mondelli, V. (2019). Markers of central inflammation in						
13	major depressive disorder: A systematic review and meta-analysis of studies						
14	examining cerebrospinal fluid, positron emission tomography and post-mortem brain						
15	tissue. Brain, Behavior, and Immunity, 81, 24-40. doi:10.1016/j.bbi.2019.06.015						
16	Gold, P. W. (2015). The organization of the stress system and its dysregulation in depressive						
17	illness. Molecular Psychiatry, 20(1), 32-47. doi:10.1038/mp.2014.163						
18	Haapakoski, R., Mathieu, J., Ebmeier, K. P., Alenius, H., & Kivimäki, M. (2015).						
19	Cumulative meta-analysis of interleukins 6 and 1 β , tumour necrosis factor α and C-						
20	reactive protein in patients with major depressive disorder. Brain, Behavior, and						
21	Immunity, 49, 206-215. doi:10.1016/j.bbi.2015.06.001						
22	Hamilton, M. (1960). A rating scale for depression. Journal of Neurology, Neurosurgery, and						
23	Psychiatry, 23, 56-62. doi:10.1136/jnnp.23.1.56						

1	Haroon, E., Raison, C. L., & Miller, A. H. (2012). Psychoneuroimmunology meets
2	neuropsychopharmacology: translational implications of the impact of inflammation
3	on behavior. Neuropsychopharmacology, 37(1), 137-162. doi:10.1038/npp.2011.205
4	Hasler, G. P., D S; Gamma, A; Milos, G; Ajdacic, V; Eich, D; Rössler, W; Angst, J. (2004).
5	The associations between psychopathology and being overweight: a 20-year
6	prospective study. Psychological Medicine, p. 1047-1057.
7	Howren, M. B., Lamkin, D. M., & Suls, J. (2009). Associations of depression with C-reactive
8	protein, IL-1, and IL-6: a meta-analysis. Psychosomatic Medicine, 71(2), 171-186.
9	doi:10.1097/PSY.0b013e3181907c1b
10	Incollingo Rodriguez, A. C., Epel, E. S., White, M. L., Standen, E. C., Seckl, J. R., &
11	Tomiyama, A. J. (2015). Hypothalamic-pituitary-adrenal axis dysregulation and
12	cortisol activity in obesity: A systematic review. Psychoneuroendocrinology, 62, 301-
13	318. doi:10.1016/j.psyneuen.2015.08.014
14	Iob, E., Kirschbaum, C., & Steptoe, A. (2020). Persistent depressive symptoms, HPA-axis
15	hyperactivity, and inflammation: the role of cognitive-affective and somatic
16	symptoms. Molecular Psychiatry, 25(5), 1130-1140. doi:10.1038/s41380-019-0501-6
17	James, S. L., Abate, D., Abate, K. H., Abay, S. M., Abbafati, C., Abbasi, N., Murray, C.
18	J. L. (2018). Global, regional, and national incidence, prevalence, and years lived with
19	disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a
20	systematic analysis for the Global Burden of Disease Study 2017. The Lancet,
21	392(10159), 1789-1858. doi:10.1016/s0140-6736(18)32279-7
22	Jantaratnotai, N., Mosikanon, K., Lee, Y., & McIntyre, R. S. (2017). The interface of
23	depression and obesity. Obesity Research & Clinical Practice, 11(1), 1-10.
24	doi:10.1016/j.orcp.2016.07.003

1	Jha, M. K., Wakhlu, S., Dronamraju, N., Minhajuddin, A., Greer, T. L., & Trivedi, M. H.
2	(2018). Validating pre-treatment body mass index as moderator of antidepressant
3	treatment outcomes: Findings from CO-MED trial. Journal of Affective Disorders,
4	234, 34-37. doi:10.1016/j.jad.2018.02.089
5	Juruena, M. F., Bocharova, M., Agustini, B., & Young, A. H. (2018). Atypical depression
6	and non-atypical depression: Is HPA axis function a biomarker? A systematic review.
7	Journal of Affective Disorders, 233, 45-67. doi:10.1016/j.jad.2017.09.052
8	Kohler, O., Krogh, J., Mors, O., & Benros, M. E. (2016). Inflammation in Depression and the
9	Potential for Anti-Inflammatory Treatment. Current Neuropharmacology, 14(7), 732-
10	742. doi:10.2174/1570159x14666151208113700
11	Kübler, U. (2013). Structured Clinical Interview for DSM-IV (SCID). In M. D. Gellman & J.
12	R. Turner (Eds.), Encyclopedia of Behavioral Medicine (pp. 1919-1920). New York,
13	NY: Springer New York.
14	Lamers, F., Vogelzangs, N., Merikangas, K. R., de Jonge, P., Beekman, A. T., & Penninx, B.
15	W. (2013). Evidence for a differential role of HPA-axis function, inflammation and
16	metabolic syndrome in melancholic versus atypical depression. Molecular Psychiatry,
17	18(6), 692-699. doi:10.1038/mp.2012.144
18	Li, Y., Zhong, X., Cheng, G., Zhao, C., Zhang, L., Hong, Y., Wang, Z. (2017). Hs-CRP
19	and all-cause, cardiovascular, and cancer mortality risk: A meta-analysis.
20	Atherosclerosis, 259, 75-82. doi:10.1016/j.atherosclerosis.2017.02.003
21	Malhi, G. S., & Mann, J. J. (2018). Depression. The Lancet, 392(10161), 2299-2312.
22	doi:10.1016/s0140-6736(18)31948-2
23	Milaneschi, Y., Lamers, F., Berk, M., & Penninx, B. (2020). Depression Heterogeneity and
24	Its Biological Underpinnings: Toward Immunometabolic Depression. Biological
25	Psychiatry. doi:10.1016/j.biopsych.2020.01.014

1	Miller, A. H., & Raison, C. L. (2016). The role of inflammation in depression: from
2	evolutionary imperative to modern treatment target. Nature Reviews Immunology,
3	16(1), 22-34. doi:10.1038/nri.2015.5
4	Nettis, M. A., Pergola, G., Kolliakou, A., O'Connor, J., Bonaccorso, S., David, A.,
5	Mondelli, V. (2019). Metabolic-inflammatory status as predictor of clinical outcome
6	at 1-year follow-up in patients with first episode psychosis.
7	Psychoneuroendocrinology, 99, 145-153. doi:10.1016/j.psyneuen.2018.09.005
8	Nigatu, Y. T., Bultmann, U., & Reijneveld, S. A. (2015). The prospective association
9	between obesity and major depression in the general population: does single or
10	recurrent episode matter? BMC Public Health, 15, 350. doi:10.1186/s12889-015-
11	1682-9
12	Nigatu, Y. T., Reijneveld, S. A., de Jonge, P., van Rossum, E., & Bultmann, U. (2016). The
13	Combined Effects of Obesity, Abdominal Obesity and Major Depression/Anxiety on
14	Health-Related Quality of Life: the LifeLines Cohort Study. PloS One, 11(2),
15	e0148871. doi:10.1371/journal.pone.0148871
16	Nikkheslat, N., McLaughlin, A. P., Hastings, C., Zajkowska, Z., Nettis, M. A., Mariani, N., .
17	. Mondelli, V. (2019). Childhood trauma, HPA axis activity and antidepressant
18	response in patients with depression. Brain, Behavior, and Immunity.
19	doi:10.1016/j.bbi.2019.11.024
20	Osimo, E. F., Baxter, L. J., Lewis, G., Jones, P. B., & Khandaker, G. M. (2019). Prevalence
21	of low-grade inflammation in depression: a systematic review and meta-analysis of
22	CRP levels. Psychological Medicine, 49(12), 1958-1970.
23	doi:10.1017/S0033291719001454

1	Ouakinin, S. R. S., Barreira, D. P., & Gois, C. J. (2018). Depression and Obesity: Integrating
2	the Role of Stress, Neuroendocrine Dysfunction and Inflammatory Pathways.
3	Frontiers in Endocrinology, 9, 431. doi:10.3389/fendo.2018.00431
4	Papakostas, G. I., Petersen, T., Iosifescu, D. V., Burns, A. M., Nierenberg, A. A., Alpert, J.
5	E., Fava, M. (2005). Obesity among outpatients with major depressive disorder.
6	The International Journal of Neuropsychopharmacology, 8(1), 59-63.
7	doi:10.1017/S1461145704004602
8	Pearson, T. A., Mensah, G. A., Alexander, R. W., Anderson, J. L., Cannon, R. O., Criqui, M.,
9	Vinicor, F. (2003). Markers of Inflammation and Cardiovascular Disease.
10	Circulation, 107(3), 499-511. doi:10.1161/01.cir.0000052939.59093.45
11	Perez-Cornago, A., de la Iglesia, R., Lopez-Legarrea, P., Abete, I., Navas-Carretero, S.,
12	Lacunza, C. I., Zulet, M. A. (2014). A decline in inflammation is associated with
13	less depressive symptoms after a dietary intervention in metabolic syndrome patients:
14	a longitudinal study. Nutrition Journal, 13, 36. doi:10.1186/1475-2891-13-36
15	Perrin, A. J., Horowitz, M. A., Roelofs, J., Zunszain, P. A., & Pariante, C. M. (2019).
16	Glucocorticoid Resistance: Is It a Requisite for Increased Cytokine Production in
17	Depression? A Systematic Review and Meta-Analysis. Frontiers in Psychiatry, 10,
18	423. doi:10.3389/fpsyt.2019.00423
19	Pruessner, J. C., Kirschbaum, C., Meinlschmid, G., & Hellhammer, D. H. (2003). Two
20	formulas for computation of the area under the curve represent measures of total
21	hormone concentration versus time-dependent change. Psychoneuroendocrinology,
22	28(7), 916-931. doi:10.1016/s0306-4530(02)00108-7
23	Rethorst, C. D., Bernstein, I., & Trivedi, M. H. (2014). Inflammation, obesity, and metabolic
24	syndrome in depression: analysis of the 2009-2010 National Health and Nutrition

1	Examination Survey (NHANES). Journal of Clinical Psychiatry, 75(12), e1428-1432.
2	doi:10.4088/JCP.14m09009
3	Ruttle, P. L., Klein, M. H., Slattery, M. J., Kalin, N. H., Armstrong, J. M., & Essex, M. J.
4	(2014). Adolescent adrenocortical activity and adiposity: differences by sex and
5	exposure to early maternal depression. Psychoneuroendocrinology, 47, 68-77.
6	doi:10.1016/j.psyneuen.2014.04.025
7	Schmidt, F. M., Weschenfelder, J., Sander, C., Minkwitz, J., Thormann, J., Chittka, T.,
8	Himmerich, H. (2015). Inflammatory cytokines in general and central obesity and
9	modulating effects of physical activity. PloS One, 10(3), e0121971.
10	doi:10.1371/journal.pone.0121971
11	Tremmel, M., Gerdtham, U. G., Nilsson, P. M., & Saha, S. (2017). Economic Burden of
12	Obesity: A Systematic Literature Review. International Journal of Environmental
13	Research and Public Health, 14(4). doi:10.3390/ijerph14040435
14	Uher, R., Mors, O., Hauser, J., Rietschel, M., Maier, W., Kozel, D., Farmer, A. (2009).
15	Body weight as a predictor of antidepressant efficacy in the GENDEP project.
16	Journal of Affective Disorders, 118(1-3), 147-154. doi:10.1016/j.jad.2009.02.013
17	Upadhyay, J., Farr, O., Perakakis, N., Ghaly, W., & Mantzoros, C. (2018). Obesity as a
18	Disease. Medical Clinics of North America, 102(1), 13-33.
19	doi:10.1016/j.mcna.2017.08.004
20	Valkanova, V., Ebmeier, K. P., & Allan, C. L. (2013). CRP, IL-6 and depression: a
21	systematic review and meta-analysis of longitudinal studies. Journal of Affective
22	Disorders, 150(3), 736-744. doi:10.1016/j.jad.2013.06.004
23	Zhang, J., Yue, Y., Thapa, A., Fang, J., Zhao, S., Shi, W., Yuan, Y. (2019). Baseline
24	serum C-reactive protein levels may predict antidepressant treatment responses in

patients with major depressive disorder. *Journal of Affective Disorders, 250*, 432-438.
 doi:10.1016/j.jad.2019.03.001

1 Table 1: Comparison of sample groups

	Groups				Group tests	
Variables	Overweight & depressed N= 69	Overweight control N= 35	Normal weight & depressed N= 55	Normal weight control N= 57	Statistic	Р
Demographics						
Median age in years (range)	38 (25 – 50)	38 (25 – 48)	32 (25 – 50)	30 (24 – 50)	<i>K</i> = 18.368	<0.001
Gender, female (%)	43 (62.3%)	17 (48.6%)	40 (72.7%)	44 (77.2%)	$x^2 =$ 9.495	0.023
BMI (range)	28.8 (25.3 - 47.8)	28.4 (25 – 35.6)	22.1 (18.1 – 24.9)	22.4 (18.4 - 24.9)	K= 161.87	< 0.001
Ethnicity, white (%)	63 (91.3%)	29 (82.9%)	45 (81.8%)	45 (78.9%)	$x^2 =$ 4.102	0.251
Smoking status, smoker (%)	11 (15.9%)	3 (8.6%)	11 (20%)	6 (10.5%)	$x^2 =$ 3.199	0.362
Clinical						
Median HAM-D total score (range)	19 (14 – 31)	0 (0-5)	18 (14 – 26)	0 (0-7)	<i>K</i> = 162.83	<0.001
Currently on antidepressants (%)	47 (68.1%)	NA	39 (70.9%)	NA	$x^2 = 0.019$	0.889
Mean age of depression onset ± SD	24.6 ± 9.8	NA	25.2 ± 9.4	NA	F = 1.617	0.206
hsCRP						
Median hsCRP (range)	2.2 (0.2 – 15.3)	1.3 (0.2 - 11.8)	0.7 (0.2 - 8.8)	0.5 (0.2 - 5.2)	K = 43.725	< 0.001
Mean log hsCRP mg/L \pm SD Cohen's <i>d</i> effect size <i>vs</i> controls	0.28 ± 0.48 1.25	$\begin{array}{c} 0.09 \pm 0.43 \\ 0.89 \end{array}$	$\begin{array}{c} -0.12\pm0.42\\ 0.35\end{array}$	-0.25 ± 0.35	F = 16.43	<0.001
hsCRP ≥3mg/L (%)	26 (37.7%)	5 (14.3%)	6 (10.9%)	3 (5.3%)	$x^2 =$ 25.96	< 0.001
Salivary cortisol						
Mean CAR AUCi (nmol min/L) ± SD	100.53 ± 248.99	75.79 ± 240.98	41.62 ± 240.43	86.46 ± 299.78	<i>F</i> = 0.473	0.701
Mean log AUCg (nmol hour/L) ± SD	1.65 ± 0.18	1.66 ± 0.23	1.72 ± 0.2	1.7 ± 0.26	F = 1.25	0.293

2

3 BMI= body mass index, HAM-D= Hamilton Depression Rating Scale 17 for depressive symptoms, hsCRP=

4 high sensitivity C-reactive protein, SD = standard deviation, CAR= cortisol awakening response, AUCi= area

5 under the curve with respect to increase, AUCg= area under the curve with respect to ground.

Figure 1 caption: Estimated marginal means of logarithmic transformed hsCRP levels for each group. A two-way ANCOVA showed a significant effect of group on log hsCRP, with adjustment for age and gender F(3, 211)=16.43, p<0.001. Significance of pairwise comparisons, with the False Discovery Rate correction set at 5%, are shown within the figure.

Estimated marginal means of log-hsCRP by group Anova, F(3,211) = 16.43, p = <0.0001, $\eta_g^2 = 0.19$



Supplementary material 1

2 3

Table 1: Comparison of sample groups with non-smoking participants only

		Group tests				
Variables	Overweight & depressed N= 58	Overweight control N= 32	Normal weight & depressed N= 44	Normal weight control N= 51	Statistic	Р
Demographics						
Median age in years (range)	38.5 (25 - 50)	37 (25 – 48)	30 (25 - 50)	30 (24 – 50)	K = 15.16	0.002
Gender, female (%)	36 (62.1%)	15 (46.9%)	33 (75%)	39 (76.5%)	$x^2 = 9.744$	0.021
BMI (range)	29.6 (25.3 – 47.8)	28.5 (25 – 35.6)	22 (18.1 – 24.9)	22.4 (18.8 – 24.9)	K = 138.57	<0.001
Ethnicity, white (%)	52 (89.7%)	26 (81.2%)	36 (81.8%)	40 (78.4%)	$x^2 = 2.711$	0.438
Clinical						
Median HAM-D total score (range)	19 (14 – 31)	0 (0-5)	18 (14 – 26)	0 (0-7)	K = 141.16	<0.001
Currently on antidepressants (%)	37 (63.8%)	NA	31 (70.5%)	NA	$x^2 = 0.245$	0.621
Mean age of depression onset \pm SD	25 ± 9.8	NA	25.4 ± 9.2	NA	F = 1.623	0.206
hsCRP						
Median hsCRP (range)	2.25 (0.2 - 15.3)	1.4 (0.2 - 11.8)	0.7 (0.2 - 8.8)	0.5 (0.2 - 5.2)	K = 30.514	< 0.001
Mean log hsCRP mg/L ± SD Cohen's <i>d</i> effect size <i>vs</i> controls	0.27 ± 0.51 1.11	0.09 ± 0.48 0.81	-0.11 ± 0.43 0.29	-0.23 ± 0.36	F = 12.07	<0.001
hsCRP ≥3mg/L (%)	23 (39.7%)	4 (12.5%)	5 (11.4%)	3 (5.9%)	$x^2 = 24.405$	< 0.001
Salivary cortisol						
Mean CAR AUCi (nmol min/L) ± SD	99.8 ± 262	87.7 ± 238	6.79 ± 228	89 ± 311	F = 0.889	0.448
Mean log AUCg (nmol hour/L) ± SD	1.66 ± 0.18	1.67 ± 0.23	1.7 ± 0.18	1.72 ± 0.23	F = 0.082	0.495

BMI= body mass index, HAM-D= Hamilton Depression Rating Scale 17 for depressive symptoms, hsCRP=

4 5 6 high sensitivity C-reactive protein, SD = standard deviation, CAR= cortisol awakening response, AUCi= area

under the curve with respect to increase, AUCg= area under the curve with respect to ground.

1

NIMA (Part 1) Consortium members

- 2
- 3 <u>Cambridge</u>
- 4 Edward T. Bullmore (MD, PI, EC)^{1,2,11}, Junaid Bhatti¹, Samuel J. Chamberlain^{1,2}, Marta M.
- 5 Correia^{1,12}, Anna L. Crofts¹, Amber Dickinson*, Andrew C. Foster*, Manfred G.
- 6 Kitzbichler¹, Clare Knight*, Mary-Ellen Lynall¹, Christina Maurice¹, Ciara O'Donnell¹,
- 7 Linda J. Pointon¹, Peter St George Hyslop^{1,13,14}, Lorinda Turner³¹, Petra Vertes¹, Barry
- 8 Widmer¹, Guy B. Williams^{1,14}
- 9 10 Cand
- 10 <u>Cardiff</u>
- 11 B. Paul Morgan (PI)¹⁵, Claire A. Leckey¹⁵, Angharad R. Morgan*, Caroline O'Hagan*,
- 12 Samuel Touchard¹⁵
- 13
- 14 <u>Glasgow</u>
- 15 Jonathan Cavanagh (PI, EC)³, Catherine Deith*, Scott Farmer¹⁶, John McClean¹⁶, Alison
- 16 McColl³, Andrew McPherson^{*}, Paul Scouller^{*}, Murray Sutherland¹⁶
- 17
- 18 Independent advisor
- 19 H.W.G.M. (Erik) Boddeke (EC)¹⁷
- 20

- 21 GSK
- 22 Jill C. Richardson (EC)¹⁸, Shahid Khan¹¹, Phil Murphy¹⁹, Christine A. Parker¹⁹, Jai Patel¹¹
- 24 Janssen
- 25 Declan Jones (EC)⁶, Peter de Boer⁴, John Kemp⁴, Wayne C. Drevets⁶, Jeffrey S. Nye
- 26 (deceased), Gayle Wittenberg⁶, John Isaac⁶, Anindya Bhattacharya⁶, Nick Carruthers⁶,
- 27 Hartmuth Kolb⁶
- 28
- 29 Kings College London
- 30 Carmine M. Pariante (PI)¹⁰, Federico Turkheimer (PI)²⁰, Gareth J. Barker²⁰, Heidi Byrom¹⁰,
- 31 Diana Cash²⁰, Annamaria Cattaneo¹⁰, Antony Gee²⁰, Caitlin Hastings¹⁰, Nicole Mariani¹⁰,
- 32 Anna McLaughlin¹⁰, Valeria Mondelli¹⁰, Maria Nettis¹⁰, Naghmeh Nikkheslat¹⁰, Karen
- 33 Randall²⁰, Hannah Sheridan*, Camilla Simmons²⁰, Nisha Singh²⁰, Victoria Van Loo*, Marta
- 34 Vicente-Rodriguez²⁰, Tobias C. Wood²⁰, Courtney Worrell*, Zuzanna Zajkowska*
- 35
- 36 <u>Lundbeck</u>
- 37 Niels Plath (EC)²¹, Jan Egebjerg²¹, Hans Eriksson²¹, Francois Gastambide²¹, Karen Husted
- 38 Adams²¹, Ross Jeggo*, Christian Thomsen²¹, Jan Torleif Pederson²¹, Brian Campbell*,
- 39 Thomas Möller*, Bob Nelson*, Stevin Zorn*
- 40
- 41 <u>University of Texas (sub-contracted to Lundbeck)</u>
- 42 Jason O'Connor²²
- 43
- 44 <u>Oxford</u>
- 45 Mary Jane Attenburrow (PI)^{7,23}, Alison Baird, Jithen Benjamin²³, Stuart Clare²⁵, Philip
- 46 Cowen⁷, I-Shu (Dante) Huang²⁴, Samuel Hurley^{*}, Helen Jones²³, Simon Lovestone⁷, (AD, PI,
- 47 EC) Francisca Mada*, Alejo Nevado-Holgado⁷, Akintayo Oladejo*, Elena Ribe⁷, Katy
- 48 Smith²³, Anviti Vyas*
- 49
- 50 <u>Pfizer</u>

- 1 Zoe Hughes*, Rita Balice-Gordon*, James Duerr*, Justin R. Piro*, Jonathan Sporn*
- 2
- 3 Southampton
- 4 V. Hugh Perry (PI)²⁷, Madeleine Cleal*, Gemma Fryatt²⁷, Diego Gomez-Nicola²⁷, Renzo
- 5 Mancuso³², Richard Reynolds²⁷
- 6
- 7 <u>Sussex</u>
- 8 Neil A. Harrison (PI, EC)²⁸, Mara Cercignani²⁸, Charlotte L. Clarke²⁸, Elizabeth Hoskins*,
- 9 Charmaine Kohn*, Rosemary Murray*, Lauren Wilcock²⁹, Dominika Wlazly³⁰
- 10
- 11 <u>University of Toronto (sub-contracted to Cambridge)</u>
- 12 Howard Mount¹³
- 13
- 14 MD = Mood disorder workpackages lead
- 15 AD = Alzheimer's disease workpackages lead
- 16 PI = Principal Investigator
- 17 EC = Executive committee member
- 18
- 19⁻¹ Department of Psychiatry, School of Clinical Medicine, University of Cambridge, CB2 0SZ,
 20 UK
- 21² Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge, CB21 5EF, UK
- ³ Sackler Centre, Institute of Health & Wellbeing, University of Glasgow, Sir Graeme Davies
 Building , Glasgow, G12 8TA, UK
- ⁴ Neuroscience, Janssen Research & Development, Janssen Pharmaceutica NV, Turnhoutseweg
 30, B-2340, Beerse, Belgium
- 26⁵ The Maurice Wohl Clinical Neuroscience Institute, Cutcombe Road, London, SE5 9RT, UK
- 27 ⁶ Neuroscience, Janssen Research & Development, LLC, Titusville, NJ, 08560, USA
- 28 ⁷ Department of Psychiatry, University of Oxford, Warneford Hospital, Oxford, OX3 7JX, UK
- 29 ⁸ Brighton & Sussex Medical School, University of Sussex, Brighton, BN1 9RR, UK
- 30 ⁹ Sussex Partnership NHS Foundation Trust, Swandean, BN13 3EP, UK
- ^{31 10} Kings College London, Institute of Psychiatry, Psychology and Neuroscience, Department of
 ³² Psychological Medicine, London, SE5 9RT, UK
- 33 ¹¹ Immuno-Psychiatry, Immuno-Inflammation Therapeutic Area Unit, GlaxoSmithKline R&D,
 34 Stevenage SG1 2NY, UK
- 35¹² MRC Cognition and Brain Sciences Unit, 15 Chaucer Road, Cambridge CB2 7EF, UK
- 36 ¹³ Tanz Centre for Research in Neurodegenerative Diseases, 60 Leonard Avenue, Toronto, ON
 37 M5T 2S8 Canada
- 38¹⁴ Department of Clinical Neurosciences, University of Cambridge, CB2 0SZ, UK
- 39¹⁵ Cardiff University, Cardiff CF10 3AT, UK
- 40 ¹⁶ NHS Greater Glasgow and Clyde, 1055 Great Western Rd, Glasgow G12 0XH, UK
- 41¹⁷ University of Groningen, 9712 CP Groningen, Netherlands
- 42¹⁸ Neurosciences Virtual PoC DPU, GlaxoSmithKline R&D, Stevenage SG1 2NY, UK
- 43¹⁹ Experimental Medicine Imaging, GlaxoSmithKline R&D, Stevenage SG1 2NY, UK
- 44²⁰ King's College London, Department of Neuroimaging Sciences, Institute of Psychiatry,
- 45 Psychology & Neuroscience, De Crespigny Park, London SE5 8AF, UK
- 46²¹ H. Lundbeck A/S Ottiliavej 9, 2500, Valby, Denmark
- 47 ²² University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Dr, San Antonio,
 48 TX 78229, USA
- 49²³ NIHR Oxford cognitive health Clinical Research Facility, Warneford Hospital, Oxford, OX3
- 50 7JX, UK

- 1²⁴ The Kennedy Institute of Rheumatology, Roosevelt Dr, Oxford OX3 7FY, UK
- 2 ²⁵ Oxford Centre for Functional MRI of the Brain, John Radcliffe Hospital, Oxford OX3 9DU,
 3 UK
- 4²⁶ Pfizer, Inc, 1 Portland Street, Cambridge MA, USA
- 5²⁷ Centre for Biological Sciences, University of Southampton, Southampton, UK
- 6²⁸ Clinical Imaging Sciences Centre (CISC), University of Sussex, Brighton, BN1 9RR, UK
- 7²⁹ Sussex Partnership NHS Foundation Trust, Nevill Avenue, Hove BN3 7HZ, UK
- 8³⁰ Brighton & Sussex University Hospitals NHS Trust, Brighton BN2 5BE, UK
- 9³¹ Department of Medicine, School of Clinical Medicine, University of Cambridge, CB2 0SZ,
- 10 UK
- 11³² VIB-KU Leuven Center for Brain & Disease Research, Campus Gasthuisberg, Herestraat 49,
- 12 bus 602, 3000 Leuven, Belgium
- 13
- 14 *Former consortium members
- 15
- 16