



King's Research Portal

DOI:

[10.1007/JHEP07\(2015\)043](https://doi.org/10.1007/JHEP07(2015)043)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Assel, B., Cassani, D., Di Pietro, L., Komargodski, Z., Lorenzen, J., & Martelli, D. (2015). The Casimir energy in curved space and its supersymmetric counterpart. *Journal of High Energy Physics*, 2015(7), 1-43. Article 43. [https://doi.org/10.1007/JHEP07\(2015\)043](https://doi.org/10.1007/JHEP07(2015)043)

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

The Casimir energy in curved space and its supersymmetric counterpart

Benjamin Assel,^a Davide Cassani,^{b,c} Lorenzo Di Pietro,^d Zohar Komargodski,^d
Jakob Lorenzen^a and Dario Martelli^a

^aDepartment of Mathematics, King's College London,
The Strand, London WC2R 2LS, U.K.

^bSorbonne Universités UPMC Paris 06, UMR 7589, LP THE,
F-75005, Paris, France

^cCNRS, UMR 7589, LP THE,
F-75005, Paris, France

^dDepartment of Particle Physics and Astrophysics, Weizmann Institute of Science,
Rehovot, 76100, Israel

E-mail: benjamin.assel@kcl.ac.uk, davide.cassani@lpthe.jussieu.fr,
lorenzo.dipietro@weizmann.ac.il, zohar.komargodski@weizmann.ac.il,
jakob.lorenzen@kcl.ac.uk, dario.martelli@kcl.ac.uk

ABSTRACT: We study d -dimensional Conformal Field Theories (CFTs) on the cylinder, $S^{d-1} \times \mathbb{R}$, and its deformations. In $d = 2$ the Casimir energy (i.e. the vacuum energy) is universal and is related to the central charge c . In $d = 4$ the vacuum energy depends on the regularization scheme and has no intrinsic value. We show that this property extends to infinitesimally deformed cylinders and support this conclusion with a holographic check. However, for $\mathcal{N} = 1$ supersymmetric CFTs, a natural analog of the Casimir energy turns out to be *scheme independent* and thus intrinsic. We give two proofs of this result. We compute the Casimir energy for such theories by reducing to a problem in supersymmetric quantum mechanics. For the round cylinder the vacuum energy is proportional to $a + 3c$. We also compute the dependence of the Casimir energy on the squashing parameter of the cylinder. Finally, we revisit the problem of supersymmetric regularization of the path integral on Hopf surfaces.

KEYWORDS: Supersymmetric gauge theory, Supersymmetric Effective Theories, Anomalies in Field and String Theories

ARXIV EPRINT: [1503.05537](https://arxiv.org/abs/1503.05537)

Contents

1	Introduction and summary	1
1.1	Outline of the derivation	5
2	Supersymmetric Casimir energy	8
2.1	Definition	8
2.2	Consequences of the supersymmetry algebra	9
2.3	Supersymmetric quantum mechanics	12
2.4	Spectrum of the Hamiltonian	15
2.5	Dimensional reduction of a 4d chiral multiplet	18
3	Supersymmetric Casimir energy on a deformed three-sphere	20
3.1	Shortening conditions on chiral multiplets	20
3.2	Reduction on deformed three-sphere	23
A	Casimir energy in CFT in $d = 2$ and $d = 4$	29
A.1	The cylinder and its infinitesimal deformation	29
A.2	The Casimir energy and holography	31
A.3	Holographic check of E_0 on a squashed cylinder	32
B	E_{susy} and the Hamiltonian	35
C	Regularization of one-loop determinant	36
C.1	Two-step regularization	37
C.2	Cut-off regularization	39

1 Introduction and summary

The quantum anomalies appearing in the trace of the energy-momentum tensor encode universal information about Conformal Field Theories (CFTs) in even space-time dimensions. In $d = 2$ the conformal anomaly c characterizes conformal field theories [1], and constrains the renormalization group (RG) flows between them [2]. In four-dimensional CFTs, there are two trace anomaly coefficients, c and a . They appear in many applications. In particular, Cardy conjectured [3] that there exists a decreasing function along RG flows, reducing to a at the fixed points [4–6].

Here we will be interested in the following question. Given a conformal field theory in \mathbb{R}^d , $d = 2n$, we can use a Weyl transformation and study the theory on $S^{d-1} \times \mathbb{R}$. This is

often referred to as “radial quantization.” Denoting the noncompact coordinate by τ , we can ask about the energy of the ground state E_0 , defined as

$$E_0 = \int_{S^{d-1}} d^{d-1}x \sqrt{g} \langle T_{\tau\tau} \rangle . \quad (1.1)$$

The expectation value of the energy-momentum tensor is taken in the ground state of the CFT on the cylinder. We refer to E_0 as the Casimir energy.

It is crucial for the analysis below to understand whether E_0 is well defined, namely, scheme independent. In $d = 2$ the only dimensionless counterterm is

$$\int d^2x \sqrt{g} R , \quad (1.2)$$

where R denotes the Ricci scalar. This vanishes on the cylinder and thus does not shift the vacuum energy.¹ In $d = 4$ there are several dimensionless counterterms. A basis is given by the Euler density $E_{(4)}$, the square of the Weyl tensor W^2 , the Pontryagin density $\text{Tr}(R \wedge R)$, and R^2 . Of those four dimensionless counterterms only $\int d^4x \sqrt{g} R^2$ does not vanish on $S^3 \times \mathbb{R}$. We could thus add this counterterm to the action with some scheme-dependent coefficient b

$$\delta S = -\frac{b}{12(4\pi)^2} \int d^4x \sqrt{g} R^2 . \quad (1.3)$$

The curvature of the three-sphere of radius r_3 is $R = \frac{6}{r_3^2}$ in our conventions. As a consequence, integrating the counterterm above we get $\delta S = -\frac{3b}{8r_3} \int d\tau$. Interpreting the coefficient of $\int d\tau$ as the ground state energy we see that E_0 is ambiguous. An equivalent way to understand this ambiguity is to note that (1.3) leads to a scheme-dependent term in the trace of the energy-momentum tensor

$$\langle T_\mu^\mu \rangle = \frac{1}{(4\pi)^2} (aE_{(4)} - cW^2 + b\Box R) . \quad (1.4)$$

Since $\langle T_\mu^\mu \rangle$ is modified, this affects the other components of $\langle T_{\mu\nu} \rangle$ and in particular the vacuum energy. By contrast, in $d = 2$ no dimension 2 term can be added to the right-hand side of $\langle T_\mu^\mu \rangle = -\frac{c}{24\pi} R$.

We see that, without additional assumptions, the ground state energy on $S^3 \times \mathbb{R}$ is not an intrinsic property of the CFT. It depends on the regularization scheme, i.e. the precise way in which the CFT is defined (different definitions lead to different values of b). On the other hand, the Casimir energy on $S^1 \times \mathbb{R}$ is an intrinsic observable that does not depend on the ultraviolet completion.

For completeness, let us quote the results for E_0 in $d = 2$ and in $d = 4$. In $d = 4$ below we use the general scheme with some b . We present a self-contained derivation of these

¹More precisely, since there is the cosmological constant counterterm $\Lambda_{UV}^2 \int d^2x \sqrt{g}$, the vacuum energy would generally have power divergent pieces $E_0 \sim \Lambda_{UV}^2 r_1$ (with r_1 the radius of the S^1 and Λ_{UV} the UV cut-off) which are non-universal. So when we discuss the scheme-independence of the vacuum energy, we always have in mind the piece that remains finite when the cut-off is removed. In $d = 2n$ dimensions, only counterterms with d -derivatives of the metric are thus relevant to us.

results in appendix A. The Casimir energy on $S^1 \times \mathbb{R}$ is

$$E_0 = -\frac{c}{12r_1}, \quad (1.5)$$

where r_1 is the radius of the circle. The Casimir energy on $S^3 \times \mathbb{R}$ is

$$E_0 = \frac{3}{4r_3} \left(a - \frac{b}{2} \right), \quad (1.6)$$

where r_3 is the radius of the three-sphere. In fact, as we show in appendix A, the result in $d = 4$ remains ambiguous even if one allows for an infinitesimal deformation of the three-sphere. By this we mean that the linear response to an infinitesimal deformation is still proportional to $a - \frac{b}{2}$. We also provide a holographic test of this claim, by checking it against the linearized holographic Casimir energy of the supergravity solution of [7].

The three main points of the present paper are

- There is a natural generalization of the notion of Casimir energy for $\mathcal{N} = 1$ supersymmetric field theories on $S^3 \times \mathbb{R}$.
- It turns out that this Casimir energy is *physical*, i.e. scheme independent. We will give two proofs of this assertion and then we will evaluate the Casimir energy exactly. One proof is based on [8] and the other is based on ideas related to Chern-Simons terms in quantum mechanics. The Casimir energy comes out proportional to the linear combination $a + 3c$ of the conformal anomalies.
- When one computes supersymmetric partition functions on manifolds with $S^3 \times S^1$ topology, the answer is given in terms of a product of the Casimir factor and the usual supersymmetric index. This leads to consistent results in various limits of the partition function.

Let us now briefly explain why it might be useful to understand the Casimir factor for four-dimensional theories. Recall that in $d = 2$, in order for the torus partition function to be modular invariant, it is necessary to include the Casimir prefactor $e^{\frac{c\beta}{12r_1}}$ (where β is the length of the thermal circle). The Casimir prefactor in four dimensions might be similarly important to manifest various relations between different path integrals such as those analogous to [9] and e.g. [10]. The Casimir prefactor might be also relevant for a better understanding of the role of the modular-like transformations in four dimensions discussed in [11, 12] (and see references therein).

So let us begin by discussing how we put supersymmetric $\mathcal{N} = 1$ theories on $S^3 \times \mathbb{R}$. For theories with an R -symmetry (such as any SCFT) one can follow the procedure of [13] whereby one couples the theory to background new minimal supergravity. New minimal supergravity [14] contains the bosonic fields $g_{\mu\nu}$, A_μ , $B_{\mu\nu}$, where A_μ is the R -gauge field and $B_{\mu\nu}$ is a two-form.² The procedure works both in Lorentzian and in Euclidean signature; here we refer to the Euclidean case for definiteness.

²One is allowed to discuss the R -symmetry in curved space without any significant modifications because its gravitational anomaly vanishes under very general assumptions [15].

It turns out that for every complex manifold with a Hermitian metric one can find appropriate A_μ and $B_{\mu\nu}$ such that one preserves at least one rigid supercharge [16, 17]. This can be viewed as a generalization of the notion of twisting [18]. Furthermore, the partition function is independent of the Hermitian metric; it only depends on the complex structure of the underlying complex manifold [19]. A particularly interesting class of complex manifolds are primary Hopf surfaces, which are topologically equivalent to $S^3 \times S^1$. In this case, the relevant branch of the moduli space of complex structures is two-complex dimensional, parameterized by complex numbers p, q .

For the special case of the round metric on $S^3 \times S^1$, the complex structure parameters are given in terms of the S^1 -length β and the S^3 -radius r_3 as $p = q = e^{-\frac{\beta}{r_3}}$. This geometry preserves in fact four supercharges and in addition to the round metric we need to activate the background fields $A_\tau = V_\tau = \frac{i}{r_3}$, where $V_\mu = \frac{1}{4}\epsilon_{\mu\nu\rho\sigma}\nabla^\nu B^{\rho\sigma}$. The two-form $B_{\mu\nu}$ couples to a redundant operator in conformal field theories and is not so important. However, the flat gauge field A along the Euclidean time direction implies that the Hamiltonian, H_{susy} , is shifted with respect to what we would get from radial quantization according to

$$H_{\text{susy}} = \Delta - \frac{1}{2r_3}R. \quad (1.7)$$

Above Δ is the time translation operator that is obtained by mapping the dilatation operator in flat space to the cylinder (equivalently, Δ is the canonical Hamiltonian for a theory on the cylinder with $A_\tau = \frac{3i}{2r_3}$, $V_\tau = \frac{i}{r_3}$). The Hamiltonian (1.7) commutes with the four supercharges on the round $S^3 \times S^1$ and so the Hilbert space is organized in representations of the appropriate superalgebra.

The Casimir energy, E_{susy} , is thus naturally defined from the large β limit of the partition function [20]

$$Z_{S^3 \times S^1_\beta}^{\text{susy}} \sim e^{-\beta E_{\text{susy}}} \quad \text{for} \quad \beta \rightarrow \infty. \quad (1.8)$$

In other words, E_{susy} is the lowest eigenvalue of the Hamiltonian (1.7). See also appendix B.

This definition can be extended to manifolds $M_3 \times S^1$, with M_3 a Seifert manifold (a Seifert manifold is, loosely speaking, a circle bundle over a Riemann surface). For primary Hopf surfaces, we will deform the three-sphere with some squashing parameter \mathfrak{b} , related to the underlying complex structure parameters p, q , and study $E_{\text{susy}}(\mathfrak{b})$. Our final result for the round sphere ($\mathfrak{b} = 1$) with radius r_3 is³

$$E_{\text{susy}} = \frac{4}{27r_3}(a + 3c), \quad (1.9)$$

in agreement with [21]. The partition function is therefore given by

$$Z_{S^3 \times S^1_\beta}^{\text{susy}} = e^{-\frac{4\beta}{27r_3}(a+3c)} \mathcal{I}_{S^3 \times S^1_\beta}, \quad (1.10)$$

with $\mathcal{I}_{S^3 \times S^1_\beta}$ being the usual supersymmetric index [22–25]. The results of [26–29] are consistent with (1.10).

³The formula also applies to non-conformal theories by replacing a and c with the appropriate linear combinations of traces over the fermion R -charges $\text{Tr}R$ and $\text{Tr}R^3$ (see (2.53)).

Since the Casimir energy is not well defined without supersymmetry, one has to develop a supersymmetric method to regularize the a priori divergent Casimir energy. As we will explain in the following subsection, the main idea is that the expectation value of (1.7) can be related to the expectation value of the R -charge in the vacuum. The latter can be thought of as a Chern-Simons term in quantum mechanics and this leads to a determination of the ordering ambiguities (see also [30]).

Let us mention that recovering (1.9) from a holographic computation remains an open problem. Recall that for four-dimensional SCFTs admitting a weakly coupled gravity dual, at leading order in the large N limit, one has $a = c$ and thus we predict $E_{\text{susy}} = \frac{16}{27r_3}a \sim \mathcal{O}(N^2)$. Therefore this should be reproduced by the on-shell action of an appropriate supergravity solution. It was noted in [7] that an obvious candidate for such a solution is AdS_5 , with the addition of an appropriate flat background gauge field; however, its renormalised on-shell action is the same as that of pure AdS_5 , which agrees with (1.6) (with $b = 0$) [31], and does not agree with (1.9). It is clearly worthwhile to revisit this problem.

In the next subsection we briefly summarize the derivation of (1.9). Section 2 contains the detailed derivation of (1.9) and section 3 contains the generalization to squashed cylinders. Appendix A contains a study (which is mostly a review) of the non-supersymmetric Casimir energy. In appendix B we include some details about the identification of E_{susy} with the VEV of the Hamiltonian. Finally, in appendix C we reconsider the full partition function on $S^3 \times S^1$. We revisit the regularization of one-loop determinants and determine the relation of the partition function and the supersymmetric index for all p, q , a study initiated in [20] (see also [32]), thereby generalizing (1.10). The correct regularization of the one-loop determinants leads to results consistent with the Cardy-like formula [26] (see also [27]), namely

$$Z_{S^3 \times S^1_\beta}^{\text{susy}} \sim \exp\left(-\frac{16\pi^2 r_3}{3\beta}(a - c)\right) \quad \text{for} \quad \beta \rightarrow 0, \quad (1.11)$$

as advertised in [26, 30].

1.1 Outline of the derivation

Let us now summarize the main points in the derivation of the supersymmetric Casimir energy for SCFTs.

In section 2 we consider an $\mathcal{N} = 1$ theory with an R -symmetry on the conformally flat background $S^3 \times \mathbb{R}$, with round metric on S^3 . We work in Lorentzian signature and denote the real time $t = i\tau$. Preserving supersymmetry requires to turn on background fields of the new minimal supergravity multiplet $A_t = V_t = \frac{1}{r_3}$. Focusing on two supercharges of opposite R -charge (out of the four preserved by the background), the supersymmetry algebra on $S^3 \times \mathbb{R}$ takes the form

$$\frac{1}{2}\{Q, Q^\dagger\} = H_{\text{susy}} - \frac{1}{r_3}(R + 2J_3), \quad [H_{\text{susy}}, Q] = [R + 2J_3, Q] = 0. \quad (1.12)$$

Here R is the generator of $U(1)_R$ symmetry while J_3 is a Cartan generator of the isometries of the three-sphere. The Casimir energy is given by the VEV of the Hamiltonian

appearing in the algebra, $E_{\text{susy}} = \langle H_{\text{susy}} \rangle$. Here, the expectation value is evaluated in the vacuum of the theory coupled to the various background fields. Since the supercharge is time-independent, we can then Wick rotate and compactify the time direction on a circle of length β , and E_{susy} gives the dominant contribution to the supersymmetric partition function for $\beta \rightarrow \infty$ as in (1.8).

Because of the separation of scales between the radius of the three-sphere and the radius of the circle, it is natural to study the reduction of the theory on the three-sphere. The result of this reduction is a *supersymmetric quantum mechanics* with infinitely many degrees of freedom. We are thus led to consider a $1d$ system with Hamiltonian H_{susy} , a global symmetry with charge Σ and a supersymmetry algebra of the form

$$\{Q, Q^\dagger\} = 2\left(H_{\text{susy}} - \frac{1}{r_3}\Sigma\right), \quad [H_{\text{susy}}, Q] = [\Sigma, Q] = 0. \quad (1.13)$$

Unbroken supersymmetry implies that $r_3\langle H_{\text{susy}} \rangle = \langle \Sigma \rangle$. When we reduce to $1d$, the effective action that computes $\langle \Sigma \rangle$ is given by

$$\mathcal{W}[A_t^\Sigma] = \langle \Sigma \rangle \int dt A_t^\Sigma, \quad (1.14)$$

where A^Σ is the background gauge field for the symmetry generated by Σ . This is a $1d$ Chern-Simons (CS) term associated to the VEV of Σ , i.e. the charge of the vacuum.

In quantum mechanics (1.14) can in principle be completed to a supersymmetric counterterm, involving the $1d$ metric g_{tt} as well as other background supergravity fields. Such a counterterm would reflect the fact that $\langle H_{\text{susy}} \rangle$ and $\langle \Sigma \rangle$ can be shifted by an arbitrary amount, preserving the relation $r_3\langle H_{\text{susy}} \rangle = \langle \Sigma \rangle$ imposed by the superalgebra (1.13). By contrast, if the quantum mechanical model arises from a local higher-dimensional model, the allowed counterterms must descend from local higher-dimensional counterterms. It is easy to see that the quantum-mechanical CS term (1.14) cannot descend from a higher-dimensional counterterm and thus it is scheme independent. As a consequence, since the vacuum energy is fixed by supersymmetry to be the same as the $1d$ CS term, the vacuum energy is physical. Furthermore, since the CS term cannot depend on continuous coupling constants, the vacuum energy is also independent of continuous coupling constants. If we further assume the existence of a weakly coupled point, we can reliably compute the Casimir energy using a free field theory.

We can then consider a free chiral multiplet. Supersymmetry implies that upon reduction on the S^3 the resulting Lagrangian and supersymmetry transformations must be organized in terms of $1d$ multiplets and Lagrangians. In particular, there are two types of multiplets, that we will call “short.” These are the *chiral* (ϕ, ψ) and the *Fermi* (λ, f) multiplets. A key point is that the combination of the two multiplets can form a reducible but indecomposable representation of supersymmetry. Thus they can join to form a “long” multiplet, with the coupling between the short components controlled by a parameter p . When $p = 0$, the short multiplets are decoupled.

The scalar fields of the $1d$ chiral multiplets arise from “Fourier modes” of the scalar fields in the $4d$ chiral multiplet. For example, on the round sphere we can use the spherical harmonics and write $\phi = \sum \phi_{\ell, m, n} Y_{\ell, m, n}$. Here ℓ, m, n are the usual $\text{SU}(2)_l \times \text{SU}(2)_r$

quantum numbers. Similarly, one can expand the two-component fermion of the $4d$ chiral multiplet in spinorial harmonics, and the $1d$ fermions ψ and λ may be identified with linear combinations of its components. The deformation parameter p that governs the shortening of the multiplets is related to the $4d$ quantum number as $p^2 = (\ell - 2m)(\ell + 2 + 2m)$, hence the shortening condition $p = 0$ is satisfied if and only if $m = \ell/2$ or $m = -\ell/2 - 1$.

We will see that in the correct renormalization scheme (whose existence we established above), the Hamiltonian of a long multiplet exactly *vanishes* on the vacuum, while for the short multiplets we find

$$\begin{aligned} \text{chiral } \left(m = \frac{\ell}{2}\right) : & \quad \langle H_{\text{chiral}} \rangle = \frac{1}{2r_3}(\ell + r), \\ \text{Fermi } \left(m = -\frac{\ell}{2} - 1\right) : & \quad \langle H_{\text{Fermi}} \rangle = -\frac{1}{2r_3}(\ell + 2 - r), \end{aligned} \quad (1.15)$$

where r is the R -charge of the $4d$ chiral multiplet. Thus, given that the long multiplets do not contribute, the expectation value of the total Hamiltonian is

$$\langle H_{\text{susy}} \rangle = \sum_{\text{chiral}} \langle H_{\text{chiral}} \rangle + \sum_{\text{Fermi}} \langle H_{\text{Fermi}} \rangle. \quad (1.16)$$

The two infinite sums can be regularized using different methods, for example using the Hurwitz zeta function, or a cut-off method, (see also [20] and [30]) yielding in any case the result⁴

$$\langle H_{\text{susy}} \rangle = E_{\text{susy}} = \frac{4}{27r_3}(a + 3c). \quad (1.17)$$

Using a similar strategy, in section 3 we compute the supersymmetric Casimir energy for an $\mathcal{N} = 1$ theory (with an R -symmetry) on a class of Hopf surfaces with $U(1)^3$ symmetry, considered in [20]. Let us describe this briefly. Again we start from the free chiral multiplet Lagrangian in four dimensions, and reduce this to an infinite set of fields in $1d$, comprising long and short multiplets. Since the quantized Hamiltonian of long multiplets vanishes on the vacuum, we can focus on the short multiplets. The shortening conditions are expressed as first-order differential equations for the eigenstates of the Hamiltonians for chiral and Fermi multiplets, which are labeled by two non-negative integers n_1, n_2 . This can be seen using the method of [34] or, equivalently, utilising the variables of [32]. Each chiral and Fermi multiplet contributes as

$$\begin{aligned} \text{chiral :} \quad & \langle H_{\text{chiral}} \rangle = \frac{1}{2r_3} \left(|\mathfrak{b}|n_1 + |\mathfrak{b}^{-1}|n_2 + \frac{r}{2}(|\mathfrak{b}| + |\mathfrak{b}^{-1}|) \right), \\ \text{Fermi :} \quad & \langle H_{\text{Fermi}} \rangle = -\frac{1}{2r_3} \left(|\mathfrak{b}|n_1 + |\mathfrak{b}^{-1}|n_2 + \frac{2-r}{2}(|\mathfrak{b}| + |\mathfrak{b}^{-1}|) \right), \end{aligned} \quad (1.18)$$

where \mathfrak{b} is a parameter deforming the complex structure of the Hopf surface. Again the two sums in (1.16) must be regulated separately. This can be done using the Barnes double

⁴In this paper we do not carry out the explicit analysis for the vector multiplets; the results would be consistent with our conclusions. Additionally, the combination $a + 3c$ has an interesting interpretation in terms of the anomaly polynomial [33].

zeta function, which is a two-parameter generalization of the Hurwitz zeta function. In the end, we obtain

$$\langle H_{\text{susy}} \rangle = \frac{2}{3r_3} (|\mathfrak{b}| + |\mathfrak{b}|^{-1}) (a - c) + \frac{2}{27r_3} (|\mathfrak{b}| + |\mathfrak{b}|^{-1})^3 (3c - 2a), \quad (1.19)$$

which is the expression (5.10) in [20] (with the redefinition $b_1 = \frac{\beta \mathfrak{b}}{2\pi r_3}$ and $b_2 = \frac{\beta \mathfrak{b}^{-1}}{2\pi r_3}$).

2 Supersymmetric Casimir energy

In this section we perform a manifestly supersymmetric analysis of the Casimir energy problem on $S^3 \times \mathbb{R}$. Our strategy is to reduce to a one-dimensional quantum mechanical problem. Using the same idea, in section 3 we will discuss the more general case leading to (1.19).

2.1 Definition

Consider an $\mathcal{N} = 1$ theory with an R -symmetry. The energy-momentum tensor can be embedded in the so-called R -multiplet (see, for instance, [35]), hence the theory can be coupled canonically to new minimal supergravity. The bosonic part of the supergravity multiplet consists of the metric $g_{\mu\nu}$ and two auxiliary fields, the R -gauge field A_μ and a conserved one-form V_μ , $\nabla_\mu V^\mu = 0$, dual to the field strength of a two-form gauge field $B_{\mu\nu}$. Gravity can then be made non-dynamical by taking a rigid limit, so that one is left with an ordinary field theory on a four-manifold M_4 . In this limit generically all the supergravity fields above are turned on, and play the role of background fields. For particular choices of background fields and manifolds M_4 one can show that the theory on M_4 preserves some rigid supersymmetry. The condition is that there exists a nontrivial solution to the generalized Killing spinor equation [13]

$$(\nabla_\mu - iA_\mu)\zeta_\alpha + iV_\mu\zeta_\alpha + iV^\nu(\sigma_{\mu\nu}\zeta)_\alpha = 0, \quad (2.1)$$

and its conjugate equation. In Lorentzian signature, these equations admit a solution if and only if the background admits a null Killing vector [36]. In Euclidean signature, solutions were classified in [16, 17]. It turns out that a solution exists if and only if the manifold is complex, with a Hermitian metric. For some special cases (and the cases we analyze in this paper are of this type) one can naturally Wick rotate from the Euclidean to the Minkowski solution. Examples of such backgrounds are $M_4 = M_3 \times S^1$, with M_3 a Seifert manifold (examples include spheres and Lens spaces).

An $\mathcal{N} = 1$ theory on a curved manifold M_4 presents the same short-distance behavior as in flat space. So in general there will be divergences in the computation of e.g. the partition function on M_4 , which need to be renormalized. Assuming that the theory is regularized in a way which respects both supersymmetry and the R -symmetry, we infer that two different regularization schemes differ by some local counterterm, namely some local action constructed using the background new minimal supergravity fields, $\int d^4x \sqrt{g} \mathcal{L}(g_{\mu\nu}, A_\mu, V_\mu)$. This has to respect supersymmetry and gauge invariance. Local terms of dimension four affect the finite part of the partition function, as they remain finite when the UV cut-off is removed. In [8] it was shown that when there exist two Killing spinors of opposite

chirality, then all local counterterms of dimension four evaluate to zero. This means that such partition functions are in fact scheme-independent.⁵

The results of [18, 19] tell us that $Z_{M_3 \times S^1}^{\text{susy}}$ is independent of coupling constants and only depends on complex structure parameters of the underlying complex four-manifold $M_3 \times S^1$. The findings of this work are consistent with that.

The partition function can be interpreted as usual as a trace over the Hilbert space $Z_{M_3 \times S^1}^{\text{susy}} = \text{Tr} [(-)^F e^{-\beta H_{\text{susy}}}]$, where H_{susy} generates the time evolution along S^1 . Then the Casimir energy is extracted from the large circle limit as

$$E_{\text{susy}} = - \lim_{\beta \rightarrow \infty} \frac{d}{d\beta} \log Z_{M_3 \times S^1}^{\text{susy}}. \quad (2.2)$$

The scheme-independence of the partition function implies that the vacuum energy is universal (below we will also give a new proof of this fact). This should be contrasted with the non-universality of the ordinary Casimir energy E_0 discussed in the introduction.

2.2 Consequences of the supersymmetry algebra

In this section we make a few observations based on the supersymmetry algebra. In Lorentzian signature, let us consider the simple case $M_4 = S^3 \times \mathbb{R}$, where S^3 is a round three-sphere of radius r_3 and \mathbb{R} is the time direction parameterized by t (this is related to the Euclidean time τ in the introduction by $t = i\tau$). By fixing the other background fields to $A_t = V_t = \frac{1}{r_3}$, with all other components vanishing, we can preserve four supercharges⁶ for any $\mathcal{N} = 1$ theory with an R -symmetry [13]. In fact one could take any constant A_t for the flat gauge field. However, only in the case that $A_t = \frac{1}{r_3}$ one gets time-independent supercharges. So we will make this choice throughout.

The superalgebra preserved by the background is [22]

$$\begin{aligned} \frac{1}{2} \{Q_\alpha, Q^{\dagger\beta}\} &= \delta^\beta_\alpha \left(H_{\text{susy}} - \frac{1}{r_3} R \right) - \frac{2}{r_3} \sigma^i{}^\beta{}_\alpha J_l^i, \\ [H_{\text{susy}}, Q_\alpha] &= 0, \quad [R, Q_\alpha] = -Q_\alpha, \quad [J_l^i, Q_\alpha] = \frac{1}{2} Q_\beta \sigma^i{}^\beta{}_\alpha, \end{aligned} \quad (2.3)$$

where H_{susy} is the generator of translations along the circle, R is the R -charge, the J_l^i , $i = 1, 2, 3$, are the generators of the $\text{SU}(2)_l \subset \text{SU}(2)_l \times \text{SU}(2)_r$ isometry of the sphere and σ^i are the Pauli matrices. The supercharges Q_α , $\alpha = 1, 2$, form a doublet of $\text{SU}(2)_l$, while the $\text{SU}(2)_r$ subgroup does not appear in the superalgebra.

A first remark is that we assume *the vacuum does not break supersymmetry*. Suppose the vacuum were not supersymmetric, in which case either Q_1 , or Q_2 (or both) would not annihilate the vacuum. Then, $Q|\text{VAC}\rangle$ is a new state with the same value of H_{susy} , but contributing with an opposite sign to the index or partition function. Therefore if

⁵As in footnote 1, one should bear in mind that there is a dimension < 4 counterterm that does not vanish in general, but this contains a positive power of the cut-off. We will use such divergent counterterm in appendix C to regularize the partition function.

⁶This requires making a special choice of the so-called κ parameter [30]. For generic choices of κ , the background preserves only two supercharges.

supersymmetry were broken, the index on $S^3 \times \mathbb{R}$ would not receive a contribution from the unit operator. In the case of SCFTs, the fact that supersymmetry is unbroken on $S^3 \times \mathbb{R}$ is a simple consequence of radial quantization. Indeed, in this case, the spectrum of H_{susy} is identical to the spectrum of $\Delta - \frac{1}{2r_3}R$, which has a gap above the unit operator.

Another simple observation is that J_l^3 annihilates the vacuum, $J_l^3|\text{VAC}\rangle = 0$. Indeed, J_l^3 appears with different signs on the right hand side of $\{Q_1, Q_1^\dagger\}$ and $\{Q_2, Q_2^\dagger\}$. Hence, it must vanish or one of the Q 's does not annihilate the vacuum. (Also, if J_l^3 were nonzero on the vacuum, the vacuum would not be unique.)

It is useful to focus on the algebra of one specific supercharge, say Q_1 . We will also rescale R and J_l^3 to reabsorb the radius r_3 of S^3 . Then the Q_1 -algebra reads

$$\begin{aligned} \{Q_1, Q_1^\dagger\} &= 2(H_{\text{susy}} - R - 2J_l^3), & Q_1^2 &= 0, \\ [H_{\text{susy}}, Q_1] &= [R + 2J_l^3, Q_1] = 0. \end{aligned} \quad (2.4)$$

From this we conclude that

$$\langle H_{\text{susy}} \rangle = \langle R \rangle. \quad (2.5)$$

However, the algebra (2.4) is invariant under shifting H_{susy} and the R -charge by some c-number ϵ so we cannot yet determine the actual expectation value of H_{susy} in the vacuum. Notice that an equivalent way to express the Ward identity (2.5) is in terms of the vacuum expectation value of Δ , defined in (1.7):

$$\langle \Delta \rangle = \frac{3}{2} \langle R \rangle. \quad (2.6)$$

Equation (2.6) takes the form of the familiar BPS relation. Here we interpret it on the cylinder, where both sides have nonzero vacuum expectation values.

We will approach the problem of determining $\langle H_{\text{susy}} \rangle$ by reducing the theory on the three-sphere. In this way we get a quantum mechanics (QM) theory with infinitely many degrees of freedom. The theory has four supercharges, Q_1, Q_2 and their Hermitian conjugates. The R -symmetry group is $\text{SU}(2)_l \times \text{U}(1)$ and the supercharges furnish the $(2, 1)$ representation. $\text{SU}(2)_r$ is a global symmetry in the quantum mechanics theory.

The generating functional of R -current connected correlators discussed in (1.14) is specifically given by

$$\mathcal{W}[A_t] = \langle R \rangle \int dt A_t, \quad (2.7)$$

where A_t is the component along S^1 of the four-dimensional R -symmetry gauge field. This term is a 1d Chern-Simons term fixing the R -charge of the vacuum. Given that $\langle H_{\text{susy}} \rangle = \langle R \rangle$, this also fixes the Casimir energy. Since the relation $\langle H_{\text{susy}} \rangle = \langle R \rangle$ is a consequence of supersymmetry, the one-dimensional local term (2.7) must be part of a supersymmetric term in a one-dimensional supergravity, obtained by dimensional reduction of 4d new minimal supergravity on S^3 . In fact, in the specific case of round $S^1 \times S^3$ that we are discussing here, it is easy to see that a plausible candidate for the completion of the CS term (2.7) to a 1d supergravity invariant is given by

$$\mathcal{W} = \langle R \rangle \int dt \left(A_t + \frac{3}{2r_3} \sqrt{|g_{tt}|} - \frac{3}{2} V_t \right), \quad (2.8)$$

where g_{tt} and V_t are simply the components of the respective background fields in four dimensions. This expression does not depend on the choice of the parameter κ and correctly reproduces (2.6) in a SCFT. However, we leave the systematic study of such terms for the future.

Equation (2.7) implies that $\langle H_{\text{susy}} \rangle$ is *physical* (i.e. scheme independent) without relying on the classification of four-derivatives counterterms of [8]. The idea is that if we just studied a quantum mechanics theory of finitely many degrees of freedom, then there would be no meaningful way to compute $\langle H_{\text{susy}} \rangle$ itself. Indeed, we could always add a counterterm of the form (2.7) with an arbitrary coefficient, shifting the values of $\langle H_{\text{susy}} \rangle$ and $\langle R \rangle$ at will, while preserving $\langle H_{\text{susy}} \rangle = \langle R \rangle$.⁷ So without additional assumptions, there is no possible way to fix $\langle H_{\text{susy}} \rangle$ itself at the level of quantum mechanics. However, it is crucial that in the present study the quantum mechanics theory derives from a four-dimensional local quantum field theory. Then, the Chern-Simons counterterm (2.7) would be admissible only if it came from a local term in four dimensions. It is easy to convince oneself that there is no way to derive (2.7) — with its specific normalization — by dimensional reduction of a four-dimensional counterterm on $S^3 \times \mathbb{R}$. Indeed to get the normalization right for the Chern-Simons term, it needs to come from a term in four dimensions of mass dimension four. But then it is straightforward to see that one cannot write anything, *regardless of supersymmetry*, that would look like $\int dt A_t$ after integrating over the sphere with radius r_3 . Therefore, the charge of the vacuum becomes physical and so does the ground state energy by the relation $\langle H_{\text{susy}} \rangle = \langle R \rangle$.

Another observation that follows directly from (2.7) is that *the Casimir energy cannot depend on continuous coupling constants (and hence on the RG scale)*. This follows from the fact that $\langle H_{\text{susy}} \rangle = \langle R \rangle$ and that to compute $\langle R \rangle$ we need to evaluate the coefficient of $\int dt A_t$. If the coefficient of A_t had depended on continuous coupling constants, we could have promoted them to time-dependent fields and lose the gauge invariance under small R -symmetry gauge transformations (and this cannot be accounted for by any anomaly). This is similar to the arguments in [26] (and references therein). This conclusion is consistent with the arguments of [18, 19].

An important consequence of this observation is that it is sufficient to calculate the Casimir energy starting from a free field theory in $4d$. We henceforth assume that such a free point exists in the space of continuous coupling constants. It would be very interesting to generalize our considerations to non-Lagrangian theories. (It is not currently clear to us how to do so.)

To summarize, by considering the supersymmetry algebra and reducing to a quantum mechanical problem, we established that what we need to compute is the coefficient of the generating functional (2.7) in quantum mechanics. We perform this computation below, after having introduced some notions of supersymmetric quantum mechanics.

⁷This freedom is constrained when the R -symmetry is compact. In this case, we can only add (2.7) with an integer coefficient. Thus, we can only change the R charge by an integer amount, such that $e^{2\pi i R} = 1$ on all the states is retained. So if the R -symmetry were compact, then the ambiguity in the vacuum energy would be only by an integer, even in quantum mechanics.

Before coming to these issues, let us comment about the case in which two supercharges are preserved instead of four. This is in fact the generic supersymmetric case, pertinent to various deformations of S^3 (these will be discussed in section 3) as well as to other topologies. In this case, some of the claims above are valid while some others are not necessarily true. An important difference is that there is no $SU(2)_I$ algebra in general, rather, there is just some $U(1)$ isometry of M_3 , generated by J_3 . Thus, a simple argument that J_3 vanishes in the vacuum does not exist. The QM algebra, inherited from the $4d$ deformed algebra [17, 20], now takes the form (1.13), so that in the vacuum we have $\langle H_{\text{susy}} \rangle = \langle \Sigma \rangle$, with Σ a QM flavor symmetry. A quantum mechanical term like (2.7) would still exist, with A_t replaced by the gauge field A_t^Σ for the flavor symmetry Σ , and the claims made below eq. (2.7) still apply. In particular, the vacuum energy is still independent of the renormalization scheme and of the coupling constants. We will use this to compute E_{susy} for $N = 1$ theories with an R-symmetry on $M_3 \times S^1$, where M_3 has S^3 topology.

2.3 Supersymmetric quantum mechanics

Let us model the situation above, governed by the two supercharges algebra (2.4), with

$$\begin{aligned} \{Q, Q^\dagger\} &= 2(H - \Sigma), & Q^2 &= 0, \\ [H, Q] &= [\Sigma, Q] = 0, \end{aligned} \quad (2.9)$$

where H generates time translations, while Σ is some Hermitian conserved charge. At the formal level, we can just redefine H by Σ . However, in order to be able to connect more easily to the reduction over S^3 , we will keep the algebra in the form (2.4). Similar supersymmetric systems were studied in [37, 38].

We can define two types of multiplets: a *chiral multiplet* (ϕ, ψ) , and a *Fermi multiplet* (λ, f) , where ϕ, f are complex and commuting while ψ, λ are complex and anti-commuting. These two multiplets have the following supersymmetry transformations

$$\begin{aligned} \text{chiral :} \quad & \delta\phi = \sqrt{2}\zeta\psi, & \delta\psi &= -\sqrt{2}i\zeta^\dagger D_t\phi, \\ \text{Fermi :} \quad & \delta\lambda = \sqrt{2}\zeta f + p\sqrt{2}\zeta^\dagger\phi, & \delta f &= -\sqrt{2}i\zeta^\dagger D_t\lambda - p\sqrt{2}\zeta^\dagger\psi, \end{aligned} \quad (2.10)$$

where on all the fields we define $D_t = \partial_t - i\sigma$, with σ the charge of the field under Σ . The complex parameter ζ is independent of time and uncharged under Σ . In the variations of the Fermi multiplet there appears a parameter p . When $p = 0$, the chiral and Fermi multiplets are independent of each other. We will refer to each of the decoupled multiplets as “short.” When instead $p \neq 0$ the two multiplets form one reducible but indecomposable representation of supersymmetry. Thus, for $p \neq 0$ we call the combined chiral and Fermi multiplets a “long” multiplet.

On each component of a multiplet with charge σ , the transformations (2.10) give

$$\{\delta_1, \delta_2\} = -2i(\zeta_1^\dagger\zeta_2 + \zeta_2^\dagger\zeta_1)D_t, \quad (2.11)$$

which is consistent with the algebra (2.9) when H is represented as $-i\partial_t$.

The supersymmetric Lagrangian of a long multiplet takes the form

$$\begin{aligned}
 L = & |D_t \phi|^2 - i\mu(\phi D_t \phi^\dagger - \phi^\dagger D_t \phi) + i\psi^\dagger D_t \psi - 2\mu\psi\psi^\dagger \\
 & + i\lambda^\dagger D_t \lambda + |f|^2 \\
 & - p^2|\phi|^2 - p(\lambda\psi^\dagger + \psi\lambda^\dagger),
 \end{aligned} \tag{2.12}$$

where μ is an additional free parameter, giving a mass to ψ . For $p = 0$, the first and the second lines are the Lagrangians of a free chiral and free Fermi multiplet, respectively, and are separately supersymmetric.⁸

We now pass to Hamiltonian formalism and quantize the theory. The canonical momenta are

$$\Pi_\phi = (D_t + i\mu)\phi^\dagger, \quad \Pi_\psi = -i\psi^\dagger, \quad \Pi_\lambda = -i\lambda^\dagger, \quad \Pi_f = 0. \tag{2.13}$$

The canonical (anti-)commutation relations are

$$[\phi, \Pi_\phi] = i, \quad \{\psi, \Pi_\psi\} \equiv -i\{\psi, \psi^\dagger\} = -i, \quad \{\lambda, \Pi_\lambda\} \equiv -i\{\lambda, \lambda^\dagger\} = -i, \tag{2.14}$$

together with their Hermitian conjugates.

The Hamiltonian reads

$$\begin{aligned}
 H = & |\Pi_\phi|^2 + i(\mu + \sigma)(\Pi_\phi \phi - \phi^\dagger \Pi_{\phi^\dagger}) + \mu^2|\phi|^2 + (\sigma + 2\mu)\psi\psi^\dagger \\
 & + \sigma\lambda\lambda^\dagger \\
 & + p^2|\phi|^2 + p(\lambda\psi^\dagger + \psi\lambda^\dagger) + \tilde{\alpha},
 \end{aligned} \tag{2.15}$$

where again when $p = 0$ the first line gives the Hamiltonian of a chiral multiplet, while the second line is the Hamiltonian of a Fermi multiplet. The field f has been set to zero by its equation of motion. Note that we have introduced a constant $\tilde{\alpha}$, parameterizing the usual ordering ambiguity.

In terms of canonical variables, the charge Σ reads

$$\Sigma = i\sigma(\Pi_\phi \phi - \phi^\dagger \Pi_{\phi^\dagger}) + \sigma(\psi\psi^\dagger + \lambda\lambda^\dagger) + \alpha, \tag{2.16}$$

where α parameterizes the ordering ambiguity in this operator. The supercharge is

$$Q = \sqrt{2}i\psi(\Pi_\phi - i\mu\phi^\dagger) + \sqrt{2}p\phi^\dagger\lambda, \tag{2.17}$$

and is free of ordering ambiguities. Evaluating $\{Q, Q^\dagger\}$ we find that (2.9) is upheld provided we take

$$\tilde{\alpha} = \alpha - 2\mu. \tag{2.18}$$

Hence supersymmetry fixes the ordering ambiguity in $H - \Sigma$. Of course, after having solved for $\tilde{\alpha}$ we still have the freedom to shift H and Σ by an equal amount, corresponding to

⁸When $p = 0$ an additional term like $\delta(\lambda W(\phi))$ can be introduced, in case the total charge under Σ vanishes. We will not need to consider this term.

the remaining parameter α . Without additional assumptions, this freedom would have remained in the framework of ordinary quantum mechanics in one dimension.

In order to explain how to fix the ordering ambiguity that is left, it is useful to recall that we are computing the coefficient of a CS term in the low-energy $1d$ effective action. This term takes the form

$$k \int dt A_t^\Sigma, \quad (2.19)$$

where A_t^Σ is the background gauge field associated to the charge Σ . A single fermion of mass M and charge q shifts the coefficient of the Chern-Simons term by $\frac{q}{2}\text{sgn}(M)$ [39]. We can think about that as if we are starting from some theory in the UV with Chern-Simons coefficient k_{uv} and then we integrate out the massive fermion leading to a Chern-Simons coefficient in the infrared k_{ir} (this interpretation was elaborated upon in [40])⁹

$$k_{\text{ir}} - k_{\text{uv}} = -\frac{q}{2}\text{sgn}(M). \quad (2.22)$$

From the point of view of the quantum mechanics, the arbitrariness in the charge of the vacuum corresponds to the arbitrariness in the UV coefficient k_{uv} . However, our theory arises from a higher-dimensional model. As already observed, it is easy to convince oneself that a term like (2.19) cannot be generated by dimensional reduction of a four-dimensional local term. So we must take

$$k_{\text{uv}} = 0, \quad (2.23)$$

i.e. no Chern-Simons contact term in the UV generating functional. This key requirement fixes the ordering ambiguity in H . Together with (2.22), this implies that multiplets containing pairs of fermions with masses of opposite sign do not contribute to the Casimir energy. We will see below that as long as the Hamiltonian is bounded from below, a long multiplet necessarily contains fermions with masses of opposite sign. As a result, the choice of the ordering coefficient must be such that H and Σ vanish in the ground state of a long multiplet. This leads to the conclusion that the correct choice of the ordering constant is

$$\alpha = -2\sigma. \quad (2.24)$$

⁹A simple way to derive (2.22) is as follows. First, from dimensional analysis and the fact that M and k are odd under charge conjugation we infer

$$k_{\text{ir}} - k_{\text{uv}} = x \text{sgn}(M), \quad (2.20)$$

where x is a coefficient, independent of M . To fix x we can consider a free fermion with mass M and charge q with a constant background gauge field A_t^Σ . This has Hamiltonian $H = (M + qA_t^\Sigma)(\psi\psi^\dagger + \hat{\alpha})$, where $\hat{\alpha}$ is an arbitrary ordering constant. The partition function is given by

$$Z = e^{-\beta(M+qA_t^\Sigma)\hat{\alpha}} \left(1 + e^{-\beta(M+qA_t^\Sigma)} \right). \quad (2.21)$$

The idea now is that we can keep the ultraviolet fixed and consider two different RG flows, one with positive M and one with negative M . By subtracting the resulting Chern-Simons terms in the infrared (which we will read out from the charge of the vacuum), we will find $2x$. If $M > 0$ then taking $M \rightarrow \infty$ we can read off the CS term (i.e. charge) in the IR to be $q\hat{\alpha} \int dt A_t^\Sigma$. On the other hand, if $M < 0$ we read out the CS term in the IR by taking the limit $M \rightarrow -\infty$ and we find $q(\hat{\alpha} + 1) \int dt A_t^\Sigma$. Subtracting these yields $2x = -q$.

We will use this choice in the following and one can verify that in all cases the results are consistent with (2.23). Incidentally, it turns out that (2.24) also corresponds to Weyl ordering for the Hamiltonian.¹⁰

2.4 Spectrum of the Hamiltonian

We now study the spectrum of the Hamiltonian and determine the vacuum state.

Long multiplet. Let us start from the bosonic sector of (2.15):

$$H_{\text{bosonic}} = |\Pi_\phi|^2 + i(\mu + \sigma)(\Pi_\phi \phi - \phi^\dagger \Pi_{\phi^\dagger}) + (\mu^2 + p^2)|\phi|^2 - \mu - \sigma, \quad (2.25)$$

where we have included half of the ordering constant appearing there (the other half will enter in the fermionic sector). This ensures Weyl ordering. We can introduce creation operators a^\dagger, b^\dagger and annihilation operators a, b via

$$\phi = \frac{(\mu^2 + p^2)^{-1/4}}{\sqrt{2}}(a + b^\dagger), \quad \Pi_\phi = \frac{i(\mu^2 + p^2)^{1/4}}{\sqrt{2}}(a^\dagger - b). \quad (2.26)$$

The canonical commutation relations between ϕ and Π_ϕ (and their Hermitian conjugates) imply that these satisfy $[a, a^\dagger] = [b, b^\dagger] = 1$, $[a, b] = [a^\dagger, b] = [a, b^\dagger] = [a^\dagger, b^\dagger] = 0$. Then the bosonic Hamiltonian can be written as

$$\begin{aligned} H_{\text{bosonic}} &= \sqrt{\mu^2 + p^2}(a^\dagger a + b^\dagger b + 1) + (\sigma + \mu)(b^\dagger b - a^\dagger a) \\ &= \frac{1}{2}\sqrt{\mu^2 + p^2}(\{a, a^\dagger\} + \{b, b^\dagger\}) + \frac{1}{2}(\sigma + \mu)(\{b, b^\dagger\} - \{a, a^\dagger\}), \end{aligned} \quad (2.27)$$

where in the second line we have emphasized that H_{bosonic} is Weyl ordered. The state annihilated by a and b has energy $\sqrt{\mu^2 + p^2}$. Acting on this with $(a^\dagger)^m(b^\dagger)^n$ (with m, n positive integers) we obtain a state with energy

$$H_{\text{bosonic}}(m, n) = \sqrt{\mu^2 + p^2} + m(\sqrt{\mu^2 + p^2} - \mu - \sigma) + n(\sqrt{\mu^2 + p^2} + \mu + \sigma). \quad (2.28)$$

We see that in order for the Hamiltonian to have a spectrum that is bounded from below we need to assume $\sqrt{\mu^2 + p^2} > |\mu + \sigma|$.¹¹ Hence the state of minimum energy in the bosonic sector is the one with $m = n = 0$.

Next we address the fermionic sector. The Hamiltonian reads

$$\begin{aligned} H_{\text{fermionic}} &= p(\lambda\psi^\dagger + \psi\lambda^\dagger) + (2\mu + \sigma)\psi\psi^\dagger + \sigma\lambda\lambda^\dagger - \mu - \sigma \\ &= \begin{pmatrix} \psi & \lambda \end{pmatrix} \begin{pmatrix} 2\mu + \sigma & p \\ p & \sigma \end{pmatrix} \begin{pmatrix} \psi^\dagger \\ \lambda^\dagger \end{pmatrix} - \mu - \sigma, \end{aligned} \quad (2.29)$$

¹⁰This explains why our final result is identical to that of [30] for the VEV of H . But, unlike [30], our result for the VEV of Σ in the vacuum manifestly respects the BPS condition $H = \Sigma$.

¹¹Allowing for $\sqrt{\mu^2 + p^2} = |\mu + \sigma|$ yields a Hamiltonian bounded from below but introduces a degenerate vacuum. Let us discard this case.

where we have kept the ordering constant that ensures Weyl ordering. We can make a unitary $U(2)$ rotation to diagonalize the above matrix. This preserves the anti-commutation relations. The eigenvalues are

$$x_{\pm} = \mu + \sigma \pm \sqrt{\mu^2 + p^2}. \quad (2.30)$$

Denoting the eigenvectors $u_+, u_-, u_+^\dagger, u_-^\dagger$, the Hamiltonian is thus

$$\begin{aligned} H_{\text{fermionic}} &= x_+ u_+ u_+^\dagger + x_- u_- u_-^\dagger - \mu - \sigma \\ &= \frac{x_+}{2} [u_+, u_+^\dagger] + \frac{x_-}{2} [u_-, u_-^\dagger], \end{aligned} \quad (2.31)$$

with $\{u_{\pm}, u_{\pm}^\dagger\} = 1$. The charge operator Σ takes the form

$$\begin{aligned} \Sigma_{\text{fermionic}} &= \sigma (u_+ u_+^\dagger + u_- u_-^\dagger - 1) \\ &= \sigma [u_+, u_+^\dagger] + \sigma [u_-, u_-^\dagger]. \end{aligned} \quad (2.32)$$

Starting with the state $|0\rangle$ which is annihilated by both u_{\pm}^\dagger , we can act with u_- , u_+ or $u_- u_+$. The spectrum therefore consists of four states with the following energy and charge:

state	$ 0\rangle$	$u_- 0\rangle$	$u_+ 0\rangle$	$u_+ u_- 0\rangle$
energy	$-\mu - \sigma$	$-\sqrt{\mu^2 + p^2}$	$\sqrt{\mu^2 + p^2}$	$\mu + \sigma$
charge	$-\sigma$	0	0	σ

(2.33)

Since we assumed $\sqrt{\mu^2 + p^2} > |\mu + \sigma|$, the state of lowest energy is $u_-|0\rangle$.

We now combine the information obtained studying the bosonic and fermionic sectors of the Hamiltonian and identify a state with minimum energy that respects supersymmetry. Adding H_{bosonic} and $H_{\text{fermionic}}$, the complete Hamiltonian is

$$\begin{aligned} H &= \sqrt{\mu^2 + p^2} (a^\dagger a + b^\dagger b + 1) + (\sigma + \mu) (b^\dagger b - a^\dagger a) \\ &\quad + x_+ u_+ u_+^\dagger + x_- u_- u_-^\dagger - \mu - \sigma. \end{aligned} \quad (2.34)$$

One can also check that the full charge operator reads

$$\Sigma = \sigma (b^\dagger b - a^\dagger a + u_+ u_+^\dagger + u_- u_-^\dagger - 1). \quad (2.35)$$

From the discussion above, the state with minimum energy is clearly

$$|\text{VAC}\rangle \equiv |m=0, n=0, x_-\rangle, \quad (2.36)$$

where $m=0, n=0$ indicates that no bosonic oscillators are excited, and by x_- we mean that we excite one fermionic oscillator with eigenvalue x_- . Its total energy is

$$H = \sqrt{\mu^2 + p^2} - \sqrt{\mu^2 + p^2} = 0, \quad (2.37)$$

and thus vanishes due to an exact cancellation between the bosonic and the fermionic contributions. Since we have just one fermionic oscillator the charge is $\Sigma = 0$, hence the relation $(H - \Sigma)|\text{VAC}\rangle = 0$ is satisfied and supersymmetry is unbroken in the vacuum, as expected.

We conclude that the long multiplets yield a vanishing contribution to the vacuum energy and charge:

$$\langle H_{\text{long}} \rangle = \langle \Sigma_{\text{long}} \rangle = 0. \quad (2.38)$$

Note that this is a consequence of our choice of ordering constant, and as argued at the end of the previous subsection this is the correct choice for a quantum mechanics arising from a higher-dimensional theory.

If we had a theory of long multiplets only, the vacuum energy would just be zero. However, if short multiplets are also present, this is not the case, as we now show.

Fermi multiplet. Consider the Fermi multiplet. Then the supercharge identically vanishes. The Hamiltonian and the charge generator take the same form,

$$H_{\text{Fermi}} = \Sigma_{\text{Fermi}} = \sigma \left(\lambda \lambda^\dagger - \frac{1}{2} \right). \quad (2.39)$$

The only two states have energy $-\frac{1}{2}\sigma$ and $+\frac{1}{2}\sigma$. The contribution of a Fermi multiplet to the vacuum energy and charge is thus

$$\langle H_{\text{Fermi}} \rangle = \langle \Sigma_{\text{Fermi}} \rangle = -\frac{|\sigma|}{2}. \quad (2.40)$$

Chiral multiplet. The bosonic sector of the chiral multiplet can be treated as we did for the long multiplet, setting $p = 0$. The full Hamiltonian and charge operator can thus be written as

$$H_{\text{chiral}} = |\mu|(a^\dagger a + b^\dagger b + 1) + (\sigma + \mu)(b^\dagger b - a^\dagger a) + (2\mu + \sigma)\psi\psi^\dagger - \mu - \frac{\sigma}{2}, \quad (2.41)$$

$$\Sigma_{\text{chiral}} = \sigma(b^\dagger b - a^\dagger a) + \sigma\psi\psi^\dagger - \frac{1}{2}\sigma. \quad (2.42)$$

Since $p = 0$, the condition for the Hamiltonian to be bound from below becomes

$$|\mu| > |\mu + \sigma|. \quad (2.43)$$

In the vacuum all bosonic oscillators are zero. Then we have two possible states:

1. the state annihilated by ψ^\dagger , with $H = |\mu| - \mu - \frac{1}{2}\sigma$ and $\Sigma = -\frac{1}{2}\sigma$;
2. the state with an oscillator ψ excited, with $H = |\mu| + \mu + \frac{1}{2}\sigma$ and $\Sigma = +\frac{1}{2}\sigma$.

Which state has minimum energy depends on the values of μ and σ . Note that (2.43) requires μ and σ to have opposite signs. If $\mu > 0$, $\sigma < 0$, then (2.43) implies $-2\mu < \sigma < 0$, and the state number 1 has minimum energy $H = -\frac{1}{2}\sigma$; since $H = \Sigma$, this state is supersymmetric, while the state 2 is non-supersymmetric. Conversely, if $\mu < 0$ and $\sigma > 0$, then from (2.43) we deduce $0 < \sigma < -2\mu$, hence the state number 1 now has higher energy and the state 2 is the supersymmetric vacuum, with $H = \frac{1}{2}\sigma$.

Thus, a chiral multiplet contributes to the vacuum energy and charge as

$$\langle H_{\text{chiral}} \rangle = \langle \Sigma_{\text{chiral}} \rangle = \frac{|\sigma|}{2}. \quad (2.44)$$

In conclusion, the analysis in supersymmetric quantum mechanics establishes that a long multiplet yields a vanishing contribution to the vacuum energy and charge, that a Fermi multiplet contributes as in (2.40) while a chiral multiplet contributes as in (2.44).

2.5 Dimensional reduction of a 4d chiral multiplet

Consider a free four-dimensional chiral multiplet (ϕ, ψ_α, F) on $S^3 \times \mathbb{R}$. The Lagrangian and supersymmetry transformations can be found in [13]. The only parameter appearing in the Lagrangian is the charge r under the background R -symmetry gauge field. Here we will restrict to $0 < r \leq 2$.¹² This range is compatible with the inequalities mentioned in the previous subsection, ensuring that the spectrum of the Hamiltonian is bounded from below. Expanding in appropriate spherical harmonics, the chiral multiplet reduces to a one-dimensional theory with infinitely many fields. These organize in one-dimensional multiplets with different values of the parameters μ, p, σ introduced above. Some have $p \neq 0$ and are thus long multiplets, while some others have $p = 0$ and are thus short multiplets, either chiral or Fermi.

More explicitly, we can expand the scalars in spherical harmonics $Y_{\ell, m, n}$ transforming in representations $(\frac{\ell}{2}, \frac{\ell}{2})$ of $SU(2)_l \times SU(2)_r$. The quantum number ℓ is a non-negative integer. For a fixed ℓ , the quantum numbers m, n of the scalar harmonic $Y_{\ell, m, n}$ range in $-\frac{\ell}{2} \leq m, n \leq \frac{\ell}{2}$. So we can write

$$\phi = \sum_{\ell, m, n} \phi_{\ell, m, n} Y_{\ell, m, n}, \quad (2.45)$$

and similarly for the auxiliary field F . The fermionic field ψ_α can be expanded in spinorial harmonics. A review of spinor spherical harmonics on S^3 can be found in [30, 41]. A single 4d fermion reduces to two infinite series of 1d fermions furnishing the representation $\sum_l (\frac{\ell-1}{2}, \frac{\ell}{2}) \oplus (\frac{\ell+1}{2}, \frac{\ell}{2})$ of $SU(2)_l \times SU(2)_r$.¹³

Integrating over S^3 and using the orthonormality of the spherical harmonics, the action of a four-dimensional chiral multiplet gives rise to a one-dimensional action for an infinite number of fields. These arrange in multiplets of supersymmetric quantum mechanics labeled by ℓ, m, n , and one can check that the Lagrangian of each of these multiplets takes the form (2.12). Here we do not need to present all details of the reduction. All we need to know is how the R -charge r and the quantum numbers ℓ, m, n map into the parameters σ, p, μ entering in (2.12) and characterizing each multiplet in supersymmetric quantum mechanics. Actually, the discussion in subsection 2.4 shows that for the purpose of determining the vacuum energy we just need to know when a multiplet is shortened (namely when $p = 0$), if it is a chiral or a Fermi multiplet, and what is the value of its charge σ .

¹²Outside this range there are complications, see [41]. Here we see additional complications, for example, the cancelation previously discussed for long multiplets would fail.

¹³The symmetry between left and right is broken by the choice of spin bundle.

By comparing the four-dimensional algebra (2.4) with (2.9), we deduce that we must identify (restoring the S^3 radius r_3) $\Sigma = \frac{1}{r_3}(R + 2J_l^3)$, and therefore

$$\sigma = \frac{1}{r_3}(r + 2m). \quad (2.46)$$

Moreover, reducing the four-dimensional Lagrangian to one dimension, one finds¹⁴

$$\begin{aligned} p^2 &= \frac{1}{r_3^2}(\ell - 2m)(\ell + 2 + 2m), \\ \mu &= -\frac{1}{r_3}(2m + 1), \end{aligned} \quad (2.47)$$

hence the shortening condition $p = 0$ is satisfied if and only if $m = \ell/2$ or $m = -\ell/2 - 1$. In the former case a chiral multiplet is obtained, with charge $\sigma = \frac{1}{r_3}(\ell + r)$. In the latter case a Fermi multiplet is obtained, with charge $\sigma = -\frac{1}{r_3}(\ell + 2 - r)$. Recalling (2.40), (2.44) we conclude that the respective contribution to the vacuum energy is:

$$\begin{aligned} \text{chiral} \quad \left(m = \frac{\ell}{2}\right) : \quad \langle H_{\text{chiral}} \rangle &= \frac{1}{2r_3}(\ell + r), \\ \text{Fermi} \quad \left(m = -\frac{\ell}{2} - 1\right) : \quad \langle H_{\text{Fermi}} \rangle &= -\frac{1}{2r_3}(\ell + 2 - r). \end{aligned} \quad (2.48)$$

The expectation value of the Hamiltonian is obtained by adding up the contributions of all chiral and Fermi multiplets:

$$\begin{aligned} \langle H_{\text{susy}} \rangle &= \sum_{\text{chiral}} \langle H_{\text{chiral}} \rangle + \sum_{\text{Fermi}} \langle H_{\text{Fermi}} \rangle \\ &= \sum_{\ell \geq 0} \frac{1}{2r_3}(\ell + 1)(\ell + r) - \sum_{\ell \geq 0} \frac{1}{2r_3}(\ell + 1)(\ell + 2 - r), \end{aligned} \quad (2.49)$$

where the $(\ell + 1)$ factor comes from the degeneracy associated with $\text{SU}(2)_r$.

To regularize the sum, we dress the terms in the sum with some decreasing weights. To do this in a supersymmetric fashion, we can decompose H as a sum of Hamiltonians acting on the Hilbert space of a single free $1d$ multiplet

$$H_{\text{susy}} = \sum_{\ell, m, n} H_{\ell, m, n}, \quad (2.50)$$

and regularize the sum with a function of the $H_{\ell, m, n}$ operators, for instance

$$H_{\text{susy}} = \sum_{\ell, m, n} H_{\ell, m, n} e^{-2tr_3|H_{\ell, m, n}|}, \quad (2.51)$$

¹⁴More generally, one could easily restore the dependence on the parameter κ . This affects only μ but not p^2 and σ . In the notation of [30] one finds that the parameter μ is related with the parameters in the four-dimensional Lagrangian as $r_3\mu = -2m - \frac{3}{2}r - \kappa(\frac{3}{2}r - \epsilon)$.

with t a positive number. This yields

$$\langle H_{\text{susy}} \rangle = \sum_{\ell \geq 0} \frac{1}{2r_3} (\ell + 1)(\ell + r) e^{-t(\ell+r)} - \sum_{\ell \geq 0} \frac{1}{2r_3} (\ell + 1)(\ell + 2 - r) e^{-t(\ell+2-r)}. \quad (2.52)$$

Taking the small t limit and dropping the diverging term in t^{-2} ,¹⁵ we obtain a regularized result for the vacuum energy that, after recalling the trace anomaly coefficients [42]

$$a = \frac{3}{32} [3(r-1)^3 - (r-1)] , \quad c = \frac{1}{32} [9(r-1)^3 - 5(r-1)] , \quad (2.53)$$

reads

$$E_{\text{susy}} = \langle H_{\text{susy}} \rangle = \frac{4}{27r_3} (a + 3c). \quad (2.54)$$

This is the result advertised in eq. (1.9) of the introduction.

One could consider a supersymmetric regularization with a different function $f(tH_{\ell,m,n})$ of the $H_{\ell,m,n}$ operators. It can be shown, using an Euler-MacLaurin expansion (see appendix C for a related application) that for all smooth functions f such that $f(0) = 1$ (and such that the series converges), one obtains the same result for the finite piece in the small t expansion. This is in agreement with the fact that the supersymmetric Casimir energy is unambiguous. The regularization using the Hurwitz zeta function [30] also reproduces the same result.

It is possible to contrast our results with several previous works in which localization techniques on $S^3 \times S^1$ were utilized. Comparing with [20] (see also [32] and [43] where similar localization techniques are used in other topologies), one finds agreement regarding the vacuum energy. However, as we will briefly discuss in appendix C, the regularization scheme of [20] in fact does not preserve supersymmetry, as it violates certain SUSY Ward identities in the small circle limit. Our result for the vacuum energy also agrees with that of [30] (this method is a Hamiltonian version of [21]), but as mentioned in footnote 10 the result of [30] does not preserve some SUSY Ward identities as well.

3 Supersymmetric Casimir energy on a deformed three-sphere

In this section we study the chiral multiplet on a supersymmetric $S^3 \times S^1$ background with more general metric and complex structure. We will use results known from the computation of the partition function on these spaces, based on localization, to implement the Hamiltonian approach to the evaluation of the Casimir energy.

3.1 Shortening conditions on chiral multiplets

We start by reviewing some results of [32], where the fermionic degrees of freedom in the chiral multiplet were conveniently redefined. This made it particularly easy to show that in backgrounds preserving two supercharges of opposite R -charge, the modes contributing

¹⁵The diverging term can be associated to the four-dimensional Einstein-Hilbert counterterm. We will discuss this more in appendix C.

to the partition function solve first-order differential equations that can be interpreted as shortening conditions.

We work in Euclidean signature and follow the conventions of [20].¹⁶ We assume the existence of at least one positive-chirality spinor ζ_α and one negative-chirality spinor $\tilde{\zeta}^{\dot{\alpha}}$ satisfying the Killing spinor equations

$$\begin{aligned}(\nabla_\mu - iA_\mu)\zeta + iV_\mu\zeta + iV^\nu\sigma_{\mu\nu}\zeta &= 0, \\ (\nabla_\mu + iA_\mu)\tilde{\zeta} - iV_\mu\tilde{\zeta} - iV^\nu\tilde{\sigma}_{\mu\nu}\tilde{\zeta} &= 0.\end{aligned}\tag{3.1}$$

These independent equations are the Euclidean version of the supersymmetry condition given in (2.1). Note that ζ has R -charge $+1$ while $\tilde{\zeta}$ has R -charge -1 . In Euclidean signature the chiral multiplet is made of (ϕ, ψ_α, F) with R -charge $(r, r-1, r-2)$ and of the independent fields $(\tilde{\phi}, \tilde{\psi}^{\dot{\alpha}}, \tilde{F})$ with R -charge $(-r, -r+1, -r+2)$. The supersymmetry transformations are

$$\begin{aligned}\delta\phi &= \sqrt{2}\zeta\psi, & \delta\tilde{\phi} &= \sqrt{2}\tilde{\zeta}\tilde{\psi}, \\ \delta\psi &= \sqrt{2}F\zeta + \sqrt{2}i\sigma^\mu\tilde{\zeta}D_\mu\phi, & \delta\tilde{\psi} &= \sqrt{2}\tilde{F}\tilde{\zeta} + \sqrt{2}i\tilde{\sigma}^\mu\zeta D_\mu\tilde{\phi}, \\ \delta F &= \sqrt{2}iD_\mu(\tilde{\zeta}\tilde{\sigma}^\mu\psi), & \delta\tilde{F} &= \sqrt{2}iD_\mu(\zeta\sigma^\mu\tilde{\psi}),\end{aligned}\tag{3.2}$$

where on a field of R -charge q the covariant derivative D_μ is defined as

$$D_\mu = \nabla_\mu - iqA_\mu,\tag{3.3}$$

with ∇_μ the Levi-Civita connection. A supersymmetric Lagrangian is

$$\begin{aligned}\mathcal{L} &= D_\mu\tilde{\phi}D^\mu\phi + V^\mu(iD_\mu\tilde{\phi}\phi - i\tilde{\phi}D_\mu\phi) + \frac{r}{4}(R + 6V_\mu V^\mu)\tilde{\phi}\phi - \tilde{F}F \\ &\quad + i\tilde{\psi}\tilde{\sigma}^\mu D_\mu\psi + \frac{1}{2}V^\mu\tilde{\psi}\tilde{\sigma}_\mu\psi,\end{aligned}\tag{3.4}$$

where R is the Ricci scalar on the four-manifold.

Following [32] (see section 5 therein), we decompose the fermion fields $\psi, \tilde{\psi}$ in anti-commuting scalars as¹⁷

$$B = \frac{1}{\sqrt{2}}\frac{\zeta^\dagger\psi}{|\zeta|^2}, \quad C = \sqrt{2}\zeta\psi \quad \Leftrightarrow \quad \psi_\alpha = \sqrt{2}\zeta_\alpha B - \frac{1}{\sqrt{2}}\frac{\varepsilon_{\alpha\beta}\zeta^{\dagger\beta}}{|\zeta|^2}C,\tag{3.5}$$

$$\tilde{B} = \frac{1}{\sqrt{2}}\frac{\tilde{\zeta}^\dagger\tilde{\psi}}{|\tilde{\zeta}|^2}, \quad \tilde{C} = \sqrt{2}\tilde{\zeta}\tilde{\psi} \quad \Leftrightarrow \quad \tilde{\psi}^{\dot{\alpha}} = \sqrt{2}\tilde{\zeta}^{\dot{\alpha}}\tilde{B} - \frac{1}{\sqrt{2}}\frac{\varepsilon^{\dot{\alpha}\dot{\beta}}\tilde{\zeta}_{\dot{\beta}}^\dagger}{|\tilde{\zeta}|^2}\tilde{C}.\tag{3.6}$$

Note that B has R -charge $r-2$ while C has R -charge r . Similarly, \tilde{B} has R -charge $-r+2$ while \tilde{C} has R -charge $-r$. We also introduce the complex vectors

$$K^\mu = \tilde{\zeta}\tilde{\sigma}^\mu\zeta, \quad \bar{K}^\mu = \frac{\tilde{\zeta}^\dagger\tilde{\sigma}^\mu\zeta^\dagger}{4|\zeta|^2|\tilde{\zeta}|^2}, \quad Y^\mu = \frac{\tilde{\zeta}^\dagger\tilde{\sigma}^\mu\zeta}{2|\zeta|^2}, \quad \bar{Y}^\mu = -\frac{\tilde{\zeta}\tilde{\sigma}^\mu\zeta^\dagger}{2|\zeta|^2},\tag{3.7}$$

¹⁶We could also work in Lorentzian signature, building on the results of [36]. One reason for choosing the Euclidean signature is that it allows to make contact with the computation of the partition function via localization.

¹⁷Note that ζ_α and $\varepsilon_{\alpha\beta}(\zeta^\dagger)^\beta$ form a basis of chiral spinors.

which define a complex frame. These satisfy $K_\mu \bar{K}^\mu = Y_\mu \bar{Y}^\mu = \frac{1}{2}$, with all other contractions vanishing. The vectors K, \bar{K} have vanishing R -charge, while Y has R -charge $+2$ and \bar{Y} has R -charge -2 . They satisfy

$$\nabla_{(\mu} K_{\nu)} = 0, \quad \nabla_\mu \bar{K}^\mu = 0, \quad D_\mu Y^\mu = 0, \quad D_\mu \bar{Y}^\mu = 0, \quad (3.8)$$

hence K is a (complex) Killing vector. Finally, we define the differential operators

$$\hat{\mathcal{L}}_K = K^\mu D_\mu, \quad \hat{\mathcal{L}}_{\bar{K}} = \bar{K}^\mu D_\mu, \quad \hat{\mathcal{L}}_Y = Y^\mu D_\mu, \quad \hat{\mathcal{L}}_{\bar{Y}} = \bar{Y}^\mu D_\mu. \quad (3.9)$$

In the new variables the supersymmetry transformations (3.2) take the form (here we distinguish between $\delta \equiv \delta_\zeta$ and $\tilde{\delta} \equiv \delta_{\tilde{\zeta}}$):

$$\begin{aligned} \delta\phi &= C, & \tilde{\delta}\phi &= 0, \\ \delta C &= 0, & \tilde{\delta}C &= 2i \hat{\mathcal{L}}_K \phi, \\ \delta B &= F, & \tilde{\delta}B &= -2i \hat{\mathcal{L}}_{\bar{Y}} \phi, \\ \delta F &= 0, & \tilde{\delta}F &= 2i (\hat{\mathcal{L}}_K B + \hat{\mathcal{L}}_{\bar{Y}} C), \end{aligned} \quad (3.10)$$

with similar transformations for the tilded fields. The Lagrangian (3.4) can then be rewritten as

$$\begin{aligned} \mathcal{L} &= 4 \mathcal{L}_{\bar{K}} \tilde{\phi} \hat{\mathcal{L}}_K \phi + 4 \mathcal{L}_Y \tilde{\phi} \hat{\mathcal{L}}_{\bar{Y}} \phi + i\kappa (\hat{\mathcal{L}}_K \tilde{\phi} \phi - \tilde{\phi} \hat{\mathcal{L}}_K \phi) - \tilde{F} F \\ &\quad + 2i \tilde{B} \hat{\mathcal{L}}_K B + 2i \tilde{C} \hat{\mathcal{L}}_{\bar{K}} C + 2i \tilde{B} \hat{\mathcal{L}}_{\bar{Y}} C - 2i \tilde{C} \hat{\mathcal{L}}_Y B - \kappa \tilde{C} C. \end{aligned} \quad (3.11)$$

Here, κ is a function describing a redundancy in the choice of the background fields [17]; it will play no important role for us as it drops from the final answer.

In [32] it was showed that in a background with two supercharges $\zeta, \tilde{\zeta}$, the partition function of a chiral multiplet reduces to

$$Z = \frac{\prod \lambda^B}{\prod \lambda^\phi}, \quad (3.12)$$

where λ^B, λ^ϕ are eigenvalues determined by the first-order differential conditions

$$\hat{\mathcal{L}}_Y B = 0, \quad i \hat{\mathcal{L}}_K B = \lambda^B B, \quad (3.13)$$

and

$$\hat{\mathcal{L}}_{\bar{Y}} \phi = 0, \quad i \hat{\mathcal{L}}_K \phi = \lambda^\phi \phi. \quad (3.14)$$

The equations on the left can be read as shortening conditions for the chiral multiplet. Indeed, $\hat{\mathcal{L}}_Y B = 0$ allows to set C (and ϕ) to zero consistently with its equation of motion. Similarly, $\hat{\mathcal{L}}_{\bar{Y}} \phi = 0$ permits to set B to zero consistently with its supersymmetry transformation in (3.10). In order to set $B = 0$ consistently with its equation of motion one also needs $\hat{\mathcal{L}}_{\bar{Y}} C = 0$; this condition is also needed to set $F = 0$ respecting its supersymmetry

variation. Similar considerations apply to the tilded fields. (In [32], the latter were related to the untilded fields via Hermitian conjugation).

These shortenings are better understood by reducing to lower dimensions, where (B, F) and (ϕ, C) define irreducible representations of supersymmetry. In [32], this was done in an $S^2 \times \mathbb{T}^2$ background by expanding in monopole harmonics on S^2 and noting that the aforementioned equations correspond to shortenings of the $(0, 2)$ supersymmetry multiplets on \mathbb{T}^2 . Therefore only modes in shortened multiplets contribute to the one-loop determinant of the chiral multiplet partition function. The long multiplets lead to paired bosonic and fermionic eigenmodes and hence yield a trivial contribution.

Similarly, in the following we consider $S^3 \times S^1$ with a general metric and reduce on S^3 to supersymmetric quantum mechanics. This allows to make contact with the approach to the computation of the Casimir energy developed in section 2.

3.2 Reduction on deformed three-sphere

We consider the class of supersymmetric backgrounds studied in [20]. These are complex manifolds with the topology of $S^3 \times S^1$ known as primary Hopf surfaces. The complex structure moduli space is described by two parameters p, q . In the notation of [20], $p = e^{-2\pi|b_1|}$, $q = e^{-2\pi|b_2|}$ and b_1, b_2 are chosen real. The circle S^1 is parameterized by $\tau \sim \tau + 2\pi$. The three-sphere is described as a torus fibration over an interval: the torus angular coordinates are $\varphi_1 \sim \varphi_1 + 2\pi$ and $\varphi_2 \sim \varphi_2 + 2\pi$, while the interval coordinate is $\rho \in [0, 1]$. The metric is

$$ds^2 = \Omega^2 d\tau^2 + ds^2(M_3) = \Omega^2 d\tau^2 + f^2 d\rho^2 + m_{IJ} d\varphi_I d\varphi_J, \quad (3.15)$$

where $I, J = 1, 2$. Supersymmetry imposes the constraint

$$\Omega^2 = b^I m_{IJ} b^J, \quad (3.16)$$

which ensures Hermiticity of the metric. f and m_{IJ} are arbitrary functions of ρ , except that m_{IJ} must be positive definite and suitable boundary conditions need to be satisfied at the extrema of the interval. As $\rho \rightarrow 0$, we require that

$$f \rightarrow f_2, \quad m_{11} \rightarrow m_{11}(0), \quad m_{22} = (f_2 \rho)^2 + \mathcal{O}(\rho^3), \quad m_{12} = \mathcal{O}(\rho^2), \quad (3.17)$$

where $f_2 > 0$ and $m_{11}(0) > 0$ are constants. Similar boundary conditions are taken at $\rho \rightarrow 1$. The background field A is given by

$$A = \frac{|b_1 b_2|}{2\Omega^2 f} (\Omega \sqrt{m})' \left(\frac{d\varphi_1}{b_1} - \frac{d\varphi_2}{b_2} \right) + \frac{1}{2} d\omega, \quad (3.18)$$

where $m = \det m_{IJ}$, a prime denotes derivative with respect to ρ , and

$$\omega = \text{sgn}(b_1) \varphi_1 + \text{sgn}(b_2) \varphi_2. \quad (3.19)$$

We will not need the expression of the remaining background field V .

The class of three-sphere metrics $ds^2(M_3)$ in (3.15) includes the elliptically squashed three-sphere S^3_b , first studied in the context of localization in [44]. This is obtained by

setting $b_1 = \frac{\beta \mathfrak{b}}{2\pi r_3}$, $b_2 = \frac{\beta}{2\pi r_3 \mathfrak{b}}$, redefining the coordinate ρ into a coordinate $\vartheta \in [0, \pi/2]$ such that $f d\rho = [\mathfrak{b}^{-2} \sin^2 \vartheta + \mathfrak{b}^2 \cos^2 \vartheta]^{1/2} d\vartheta$, and taking $m_{11} = \mathfrak{b}^{-2} \cos^2 \vartheta$, $m_{22} = \mathfrak{b}^2 \sin^2 \vartheta$, $m_{12} = 0$. With these choices, $\Omega = (\frac{\beta}{2\pi r_3})^2$ and the expressions of the background fields also simplify. Everything that we will say applies to this particular case.

The complex vectors K and Y defined in (3.7) take the form

$$K = \frac{1}{2} \left[b_1 \frac{\partial}{\partial \varphi_1} + b_2 \frac{\partial}{\partial \varphi_2} - i \frac{\partial}{\partial \tau} \right], \quad (3.20)$$

$$Y = -\frac{e^{i\omega}}{2} \left[\frac{1}{f} \frac{\partial}{\partial \rho} + i \frac{\text{sgn}(b_1 b_2)}{\Omega \sqrt{m}} b_I m_{IJ} \varepsilon^{JK} \frac{\partial}{\partial \varphi_K} \right], \quad (3.21)$$

where ε^{JK} is the antisymmetric symbol, with $\varepsilon^{12} = +1$. In order to evaluate the differential operators (3.9) it will be useful to record that

$$Y^\mu A_\mu = \frac{i}{4} e^{i\omega} \left[\frac{1}{f} (\log(\Omega \sqrt{m}))' - \frac{\text{sgn}(b_1 b_2)}{\Omega \sqrt{m}} b_I m_{IJ} \varepsilon^{JK} \text{sgn}(b_K) \right], \quad (3.22)$$

$$K^\mu A_\mu = \frac{1}{2} K^\mu \partial_\mu \omega = \frac{|b_1| + |b_2|}{4}. \quad (3.23)$$

In principle, the action of the 4d chiral multiplet can be reduced to one dimension by expanding all the fields in an orthonormal basis of eigenfunctions of the relevant differential operators, and performing the integral over the three-manifold M_3 . This yields a 1d action for infinitely many 1d multiplets. In practice, carrying out this reduction on a general background such as the one above is not feasible, as the eigenfunctions are not known. However, for the purpose of computing the Casimir energy there is no need to perform the complete reduction. Indeed, comparing the supersymmetry transformations (3.10) with the one-dimensional ones in (2.10), it is clear that the modes that do not satisfy the shortening conditions in (3.13), (3.14) are going to produce long multiplets in 1d. As explained in section 2.4, these do not contribute to the Casimir energy. Therefore we can focus our attention on the subsector that does satisfy either one of the shortening conditions in (3.13), (3.14). Let us study the two cases in turn.

Reduction to 1d Fermi multiplets. We start analyzing the conditions (3.13), which lead to 1d Fermi multiplets. We Fourier expand the dependence of B on the torus coordinates φ_1, φ_2 as

$$B(\rho, \varphi_1, \varphi_2, \tau) = \sum_{n_1, n_2} b_{n_1, n_2}(\tau) B_{n_1, n_2}(\rho) e^{-in_1 \varphi_1 - in_2 \varphi_2}, \quad (3.24)$$

where n_1, n_2 are integer. Recalling (3.21), (3.22) and that B has R -charge $r - 2$, one can see that the condition $\hat{\mathcal{L}}_Y B = 0$ yields for each choice of n_1, n_2 :

$$\frac{1}{f} B'_{n_1, n_2} = \left[\frac{r-2}{2} \frac{(\log(\Omega \sqrt{m}))'}{f} - \frac{\text{sgn}(b_1 b_2)}{\Omega \sqrt{m}} b_I m_{IJ} \varepsilon^{JK} \left(n_K + \frac{r-2}{2} \text{sgn}(b_K) \right) \right] B_{n_1, n_2}. \quad (3.25)$$

This differential equation determines $B_{n_1, n_2}(\rho)$. The actual solution depends on the form of the metric functions and is not important. For our purposes it is sufficient to impose

that the solution is normalizable, and thus find a restriction for the allowed values of n_1, n_2 . To this end, it is sufficient to study the equation near the extrema of the interval parameterized by $\rho \in [0, 1]$ and make sure that it does not develop singularities. Recalling the boundary conditions (3.17), we find that near $\rho = 0$ the equation is solved by

$$B_{n_1, n_2} = k \rho^{-n_2 \text{sgn}(b_2)} + \dots \quad (3.26)$$

for some constant k . This is normalizable when $n_2 \text{sgn}(b_2) \leq 0$. A similar analysis of the behavior near $\rho = 1$ leads to require $n_1 \text{sgn}(b_1) \leq 0$. It is thus convenient to redefine $n_1 \rightarrow -n_1 \text{sgn}(b_1)$, $n_2 \rightarrow -n_2 \text{sgn}(b_2)$. After this redefinition, the integers n_1, n_2 must take non-negative values, and the expansion (3.24) becomes

$$B = \sum_{n_1, n_2 \geq 0} b_{n_1, n_2}(\tau) B_{n_1, n_2}(\rho) e^{in_1 \text{sgn}(b_1) \varphi_1 + in_2 \text{sgn}(b_2) \varphi_2}. \quad (3.27)$$

Recalling (3.20), (3.23), we can also compute

$$\hat{\mathcal{L}}_K B = -\frac{i}{2} \sum_{n_1, n_2 \geq 0} D_\tau b_{n_1, n_2}(\tau) B_{n_1, n_2}(\rho) e^{in_1 \text{sgn}(b_1) \varphi_1 + in_2 \text{sgn}(b_2) \varphi_2}, \quad (3.28)$$

with

$$D_\tau b_{n_1, n_2} = (\partial_\tau + \lambda_{n_1, n_2}^B) b_{n_1, n_2}, \quad (3.29)$$

and

$$\lambda_{n_1, n_2}^B = -|b_1|n_1 - |b_2|n_2 + \frac{r-2}{2}(|b_1| + |b_2|). \quad (3.30)$$

We are now ready to perform the reduction to one dimension. Setting $\hat{\mathcal{L}}_Y B = 0$ and $\phi = C = 0$, the chiral multiplet Lagrangian (3.11) becomes

$$\mathcal{L} = 2i\tilde{B}\hat{\mathcal{L}}_K B - \tilde{F}F, \quad (3.31)$$

and the supersymmetry variations simplify to

$$\begin{aligned} \delta B &= F, & \tilde{\delta} B &= 0, \\ \delta F &= 0, & \tilde{\delta} F &= 2i\hat{\mathcal{L}}_K B. \end{aligned} \quad (3.32)$$

We can expand F as

$$F = \sum_{n_1, n_2 \geq 0} f_{n_1, n_2}(\tau) B_{n_1, n_2}(\rho) e^{in_1 \text{sgn}(b_1) \varphi_1 + in_2 \text{sgn}(b_2) \varphi_2} \quad (3.33)$$

(note that the dependence on $\rho, \varphi_1, \varphi_2$ is chosen the same as that of B , so F also satisfies $\hat{\mathcal{L}}_Y F = 0$). Similar expansions hold for \tilde{B} and \tilde{F} . The action associated with the Lagrangian (3.31) reduces to an action for an infinite set of Fermi multiplets in one dimension:

$$S = \int d\tau \int_{M_3} d^3x \sqrt{g} \mathcal{L} = \int d\tau \sum_{n_1, n_2 \geq 0} \left(\tilde{b}_{n_1, n_2} D_\tau b_{n_1, n_2} - \tilde{f}_{n_1, n_2} f_{n_1, n_2} \right). \quad (3.34)$$

This is an infinite set of decoupled 1d multiplets, labeled by n_1, n_2 . It is also straightforward to see that the 4d supersymmetry variations decompose into a set of 1d variations for the Fermi multiplets:

$$\begin{aligned}\delta b_{n_1, n_2} &= f_{n_1, n_2}, & \tilde{\delta} b_{n_1, n_2} &= 0, \\ \delta f_{n_1, n_2} &= 0, & \tilde{\delta} f_{n_1, n_2} &= D_\tau b_{n_1, n_2}.\end{aligned}\quad (3.35)$$

From the form of the covariant derivative in (3.29), we see that λ_{n_1, n_2}^B should be identified with the charge σ in section 2. More precisely, $\sigma_{\text{Fermi}} = \frac{2\pi}{\beta} \lambda_{n_1, n_2}^B$.¹⁸ Recalling eq. (2.40), the contribution of the infinitely many Fermi multiplets to the vacuum expectation values of Σ and H is thus

$$\langle H_{\text{Fermi}} \rangle = \langle \Sigma_{\text{Fermi}} \rangle = \frac{\pi}{\beta} \sum_{n_1, n_2 \geq 0} \lambda_{n_1, n_2}^B. \quad (3.36)$$

Reduction to 1d chiral multiplets. In the same way we can study the conditions (3.14), which define a reduction to 1d chiral multiplets. With some foresight, we expand

$$\phi(\rho, \varphi_1, \varphi_2, \tau) = \sum_{n_1, n_2} \phi_{n_1, n_2}(\tau) \Phi_{n_1, n_2}(\rho) e^{-in_1 \text{sgn}(b_1) \varphi_1 - in_2 \text{sgn}(b_2) \varphi_2}. \quad (3.37)$$

Recalling (3.21), (3.22), the condition $\hat{\mathcal{L}}_{\bar{Y}} \phi = 0$ yields for each choice of n_1, n_2 :

$$\frac{1}{f} \Phi'_{n_1, n_2} = \left[-\frac{r}{2} \frac{(\log(\Omega \sqrt{m}))'}{f} + \frac{\text{sgn}(b_1 b_2)}{\Omega \sqrt{m}} b_I m_{IJ} \varepsilon^{JK} \left(n_K \text{sgn}(b_K) + \frac{r}{2} \text{sgn}(b_K) \right) \right] \Phi_{n_1, n_2}. \quad (3.38)$$

This determines $\Phi_{n_1, n_2}(\rho)$. By studying the equation near $\rho = 0$ and $\rho = 1$, we see that normalizability requires $n_1 \geq 0, n_2 \geq 0$. We also compute

$$\hat{\mathcal{L}}_K \phi = -\frac{i}{2} \sum_{n_1, n_2 \geq 0} D_\tau \phi_{n_1, n_2}(\tau) \Phi_{n_1, n_2}(\rho) e^{-in_1 \text{sgn}(b_1) \varphi_1 - in_2 \text{sgn}(b_2) \varphi_2}, \quad (3.39)$$

with

$$D_\tau \phi_{n_1, n_2} = \left(\partial_\tau + \lambda_{n_1, n_2}^\phi \right) \phi_{n_1, n_2}, \quad (3.40)$$

and

$$\lambda_{n_1, n_2}^\phi = |b_1| n_1 + |b_2| n_2 + \frac{r}{2} (|b_1| + |b_2|). \quad (3.41)$$

Setting $\hat{\mathcal{L}}_{\bar{Y}} \phi = \hat{\mathcal{L}}_{\bar{Y}} C = B = F = 0$, the 4d Lagrangian (3.11) becomes

$$\mathcal{L} = 4 \mathcal{L}_{\bar{K}} \tilde{\phi} \hat{\mathcal{L}}_K \phi + i\kappa (\hat{\mathcal{L}}_K \tilde{\phi} \phi - \tilde{\phi} \hat{\mathcal{L}}_K \phi) + 2i \tilde{C} \hat{\mathcal{L}}_{\bar{K}} C - \kappa \tilde{C} C, \quad (3.42)$$

with supersymmetry variations

$$\begin{aligned}\delta \phi &= C, & \tilde{\delta} \phi &= 0, \\ \delta C &= 0, & \tilde{\delta} C &= 2i \hat{\mathcal{L}}_K \phi.\end{aligned}\quad (3.43)$$

¹⁸In order to compare with section 2, we must first rescale the S^1 coordinate as $\tau^{\text{new}} = \frac{\beta}{2\pi} \tau^{\text{old}}$, so that it has period β . Then we implement a Wick rotation $t = i\tau^{\text{new}}$. Overall this gives $(\partial_\tau + \lambda) = i(\partial_t - \frac{2\pi i}{\beta} \lambda)$.

By expanding C as

$$C = \sum_{n_1, n_2 \geq 0} c_{n_1, n_2}(\tau) \Phi_{n_1, n_2}(\rho) e^{-in_1 \text{sgn}(b_1) \varphi_1 - in_2 \text{sgn}(b_2) \varphi_2}, \quad (3.44)$$

performing similar expansions for $\tilde{\phi}$, \tilde{C} and integrating over the four-manifold, we obtain an infinite set of decoupled one-dimensional chiral multiplets. However, the essential information that we need is more straightforwardly extracted from the supersymmetry transformations. For any choice of n_1, n_2 , these read

$$\begin{aligned} \delta \phi_{n_1, n_2} &= c_{n_1, n_2}, & \tilde{\delta} \phi_{n_1, n_2} &= 0, \\ \delta c_{n_1, n_2} &= 0, & \tilde{\delta} c_{n_1, n_2} &= D_\tau \phi_{n_1, n_2}. \end{aligned} \quad (3.45)$$

By comparing with the supersymmetry variations given in (2.10), we see that λ_{n_1, n_2}^ϕ is proportional to the charge σ of the 1d chiral multiplet: $\sigma_{\text{chiral}} = \frac{2\pi}{\beta} \lambda_{n_1, n_2}^\phi$. Then using (2.44) the contribution of the 1d chiral multiplets to the vacuum expectation value of Σ and H is

$$\langle H_{\text{chiral}} \rangle = \langle \Sigma_{\text{chiral}} \rangle = \frac{\pi}{\beta} \sum_{n_1, n_2 \geq 0} \lambda_{n_1, n_2}^\phi. \quad (3.46)$$

The vacuum energy. The VEV of the full one-dimensional Hamiltonian is given by

$$\langle H_{\text{susy}} \rangle = \frac{\pi}{\beta} \sum_{n_1, n_2 \geq 0} \lambda_{n_1, n_2}^\phi + \frac{\pi}{\beta} \sum_{n_1, n_2 \geq 0} \lambda_{n_1, n_2}^B, \quad (3.47)$$

with

$$\begin{aligned} \lambda_{n_1, n_2}^\phi &= |b_1|n_1 + |b_2|n_2 + \frac{r}{2}(|b_1| + |b_2|), \\ \lambda_{n_1, n_2}^B &= -|b_1|n_1 - |b_2|n_2 + \frac{r-2}{2}(|b_1| + |b_2|). \end{aligned} \quad (3.48)$$

These infinite sums are divergent and require regularization. As in subsection 2.5, we will regularize the two sums separately, using the Barnes double zeta function. For simplicity in the next expressions we assume $b_1 > 0$, $b_2 > 0$; the formulae however hold more generally, with b_1, b_2 replaced by $|b_1|, |b_2|$.

The sum (3.47) can be written as

$$\langle H_{\text{susy}} \rangle = \frac{1}{2} \zeta_2 \left(-1; b_1, b_2, \frac{r}{2}(b_1 + b_2) \right) - \frac{1}{2} \zeta_2 \left(-1; b_1, b_2, \frac{2-r}{2}(b_1 + b_2) \right), \quad (3.49)$$

where ζ_2 is the Barnes double zeta function, defined as

$$\zeta_2(s; b_1, b_2, x) = \sum_{n_1, n_2 \geq 0} (b_1 n_1 + b_2 n_2 + x)^{-s}, \quad (3.50)$$

with b_1, b_2, x real and positive. At $s = -1$, it evaluates to (see e.g. [45])

$$\zeta_2(-1; b_1, b_2, x) = -\frac{b_1 + b_2}{24} + \left(3 + \frac{b_1}{b_2} + \frac{b_2}{b_1} \right) \frac{x}{12} - \left(\frac{1}{b_1} + \frac{1}{b_2} \right) \frac{x^2}{4} + \frac{x^3}{6b_1 b_2}. \quad (3.51)$$

This gives for the vacuum energy

$$\langle H_{\text{susy}} \rangle = \frac{4\pi}{3\beta} (b_1 + b_2) (a - c) + \frac{4\pi}{27\beta} \frac{(b_1 + b_2)^3}{b_1 b_2} (3c - 2a). \quad (3.52)$$

This is the same expression appearing in eq. (5.10) of [20]. Redefining $b_1 = \frac{\beta \mathfrak{b}}{2\pi r_3}$ and $b_2 = \frac{\beta \mathfrak{b}^{-1}}{2\pi r_3}$, where β coincides with the length of S^1 , gives the result (1.19) advertised in the introduction.

Back to the round case. Let us conclude by coming back to the $S^3 \times S^1$ background with $\text{SU}(2)_l \times \text{SU}(2)_r \times \text{U}(1)$ symmetry discussed in subsection 2.5. The round sphere is just $S^3_{\mathfrak{b}}$ with $\mathfrak{b} = 1$. In this case $|b_1| = |b_2| = \frac{\beta}{2\pi r_3}$ and the result above for the Casimir energy simplifies to the result in subsection 2.5. In more detail, the sum (3.47) becomes

$$\begin{aligned} \langle H_{\text{susy}} \rangle &= \frac{1}{2r_3} \sum_{n_1, n_2 \geq 0} (n_1 + n_2 + r) + \frac{1}{2r_3} \sum_{n_1, n_2 \geq 0} (-n_1 - n_2 + r - 2) \\ &= \frac{1}{2r_3} \sum_{\ell \geq 0} (\ell + 1) (\ell + r) - \frac{1}{2r_3} \sum_{\ell \geq 0} (\ell + 1) (\ell + 2 - r), \end{aligned} \quad (3.53)$$

where in the second line we have defined $\ell = n_1 + n_2$. So the regularization used above and the one used in subsection 2.5 are compatible.

It is also straightforward to see that the shortening conditions match. When expanding the chiral multiplet (ϕ, C, B, F) in scalar spherical harmonics, the differential operators discussed above become

$$\hat{\mathcal{L}}_Y = iL_l^-, \quad \hat{\mathcal{L}}_K = -\frac{i}{2} \left(L_l^3 + \nabla_\tau + \frac{q_R}{2} \right), \quad (3.54)$$

where q_R is the charge of the field the operator is acting on, and we introduced the $\text{SU}(2)_l$ generators L_l^\pm, L_3 . Then the condition $\hat{\mathcal{L}}_Y \phi = 0$ is nothing but $L_l^+ Y_{\ell, m, n} = 0$, which selects the harmonics with highest quantum number m , namely $Y_{\ell, \frac{\ell}{2}, n}$. The condition $\hat{\mathcal{L}}_Y B = 0$ translates into $L_l^- Y_{\ell, m, n} = 0$ and thus selects $Y_{\ell, -\frac{\ell}{2}, n}$.

Acknowledgments

We would like to thank H.-C. Kim, S. Kim, S. Minwalla, and A. Schwimmer for useful discussions. B.A., J.L., and D.M. acknowledge support by the ERC Starting Grant N. 304806, “The Gauge/Gravity Duality and Geometry in String Theory.” D.C. is supported by an European Commission Marie Curie Fellowship under the contract PIEF-GA-2013-627243. Z.K. and L.D.P are supported by the ERC STG grant 335182, by the Israel Science Foundation under grant 884/11, by the United States-Israel Binational Science Foundation (BSF) under grant 2010/629, as well as by the I-CORE Program of the Planning and Budgeting Committee and by the Israel Science Foundation center for excellence grant (grant no. 1989/14). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the funding agencies.

A Casimir energy in CFT in $d = 2$ and $d = 4$

In this appendix we consider a CFT, not necessarily supersymmetric, in d dimensions coupled to a background metric $g_{\mu\nu}$ with Euclidean signature. We denote the generating functional of connected correlation functions by $\mathcal{W}[g] = -\log Z[g]$. Derivatives with respect to the metric give insertions of the energy-momentum tensor

$$\langle \sqrt{g} T^{\mu\nu} \rangle = 2 \frac{\delta \mathcal{W}}{\delta g_{\mu\nu}} . \quad (\text{A.1})$$

Under an infinitesimal Weyl transformations $g_{\mu\nu} \rightarrow (1 + 2\sigma)g_{\mu\nu}$, the transformation of \mathcal{W} is given by the Weyl anomaly. In $d = 2$ and $d = 4$ respectively we get

$$\delta_\sigma \mathcal{W} = \int d^2x \sqrt{g} \sigma \left(-\frac{c}{24\pi} R \right) , \quad (\text{A.2})$$

$$\delta_\sigma \mathcal{W} = \frac{1}{(4\pi)^2} \int d^4x \sqrt{g} \sigma (aE_{(4)} - cW^2) . \quad (\text{A.3})$$

We have set the coefficient b that appears in (1.3) to zero. The above infinitesimal Weyl transformations can be integrated to the so-called Dilaton action,¹⁹ see e.g. [46–48]:

$$\mathcal{W}[e^{2\sigma}g] - \mathcal{W}[g] = -S_D[-\sigma, g] = S_D[\sigma, e^{2\sigma}g] , \quad (\text{A.4})$$

where S_D has the following explicit expressions in $d = 2$ and $d = 4$

$$S_D^{d=2}[\sigma, g_{\mu\nu}] = \frac{c}{24\pi} \int d^2x \sqrt{g} (-\sigma R + (\partial_\mu \sigma)^2) , \quad (\text{A.5})$$

$$\begin{aligned} S_D^{d=4}[\sigma, g_{\mu\nu}] &= \frac{a}{(4\pi)^2} \int d^4x \sqrt{g} \left(\sigma E_4 + 4\partial_\mu \sigma \partial_\nu \sigma \left(R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} R \right) - 4(\partial\sigma)^2 \square\sigma + 2(\partial\sigma)^4 \right) \\ &\quad - \frac{c}{(4\pi)^2} \int d^4x \sqrt{g} \sigma W^2 . \end{aligned} \quad (\text{A.6})$$

Taking a derivative of (A.4) with respect to the metric, we find

$$e^{d\sigma} \langle T^\mu_\nu \rangle_{e^{2\sigma}g} - \langle T^\mu_\nu \rangle_g = \frac{2}{\sqrt{g}} g_{\nu\lambda} \frac{\delta(-S_D)}{\delta g_{\mu\lambda}} [-\sigma, g] . \quad (\text{A.7})$$

We see that the change in the VEV of the energy-momentum tensor under Weyl rescaling is fixed by the Dilaton action.

A.1 The cylinder and its infinitesimal deformation

In the case of a conformally flat geometry with metric $e^{2\sigma} \delta_{\mu\nu}$ (such as $S^{d-1} \times \mathbb{R}$) we can use the fact that the VEV in flat space is unambiguously fixed to vanish, i.e. $\langle T^\mu_\nu \rangle_{\mathbb{R}^d} = 0$. One then finds for the energy-momentum tensor on this space

$$\langle T^\nu_\mu \rangle_{e^{2\sigma}\delta}^{(d=2)} = \frac{c}{12\pi} e^{-2\sigma} \left(\square\sigma \delta^\nu_\mu - \partial^\nu \partial_\mu \sigma + \partial^\nu \sigma \partial_\mu \sigma - \frac{1}{2} \delta^\nu_\mu (\partial\sigma)^2 \right) , \quad (\text{A.8})$$

¹⁹We thank Adam Schwimmer for discussions.

$$\langle T^\nu_\mu \rangle_{e^{2\sigma}\delta}^{(d=4)} = -\frac{a}{(4\pi)^2} \left[\delta^\nu_\mu \left(R^{\rho\tau} R_{\rho\tau} - \frac{1}{2} R^2 \right) - 2R^{\nu\rho} R_{\rho\mu} + \frac{4}{3} R R^\nu_\mu \right] \Big|_{e^{2\sigma}\delta}. \quad (\text{A.9})$$

These results were first derived in [49] using a different method.

Consider in particular the round cylinder $S^{d-1} \times \mathbb{R}$ with the conformally flat metric

$$ds^2_{S^{d-1} \times \mathbb{R}} = d\tau^2 + r_{d-1}^2 d\Omega_{d-1}^2 = \left(\frac{r_{d-1}}{r} \right)^2 (dr^2 + r^2 d\Omega_{d-1}^2). \quad (\text{A.10})$$

Here $d\Omega_{d-1}^2$ denotes the metric on a $(d-1)$ -sphere with unit radius, and r_{d-1} is the radius of the sphere. Evaluating the energy-momentum tensor in this case, one finds the following non-zero components of $\langle T^\mu_\nu \rangle$ in $d=2$:

$$T^\tau_\tau = -\frac{c}{24\pi r_1^2}, \quad T^\theta_\theta = \frac{c}{24\pi r_1^2}, \quad (\text{A.11})$$

and in $d=4$:

$$T^\tau_\tau = \frac{3a}{8\pi^2 r_3^4}, \quad T^i_j = -\frac{a}{8\pi^2 r_3^4} \delta^i_j. \quad (\text{A.12})$$

Note that $\langle T^\mu_\mu \rangle_{S^3 \times \mathbb{R}} = 0$, because the Weyl anomalies evaluate to zero on the cylinder. We can easily reintroduce the b dependence of the result in $d=4$ by taking a functional derivative of the associated counterterm $-\frac{b}{12(4\pi)^2} \int d^4x \sqrt{g} R^2$. This results in a shift of the coefficient $a \rightarrow a - \frac{b}{2}$.

Integrating $\langle T^\tau_\tau \rangle$ over S^1 and S^3 respectively gives the Casimir energies quoted in the text (1.5) and (1.6).

Above we have reviewed how to compute the Casimir energy on a conformally flat background $e^{2\sigma}\delta_{\mu\nu}$. We will now slightly extend our considerations by taking a geometry that is a generic small perturbation of a conformally flat one, i.e. $g_{\mu\nu} = e^{2\sigma}\delta_{\mu\nu} + h_{\mu\nu}$. Let us compute the change in the ground state energy to first order in $h_{\mu\nu}$. The idea is that we can infer the ground state energy from the change in the partition function \mathcal{W} as a result of the perturbation.

This can be approached as follows: from the definition (A.1) we have

$$\delta\mathcal{W} = \frac{1}{2} \int d^4x \sqrt{g} \langle T_{\mu\nu} \rangle_{e^{2\sigma}\delta} h^{\mu\nu} + \mathcal{O}(h^2), \quad (\text{A.13})$$

so we just need to know $\langle T_{\mu\nu} \rangle_{e^{2\sigma}\delta}$, i.e. the VEV in the conformally flat original space.

For instance, consider the round cylinder $S^3 \times \mathbb{R}$ with a time-independent perturbation h_{ij} of the metric on the three-sphere. The result for the VEV of the energy-momentum tensor on the cylinder was given in equation (A.12). Therefore we find that

$$\delta\mathcal{W} = -\frac{(2a-b)}{32\pi^2 r_3^4} \int d^4x \sqrt{g} h^i_i. \quad (\text{A.14})$$

Since the metric perturbation is assumed time-independent, we can interpret this integral as

$$-\frac{(2a-b)}{32\pi^2 r_3^4} \int d^4x \sqrt{g} h^i_i = -\frac{(2a-b)}{32\pi^2 r_3^4} \int d^3x \sqrt{g} h^i_i \int d\tau \quad (\text{A.15})$$

and the coefficient of $\int d\tau$ gives the correction to the energy of the ground state. The final result for the Casimir energy is

$$E_0 = (2a - b) \left(\frac{3}{8r_3} - \frac{1}{32\pi^2 r_3^4} \int d^3x \sqrt{g} h^i_i \right) + \mathcal{O}(h^2) . \quad (\text{A.16})$$

We see that to order $\mathcal{O}(h)$ the Casimir energy is still scheme dependent, being proportional to the same combination $2a - b$ as the leading order result. However, the ratio of the leading term and the subleading term is unambiguous (in fact, the ratio is fixed by imagining a small perturbation that only changes the radius of the three-sphere).

One can speculate that going to second order in the perturbation $\mathcal{O}(h^2)$ the c -anomaly will also appear and that some terms will be scheme independent. Notice, however, that at this order a new counterterm, proportional to $\int d^4x \sqrt{g} W^2$, could contribute.

A.2 The Casimir energy and holography

We now summarize the status of the Casimir energy in the context of the gauge/gravity duality. We will mainly focus on four-dimensional CFTs admitting a dual description in terms of solutions of type IIB supergravity of the type $\text{AdS}_5 \times M_5$, and their deformations. Similar considerations can be made for six-dimensional CFTs with AdS_7 gravity duals.

The comparison of the Casimir energy of a four-dimensional CFT on $S^3 \times \mathbb{R}$ with the expectation value of the holographic energy-momentum tensor was first made in [50], although in this reference the authors focussed on $\mathcal{N} = 4$ SYM and its $\text{AdS}_5 \times S^5$ dual, and some of the comments made there apply only to this example. Let us review the discussion of [50]. Using the holographic energy-momentum tensor, computed in global AdS_5 , one obtains for the Casimir energy, defined exactly as in (1.1), the expression²⁰

$$E_0^{\text{holo}} = \frac{3N^2}{16\ell} . \quad (\text{A.17})$$

It is important to note that this result is obtained using a minimal holographic renormalization scheme, where there are no finite counterterms ΔS_{ct} added to the on-shell action. This corresponds to the absence of the $\square R$ term in the holographic trace anomaly $\langle T^\mu_\mu \rangle$. In other words, it corresponds to the scheme $b = 0$.

The expression (A.17) is compared with a Casimir energy in a free CFT, comprising n_0 real scalars, $n_{1/2}$ Weyl fermions, and n_1 Abelian gauge fields. In particular, the scalar fields considered are conformally coupled, and the fermions are massless. This results in the expression

$$E_0^{\text{free}} = \frac{1}{960r_3} (4n_0 + 17n_{1/2} + 88n_1) , \quad (\text{A.18})$$

obtained summing up the contributions of the single fields, which in turn are regularised using zeta function. For the specific case of the $\mathcal{N} = 4$ SYM theory, we have

$$E_0^{\text{free}} = \frac{3(N^2 - 1)}{16r_3} = \frac{3}{4r_3} a \quad \text{for } \mathcal{N} = 4 \text{ SYM}, \quad (\text{A.19})$$

which agrees with (A.17) at leading order in N [50].

²⁰This was referred to as “mass of global AdS_5 ” in [50]. We have used the standard formula for the Newton constant $G_5^{-1} = \frac{2N^2}{\ell^3} \pi^4 \text{vol}'(M_5)$, where ℓ is the AdS radius and in the specific case $\text{vol}'(S^5) = \pi^3$.

However, the agreement of E_0^{holo} with E_0^{free} is, in some sense, accidental, and may be misleading. Indeed for generic n_0 , $n_{1/2}$, n_1 , the expression (A.18) for E_0^{free} is *not* proportional to the a anomaly: it also includes a contribution from b . In fact, the scheme leading to (A.18) generically also gives a $\square R$ contribution to the trace anomaly. Once the particular matter content of $\mathcal{N} = 4$ SYM is specified, the coefficient b happens to vanish (see e.g. [46]) and the scheme leading to (A.18) effectively coincides with the scheme $b = 0$ which was used in the holographic computation.

More generally, one can ask whether the computation of the Casimir energy in a deformed cylinder can be reproduced by a five-dimensional supergravity solution deforming AdS_5 . At least at leading order in the deformation, there exists a holographic counterpart to the formula (A.13). Consider a one-parameter family of deformations of the *boundary* metric, with parameter μ . Then by simply applying the chain rule to the renormalised on-shell action one obtains²¹

$$\frac{d}{d\mu} S_{\text{ren}} = \int_{\partial M_5} d^4x \sqrt{g} \left(-\frac{1}{2} \langle T_{\mu\nu} \rangle \frac{dg^{\mu\nu}}{d\mu} \right), \quad (\text{A.20})$$

where $g_{\mu\nu}$ is the finite metric on the boundary, and $\langle T_{\mu\nu} \rangle$ is the holographic energy-momentum tensor. Setting $g = g_{(0)} + \mu h$, with μ infinitesimal, and expanding (A.20) at first order in μ leads exactly to the holographic version of the formula (A.13). Thus, the holographic Casimir energy on an infinitesimally squashed cylinder is guaranteed to agree with the field theory result. In the next subsection, we verify this in an explicit example.

For the supersymmetric version of the Casimir energy the situation is different. One of the main points emphasized in this paper is that on the field theory side, computing in the free limit gives a reliable result for E_{susy} , valid also at strongly coupled points. However, how to reproduce this in the gravity side remains an open problem. In [7] it was suggested that a supersymmetric treatment of holographic renormalization might reveal the existence of new boundary terms, that would lead to a matching of the on-shell action of existing solutions with E_{susy} . Another possibility is that there exist other (Euclidean) supersymmetric solutions, whose on-shell action would match with E_{susy} , using the standard holographic renormalization technology. In light of the results of the present paper, it will be very interesting to revisit this problem.

Let us also mention that there exists a similar open problem in the context of SCFTs on the six-dimensional cylinder $S^5 \times \mathbb{R}$. A supersymmetric version of the Casimir energy was discussed in [21], where it was also noted that it does not match the standard holographic Casimir energy in AdS_7 . It may be useful to prove that this quantity is physical in 6d SCFTs by performing an analysis like the one in the present paper, and to investigate the seven-dimensional holographic dual.

A.3 Holographic check of E_0 on a squashed cylinder

Below we will compare the result of the first-order correction to the ordinary Casimir energy due to a non-conformally flat geometry (A.16), with a corresponding holographic

²¹In [7] this identity was written to include the variation of the boundary gauge field. However, if we are not interested in supersymmetry, we can vary the metric independently.

result, that can be easily extracted from the gravity solution presented in [7]. Although the solution in [7] is supersymmetric, we can obtain from this an expression for the Casimir energy, that is valid independently of supersymmetry. In order to make the discussion as self-contained as possible, we will begin recalling relevant aspects of the solution in [7], referring the reader to this reference for more details.

This is a one-parameter family of supersymmetric solutions deforming AdS_5 , constructed as an asymptotically locally anti de Sitter (AlAdS) solution of five-dimensional minimal gauged supergravity.

An analytic continuation to Euclidean signature yields the gravity dual to a class of four-dimensional $\mathcal{N} = 1$ supersymmetric gauge theories on a curved manifold with topology $S^3 \times S^1$. In particular, the boundary metric is that of a squashed three-sphere preserving $\text{SU}(2) \times \text{U}(1)$ isometry, and there is a non-trivial background gauge field coupling to the R -symmetry current. So, this is an instance of a Hopf surface, as discussed in the main part of the text [7].

In a coordinate system, the five-dimensional metric and the graviphoton field take the following asymptotic form near to the boundary, corresponding to $\rho \rightarrow \infty$:

$$\begin{aligned} ds_{5d}^2 &= d\rho^2 + e^{2\rho/\ell} ds_{\text{bdry}}^2 + \dots, \\ A_{5d} &= A_{\text{bdry}} + \mathcal{O}(e^{-\rho}), \end{aligned} \tag{A.21}$$

where the boundary values read

$$ds_{\text{bdry}}^2 = (2a_0)^2 \left[-\frac{1}{v^2} dt^2 + \frac{\ell^2}{4} (\sigma_1^2 + \sigma_2^2 + v^2 \sigma_3^2) \right], \tag{A.22}$$

and

$$A_{\text{bdry}} = \frac{1}{2\sqrt{3}} \left[\frac{dt}{\ell} + (v^2 - 1)\sigma_3 \right], \tag{A.23}$$

respectively. Here ℓ is the AdS radius, that can be identified with the radius of the S^3 , and the left-invariant one-forms σ_a 's are defined as usual (see [7]). The full five-dimensional solution is determined in terms of the single parameter v , measuring the squashing of the boundary metric. For $v^2 = 1$ the solution reduces to AdS_5 and the boundary metric is conformally flat. The parameter $2a_0$ is an overall scale, that can be set to any value by simply shifting the radial coordinate ρ .²² Notice that the background gauge field in (A.23) comprises a constant part along dt , that is necessary to have well defined supercharges in the compactified geometry, as discussed around (2.3). However, in order to compare with the result above, which is valid in the *absence* of such term²³ (and independently of supersymmetry), we should remove this by shifting the graviphoton field as $A_{5d} \rightarrow A_{5d} - \frac{1}{2\sqrt{3}\ell} dt$. It is simple to check that the solution with this new gauge field is still non-singular, and its relevant properties may be extracted from [7], by following through this simple change of gauge.

²²In [7] the parameter a_0 is chosen to be a convenient function of v (so that the solution ends at $\rho = 0$). However, this is irrelevant for the present discussion.

²³This term maps to a gauge field $A \sim \frac{dx}{r}$ that is *singular* at the origin of \mathbb{R}^4 ; this is why we do not include it in our treatment.

After Wick rotating to Euclidean signature, and setting $v = 1 + \epsilon$, we have at leading order the following boundary metric

$$ds_{\text{bdry}}^2 = ds_{(0)}^2 + ds_{(1)}^2 + \mathcal{O}(\epsilon^2), \quad (\text{A.24})$$

with

$$\begin{aligned} ds_{(0)}^2 &= g_{\mu\nu}^{(0)} dx^\mu dx^\nu = d\tau^2 + \frac{\ell^2}{4} (\sigma_1^2 + \sigma_2^2 + \sigma_3^2), \\ ds_{(1)}^2 &= h_{\mu\nu} dx^\mu dx^\nu = \epsilon \left[-2d\tau^2 + \frac{\ell^2}{2} \sigma_3^2 \right], \end{aligned} \quad (\text{A.25})$$

and background gauge field

$$A_{\text{bdry}} = \frac{\epsilon}{\sqrt{3}} \sigma_3 + \mathcal{O}(\epsilon^2). \quad (\text{A.26})$$

In particular, this implies that the contribution of the background R -symmetry current to the Casimir energy through a term of the type $\int_{S^3} d^3x \sqrt{g} \langle J_R^\mu \rangle A_\mu$ can only affect the result at order $\mathcal{O}(\epsilon^2)$. Therefore at linearized order we can neglect the background gauge field.

Using (A.12), (A.13) we can evaluate the correction to the Casimir energy due to the perturbation in (A.25). We obtain

$$\delta E_0 = \frac{1}{2} (2\pi^2 \ell^3) \left(-\epsilon \frac{a}{\pi^2 \ell^4} \right) = -\epsilon \frac{a}{\ell}, \quad (\text{A.27})$$

where the factor $2\pi^2 \ell^3$ comes from the integration over the three-sphere. Thus, the Casimir energy at first order in the squashing reads

$$E_0 = \left(\frac{3}{4} - \epsilon \right) \frac{a}{\ell}. \quad (\text{A.28})$$

This can be compared with the result of the holographic computation. The Euclidean on-shell action of five-dimensional supergravity was computed exactly as a function of the parameter v in [7], and reads

$$S = \frac{8a}{\ell} \left(\frac{2}{27v^2} + \frac{2}{27} - \frac{13}{108} v^2 + \frac{19}{288} v^4 \right) \int d\tau, \quad (\text{A.29})$$

where we rewrote the five-dimensional Newton constant in terms of the a anomaly of the dual CFT. This was obtained in a gauge such that $A_\tau^{\text{bdry}} \neq 0$, so we should shift the gauge to $A_\tau^{\text{bdry}} = 0$ and correspondingly the on-shell action using eq. (4.14) therein. In fact, the shift due to the change of gauge is proportional to ϵ^2 , so at linear order, this shift is immaterial. In any case, after doing this shift, we obtain exactly the expression appearing in (4.35) therein for E_0 , times $\int d\tau$. The on-shell action (A.29) expanded at first order in ϵ , gives

$$S = \left(\frac{3}{4} - \epsilon \right) \frac{a}{\ell} \int d\tau, \quad (\text{A.30})$$

thus we get perfect agreement with the dual field theory result (A.28). Note that (A.29) has been evaluated using a holographic renormalization scheme without the $\int d^4x \sqrt{g} R^2$ counterterm, precisely as in the conformally flat background.

B E_{susy} and the Hamiltonian

In this appendix we show that the supersymmetric vacuum energy E_{susy} , defined within the path integral approach as

$$E_{\text{susy}} = - \lim_{\beta \rightarrow \infty} \frac{d}{d\beta} \log Z_{M_3 \times S^1_\beta}^{\text{susy}}, \quad (\text{B.1})$$

coincides with the VEV of the charge associated to the time translation symmetry. In other words, E_{susy} is the VEV of the Hamiltonian, $E_{\text{susy}} = \langle H_{\text{susy}} \rangle$. This is well-known, in particular in the context of quantum field theories at non-zero temperature. Nevertheless in the presence of additional non-dynamical background fields it may be useful to spell out some details.

E_{susy} can be expressed in terms of the energy-momentum tensor and the currents appearing in the R -multiplet in the following way. Consider a background $M_3 \times S^1_\beta$ as in the main text, then perform a trivial rescaling of the S^1 coordinate τ so that $\tau \sim \tau + 1$ and the background fields $g_{\tau\tau}$, A_τ and V_τ acquire a dependence on β . Applying the chain rule to the variation of $-\log Z$ with respect to β , E_{susy} can be written as

$$\begin{aligned} E_{\text{susy}} &= \lim_{\beta \rightarrow \infty} \left\langle \int d^4x \sqrt{g} \left(-\frac{1}{2} T_{\mu\nu} \frac{dg^{\mu\nu}}{d\beta} + J_R^\mu \frac{dA_\mu}{d\beta} - \frac{3}{2} J_{\text{FZ}}^\mu \frac{dV_\mu}{d\beta} \right) \right\rangle \\ &= \lim_{\beta \rightarrow \infty} \frac{1}{\beta} \left\langle \int d^4x \sqrt{g} \left(T^\tau{}_\tau + J_R^\tau A_\tau - \frac{3}{2} J_{\text{FZ}}^\tau V_\tau \right) \right\rangle, \end{aligned} \quad (\text{B.2})$$

where the energy-momentum tensor $T_{\mu\nu}$, the R -current J_R^μ and the Ferrara-Zumino current J_{FZ}^μ are defined as

$$T_{\mu\nu} = -\frac{2}{\sqrt{g}} \frac{\delta S}{\delta g^{\mu\nu}}, \quad J_R^\mu = \frac{1}{\sqrt{g}} \frac{\delta S}{\delta A_\mu}, \quad -\frac{3}{2} J_{\text{FZ}}^\mu = \frac{1}{\sqrt{g}} \frac{\delta S}{\delta V_\mu}. \quad (\text{B.3})$$

These are components of the R -multiplet, which at the linear level have canonical couplings to the metric, to A and to V , respectively. However, we remark that we do not set A and V to zero after having taken the variation, so these are currents in the presence of sources.

The expression in (B.2) shows that E_{susy} receives a contribution from the current in the R -multiplet in addition to the one from the temporal component of the energy-momentum tensor. This is an alternate way to see that E_{susy} is different from the ordinary Casimir energy E_0 defined in eq. (1.1).

We now construct the charge associated with the time translation symmetry and compare it with (B.2). Recall that in the presence of background fields other than the metric, the energy-momentum tensor $T_{\mu\nu}$ is in general not conserved, $\nabla^\mu T_{\mu\nu} \neq 0$. This can be easily seen by revisiting the standard conservation proof (see e.g. appendix E of [51]) allowing for fields that do not satisfy their Euler-Lagrange equation. In particular, this applies to a supersymmetric field theory defined via the rigid limit of new minimal supergravity. One can see that the non-conservation equation of the energy-momentum tensor reads

$$\nabla^\mu T_{\mu\nu} = (dA)_{\nu\mu} J_R^\mu - \frac{3}{2} (dV)_{\nu\mu} J_{\text{FZ}}^\mu + \frac{3}{2} V_\nu \nabla_\mu J_{\text{FZ}}^\mu, \quad (\text{B.4})$$

where we used that the R -current is conserved, $\nabla_\mu J_R^\mu = 0$, while generically $\nabla_\mu J_{\text{FZ}}^\mu \neq 0$.

Although $T_{\mu\nu}$ is not conserved, when the background admits a Killing symmetry generated by a vector ξ^μ one can introduce a modified energy-momentum current

$$Y_\xi^\mu = \xi^\nu \left(T_\nu^\mu + J_R^\mu A_\nu - \frac{3}{2} J_{FZ}^\mu V_\nu \right), \quad (\text{B.5})$$

that is conserved, $\nabla_\mu Y_\xi^\mu = 0$. This is easily seen using $\mathcal{L}_\xi g = \mathcal{L}_\xi A = \mathcal{L}_\xi V = 0$. One can show that Y_ξ^μ is just the canonical Noether current associated to the symmetry generated by ξ . Thus a conserved charge can be defined in the usual way.

We recall that two such Killing vectors exist in Euclidean backgrounds preserving two supercharges of opposite R -charge [16, 17]. In Lorentzian signature, at least one null Killing vector exists [36]. Our $M_3 \times S_\beta^1$ background admits a Killing vector generating (Euclidean) time translations, $\xi = \frac{\partial}{\partial \tau}$. The associated conserved charge, to be identified with the Hamiltonian, is

$$\begin{aligned} H &= \int_{M_3} d^3x \sqrt{g_{(3)}} u_\mu Y_{\xi = \frac{\partial}{\partial \tau}}^\mu \\ &= \frac{1}{\beta} \int_{M_4} d^4x \sqrt{g_{(4)}} \left(T^\tau{}_\tau + J_R^\tau A_\tau - \frac{3}{2} J_{FZ}^\tau V_\tau \right), \end{aligned} \quad (\text{B.6})$$

where u^μ is the unit time-like vector, that in the second line we expressed as $u = u_\mu dx^\mu = \sqrt{g_{\tau\tau}} d\tau$ (in the second line we also multiplied the three-dimensional integral by $1 = \frac{1}{\beta} \int_{S^1} d\tau$; since the charge is constant, this can be rewritten as a four-dimensional integral). The VEV of the final expression in (B.6) for $\beta \rightarrow \infty$ is immediately recognized as the E_{susy} given in (B.2), that is what we wanted to show.

Finally, note that the Hamiltonian in (B.6) coincides with the supersymmetric H_{susy} used in section 2 only for the special choice $A_\tau = i/r_3$, which guarantees time-independent supersymmetry.

C Regularization of one-loop determinant

In this appendix we reconsider the regularization of the one-loop determinant for a free chiral multiplet on the Hopf surface $M^3 \times S^1$ (with $M^3 \simeq S^3$) computed in [20]. It was found in [26] that the partition function on the Hopf surface has a universal behaviour in the small β limit, where β is the length of S^1 . In the case of the round $S^3 \times S^1$, this takes the form²⁴

$$\log Z_{S^3 \times S_\beta^1}^{\text{susy}} = \frac{16\pi^2 r_3}{3\beta} (c - a) + \mathcal{O}(\beta^0), \quad (\text{C.1})$$

where r_3 is the radius of S^3 . The result found in [20] are incompatible with this expansion, as in [20] the order $\mathcal{O}(\beta^{-1})$ vanishes instead. We propose here an alternative method to regularize one-loop determinant which on one hand agrees with the small beta expansion of [26] and on the other hand reproduces the large β behaviour leading to the Casimir energy (1.19). Moreover, we show that this regularization method is equivalent to a

²⁴Subleading terms in the small β expansion were worked out in [29].

cut-off regularization which manifestly preserves supersymmetry. The main difference between the procedure discussed below and that in [20] is that here all the Kaluza-Klein (KK) modes on S^1 are dealt with in a manifestly symmetric way, while in [20] these were somewhat artificially split and combined with the Fourier modes on the three-sphere, in the triple gamma functions. However, the precise reason for the discrepancy of the two methods remains unclear. While we were writing up this paper, [29] appeared, presenting a similar regularization method of one-loop determinants, leading to the result (1.19) for the general deformed sphere.

One-loop determinant regularizations are based on the use of multiple gamma functions [52] and generalized zeta functions. The main mathematical tools that we will use are presented in a convenient way in appendices A and B of [53]. We will also rely on [54].

The one-loop determinant suffers from UV divergences. It is given for a free chiral multiplet²⁵ by the formal expression

$$Z_{1\text{-loop}}^{\text{chiral}} \equiv Z = \prod_{n_0 \in \mathbb{Z}} \prod_{n_1, n_2 \geq 0} \frac{-\frac{r}{2}(a_1 + a_2) - n_0 + (n_1 + 1)a_1 + (n_2 + 1)a_2}{\frac{r}{2}(a_1 + a_2) + n_0 + n_1 a_1 + n_2 a_2}, \quad (\text{C.2})$$

where $a_1 = ib_1$, $a_2 = ib_2$ in the notations of [20] and here we consider $b_1 > 0, b_2 > 0$. r is the R -charge of the chiral multiplet. The analysis of this paper allows to re-interpret each factor appearing in the numerator of (C.2) as the contribution from a single Fermi multiplet to the one-loop determinant and each factor in the denominator as the contribution of a single chiral multiplet. A crucial difference with the Hamiltonian quantization analysis is that now we have a tower of KK modes on S^1 with KK level parameterized by $n_0 \in \mathbb{Z}$.

C.1 Two-step regularization

Here we proceed by first regularizing the sum over $n_1, n_2 \geq 0$ with double gamma functions for a fixed n_0 , which corresponds to the one-loop determinant on M_3 of the n_0 -th KK mode along S^1 , and then regularizing the infinite sum over $n_0 \in \mathbb{Z}$. We start with

$$\begin{aligned} Z &= \prod_{n_0 \in \mathbb{Z}} F[u + n_0] \\ F[v] &= \prod_{n_1, n_2 \geq 0} \frac{-v + (n_1 + 1)a_1 + (n_2 + 1)a_2}{v + n_1 a_1 + n_2 a_2} = \frac{\Gamma_2(v, a_1, a_2)}{\Gamma_2(a_1 + a_2 - v, a_1, a_2)} \\ &= \Gamma_h(v, a_1, a_2), \end{aligned} \quad (\text{C.3})$$

where $u = \frac{r}{2}(a_1 + a_2)$ and the hyperbolic gamma function is defined by

$$\begin{aligned} \Gamma_h(v, a_1, a_2) &= e^{\frac{\pi i}{2} B(v, a_1, a_2)} \frac{(e^{2\pi i(a_1 - v)/a_2}; e^{2\pi i a_1/a_2})_\infty}{(e^{-2\pi i v/a_1}; e^{-2\pi i a_2/a_1})_\infty} \\ (x, q)_\infty &= \prod_{k \geq 0} (1 - x q^k), \quad B(v, a_1, a_2) = \frac{1}{a_1 a_2} \left(\left(v - \frac{a_1 + a_2}{2} \right)^2 - \frac{a_1^2 + a_2^2}{12} \right). \end{aligned} \quad (\text{C.4})$$

²⁵ Adding a gauge interaction with a flat gauge field as in the localization computation of [20] is straightforward.

The hyperbolic gamma function admits also the following representation [53]

$$\Gamma_h(v, a_1, a_2) = e^{-\frac{\pi i}{2} B(v, a_1, a_2)} \frac{(e^{2\pi i(v-a_2)/a_1}; e^{-2\pi i a_2/a_1})_\infty}{(e^{2\pi i v/a_2}; e^{2\pi i a_1/a_2})_\infty}. \quad (\text{C.5})$$

Using (C.4) for $n_0 \geq 1$ and (C.5) for $n_0 \leq 0$, we obtain

$$Z = e^{\frac{\pi i}{2} C(u, a_1, a_2)} \frac{\Gamma_e\left(\frac{u}{a_2}, -\frac{1}{a_2}, \frac{a_1}{a_2}\right)}{\Gamma_e\left(\frac{u-a_2}{a_1}, -\frac{1}{a_1}, -\frac{a_2}{a_1}\right)} = e^{\frac{\pi i}{2} C(u, a_1, a_2)} e^{\frac{\pi i}{2} D(u, a_1, a_2)} \Gamma_e(u, a_1, a_2), \quad (\text{C.6})$$

where the elliptic gamma function Γ_e is defined by

$$\Gamma_e(x, \tau, \sigma) = \prod_{n_1, n_2 \geq 0} \frac{1 - e^{2\pi i(-x + (n_1+1)\tau + (n_2+1)\sigma)}}{1 - e^{2\pi i(x + n_1\tau + n_2\sigma)}}. \quad (\text{C.7})$$

The last equality in (C.6) is obtained from the modular properties of the elliptic gamma function [53, 54] and the functions C, D are given by

$$D(u, a_1, a_2) = \frac{1}{a_1 a_2} \left(\frac{2}{3} (u')^3 + (u')^2 + \frac{2 - a_1^2 - a_2^2}{6} u' - \frac{a_1^2 + a_2^2}{12} \right), \quad (\text{C.8})$$

$$C(u, a_1, a_2) = \frac{1}{a_1 a_2} \left(\sum_{n \geq 1} \left[(u' + n)^2 - \frac{a_1^2 + a_2^2}{12} \right] - \sum_{n \geq 0} \left[(-u' + n)^2 - \frac{a_1^2 + a_2^2}{12} \right] \right), \quad (\text{C.9})$$

with $u' = u - \frac{a_1 + a_2}{2} = \frac{r-1}{2}(a_1 + a_2)$. Note that $D(u, a_1, a_2)$ is a well defined function, but $C(u, a_1, a_2)$ involves an infinite sum over n , which needs to be regularized. We do this using the standard Riemann zeta function,²⁶ which is compatible with the partial cancellation of terms between the two infinite sums:

$$C(u, a_1, a_2) = \frac{1}{a_1 a_2} \left(-(u')^2 + \frac{a_1^2 + a_2^2}{12} + 4u' \sum_{n \geq 1} n \right) = \frac{1}{a_1 a_2} \left(-(u')^2 + \frac{a_1^2 + a_2^2}{12} - \frac{u'}{3} \right), \quad (\text{C.10})$$

where we used $\sum_{n \geq 1} n = \zeta(-1) = -\frac{1}{12}$. Notice that despite the formal similarity of the infinite sums in (C.9) and (2.49), the regularization of the two sums is performed using two *different* prescriptions. The final result is

$$Z = e^{i\pi \Psi(u, a_1, a_2)} \Gamma_e(u, a_1, a_2), \quad \Psi(u, a_1, a_2) = \frac{(u')^3}{3a_1 a_2} - \frac{a_1^2 + a_2^2}{12a_1 a_2} u'. \quad (\text{C.11})$$

This reproduces the supersymmetric Casimir energy (1.19) and it can be shown that is compatible with the small β limit of [26], which is obtained by setting $a_1 = i \frac{\beta}{2\pi r_3} \mathfrak{b}$, $a_2 = i \frac{\beta}{2\pi r_3} \mathfrak{b}^{-1}$ and keeping \mathfrak{b} fixed in the limit. The difference with the result of [20] is only the absence of the term $\frac{u'}{6a_1 a_2} = \frac{r-1}{12} \frac{a_1 + a_2}{a_1 a_2}$ in $\Psi(u, a_1, a_2)$, which affects the small β limit, but not the large β limit.

²⁶This is similar to the regularization presented in appendix B of [55].

C.2 Cut-off regularization

We present now a different method of regularization which uses a cut-off on the momentum modes. The treatment of the KK modes in this method is more transparent and the result agrees with (C.11), thus corroborating it.

A reliable way of regularizing the one-loop determinant is to introduce a cut-off on the momentum of the modes or more generally a smooth truncation of the modes above a momentum scale Λ (this means that the contributions of modes above Λ are counted with decreasing weights). Note that this preserves supersymmetry since each momentum in the product carries the contribution of a specific multiplet in supersymmetric quantum mechanics. The (logarithm of the) regularized determinant can be expanded in powers of the cut-off at large Λ . The diverging contributions are expected to be removed by local supersymmetric counterterms, while the finite piece yields the regularized result, possibly subject to ambiguities. In the case at hand we compute the partition function on a curved manifold admitting two Killing spinors of new minimal supergravity of opposite chirality and it was shown in [8] that there is no ambiguity in the finite result.

For simplicity, we focus here on the case of the round $S^3 \times S^1$, which amounts to setting $a_1 = a_2 = a \in i\mathbb{R}_{>0}$. In this case, the chiral multiplet one-loop determinant simplifies to

$$\begin{aligned} Z &= \prod_{n_0 \in \mathbb{Z}} \prod_{n_1, n_2 \geq 0} \frac{-u - n_0 + (n_1 + n_2 + 2)a}{u + n_0 + (n_1 + n_2)a} = \prod_{n_0 \in \mathbb{Z}} \prod_{m \geq 0} \left(\frac{-u - n_0 + (m + 2)a}{u + n_0 + ma} \right)^{m+1} \\ &= \prod_{n_0 \in \mathbb{Z}} F[u + n_0], \end{aligned} \quad (\text{C.12})$$

with $u = ra$ and

$$\log F[v] = \sum_{m \geq 0} (m + 1) [\log(-v + (m + 2)a) - \log(v + ma)]. \quad (\text{C.13})$$

We regularize the sum by introducing a cut-off $\frac{\beta}{2\pi}\Lambda$ on the S^1_β KK momentum level n_0 and $\frac{\beta}{2\pi|a|}\Lambda$ on the S^3 KK momentum level m .²⁷ To simplify slightly the notations we simply set $\beta = 2\pi$ in the following. The regularization is done by using any smooth decreasing function f such that $f(0) = 1$ and going to zero at infinity sufficiently fast to make the sum converge. The regularized sum over S^3 modes is

$$\log F[v] = \sum_{m \geq 0} (m + 1) [\log(-v + (m + 2)a) - \log(v + ma)] f\left(\frac{m}{\Lambda|a^{-1}|}\right). \quad (\text{C.14})$$

We make use of the Euler-MacLaurin formula²⁸ to work out the large Λ expansion

$$\begin{aligned} \log F[v] &= 2\Lambda|a^{-1}|(1 - va^{-1}) \int_0^\infty f + \mathcal{O}(\Lambda^0) \\ &= c_1 \Lambda a^{-2}(v - a) + \mathcal{O}(\Lambda^0), \end{aligned} \quad (\text{C.16})$$

²⁷The relative factor $|a| = \frac{\beta}{2\pi r_3}$ takes into account the ratio of scales between S^1_β and S^3 KK modes.

²⁸The Euler-MacLaurin formula for a convergent sum reads:

$$\sum_{n \geq 0} g(n) = \int_0^\infty g(x) dx + \frac{1}{2}g(0) + \sum_{k \geq 1} \frac{\zeta(1 - 2k)}{(2k - 1)!} g^{(2k-1)}(0). \quad (\text{C.15})$$

with $c_1 = 2i \int_0^\infty f$. The finite part of the large Λ expansion is complicated as it receives an infinite number of contributions from the Euler-MacLaurin expansion, however it is easy to see that it is independent of the function f and we can trust that it will reproduce the (logarithm of the) hyperbolic gamma function at $a_1 = a_2$, since this is the well-known result for the 3d chiral multiplet one-loop determinant. We obtain

$$F[u + n_0] \simeq e^{c_1 \Lambda a^{-2}(u-a+n_0)} \Gamma_h(u + n_0, a, a), \quad (\text{C.17})$$

where \simeq indicates that we dropped the term $\mathcal{O}(\Lambda^{-1})$ in $\log F[u + n_0]$. The same manipulations as in the regularization above yield the result

$$Z = e^{\frac{\pi i}{2} \tilde{C}(u, a, a)} e^{\frac{\pi i}{2} D(u, a, a)} \Gamma_e(u, a, a), \quad (\text{C.18})$$

with

$$\begin{aligned} \tilde{C}(u, a, a) = & \frac{1}{a^2} \sum_{n \geq 1} \left[(u' + n)^2 - \frac{a^2}{6} + c_1 \Lambda (u' + n) \right] \hat{f}\left(\frac{n}{\Lambda}\right) \\ & + \frac{1}{a^2} \sum_{n \geq 0} \left[-(u' - n)^2 + \frac{a^2}{6} + c_1 \Lambda (u' - n) \right] \hat{f}\left(\frac{n}{\Lambda}\right), \end{aligned} \quad (\text{C.19})$$

where $u' = u - a = (r - 1)a$ and \hat{f} is a second regulating function implementing the smooth cut-off on S^1 KK modes. Applying again the Euler-MacLaurin formula we obtain

$$\tilde{C}(u, a, a) = \frac{1}{a^2} \left(c_2 \Lambda^2 u' - u'^2 - \frac{u'}{3} + \frac{a^2}{6} \right), \quad (\text{C.20})$$

with $c_2 = 4 \int_0^\infty y \hat{f}(y) dy + \frac{8c_1}{i\pi} \int_0^\infty \hat{f}(y) dy$. The finite piece reproduces the result (C.10) for $a_1 = a_2 = a$, so that the cut-off regularization result Z_{reg} will match the regularization described above. The diverging piece should be removed with a dimension two supersymmetric counterterm constructed with new minimal supergravity background fields [8]. We have obtained

$$\log Z \simeq \frac{i\pi}{2} c_2 \Lambda^2 (r - 1) a^{-1} + \log Z_{\text{reg}}. \quad (\text{C.21})$$

There is a single supergravity term of mass dimension two in new minimal supergravity [8]; it is the usual Einstein-Hilbert new minimal supergravity action, that we can take with coefficient Λ^2 . Its bosonic part is given by

$$S_R^{(4)} = \frac{\Lambda^2}{2} \int d^4x \sqrt{g} (R + 6V_\mu V^\mu - 8A_\mu V^\mu). \quad (\text{C.22})$$

Its evaluation on the $S^3 \times S^1$ background [17] yields²⁹

$$S_R^{(4)} \propto \Lambda^2 a^{-2} \kappa, \quad (\text{C.23})$$

where the constant κ parametrizes a freedom in the choice of background. The choice $\kappa = a$ is the most natural, because it preserves the $\text{SU}(2)_l \times \text{SU}(2)_r$ isometries of S^3 . This is precisely the term needed to remove the divergent piece in (C.21). We conclude that the cut-off regularization further validates the regularization method presented above.

²⁹Note that the result is in units where the S^1 radius is fixed to one. To reinstate the β dependence, one can shift $\Lambda \rightarrow \frac{\beta}{2\pi} \Lambda$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](#)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] J. Cardy, *The ubiquitous ‘c’: from the Stefan-Boltzmann law to quantum information*, *J. Stat. Mech.* **1010** (2010) P10004 [[arXiv:1008.2331](#)] [[INSPIRE](#)].
- [2] A.B. Zamolodchikov, *Irreversibility of the flux of the renormalization group in a 2D field theory*, *JETP Lett.* **43** (1986) 730 [*Pisma Zh. Eksp. Teor. Fiz.* **43** (1986) 565] [[INSPIRE](#)].
- [3] J.L. Cardy, *Is there a c theorem in four-dimensions?*, *Phys. Lett. B* **215** (1988) 749 [[INSPIRE](#)].
- [4] I. Jack and H. Osborn, *Analogues for the c theorem for four-dimensional renormalizable field theories*, *Nucl. Phys. B* **343** (1990) 647 [[INSPIRE](#)].
- [5] Z. Komargodski and A. Schwimmer, *On renormalization group flows in four dimensions*, *JHEP* **12** (2011) 099 [[arXiv:1107.3987](#)] [[INSPIRE](#)].
- [6] Z. Komargodski, *The constraints of conformal symmetry on RG flows*, *JHEP* **07** (2012) 069 [[arXiv:1112.4538](#)] [[INSPIRE](#)].
- [7] D. Cassani and D. Martelli, *The gravity dual of supersymmetric gauge theories on a squashed $S^1 \times S^3$* , *JHEP* **08** (2014) 044 [[arXiv:1402.2278](#)] [[INSPIRE](#)].
- [8] B. Assel, D. Cassani and D. Martelli, *Supersymmetric counterterms from new minimal supergravity*, *JHEP* **11** (2014) 135 [[arXiv:1410.6487](#)] [[INSPIRE](#)].
- [9] S. Pasquetti, *Factorisation of $N = 2$ theories on the squashed 3-sphere*, *JHEP* **04** (2012) 120 [[arXiv:1111.6905](#)] [[INSPIRE](#)].
- [10] Y. Yoshida, *Factorization of 4d $N = 1$ superconformal index*, [arXiv:1403.0891](#) [[INSPIRE](#)].
- [11] V.P. Spiridonov and G.S. Vartanov, *Elliptic hypergeometric integrals and ’t Hooft anomaly matching conditions*, *JHEP* **06** (2012) 016 [[arXiv:1203.5677](#)] [[INSPIRE](#)].
- [12] S.S. Razamat, *On a modular property of $N = 2$ superconformal theories in four dimensions*, *JHEP* **10** (2012) 191 [[arXiv:1208.5056](#)] [[INSPIRE](#)].
- [13] G. Festuccia and N. Seiberg, *Rigid supersymmetric theories in curved superspace*, *JHEP* **06** (2011) 114 [[arXiv:1105.0689](#)] [[INSPIRE](#)].
- [14] M.F. Sohnius and P.C. West, *An alternative minimal off-shell version of $N = 1$ supergravity*, *Phys. Lett. B* **105** (1981) 353 [[INSPIRE](#)].
- [15] D. Cassani and D. Martelli, *Supersymmetry on curved spaces and superconformal anomalies*, *JHEP* **10** (2013) 025 [[arXiv:1307.6567](#)] [[INSPIRE](#)].
- [16] C. Klare, A. Tomasiello and A. Zaffaroni, *Supersymmetry on curved spaces and holography*, *JHEP* **08** (2012) 061 [[arXiv:1205.1062](#)] [[INSPIRE](#)].
- [17] T.T. Dumitrescu, G. Festuccia and N. Seiberg, *Exploring curved superspace*, *JHEP* **08** (2012) 141 [[arXiv:1205.1115](#)] [[INSPIRE](#)].
- [18] C. Closset, T.T. Dumitrescu, G. Festuccia and Z. Komargodski, *From rigid supersymmetry to twisted holomorphic theories*, *Phys. Rev. D* **90** (2014) 085006 [[arXiv:1407.2598](#)] [[INSPIRE](#)].

- [19] C. Closset, T.T. Dumitrescu, G. Festuccia and Z. Komargodski, *The geometry of supersymmetric partition functions*, *JHEP* **01** (2014) 124 [[arXiv:1309.5876](#)] [[INSPIRE](#)].
- [20] B. Assel, D. Cassani and D. Martelli, *Localization on Hopf surfaces*, *JHEP* **08** (2014) 123 [[arXiv:1405.5144](#)] [[INSPIRE](#)].
- [21] H.-C. Kim and S. Kim, *M5-branes from gauge theories on the 5-sphere*, *JHEP* **05** (2013) 144 [[arXiv:1206.6339](#)] [[INSPIRE](#)].
- [22] C. Romelsberger, *Counting chiral primaries in $N = 1$, $D = 4$ superconformal field theories*, *Nucl. Phys. B* **747** (2006) 329 [[hep-th/0510060](#)] [[INSPIRE](#)].
- [23] J. Kinney, J.M. Maldacena, S. Minwalla and S. Raju, *An index for 4 dimensional super conformal theories*, *Commun. Math. Phys.* **275** (2007) 209 [[hep-th/0510251](#)] [[INSPIRE](#)].
- [24] C. Romelsberger, *Calculating the superconformal index and Seiberg duality*, [arXiv:0707.3702](#) [[INSPIRE](#)].
- [25] F.A. Dolan and H. Osborn, *Applications of the superconformal index for protected operators and q -hypergeometric identities to $N = 1$ dual theories*, *Nucl. Phys. B* **818** (2009) 137 [[arXiv:0801.4947](#)] [[INSPIRE](#)].
- [26] L. Di Pietro and Z. Komargodski, *Cardy formulae for SUSY theories in $d = 4$ and $d = 6$* , *JHEP* **12** (2014) 031 [[arXiv:1407.6061](#)] [[INSPIRE](#)].
- [27] A.A. Ardehali, J.T. Liu and P. Szepietowski, *c -a from the $N = 1$ superconformal index*, *JHEP* **12** (2014) 145 [[arXiv:1407.6024](#)] [[INSPIRE](#)].
- [28] A.A. Ardehali, J.T. Liu and P. Szepietowski, *Central charges from the $N = 1$ superconformal index*, *Phys. Rev. Lett.* **114** (2015) 091603 [[arXiv:1411.5028](#)] [[INSPIRE](#)].
- [29] A.A. Ardehali, J.T. Liu and P. Szepietowski, *High-temperature expansion of supersymmetric partition functions*, [arXiv:1502.07737](#) [[INSPIRE](#)].
- [30] J. Lorenzen and D. Martelli, *Comments on the Casimir energy in supersymmetric field theories*, [arXiv:1412.7463](#) [[INSPIRE](#)].
- [31] C.P. Herzog and K.-W. Huang, *Stress tensors from trace anomalies in conformal field theories*, *Phys. Rev. D* **87** (2013) 081901 [[arXiv:1301.5002](#)] [[INSPIRE](#)].
- [32] C. Closset and I. Shamir, *The $N = 1$ chiral multiplet on $T^2 \times S^2$ and supersymmetric localization*, *JHEP* **03** (2014) 040 [[arXiv:1311.2430](#)] [[INSPIRE](#)].
- [33] N. Bobev, M. Bullimore and H.-C. Kim, in progress.
- [34] L.F. Alday, D. Martelli, P. Richmond and J. Sparks, *Localization on three-manifolds*, *JHEP* **10** (2013) 095 [[arXiv:1307.6848](#)] [[INSPIRE](#)].
- [35] Z. Komargodski and N. Seiberg, *Comments on supercurrent multiplets, supersymmetric field theories and supergravity*, *JHEP* **07** (2010) 017 [[arXiv:1002.2228](#)] [[INSPIRE](#)].
- [36] D. Cassani, C. Klare, D. Martelli, A. Tomasiello and A. Zaffaroni, *Supersymmetry in Lorentzian curved spaces and holography*, *Commun. Math. Phys.* **327** (2014) 577 [[arXiv:1207.2181](#)] [[INSPIRE](#)].
- [37] A.V. Smilga, *Weak supersymmetry*, *Phys. Lett. B* **585** (2004) 173 [[hep-th/0311023](#)] [[INSPIRE](#)].
- [38] E. Ivanov and S. Sidorov, *Deformed supersymmetric mechanics*, *Class. Quant. Grav.* **31** (2014) 075013 [[arXiv:1307.7690](#)] [[INSPIRE](#)].

- [39] S. Elitzur, Y. Frishman, E. Rabinovici and A. Schwimmer, *Origins of global anomalies in quantum mechanics*, *Nucl. Phys. B* **273** (1986) 93 [[INSPIRE](#)].
- [40] C. Closset, T.T. Dumitrescu, G. Festuccia, Z. Komargodski and N. Seiberg, *Comments on Chern-Simons contact terms in three dimensions*, *JHEP* **09** (2012) 091 [[arXiv:1206.5218](#)] [[INSPIRE](#)].
- [41] E. Gerchkovitz, *Constraints on the R-charges of free bound states from the Römelsberger index*, *JHEP* **07** (2014) 071 [[arXiv:1311.0487](#)] [[INSPIRE](#)].
- [42] D. Anselmi, D.Z. Freedman, M.T. Grisaru and A.A. Johansen, *Nonperturbative formulas for central functions of supersymmetric gauge theories*, *Nucl. Phys. B* **526** (1998) 543 [[hep-th/9708042](#)] [[INSPIRE](#)].
- [43] T. Nishioka and I. Yaakov, *Generalized indices for $N = 1$ theories in four-dimensions*, *JHEP* **12** (2014) 150 [[arXiv:1407.8520](#)] [[INSPIRE](#)].
- [44] N. Hama, K. Hosomichi and S. Lee, *SUSY gauge theories on squashed three-spheres*, *JHEP* **05** (2011) 014 [[arXiv:1102.4716](#)] [[INSPIRE](#)].
- [45] M. Spreafico, *On the Barnes double zeta and Gamma functions*, *J. Number Theor.* **129** (2009) 2035.
- [46] E.S. Fradkin and A.A. Tseytlin, *Conformal anomaly in Weyl theory and anomaly free superconformal theories*, *Phys. Lett. B* **134** (1984) 187 [[INSPIRE](#)].
- [47] M.R. Brown and A.C. Ottewill, *Effective actions and conformal transformations*, *Phys. Rev. D* **31** (1985) 2514 [[INSPIRE](#)].
- [48] A. Cappelli and A. Coste, *On the stress tensor of conformal field theories in higher dimensions*, *Nucl. Phys. B* **314** (1989) 707 [[INSPIRE](#)].
- [49] L.S. Brown and J.P. Cassidy, *Stress tensors and their trace anomalies in conformally flat space-times*, *Phys. Rev. D* **16** (1977) 1712 [[INSPIRE](#)].
- [50] V. Balasubramanian and P. Kraus, *A stress tensor for anti-de Sitter gravity*, *Commun. Math. Phys.* **208** (1999) 413 [[hep-th/9902121](#)] [[INSPIRE](#)].
- [51] R.M. Wald, *General relativity*, University of Chicago Press, Chicago U.S.A. (1984) [[INSPIRE](#)].
- [52] S. Ruijsenaars, *On Barnes' multiple zeta and gamma functions*, *Adv. Math.* **156** (2000) 107.
- [53] V.P. Spiridonov, *Elliptic beta integrals and solvable models of statistical mechanics*, *Contemp. Math.* **563** (2012) 181 [[arXiv:1011.3798](#)] [[INSPIRE](#)].
- [54] G. Felder and A. Varchenko, *The elliptic gamma function and $SL(3, Z) \ltimes Z^3$* , *Adv. Math.* **156** (2000) 44 [[math/9907061](#)].
- [55] O. Aharony, S.S. Razamat, N. Seiberg and B. Willett, *3d dualities from 4d dualities*, *JHEP* **07** (2013) 149 [[arXiv:1305.3924](#)] [[INSPIRE](#)].