



King's Research Portal

DOI:

[10.1002/ajmg.b.32862](https://doi.org/10.1002/ajmg.b.32862)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Luo, M., Meehan, A. J., Walton, E., Röder, S., Herberth, G., Zenclussen, A. C., Cosín, M., Sunyer, J., Mulder, R. H., Cortes Hidalgo, A. P., Bakermans-Kranenburg, M. J., Felix, J. F., Relton, C., Suderman, M., Pappa, I., Kok, R., Tiemeier, H., van IJzendoorn, M. H., Barker, E. D., & Cecil, C. A. M. (2021). Neonatal DNA methylation and childhood low prosocial behavior: An epigenome-wide association meta-analysis. *American Journal of Medical Genetics. Part B: Neuropsychiatric Genetics*, 186(4), 228-241. <https://doi.org/10.1002/ajmg.b.32862>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

ORIGINAL ARTICLE



WILEY

Neonatal DNA methylation and childhood low prosocial behavior: An epigenome-wide association meta-analysis

Mannan Luo^{1,2} | Alan J. Meehan^{3,4} | Esther Walton^{5,6} | Stefan Röder⁷ | Gunda Herberth⁷ | Ana C. Zenciusen⁷ | Marta Cosín-Tomás^{8,9,10} | Jordi Sunyer^{8,9,10,11} | Rosa H. Mulder^{2,12} | Andrea P. Cortes Hidalgo^{2,12} | Marian J. Bakermans-Kranenburg¹³ | Janine F. Felix^{2,14} | Caroline Relton⁵ | Matthew Suderman⁵ | Irene Pappa^{2,12} | Rianne Kok¹ | Henning Tiemeier^{12,15} | Marinus H. van IJzendoorn^{1,16} | Edward D. Barker¹⁷ | Charlotte A. M. Cecil^{12,17,18,19}

¹Department of Psychology, Education and Child Studies, Erasmus University Rotterdam, Rotterdam, The Netherlands

²Generation R Study Group, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

³Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

⁴Yale Child Study Center, Yale School of Medicine, New Haven, USA

⁵MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

⁶Department of Psychology, University of Bath, Bath, UK

⁷Department for Environmental Immunology, Helmholtz Centre for Environmental Research—UFZ, Leipzig, Germany

⁸ISGlobal, Barcelona, Catalonia, Spain

⁹Universitat Pompeu Fabra, Barcelona, Catalonia, Spain

¹⁰Centro de Investigación Biomédica en Red en Epidemiología y Salud Pública (CIBERESP), Madrid, Spain

¹¹IMIM Parc Salut Mar, Barcelona, Catalonia, Spain

¹²Department of Child and Adolescent Psychiatry/Psychology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

¹³Department of Clinical Child and Family Studies, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

¹⁴Department of Pediatrics, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

¹⁵Department of Social and Behavioral Science, Harvard TH Chan School of Public Health, Boston, Massachusetts, USA

¹⁶Department of Clinical, Educational and Health Psychology, Faculty of Brain Sciences, University College London, London, UK

¹⁷Department of Psychology, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

¹⁸Department of Epidemiology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands

¹⁹Molecular Epidemiology, Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, The Netherlands

Correspondence

Edward D. Barker, Department of Psychology, Institute of Psychiatry, Psychology & Neuroscience, King's College London, 16 De Crespigny Park, London SE5 8AF, UK.
Email: ted.barker@kcl.ac.uk

Abstract

Low prosocial behavior in childhood has been consistently linked to later psychopathology, with evidence supporting the influence of both genetic and environmental factors on its development. Although neonatal DNA methylation (DNAm) has been

Mannan Luo and Alan J. Meehan should be considered joint first authors.

Edward D. Barker and Charlotte A. M. Cecil should be considered joint senior authors.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics* published by Wiley Periodicals LLC.

Charlotte A. M. Cecil, Department of Child and Adolescent Psychiatry/Psychology Erasmus MC, University Medical Center Rotterdam, P.O. Box 2060, Rotterdam, CB 3000, The Netherlands.
Email: c.cecil@erasmusmc.nl

Funding information

Agence Nationale de Sécurité Sanitaire de l'Alimentation de l'Environnement et du Travail, Grant/Award Number: 1262C0010; Agència de Gestió d'Ajuts Universitaris i de Recerca, Grant/Award Numbers: 2009 SGR 501, 2014 SGR 822; China Scholarship Council, Grant/Award Number: 201706990036; Comissió Interdepartamental de Recerca i Innovació Tecnològica, Grant/Award Number: 1999SGR 00241; Deutsches Krebsforschungszentrum; Economic and Social Research Council, Grant/Award Numbers: ES/J500057/1, ES/R005516/1; Eunice Kennedy Shriver National Institute of Child Health and Human Development, Grant/Award Number: R01HD068437; European Commission, Grant/Award Numbers: 261357, 308333, 603794, 634453; Fundació la Marató de TV3, Grant/Award Number: 090430; H2020 European Research Council, Grant/Award Numbers: 268479, 669249; H2020 Marie Skłodowska-Curie Actions, Grant/Award Numbers: 707404, 848158; Helmholtz-Zentrum für Umweltforschung; Instituto de Salud Carlos III, Grant/Award Numbers: CB06/02/0041, CP13/00054, CP16/00128, CP18/00018, G03/176, PI041436, PI081151, PI12/01890, PI15/00118, PI16/00118, PI16/00261, PI18/00547; Medical Research Council, Grant/Award Number: MC_UU_00011/5; Ministerio de Ciencia e Innovación, Grant/Award Number: FJC2018-036335-I; National University of Ireland; Nederlandse Organisatie voor Wetenschappelijk Onderzoek, Grant/Award Numbers: 024-001-003, 050-060-810; Spanish Ministry of Economy and Competitiveness, Grant/Award Number: SAF2012-32991; Spanish Ministry of Health, Grant/Award Numbers: FIS-FEDER-PI03-1615, FIS-FEDER-PI06/0867, FIS-PI04/1436, FIS-PI08/1151, FIS-PI11/00610

found to prospectively associate with a range of psychological traits in childhood, its potential role in prosocial development has yet to be investigated. This study investigated prospective associations between cord blood DNAm at birth and low prosocial behavior within and across four longitudinal birth cohorts from the Pregnancy And Childhood Epigenetics (PACE) Consortium. We examined (a) developmental trajectories of “chronic-low” versus “typical” prosocial behavior across childhood in a case-control design ($N = 2,095$), and (b) continuous “low prosocial” scores at comparable cross-cohort time-points ($N = 2,121$). Meta-analyses were performed to examine differentially methylated positions and regions. At the cohort-specific level, three CpGs were found to associate with chronic low prosocial behavior; however, none of these associations was replicated in another cohort. Meta-analysis revealed no epigenome-wide significant CpGs or regions. Overall, we found no evidence for associations between DNAm patterns at birth and low prosocial behavior across childhood. Findings highlight the importance of employing multi-cohort approaches to replicate epigenetic associations and reduce the risk of false positive discoveries.

KEYWORDS

cord blood, DNA methylation, epigenome-wide association study, meta-analysis, prosocial behavior

1 | INTRODUCTION

Prosocial behavior, defined as voluntary behavior intended to benefit others (e.g., helping, sharing, and comforting), provides a foundation for social competence and moral development (Eisenberg, Fabes, & Spinrad, 2006). Typically emerging early in childhood, higher levels of expressed prosociality are consistently linked with greater self-esteem, peer acceptance, and academic success in young people (Caprara, Barbaranelli, Pastorelli, Bandura, & Zimbardo, 2000; Layous, Nelson, Oberle, Schonert-Reichl, & Lyubomirsky, 2012). However, levels of prosocial expression vary considerably between individuals. Notably, at the opposite end of the prosocial continuum, young people who follow developmental trajectories denoted by persistently

low levels of prosociality display elevated levels of externalizing behaviors, such as aggression and delinquency (Flynn, Ehrenreich, Beron, & Underwood, 2015; Kokko, Tremblay, Lacourse, Nagin, & Vitaro, 2006), and—albeit less consistently—internalizing problems (Nantel-Vivier, Pihl, Cote, & Tremblay, 2014).

Etiological research consistently reports both genetic and environmental contributions to prosocial development (Knafo-Noam, Versteiger, & Israel, 2018). From twin designs, heritability estimates for observed prosocial actions range between 23 and 69%, with most studies also identifying substantial shared environment effects (Gregory, Light-Häusermann, Rijdsdijk, & Eley, 2009; Knafo-Noam, Israel, & Ebstein, 2011; Knafo-Noam, Uzefovsky, Israel, Davidov, & Zahn-Waxler, 2015). These heritability estimates have also been

shown to increase during childhood, while shared environmental influences diminish over time (Knafo-Noam & Plomin, 2006). However, efforts to elucidate the specific biological processes underpinning this heritability by examining associations between specific candidate genes and various domains of prosocial behavior have yielded very small meta-analytic effect sizes (Bakermans-Kranenburg & Van IJzendoorn, 2014). Furthermore, no hypothesis-free study of genetic variation across the entire genome has been published for this phenotype to date (Knafo-Noam et al., 2018). At the same time, exposure to prenatal (e.g., maternal stress, androgen exposure) and postnatal adversity (e.g., parenting, family functioning, neighborhood deprivation) has been linked to less prosocial behavior in children, suggesting some etiological role for the early environment (Horn, Hungerlander, Windhager, Bugnyar, & Massen, 2018; Jambon, Madigan, Plamondon, & Jenkins, 2019; Loomans et al., 2011; Safra et al., 2016).

Although more research has begun to consider the potential interplay between genetic and environmental influences on prosocial development (Knafo-Noam et al., 2011; Sasaki et al., 2013), the underlying molecular mechanisms remain unclear. In recent years, DNA methylation (DNAm), an epigenetic process that regulates gene expression in response to both genetic and environmental signals, has emerged as a potential mechanism of interest through which the early environment, in combination with genetic predispositions, can increase biological vulnerability for psychopathology (Barker, Walton, & Cecil, 2018). Specifically, DNAm involves the addition of a methyl group to DNA base pairs—mainly, cytosine–guanine (CpG) dinucleotides (McGowan & Roth, 2015)—which can disrupt the binding of transcription factors (Zhang & Meaney, 2010). DNAm has been shown to regulate numerous neurobiological and developmental processes, with alterations in DNAm linked to psychological and psychiatric outcomes (Barker, Walton, Cecil, Rowe, et al., 2018; Cecil et al., 2014; Rijlaarsdam et al., 2017; Walton et al., 2017). It has also been suggested that DNAm may act as a noncausal biomarker indexing vulnerability for psychopathology, in a similar way to the approach that has been used in cancer risk and prognosis (Ladd-Acosta & Fallin, 2016). Thus, it may be possible to identify a consistent profile of DNAm that characterizes persistently “low prosocial” youth.

To date, no study has examined DNAm patterns associated with low prosocial behavior, although some research has been carried out on closely related phenotypes. For example, callous-unemotional (CU) traits are defined by a lack of prosocial emotions such as empathy and remorse, and represent a significant risk factor for aggressive and antisocial behaviors (Marsh, 2019). Within clinical samples of boys with conduct problems, higher CU traits have been associated with increased DNAm in *OXTR* gene promoter (Dadds et al., 2014) and reduced DNAm of the promoter region of the serotonin 1B receptor gene (*HTR1B*; Moul, Dobson-Stone, Brennan, Hawes, & Dadds, 2015). However, these approaches are limited in two main ways. First, single candidate genes are not likely to explain the majority of variance of more complex, multi-determined psychiatric traits and disorders, whose exact pathophysiology is not yet known (Salvatore & Dick, 2018). Second, these studies only examined DNAm for specific

genes selected based on hypothesized biological or functional relevance, precluding their ability to detect novel biological associations.

In light of these research gaps, this study drew on four independent yet highly compatible birth cohorts to investigate epigenome-wide, prospective associations between DNAm at birth and low prosocial behavior during childhood. Focusing on methylation levels at birth (i.e., before manifestation of later behavioral or psychiatric symptoms) enabled us to separate the direction of the association, an important step for establishing whether detected epigenetic modifications may serve as potential predictive markers, as opposed to secondary effects of the “disease” process itself (Rakyan, Down, Balding, & Beck, 2011). Moreover, existing longitudinal studies comparing the stability of DNAm associations across childhood have found that DNAm patterns are (a) highly dynamic over time (Mulder et al., 2021), and (b) more strongly predictive (at birth) of neurodevelopmental and behavioral outcomes related to prosocial behavior in children, including conduct problems (Cecil et al., 2018), CU traits (Cecil et al., 2014), and social communication deficits (Rijlaarsdam, Cecil, Relton, & Barker, 2021), compared to DNAm patterns examined later in childhood. However, it is not known whether such findings at birth extend to prosocial behavior itself. First, in line with previous research (Nantel-Vivier et al., 2014), we identified developmental trajectories characterized by persistently low prosocial behavior across childhood. We then ran an epigenome-wide association study (EWAS) to identify DNAm sites within each cohort that differed between trajectories. Second, as all cohorts featured comparable time-points and identical measures of prosocial behavior, we meta-analyzed results to maximize our power to detect small effects and reduce the risk of false positives. Furthermore, we used region discovery methods to identify potentially differentially methylated regions (DMRs) associated with low prosocial behavior across cohorts. Finally, previous studies of other psychiatric traits suggest that the influence of genetic and environmental risk factors may be best observed across a continuum of severity (Thapar, Langley, O'Donovan, & Owen, 2006). Thus, we adopted a dimensional approach to complement our trajectory-based analyses, repeating all steps using continuous prosocial scores assessed at comparable time-points (6–7 years of age) within each cohort.

2 | METHODS

2.1 | Participants

Four prospective birth cohorts within the Pregnancy And Childhood Epigenetics (PACE) Consortium (Felix et al., 2018) had information on DNAm in cord blood and prosocial behavior in childhood: the Avon Longitudinal Study of Parents and Children (ALSPAC; Fraser et al., 2013), Generation R (GENR; Kooijman et al., 2016), Infancia y Medio Ambiente (INMA; Guxens et al., 2012), and the Lifestyle and environmental factors and their Influence on Newborns Allergy risk (LINA; Herberth et al., 2011). Full cohort descriptions are provided in Supporting Information Methods. In brief, all included participants

were of European ancestry, with combined sample sizes of 2,095–2,121 in the current study. Ethical approval for each study was obtained by local committees and consent to use their data was obtained for all participants.

2.2 | Measures

2.2.1 | DNA methylation

DNA_m in cord blood was measured using the Illumina Infinium HumanMethylation450K BeadChip (Illumina Inc., San Diego). Full information on sample processing, quality control, and normalization procedures within each cohort is described in Appendix S1 of Supporting Information. Methylation levels were characterized by a “beta” (β) value ranging from 0 (no methylation) to 1 (full methylation). CpG sites with values outside the 25th percentile $- 3 \times$ interquartile range (IQR) and the 75th percentile $+ 3 \times$ IQR of the distribution were identified as outliers and winsorized.

2.2.2 | Prosocial behavior

In all cohorts, prosocial behavior was repeatedly assessed using maternal ratings on the “prosocial” subscale of the Strengths and Difficulties Questionnaire (SDQ), a widely used screening instrument with established reliability and validity (Goodman, 2001). Five items assessed the presence of the following behaviors “in the past six months” along a three-point scale (0 = “not true”; 1 = “somewhat true”; 2 = “certainly true”): (a) “considerate of other’s feelings”; (b) “shares readily with other children”; (c) “helpful if someone is hurt, upset, or ill”; (d) “kind to younger children”; and (e) “volunteers to help others”. Prosocial behavior was assessed in ALSPAC when children were aged 4, 7, 8, 10, 12, and 13 years, in GENR when children were aged 6 and 9 years, in INMA when children were aged 7 and 11 years, and in LINA when children were aged 7 and 10 years.

Latent profile analysis (LPA) was performed using Mplus version 7.11 (Muthén & Muthén, 2012) to classify ALSPAC children into developmental trajectories based on their levels of prosocial behavior across childhood. A type of latent variable mixture model, LPA is a person-centered technique that identifies discrete subgroups within a population based on a series of observed indicators—in this case, repeated measures of the SDQ prosocial subscale at six time-points (Oberski, 2016). We estimated two-class to five-class solutions and used several fit indices (Bayesian Information Criterion; Akaike Information Criterion; Lo–Mendell–Rubin test; entropy) to determine the optimal number of classes (see Table S1, Panel A and accompanying note for details). Based on these metrics, a three-class solution provided the best fit for these data. Within this three-class model, one trajectory subgroup, termed “chronic-low” ($n = 75$, bottom 10% of sample), identified children with consistently low levels of prosocial behavior from age 4–13 years. The two remaining trajectories differed

significantly from this chronic-low trajectory, as well as from each other. However, average levels of prosocial behavior among these two trajectory groups ($M = 9.0$ and 7.3) fell well within the “normative” or “average” scoring band (≥ 6), based on the original cut-offs for the parent-reported SDQ (Goodman, 1997). This contrasted with the chronic-low prosocial group, where the grand mean for prosocial behavior across all six time-points fell within the clinical threshold for “low” prosocial behavior ($M = 5.6$). As the present study sought to compare persistently low prosocial youth with more typical levels of prosociality in a case–control approach, we combined the two remaining trajectories, hereafter referred to as the “typical” group ($n = 673$, 89.96%). The resulting dichotomous variable (i.e., “typical” vs. “chronic-low”) was found to significantly associate with several established correlates of prosocial behavior, including empathy, social-cognitive difficulties, and CU traits ($r_s = -0.27$ – 0.32 , $p < .001$; see Table S1, Panel B).

To enable cross-cohort comparison and meta-analysis, we sought to establish an equivalent case–control design for all cohorts by replicating the clinically significant “bottom 10%” cut-off observed from trajectory-based analyses in ALSPAC. However, as GENR, INMA, and LINA each had only two repeated measures of prosocial behavior, which was insufficient to perform LPA, we instead standardized prosocial scores for each age and then calculated an average over both ages. Values below the 10th percentile for this average score (i.e., “bottom 10%” cut-off) were classed as “chronic-low” prosocial behavior across both ages, in contrast to the remainder of the sample (i.e., typical group). Having identified these groups in children who had available prosocial scores at both ages, our final sample within each cohort consisted of those who also had DNA_m data available at birth. Table 1 presents the number and proportion of children in each of these chronic-low trajectories. In all three cohorts, “grand mean” prosocial scores in the defined chronic-low (GENR: $M = 5.4$; $SD = 1.7$; INMA: $M = 5.3$; $SD = 1.2$; LINA: $M = 5.5$; $SD = 1.3$) and typical groups (GENR: $M = 8.8$; $SD = 1.3$; INMA: $M = 8.7$; $SD = 1.4$; LINA: $M = 8.5$; $SD = 1.3$) were comparable to those observed within the two ALSPAC trajectory groups (chronic-low: $M = 5.6$; typical: $M = 8.2$) and thus in line with clinical definitions for “low” and “average” levels of prosocial behavior, respectively.

In a complementary strategy to account for the dimensionality of prosocial behavior and identify DNA_m differences associated with the severity (as opposed to chronicity) of low prosociality, we repeated all analyses using continuous measures of low prosocial behavior (rather than trajectory groups) at the closest-corresponding time-points at age 6–7 years across all cohorts. These continuous scores were negatively skewed (i.e., indicating that most children exhibit higher levels of prosocial behavior) in all cohorts, which is a common occurrence in general-population samples. As square root and log transformation of the scores did not correct for this skewness, as well as the fact that transformed scores would complicate interpretation of the results, robust linear regression analyses were performed. Scale scores were reversed to be consistent with trajectory coding and facilitate interpretation, with higher scores representing lower prosocial behavior.

TABLE 1 Descriptive statistics across the four included cohorts ($N_{total} = 2,095\text{--}2,121$)

	Cohort (sample size) ^a			
	Avon Longitudinal Study of Parents and Children (ALSPAC; $n = 748$)	Generation R (GENR; $n = 891$)	Infancia y Medio Ambiente (INMA; $n = 270$)	Lifestyle and environmental factors and their Influence on Newborns Allergy risk (LINA; $n = 186$)
Child characteristics				
Sex, % female	48.4	49.6	48.9	48.1
Gestational age in weeks, mean (SD)	39.6 (1.5)	40.2 (1.4)	39.8 (1.3)	39.8 (1.4)
Child prosocial behavior				
Chronic-low prosocial trajectory, n (%)	75 (10.0)	100 (11.2)	31 (11.5)	20 (10.8)
Continuous prosocial score (age 6–7 years), mean (SD)	1.8 (1.7)	1.7 (1.8)	1.7 (1.7)	1.9 (1.7)
Age in years, mean (SD)	6.7 (0.1)	6.0 (0.3)	6.7 (0.4)	7.1 (0.3)
Maternal characteristics				
Age at intake/delivery in years, mean (SD)	29.6 (4.4)	32.1 (3.9)	30.6 (4)	30.8 (4.6)
Sustained smoking during pregnancy, %	9.7	9.8	12.6	1.6
Educational level, %				
Lower (below university degree)	79.2	29.8	68.1	36.5
Higher (university degree and above)	20.8	70.3	31.9	63.5
Psychopathology, mean (SD)	14.3, % yes ^b	0.1 (0.2)	0.7 (0.5)	2.1 (0.5)

^aReported sample sizes are drawn from prosocial trajectory analyses. For analyses of continuous “low prosocial” scores at age 6–7 years, sample sizes were as follows: $N_{ALSPAC} = 689$; $N_{GENR} = 976$; $N_{INMA} = 270$; $N_{LINA} = 186$.

^bIn ALSPAC, the presence of maternal anxiety/depression was coded as a binary variable (1 = yes vs. 0 = no).

2.2.3 | Covariates

We adjusted for child sex, maternal age at delivery (in years), smoking during pregnancy (binary categorization of “no smoking/quit in early pregnancy” vs. “smoked throughout pregnancy”), gestational age at delivery (in weeks), child age at the outcome assessment (in years), maternal education (binary categorization of “below university degree” vs. “university degree and above”), and technical covariates (e.g., batch or surrogate variables). We also adjusted for estimated cell type proportions calculated from a cord blood cell type reference panel (Gervin et al., 2019). Finally, we included mothers' experience of depression and anxiety as a covariate (see Appendix S1 of Supporting Information for full details of measures used), as a caregiver experiencing these symptoms may perceive and rate their child's behavior as more problematic than one who is not, potentially biasing observed associations between methylation and low prosociality.

2.3 | Statistical Analyses

Analyses proceeded in two main steps:

2.3.1 | Step 1: Epigenome-wide association study (EWAS) and meta-analysis

Based on a predefined analysis plan, cohort-level EWAS between DNAm at birth and prosocial trajectories during childhood was

performed using linear regression models. Cohorts excluded individuals with known chromosomal abnormalities, multiple births, and one random sibling per sibling pair.

After performing quality control on cohort-specific summary statistics, we combined results in a fixed-effects inverse variance-weighted meta-analysis using the *metafor* R package (Viechtbauer, 2010). We excluded probes mapped to X/Y chromosomes, polymorphic CpGs (overlapping with known single-nucleotide-polymorphisms; Min, Hemani, Smith, Relton, & Suderman, 2018), and control or cross-reactive probes (targeting repetitive sequences/co-hybridizing to alternate sequences; Chen et al., 2013; McCartney et al., 2016), resulting in 364,659 autosomal CpG sites in the meta-analysis. Probes were annotated using the *meffil* R package (Min et al., 2018), enhanced by use of the University of California Santa Cruz Genome Browser (including data from the RefSeq and Ensembl databases). All annotations were based on genome build hg19. Genome-wide significance was defined based on a looser 450k array p -value threshold of $p < 2.4 \times 10^{-7}$ (Saffari et al., 2018), suggestive significance at $p < 5 \times 10^{-5}$, and nominal significance at $p < .05$.

We performed several additional analyses. First, we repeated analyses using “low prosocial” continuous scores at age 6–7 years, using robust linear regression in the *MASS* R package. Second, a candidate gene follow-up analysis was performed on results extracted from the meta-analysis for CpG sites annotated to *OXTR*, in order to examine associations with DNAm at this previously implicated gene. The significance threshold was set at a Bonferroni gene-level corrected p -value of 4.17×10^{-3} ($n = 12$ CpGs).

2.3.2 | Step 2: Differentially methylated regions (DMRs) and meta-analysis

To account for the correlated structure of DNAm patterns, we next performed a regional analysis to identify regions that are differentially methylated in relation to low prosocial behavior. In contrast to site-specific EWAS analyses, which focus on individual CpG sites, regional analyses attenuate the burden of multiple testing and can also detect weaker signals that may be spread over wider regions. Regional analyses were first performed in each cohort and then meta-analyzed using the *dmrffR* package (Suderman et al., 2018). First, this method selects potentially DMRs by identifying genomic regions spanned by nominally significant CpG sites at most 500 bp apart. Second, EWAS summary statistics of all CpGs in the region are combined, while adjusting for correlations between sites in order to avoid inflating regional statistics. DMRs are then defined as having at least two spatially contiguous CpG sites. *p*-values for each region are adjusted for multiple tests using Bonferroni adjusted $p < 0.05$. As for the site-specific EWAS analyses, regional analyses were also repeated using “low prosocial” continuous scores at age 6–7 years.

3 | RESULTS

3.1 | Sample characteristics

In the main analyses, complete data for DNAm and the prosocial trajectories were available for 748 children (48.4% female) in ALSPAC, 891 children (49.6% female) in Generation R, 270 children (48.9% female) in INMA, and 186 children (48.1% female) in LINA. Detailed sample characteristics are presented in Table 1.

3.2 | Epigenome-wide association analysis and meta-analysis

Meta-analysis of the four cohorts ($N_{\text{total}} = 2,095$) did not identify significant associations between DNAm of 364,659 CpG sites in cord blood and chronic-low prosocial trajectories ($p < 2.4 \times 10^{-7}$; Figure 1). The most significant association ($\beta = -0.004$, $SE = 0.001$, $p = 8.37 \times 10^{-7}$) was found for cg03160045 annotated to *RBX1*, a gene previously associated with cognitive ability (Lee et al., 2018) and psychiatric disorders including anxiety, depression, and schizophrenia (Baselmans et al., 2019; Goes et al., 2015; Wu et al., 2020). In Table 2, we list the 24 CpG sites that were most strongly associated with prosocial trajectories at the suggestive level ($p < 5 \times 10^{-5}$). The direction of effect estimates was consistent across all cohorts for 20 of these sites. There was evidence for some genomic inflation ($\lambda = 1.25$; see Figure 1b for quantile–quantile plot).

Table S3, Panel A shows the lambda and number of cohort-level, genome-wide significant ($p < 2.4 \times 10^{-7}$) hits for typical versus chronic-low trajectories. We observed three significant cohort-level associations between DNAm and prosocial trajectories. These sites (cg22599122 [*HMGB4/CSMD2*] in INMA; cg03436478 [*SGCE/PEG10*]




FIGURE 1 Manhattan (a) and quantile–quantile (QQ) plots (b) showing meta-analytic associations between cord blood DNA methylation and chronic-low prosocial trajectory membership ($N_{\text{total}} = 2,095$). For Panel (a), the horizontal line indicates a genome-wide significance threshold of 2.4×10^{-7} . For Panel (b), $\lambda = 1.25$ [Color figure can be viewed at wileyonlinelibrary.com]

in ALSPAC; cg21552290 [*GABBR1*] in LINA) mapped to genes associated with neurodevelopmental and mental health-related phenotypes (see Table S3 and accompanying note for full details); however, none of the three hits overlapped with other cohorts.

Similar to the main trajectory-based analyses, no genome-wide associations were observed in meta-analysis of continuous “low prosocial” scores at age 6–7 years, with the most significant association ($\beta = 5.80$, $SE = 1.22$, $p = 1.92 \times 10^{-6}$) found for cg13899097 on the *RARRES3* gene (see Table S2), which is differentially expressed in autism (Seno et al., 2011). We found seven cohort-level associations between DNAm and continuous “low prosocial” scores (Table S3,

TABLE 2 Cord blood CpG sites most strongly associated with prosocial trajectories (typical vs. chronic-low) within meta-analysis

CpG site	Chr	Position	Nearest gene	Effect	SE	p-value	Direction	I ²	Heterogeneity p-value
cg03160045	22	41,345,978	RBX1	−0.004	0.001	8.37E-07	-----	42	.16
cg00979307	3	47,889,555	DHX30; MIR1226	−0.008	0.002	2.37E-06	-----	43	.15
cg01402746	6	30,616,468	C6orf136	0.004	0.001	2.62E-06	+++−	0	.66
cg14931071	6	12,717,776	PHACTR1	−0.008	0.002	4.20E-06	-----	54	.09
cg11495494	4	140,358,384	RAB33B ^a	−0.004	0.001	5.71E-06	-----	0	.69
cg16811988	3	42,103,138	TRAK1 ^a	−0.004	0.001	8.43E-06	-----	0	.72
cg18278596	14	105,915,860	MTA1	0.004	0.001	9.95E-06	+++−	32	.22
cg22117637	2	223,285,495	SGPP2 ^a	−0.007	0.002	1.02E-05	-----	25	.26
cg07572052	1	149,821,168	HIST2H2AA4; HIST2H2AA3	0.014	0.003	1.23E-05	++++	1	.39
cg19508437	13	35,517,410	NBEA	−0.001	0.000	1.38E-05	---+	0	.63
cg10182355	2	101,768,491	TBC1D8	0.003	0.001	1.76E-05	++++	0	.67
cg17394129	12	71,829,267	LGR5 ^a	−0.011	0.002	1.83E-05	-----	0	.51
cg00932677	4	187,776,068	FAT1 ^a	−0.008	0.002	2.17E-05	-----	0	.66
cg02007434	10	98,210,804	TLL2	−0.010	0.002	2.32E-05	-----	38	.19
cg20397078	7	28,966,605	CREB, TRIL, CPVL ^a	0.004	0.001	2.39E-05	++++	0	.79
cg07129067	2	180,451,222	ZNF385B	−0.005	0.001	2.41E-05	-----	0	.62
cg20313969	2	112,939,562	FBLN7	−0.012	0.003	3.05E-05	-----	56	.08
cg01825355	6	168,812,152	SMOC2 ^a	0.005	0.001	3.48E-05	−+++	27	.25
cg06850924	7	129,691,279	ZC3HC1	−0.002	0.000	3.48E-05	-----	20	.29
cg02074728	5	50,259,086	PARP8 ^a	0.003	0.001	3.52E-05	++++	0	.80
cg21165255	9	93,712,460	SYK ^a	−0.009	0.002	3.78E-05	-----	0	.42
cg14818546	7	69,296,381	AUTS2	−0.005	0.001	3.82E-05	-----	0	.74
cg24341770	6	105,148,882	HACE1 ^a	−0.009	0.002	4.05E-05	-----	0	.52
cg15727390	3	149,700,905	PFN2, LOC646903 ^a	0.010	0.002	4.75E-05	++++	0	.71

Note: Maximum $N = 2,095$; meta-analytic p -value $< 5 \times 10^{-5}$. The full model is adjusted for batch effects, estimated cell-type proportions, child sex, gestational age, maternal age in take/delivery, maternal smoking during pregnancy, maternal education, and maternal psychopathology. Chr: chromosome; Direction: direction of effect per cohort, where + and − signify positive and negative effect estimates, respectively (cohorts are listed in alphabetical order: Avon Longitudinal Study of Parents and Children [ALSPAC]; Generation R [GENR]; Infancia y Medio Ambiente [INMA]; Lifestyle and environmental factors and their [LINA]); I²: heterogeneity statistic describing variation attributable to heterogeneity across studies; high I² values suggest high heterogeneity.

^aAnnotation based on University of California Santa Cruz Known Gene fills in the nearest gene within 10 MB.

Panel C); three of which (cg06643156, cg09509433, and cg21144158) showed a consistent direction of effects across all cohorts. We found no evidence of genomic inflation for continuous analyses ($\lambda = 1.06$; quantile–quantile plot in Figure S1).

In candidate gene follow-up analysis, none of the CpGs annotated to *OXTR* reached gene-level Bonferroni significance (Table S4). Of note, although effect sizes were trivial, most CpG sites showed the same directions of effect as previous studies examining related phenotypes (e.g., CU traits), in that higher *OXTR* methylation at birth was associated with lower prosociality.

3.3 | Differentially methylated regions (DMRs) analysis

We identified no DMRs associated with “chronic-low prosocial” trajectory membership in the meta-analysis (Table 3). The top DMR associated with the chronic-low trajectory was located on chromosome

19:35,630,106–35,630,355 ($\beta = -.01$, $SE = 0.002$, $p = 2.5 \times 10^{-6}$), nearest to genes *FXYD1* and *LGI4*. We repeated the same analyses with “low prosocial” continuous scores. Again, no DMRs reached epigenome-wide significance in the meta-analysis, with a top DMR located on chromosome 9:132,805,739–132,805,979 ($\beta = -6.85$, $SE = 1.45$, $p = 2.3 \times 10^{-6}$; nearest gene *FBNP1*). DMRs that were suggestively associated with the severity of low prosocial behavior (based on continuous scores) are presented in Table S5.

3.4 | Power calculation

We performed a power calculation with a maximum effect estimate of 0.02 and alpha set at the 450k array significance threshold, $p < 2.4 \times 10^{-7}$. Results suggested that we had poor power (0.37) to detect the effect size observed given the current sample size ($N = 2,095$). Thus, it appears that larger sample sizes are needed to detect CpG sites associated with low prosocial trajectory membership.

TABLE 3 Differentially methylated regions associated with prosocial trajectories (typical vs. chronic-low) based on a suggestive threshold ($p < 5 \times 10^{-5}$)

Chr	Start	End	N of sites	Effect	SE	p-value	Nearest gene
19	35,630,106	35,630,355	5	−0.008	0.002	2.50E-06	<i>FXYP1, LIG4</i>
2	180,451,203	180,451,222	2	−0.006	0.001	8.03E-06	<i>ZNF385B</i>
11	3,876,513	3,876,808	7	0.000	0.000	1.92E-05	<i>STIM1</i>
11	3,647,365	3,647,654	3	−0.006	0.001	3.05E-05	<i>TRPC2</i>
8	142,316,216	142,316,861	3	−0.004	0.001	3.99E-05	<i>SLC45A4</i>
19	55,996,543	55,996,566	2	0.000	0.000	4.37E-05	<i>NAT14</i>

Abbreviation: Chr, chromosome.

Our power calculation also indicated that, based on the effect sizes and significance thresholds in the current study, a minimum sample size of 2,985 would be required to achieve 80% power. We did not perform this power calculation for the continuous score analysis as these power estimates would have been inflated by the large coefficient estimates of DNAm that were weighted from robust regression (the maximum effect size was 18.82).

4 | DISCUSSION

The current study examined whether DNAm patterns at birth prospectively associate with persistently low levels of prosocial behavior across childhood, using highly comparable data from four independent cohort studies. Although we identified three CpG sites from cord blood that differentiated persistently “low prosocial” children from their more typically developing peers within individual cohorts, none of these associations were replicated in another sample. Moreover, our EWAS meta-analysis ($N = 2,095$) did not identify any significant associations across the four cohorts at a 450k genome-wide level. This was true when examining both developmental trajectories of chronic-low versus typical prosocial behavior, as well as dimensional scores at similar time-points for all cohorts. We confirmed our findings using follow-up regional analyses to address non-independence between CpG sites, which yielded no meta-analytical DMRs following multiple testing corrections. Overall, our findings do not support an association between neonatal DNAm and low prosocial behavior in childhood.

It is important to consider the negative findings of the present study in the context of its considerable methodological strengths. First, our analyses drew on four independent large-scale cohorts with prospective designs. Second, availability of repeated measures of prosocial behavior enabled us to characterize participants based on sustained levels of low prosociality across childhood, capturing a discrete group of children at elevated risk for a range of psychiatric disorders (Flynn et al., 2015; Nantel-Vivier et al., 2014). This longitudinal approach was supplemented with dimension-based analyses (i.e., total scores) to explore associations across the entire continuum of prosocial behavior in the general population. Third, comprehensive harmonization between the four cohorts, in terms of both

normalization of DNAm data and phenotypic measurement, was performed to reduce study heterogeneity and maximize comparability for replication efforts and the meta-analysis. Fourth, our analytic strategy featured both site-specific and regional analyses, which can improve power and detect functionally relevant findings across genomic regions (Michels et al., 2013).

Despite the strengths of our design and analysis, we did not identify any genome-wide associations between neonatal DNAm and childhood low prosocial behavior. We offer several potential explanations for our negative findings, which also consider a number of key study limitations. First, it appears from our power analysis that, even with the large sample size offered by these combined datasets, our analyses were underpowered to detect what appear to be reasonably small associations, particularly given the relatively low numbers of participants classified as “chronic-low prosocial.” However, our dimensional analyses involving continuous low prosocial scores across the entire sample showed similarly negative results. More generally, there may be insufficient individual variation within our data to detect differences in DNAm patterns in relation to prosocial behavior. For example, the interquartile ranges of low prosocial behavior at age 6–7 years were identical in all cohorts (median = 1; inter-quartile range: 0–3), indicating that most individuals show moderate to high levels of prosocial behavior (Figure S2). However, we found that these chronic-low and typical groups did differ on many behavioral outcomes (i.e., empathy, social-cognitive difficulties, and CU traits) that are known to associate with prosociality.

Second, the lack of associations could be the result of methodological differences across cohorts, despite the application of stringent harmonization standards to both phenotypic and DNAm data. Specifically, low prosocial behavior was assessed by the same reporting source (i.e., mothers), with the same instrument, and at comparable time-points in all samples. Nonetheless, we cannot rule out perceptual biases due to single informants or situation-specificity that may have influenced the results of this study. The use of mother reports allowed us to model prosocial behavior starting in early childhood, when child self-reports are not reliable. However, the inclusion of multiple informants (e.g., teacher report, child self-report) in future research may help to provide greater differentiation of the child's prosocial behavior across different contexts. Moreover, differences in quality control, environmental exposures, and other unknown factors

could still contribute to cohort-specific methylation patterns and consequently dilute potential associations during meta-analysis (Joubert et al., 2016).

Third, our operationalization of prosocial behavior using a relatively brief measure may be overly broad. Our use of trajectory-based and dimensional approaches enabled us to examine associations for both a discrete group of “persistently low prosocial” children, as well as across a broader continuum of severity. Although the SDQ’s prosocial scale has been shown to robustly associate with psychosocial and functional outcomes (Meehan, Maughan, & Barker, 2019), it may be too nonspecific to reliably detect a unified underlying biological substrate. Moreover, prosocial behavior is increasingly viewed as a multi-faceted phenotype (Van IJzendoorn & Bakermans-Kranenburg, 2014) that incorporates, and interacts with, a range of motivational, cognitive, and affective processes, including altruism, perspective-taking, Theory of Mind, and empathic concern, as well as contextual factors (e.g., Imuta, Henry, Slaughter, Selcuk, & Ruffman, 2016; Preckel, Kanske, & Singer, 2018; Van der Graaff, Carlo, Crocetti, Koot, & Branje, 2018). Although it has been suggested that genetic associations with broader measures of constructs tend to be stronger (Dawson et al., 2002; Thapar et al., 2006), phenotypic heterogeneity within the measure, such as helping versus comforting (Paulus, Kuhn-Popp, Licata, Sodian, & Meinhardt, 2013), may differ in their developmental course and epigenetic patterns. Therefore, a measure capturing more granular components of this multidimensional construct may better elucidate specific associations with individual biological markers.

Fourth, we chose to focus on the potential effects of DNAm at birth (cord blood). This was informed by several previous epigenetic studies in birth cohorts, including ALSPAC and Generation R, which have found prospective associations between DNAm at birth (vs. later in childhood) and outcomes related to prosocial behavior such as CU traits (Cecil et al., 2014), conduct problems (Cecil et al., 2018), and social communication deficits (Rijlaarsdam et al., 2021). Furthermore, DNAm patterns at birth—but not in childhood—may mediate risk for psychiatric problems in later life; for example, DNAm alterations at birth may impact early neurodevelopment and downstream phenotypes that persist despite changes in DNAm patterns *per se*. Given our prospective study design, this focus on birth DNAm also allowed us to disentangle the directionality of observed effects (i.e., examining DNAm pre-manifestation of prosocial behavior), while also maximizing the available analytic sample. However, we cannot exclude the possibility of time-specific associations later in life. Of particular note, prosocial development is influenced by environmental processes in early childhood, most notably parental socialization (Eisenberg, Spinrad, & Knafo-Noam, 2015; Paulus, 2014), whose effects may not be reflected in DNAm at birth, but could engender changes in a child’s epigenetic profile that are only evidenced later in development. For example, Barker, Cecil, et al. (2018) found that postnatal environmental adversity was associated with age-7 DNAm, which then affected a subsequent vulnerability for internalizing problems in late childhood. It will be of interest in the future to utilize repeated

measures of DNAm to explore potential associations with postnatal environmental influences and capture dynamic changes across the lifespan.

DNAm is also highly tissue-specific (Davies et al., 2012; Dempster et al., 2014) and, like other epigenetic studies, our ability to detect effects for brain-related behavior traits may therefore be limited by a reliance on blood-based samples. Although blood samples can be reflective of DNAm in other tissue-relevant samples, including the brain (Hannon, Lunnon, Schalkwyk, & Mill, 2015; Qi et al., 2018), with some epigenetic findings successfully replicated across tissues (Kaminsky et al., 2012), the extent to which these reflect changes in the brain, as well as the functional consequences of these changes for gene expression, remains unclear. Incorporating additional tissue sources (e.g., blood, brain, and saliva) may better reveal common and differential patterns of methylation in relation to low prosociality; however, this is often not feasible in population-based cohorts. Finally, the 450k array only covers 1.7% of the total number of CpGs in the human genome, and selected CpG sites are substantially biased toward promoter and gene body regions, which limits the capacity to detect potential associations between DNAm at other, unmeasured, CpGs with prosocial behavior.

Although not replicated in other cohorts, we note that one site (cg03436478), annotated to *SGCE* and *PEG10* genes, showed significant differences in birth DNAm between chronic-low and typical prosocial trajectories in ALSPAC, as well as the same direction of effect across all cohorts. Both *SGCE* and *PEG10*, as imprinted genes, are highly expressed during embryonic development, which is a crucial period for establishing and maintaining methylation profiles (Kainz et al., 2007). With specific regard to mental health, *SGCE* has previously been linked to anxiety, mood disorders, and schizophrenia in genetic studies (Lam et al., 2019; Peall et al., 2013), while DNAm in the vicinity of *PEG10* has been related to maternal depression during pregnancy (Liu et al., 2012) and increased risk for child maltreatment (Yang et al., 2013). Furthermore, differential methylation in these two imprinted genes has been associated with neurobehavioral outcomes in a sex-specific manner: increased methylation of *SGCE/PEG10* was associated with increased risk of atypical behavior in boys, but decreased risk in girls (House et al., 2018). In this study, we did find evidence of sex differences between prosocial trajectories; specifically, boys were more prevalent in the chronic-low prosocial groups and showed lower prosocial behavior at age 6–7 years. However, due to the size of our analytic sample(s), we had insufficient power to perform epigenome-wide stratified and interaction analyses by sex. In future research with larger samples, it would be worthwhile to analyze whether DNAm was differentially associated with prosocial behavior between boys and girls.

Finally, our findings did not replicate candidate genes previously implicated in prosocial-related phenotypes, namely *OXTR*. One previous ALSPAC study reported prospective associations between a higher level of *OXTR* DNAm at birth and adolescent CU traits at age 13 (Cecil et al., 2014). Of note, while CU traits are strongly correlated with lower levels of prosocial behavior (Barker, Oliver, Viding, Salekin, & Maughan, 2011), recent factor analyses have shown that

the broader “low prosocial” phenotype denotes some unique variance over and above that shared with CU traits, and is therefore likely to be characterized by a distinct pathophysiological profile (Meehan, Hawes, Salekin, & Barker, 2019).

In summary, this study is the first to examine the association between 450k genome-wide DNAm at birth and childhood prosocial behavior, using data from four independent longitudinal birth cohorts. We also found no differentially methylated positions or regions that were associated with low prosocial behavior at an epigenome-wide significant threshold (based on examination of longitudinal trajectories and dimensional “low prosocial” scores at a single time-point). Together these findings suggest that, at least at birth and based on the current sample size, there is no robust evidence for a prospective association between birth DNAm and childhood prosocial behavior. Overall, the current study highlights the importance of multi-cohort approaches to replicate epigenetic associations and reduce the risk of false positive discoveries (Rijlaarsdam et al., 2016). Future research with larger sample sizes, more detailed phenotype definitions, and additional epigenetic time-points will help to clarify the relation between DNAm and low prosocial behavior in childhood.

ACKNOWLEDGEMENTS

ALSPAC: We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses. The ALSPAC website contains details of all available data through a data dictionary and variable search tool: <http://www.bristol.ac.uk/alspac/researchers/our-data/>. Informed consent was obtained from all participants and ethical approval was obtained from the ALSPAC Law and Ethics Committee and Local Research Committees.

The UK Medical Research Council (MRC) and Wellcome Trust (102215/2/13/2) and the University of Bristol provide core support for ALSPAC. A comprehensive list of grants funding is available on the ALSPAC website (<http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf>). Analysis of ALSPAC data was funded by the UK Economic and Social Research Council (ESRC; ES/N000498/1). ARIES was funded by the Biotechnology and Biological Sciences Research Council (BBSRC; BBI025751/1 and BB/I025263/1). Supplementary funding to generate DNA methylation data which are (or will be) included in ARIES has been obtained from the MRC, ESRC, National Institutes of Health, and other sources. ARIES is maintained under the auspices of the MRC Integrative Epidemiology Unit at the University of Bristol (MC_UU_00011/4 and MC_UU_00011/5). The ALSPAC methylation component of this article was supported by a Eunice Kennedy Shriver National Institute of Child and Human Development grant to EDB (R01HD068437) with further funding from the ESRC (ES/R005516/1). AJM was supported by the ESRC (ES/J500057/1) and National University of Ireland. This publication is the work of the authors, and EDB and CAMC serve as guarantors for the contents of this article.

Generation R: The Generation R Study is conducted by the Erasmus Medical Center in close collaboration with the School of Law and Faculty of Social Sciences at Erasmus University Rotterdam, the Municipal Health Service Rotterdam area, the Rotterdam Homecare Foundation, and the Stichting Trombosedienst & Artsenlaboratorium Rijnmond (STAR-MDC), Rotterdam. We gratefully acknowledge the contribution of children and parents, general practitioners, hospitals, midwives, and pharmacies in Rotterdam. The Generation R website contains details of ongoing data collection: <http://generationr.nl/researchers/data-collection/>. Written informed consent was obtained for all participants, and the study protocol was approved by the Medical Ethical Committee of the Erasmus University Medical Center, Rotterdam. Generation and management of the Illumina 450K methylation array for EWAS data in Generation R was executed by the Human Genotyping Facility of the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, The Netherlands. We thank Mr. Michael Verbiest, Ms. Mila Jhamai, Ms. Sarah Higgins, Mr. Marijn Verkerk and Dr. Lisette Stolk for their help in creating the EWAS database. We thank Dr. A. Teumer for his work on the quality control and normalization scripts.

The general design of the Generation R Study is made possible by financial support from the Erasmus Medical Center Rotterdam, Erasmus University Rotterdam, the Netherlands Organization for Health Research and Development and the Ministry of Health, Welfare and Sport. The EWAS data was funded by a grant from the Netherlands Genomics Initiative (NGI)/Netherlands Organization for Scientific Research (NWO) Netherlands Consortium for Healthy Aging (NCHA; project nr. 050-060-810), funds from the Genetic Laboratory of the Department of Internal Medicine, Erasmus MC, and a grant from the National Institute of Child and Human Development (R01HD068437). This project received funding from the European Union's Horizon 2020 research and innovation program (733,206, LifeCycle; 848,158, EarlyCause) and from the European Joint Programming Initiative “A Healthy Diet for a Healthy Life” (JPI HDHL, NutriPROGRAM project, ZonMw the Netherlands no.529051022 and PREcisE project no. P75416). CAMC has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 707404. CAMC and EW are supported by grant agreement No 848158 (EarlyCause). ML is supported by the scholarship from the China Scholarship Council (201706990036). MHvIJ and MJB-K are supported by the Dutch Ministry of Education, Culture, and Science and the Netherlands Organization for Scientific Research (NWO grant No. 024.001.003, Consortium on Individual Development). MJB-K was supported by the European Research Council (ERC AdG 669,249). MHvIJ and APCH are also supported by a Spinoza Award of the Netherlands Organization for Scientific Research to MHvIJ.

INMA: The epigenetic studies in INMA were mainly funded by grants from Instituto de Salud Carlos III (Red INMA G03/176, CB06/02/0041, CP18/00018), Spanish Ministry of Health (FIS-PI04/1436, FIS-PI08/1151 including FEDER funds, FIS-PI11/00610, FIS-FEDER-PI06/0867, FIS-FEDER-PI03-1615), Generalitat de Catalunya-CIRIT 1999SGR 00241, Fundació La marató de TV3

(090430), EU Commission (261357-MeDALL: Mechanisms of the Development of ALLergy), and European Research Council (268479-BREATHE: BRain dEvelopment and Air polluTion ultrafine particles in scHool childrEn). MCT is funded by a Juan de la Cierva—Formación Postdoctoral Contract awarded by Ministry of Science and Innovation (FJC2018-036335-I). The 4, 7, and 11 year follow-up visits was funded by grants from Instituto de Salud Carlos III (Red INMA G03/176; CB06/02/0041; PI041436; PI081151 incl. FEDER funds; PI12/01890 incl. FEDER funds; CP13/00054 incl. FEDER funds; PI15/00118 incl. FEDER funds; CP16/00128 incl. FEDER funds; PI16/00118 incl. FEDER funds; PI16/00261 incl. FEDER funds; PI18/00547 incl. FEDER funds), CIBERESP, Generalitat de Catalunya-CIRIT 1999SGR 00241, Generalitat de Catalunya-AGAUR (2009 SGR 501, 2014 SGR 822), Fundació La marató de TV3 (090430), Spanish Ministry of Economy and Competitiveness (SAF2012-32991 incl. FEDER funds), Agence Nationale de Securite Sanitaire de l'Alimentation de l'Environnement et du Travail (1262C0010; EST-2016 RF-21), EU Commission (261357, 308333, 603794 and 634453). We acknowledge support from the Spanish Ministry of Science and Innovation and State Research Agency through the “Centro de Excelencia Severo Ochoa 2019-2023” Program (CEX2018-000806-S), and support from the Generalitat de Catalunya through the CERCA Program.

LINA: Core funding for the LINA study is provided by the Helmholtz Center for Environmental Research—UFZ / Department of Environmental Immunology, with methylation analysis supported by the German Cancer Research Centre—DKFZ.

CONFLICT OF INTEREST

All named authors report no conflict of interests.

AUTHOR CONTRIBUTIONS

Mannan Luo, Alan J. Meehan, Esther Walton, Irene Pappa, Henning Tiemeier, Marinus H. van IJzendoorn, Edward D. Barker, Charlotte A. M. Cecil: Study conception and design. **Mannan Luo, Alan J. Meehan, Esther Walton, Stefan Röder, Gunda Herberth, Ana C. Zencussen, Marta Cosín-Tomás, Jordi Sunyer, Rosa H. Mulder, Janine F. Felix, Caroline Relton, Matthew Suderman, Edward D. Barker, Charlotte A. M. Cecil:** Data acquisition, preparation, and/or quality control. **Mannan Luo, Alan J. Meehan, Esther Walton, Stefan Röder, Marta Cosín-Tomás, Andrea P. Cortes Hidalgo:** Statistical analysis. **Mannan Luo, Alan J. Meehan, Edward D. Barker, Charlotte A. M. Cecil:** Drafting of the article. **Edward D. Barker, Charlotte A. M. Cecil:** Study supervision. All authors contributed to the interpretation of results, provided critical revisions of the article, and approved the final version.

DATA AVAILABILITY STATEMENT

All summary statistics from the EWAS meta-analysis presented in this study will be made available on Zenodo/EWAS catalog.

ORCID

Alan J. Meehan  <https://orcid.org/0000-0001-7169-997X>

Esther Walton  <https://orcid.org/0000-0002-0935-2200>

Stefan Röder  <https://orcid.org/0000-0001-9367-663X>

Gunda Herberth  <https://orcid.org/0000-0003-0212-3509>

Ana C. Zencussen  <https://orcid.org/0000-0003-3544-4552>

Marta Cosín-Tomás  <https://orcid.org/0000-0001-5012-8983>

Jordi Sunyer  <https://orcid.org/0000-0002-2602-4110>

Rosa H. Mulder  <https://orcid.org/0000-0002-2382-1704>

Andrea P. Cortes Hidalgo  <https://orcid.org/0000-0002-7426-6181>

Marian J. Bakermans-Kranenburg  <https://orcid.org/0000-0001-7763-0711>

Janine F. Felix  <https://orcid.org/0000-0002-9801-5774>

Caroline Relton  <https://orcid.org/0000-0003-2052-4840>

Matthew Suderman  <https://orcid.org/0000-0002-2715-9930>

Irene Pappa  <https://orcid.org/0000-0003-4936-5791>

Rianne Kok  <https://orcid.org/0000-0003-3620-7313>

Henning Tiemeier  <https://orcid.org/0000-0002-4395-1397>

Marinus H. van IJzendoorn  <https://orcid.org/0000-0003-1144-454X>

Edward D. Barker  <https://orcid.org/0000-0002-9914-8958>

Charlotte A. M. Cecil  <https://orcid.org/0000-0002-2389-5922>

REFERENCES

- Bakermans-Kranenburg, M. J., & Van IJzendoorn, M. H. (2014). A sociability gene? Meta-analysis of oxytocin receptor genotype effects in humans. *Psychiatric Genetics*, 24(2), 45–51. <https://doi.org/10.1097/YPG.0b013e3283643684>
- Barker, E. D., Cecil, C. A. M., Walton, E., Houtepen, L. C., O'Connor, T. G., Danese, A., ... Roberts, S. (2018). Inflammation-related epigenetic risk and child and adolescent mental health: A prospective study from pregnancy to middle adolescence. *Development and Psychopathology*, 30(3), 1145–1156. <https://doi.org/10.1017/S0954579418000330>
- Barker, E. D., Oliver, B. R., Viding, E., Salekin, R. T., & Maughan, B. (2011). The impact of prenatal maternal risk, fearless temperament and early parenting on adolescent callous-unemotional traits: A 14-year longitudinal investigation. *Journal of Child Psychology and Psychiatry*, 52(8), 878–888. <https://doi.org/10.1111/j.1469-7610.2011.02397.x>
- Barker, E. D., Walton, E., & Cecil, C. A. M. (2018). Annual research review: DNA methylation as a mediator in the association between risk exposure and child and adolescent psychopathology. *Journal of Child Psychology and Psychiatry*, 59(4), 303–322. <https://doi.org/10.1111/jcpp.12782>
- Barker, E. D., Walton, E., Cecil, C. A. M., Rowe, R., Jaffee, S. R., Maughan, B., ... Gaunt, T. R. (2018). A methylome-wide association study of trajectories of oppositional defiant behaviors and biological overlap with attention deficit hyperactivity disorder. *Child Development*, 89(5), 1839–1855. <https://doi.org/10.1111/cdev.12957>
- Baselmans, B. M. L., Jansen, R., Ip, H. F., van Dongen, J., Abdellaoui, A., van de Weijer, M. P., ... Bartels, M. (2019). Multivariate genome-wide analyses of the well-being spectrum. *Nature Genetics*, 51(3), 445–451. <https://doi.org/10.1038/s41588-018-0320-8>
- Caprara, G. V., Barbaranelli, C., Pastorelli, C., Bandura, A., & Zimbardo, P. G. (2000). Prosocial foundations of children's academic achievement. *Psychological Science*, 11(4), 302–306. <https://doi.org/10.1111/1467-9280.00260>
- Cecil, C. A. M., Lysenko, L. J., Jaffee, S. R., Pingault, J. B., Smith, R. G., Relton, C. L., ... Barker, E. D. (2014). Environmental risk, oxytocin receptor gene (OXTR) methylation and youth callous-unemotional traits: A 13-year longitudinal study. *Molecular Psychiatry*, 19(10), 1071–1077. <https://doi.org/10.1038/mp.2014.95>
- Cecil, C. A. M., Walton, E., Jaffee, S. R., O'Connor, T., Maughan, B., Relton, C. L., ... Barker, E. D. (2018). Neonatal DNA methylation and

- early-onset conduct problems: A genome-wide, prospective study. *Development and Psychopathology*, 30(2), 383–397. <https://doi.org/10.1017/S095457941700092x>
- Chen, Y. A., Lemire, M., Choufani, S., Butcher, D. T., Grafodatskaya, D., Zanke, B. W., ... Weksberg, R. (2013). Discovery of cross-reactive probes and polymorphic CpGs in the Illumina Infinium Human Methylation450 microarray. *Epigenetics*, 8(2), 203–209. <https://doi.org/10.4161/epi.23470>
- Dadds, M. R., Moul, C., Cauchi, A., Dobson-Stone, C., Hawes, D. J., Brennan, J., & Ebstein, R. E. (2014). Methylation of the oxytocin receptor gene and oxytocin blood levels in the development of psychopathy. *Development and Psychopathology*, 26(1), 33–40. <https://doi.org/10.1017/S0954579413000497>
- Davies, M. N., Volta, M., Pidsley, R., Lunnon, K., Dixit, A., Lovestone, S., ... Mill, J. (2012). Functional annotation of the human brain methylome identifies tissue-specific epigenetic variation across brain and blood. *Genome Biology*, 13(6), R43. <https://doi.org/10.1186/gb-2012-13-6-r43>
- Dawson, G., Webb, S., Schellenberg, G. D., Dager, S., Friedman, S., Aylward, E., & Richards, T. (2002). Defining the broader phenotype of autism: Genetic, brain, and behavioral perspectives. *Development and Psychopathology*, 14(3), 581–611. <https://doi.org/10.1017/S0954579402003103>
- Dempster, E. L., Wong, C. C. Y., Lester, K. J., Burrage, J., Gregory, A. M., Mill, J., & Eley, T. C. (2014). Genome-wide methylomic analysis of monozygotic twins discordant for adolescent depression. *Biological Psychiatry*, 76(12), 977–983. <https://doi.org/10.1016/j.biopsych.2014.04.013>
- Eisenberg, N., Fabes, R. A., & Spinrad, T. L. (2006). Prosocial development. In N. Eisenberg & W. Damon (Eds.), *Handbook of child psychology: Vol. 3. Social, emotional, and personality development* (6th ed., pp. 646–718). New York: Wiley.
- Eisenberg, N., Spinrad, T. L., & Knafo-Noam, A. (2015). Prosocial development. In R. M. Lerner (Ed.), *Handbook of child psychology and developmental science* (Vol. 3, 7th ed., pp. 610–656). New York: Wiley.
- Felix, J. F., Joubert, B. R., Baccarelli, A. A., Sharp, G. C., Almqvist, C., Annesi-Maesano, I., ... London, S. J. (2018). Cohort profile: Pregnancy and childhood epigenetics (PACE) consortium. *International Journal of Epidemiology*, 47(1), 22–23u. <https://doi.org/10.1093/ije/dyx190>
- Flynn, E., Ehrenreich, S. E., Beron, K. J., & Underwood, M. K. (2015). Prosocial behavior: Long-term trajectories and psychosocial outcomes. *Social Development*, 24(3), 462–482. <https://doi.org/10.1111/sode.12100>
- Fraser, A., Macdonald-Wallis, C., Tilling, K., Boyd, A., Golding, J., Smith, G. D., ... Lawlor, D. A. (2013). Cohort profile: The Avon longitudinal study of parents and children: ALSPAC mothers cohort. *International Journal of Epidemiology*, 42(1), 97–110. <https://doi.org/10.1093/ije/dys066>
- Gervin, K., Salas, L. A., Bakulski, K. M., van Zelm, M. C., Koestler, D. C., Wiencke, J. K., ... Jones, M. J. (2019). Systematic evaluation and validation of reference and library selection methods for deconvolution of cord blood DNA methylation data. *Clinical Epigenetics*, 11(1), 125. <https://doi.org/10.1186/s13148-019-0717-y>
- Goes, F. S., McGrath, J., Avramopoulos, D., Wolyniec, P., Pirooznia, M., Ruczinski, I., ... Pulver, A. E. (2015). Genome-wide association study of schizophrenia in Ashkenazi Jews. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 168(8), 649–659. <https://doi.org/10.1002/ajmg.b.32349>
- Goodman, R. (1997). The strengths and difficulties questionnaire: A research note. *Journal of Child Psychology and Psychiatry*, 38(5), 581–586. <https://doi.org/10.1111/j.1469-7610.1997.tb01545.x>
- Goodman, R. (2001). Psychometric properties of the strengths and difficulties questionnaire. *Journal of the American Academy of Child and Adolescent Psychiatry*, 40(11), 1337–1345. <https://doi.org/10.1097/00004583-200111000-00015>
- Gregory, A. M., Light-Häusermann, J. H., Rijdsdijk, F., & Eley, T. C. (2009). Behavioral genetic analyses of prosocial behavior in adolescents. *Developmental Science*, 12(1), 165–174. <https://doi.org/10.1111/j.1467-7687.2008.00739.x>
- Guxens, M., Ballester, F., Espada, M., Fernandez, M. F., Grimalt, J. O., Ibarluzea, J., ... Project, I. (2012). Cohort profile: The INMA-INfancia y Medio Ambiente-(environment and childhood) project. *International Journal of Epidemiology*, 41(4), 930–940. <https://doi.org/10.1093/ije/dyr054>
- Hannon, E., Lunnon, K., Schalkwyk, L., & Mill, J. (2015). Interindividual methylomic variation across blood, cortex, and cerebellum: Implications for epigenetic studies of neurological and neuropsychiatric phenotypes. *Epigenetics*, 10(11), 1024–1032. <https://doi.org/10.1080/15592294.2015.1100786>
- Herberth, G., Hinz, D., Roder, S., Schlink, U., Sack, U., Diez, U., ... Lehmann, I. (2011). Maternal immune status in pregnancy is related to offspring's immune responses and atopy risk. *Allergy*, 66(8), 1065–1074. <https://doi.org/10.1111/j.1398-9995.2011.02587.x>
- Horn, L., Hungerlander, N. A., Windhager, S., Bugnyar, T., & Massen, J. J. M. (2018). Social status and prenatal testosterone exposure assessed via second-to-fourth digit ratio affect 6-9-year-old children's prosocial choices. *Scientific Reports*, 8, 8–11. <https://doi.org/10.1038/s41598-018-27468-0>
- House, J. S., Mendez, M., Maguire, R. L., Gonzalez-Nahm, S., Huang, Z. Q., Daniels, J., ... Hoyo, C. (2018). Periconceptional maternal Mediterranean diet is associated with favorable offspring behaviors and altered CpG methylation of imprinted genes. *Frontiers in Cell and Developmental Biology*, 6, 107. <https://doi.org/10.3389/fcell.2018.00107>
- Imuta, K., Henry, J. D., Slaughter, V., Selcuk, B., & Ruffman, T. (2016). Theory of mind and prosocial behavior in childhood: A meta-analytic review. *Developmental Psychology*, 52(8), 1192–1205. <https://doi.org/10.1037/dev0000140>
- Jambon, M., Madigan, S., Plamondon, A., & Jenkins, J. (2019). Developmental trajectories of physical aggression and prosocial behavior in early childhood: Family antecedents and psychological correlates. *Developmental Psychology*, 55(6), 1211–1225. <https://doi.org/10.1037/dev0000714>
- Joubert, B. R., Felix, J. F., Yousefi, P., Bakulski, K. M., Just, A. C., Breton, C., ... London, S. J. (2016). DNA methylation in newborns and maternal smoking in pregnancy: Genome-wide consortium meta-analysis. *American Journal of Human Genetics*, 98(4), 680–696. <https://doi.org/10.1016/j.ajhg.2016.02.019>
- Kainz, B., Shehata, M., Bilban, M., Kienle, D., Heintel, D., Krömer-Holinger, E., ... Jäger, U. (2007). Overexpression of the paternally expressed gene 10 (PEG10) from the imprinted locus on chromosome 7q21 in high-risk B-cell chronic lymphocytic leukemia. *International Journal of Cancer*, 121(9), 1984–1993. <https://doi.org/10.1002/ijc.22929>
- Kaminsky, Z., Tochigi, M., Jia, P., Pal, M., Mill, J., Kwan, A., ... Petronis, A. (2012). A multi-tissue analysis identifies HLA complex group 9 gene methylation differences in bipolar disorder. *Molecular Psychiatry*, 17(7), 728–740. <https://doi.org/10.1038/mp.2011.64>
- Knafo-Noam, A., Israel, S., & Ebstein, R. P. (2011). Heritability of children's prosocial behavior and differential susceptibility to parenting by variation in the dopamine receptor D4 gene. *Development and Psychopathology*, 23(1), 53–67. <https://doi.org/10.1017/S0954579410000647>
- Knafo-Noam, A., & Plomin, R. (2006). Prosocial behavior from early to middle childhood: Genetic and environmental influences on stability and change. *Developmental Psychology*, 42(5), 771–786. <https://doi.org/10.1037/0012-1649.42.5.771>
- Knafo-Noam, A., Uzevovsky, F., Israel, S., Davidov, M., & Zahn-Waxler, C. (2015). The prosocial personality and its facets: Genetic and environmental architecture of mother-reported behavior of 7-year-old twins. *Frontiers in Psychology*, 6, 112. <https://doi.org/10.3389/fpsyg.2015.00112>

- Knafo-Noam, A., Vertsberger, D., & Israel, S. (2018). Genetic and environmental contributions to children's prosocial behavior: Brief review and new evidence from a reanalysis of experimental twin data. *Current Opinion in Psychology*, 20, 60–65. <https://doi.org/10.1016/j.copsyc.2017.08.013>
- Kokko, K., Tremblay, R. E., Lacourse, E., Nagin, D. S., & Vitaro, F. (2006). Trajectories of prosocial behavior and physical aggression in middle childhood: Links to adolescent school dropout and physical violence. *Journal of Research on Adolescence*, 16(3), 403–428. <https://doi.org/10.1111/j.1532-7795.2006.00500.x>
- Kooijman, M. N., Kruithof, C. J., van Duijn, C. M., Duijts, L., Franco, O. H., van IJzendoorn, M. H., ... Jaddoe, V. W. V. (2016). The Generation R Study: Design and cohort update 2017. *European Journal of Epidemiology*, 31(12), 1243–1264. <https://doi.org/10.1007/s10654-016-0224-9>
- Ladd-Acosta, C., & Fallin, M. D. (2016). The role of epigenetics in genetic and environmental epidemiology. *Epigenomics*, 8(2), 271–283. <https://doi.org/10.2217/epi.15.102>
- Lam, M., Chen, C. Y., Li, Z. Q., Martin, A. R., Bryois, J., Ma, X. X., ... neT-work-, G. R. S. (2019). Comparative genetic architectures of schizophrenia in East Asian and European populations. *Nature Genetics*, 51(12), 1670–1678. <https://doi.org/10.1038/s41588-019-0512-x>
- Layous, K., Nelson, S. K., Oberle, E., Schonert-Reichl, K. A., & Lyubomirsky, S. (2012). Kindness counts: Prompting prosocial behavior in preadolescents boosts peer acceptance and well-being. *PLoS One*, 7(12), e51380. <https://doi.org/10.1371/journal.pone.0051380>
- Lee, J. J., Wedow, R., Okbay, A., Kong, E., Maghzian, O., Zacher, M., ... Cesarini, D. (2018). Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nature Genetics*, 50(8), 1112–1121. <https://doi.org/10.1038/s41588-018-0147-3>
- Liu, Y., Murphy, S. K., Murtha, A. P., Fuemmeler, B. F., Schildkraut, J., Huang, Z. Q., ... Hoyo, C. (2012). Depression in pregnancy, infant birth weight and DNA methylation of imprint regulatory elements. *Epigenetics*, 7(7), 735–746. <https://doi.org/10.4161/epi.20734>
- Loomans, E. M., van der Stelt, O., van Eijsden, M., Gemke, R. J. B. J., Vrijlkotte, T., & Van den Bergh, B. R. H. (2011). Antenatal maternal anxiety is associated with problem behaviour at age five. *Early Human Development*, 87(8), 565–570. <https://doi.org/10.1016/j.earlhumdev.2011.04.014>
- Marsh, A. A. (2019). The caring continuum: Evolved hormonal and proximal mechanisms explain prosocial and antisocial extremes. *Annual Review of Psychology*, 70(1), 347–371. <https://doi.org/10.1146/annurev-psych-010418-103010>
- McCartney, D. L., Walker, R. M., Morris, S. W., McIntosh, A. M., Porteous, D. J., & Evans, K. L. (2016). Identification of polymorphic and off-target probe binding sites on the Illumina Infinium MethylationEPIC BeadChip. *Genomics Data*, 9, 22–24. <https://doi.org/10.1016/j.gdata.2016.05.012>
- McGowan, P. O., & Roth, T. L. (2015). Epigenetic pathways through which experiences become linked with biology. *Development and Psychopathology*, 27(2), 637–648. <https://doi.org/10.1017/S0954579415000206>
- Meehan, A. J., Hawes, D. J., Salekin, R. T., & Barker, E. D. (2019). Shared and unique variances of interpersonal callousness and low prosocial behavior. *Psychological Assessment*, 31(3), 376–388. <https://doi.org/10.1037/pas0000675>
- Meehan, A. J., Maughan, B., & Barker, E. D. (2019). Health and functional outcomes for shared and unique variances of interpersonal callousness and low prosocial behavior. *Journal of Psychopathology and Behavioral Assessment*, 41(3), 353–365. <https://doi.org/10.1007/s10862-019-09756-9>
- Michels, K. B., Binder, A. M., Dedeurwaerder, S., Epstein, C. B., Greally, J. M., Gut, I., ... Irizarry, R. A. (2013). Recommendations for the design and analysis of epigenome-wide association studies. *Nature Methods*, 10(10), 949–955. <https://doi.org/10.1038/nmeth.2632>
- Min, J. L., Hemani, G., Smith, G. D., Relton, C., & Suderman, M. (2018). Meffil: Efficient normalization and analysis of very large DNA methylation datasets. *Bioinformatics*, 34(23), 3983–3989. <https://doi.org/10.1093/bioinformatics/bty476>
- Moul, C., Dobson-Stone, C., Brennan, J., Hawes, D. J., & Dadds, M. R. (2015). Serotonin 1B receptor gene (HTR1B) methylation as a risk factor for callous-unemotional traits in antisocial boys. *PLoS One*, 10(5), e0126903. <https://doi.org/10.1371/journal.pone.0126903>
- Mulder, R. H., Neumann, A., Cecil, C. A. M., Walton, E., Houtepen, L. C., Simpkin, A. J., ... Suderman, M. (2021). Epigenome-wide change and variation in DNA methylation in childhood: Trajectories from birth to late adolescence. *Human Molecular Genetics*, 30(1), 119–134. <https://doi.org/10.1093/hmg/ddaa280>
- Muthén, L. K., & Muthén, B. O. (2012). *Mplus Version 7 user's guide*. Los Angeles, CA: Muthén & Muthén.
- Nantel-Vivier, A., Pihl, R. O., Cote, S., & Tremblay, R. E. (2014). Developmental association of prosocial behaviour with aggression, anxiety and depression from infancy to preadolescence. *Journal of Child Psychology and Psychiatry*, 55(10), 1135–1144. <https://doi.org/10.1111/jcpp.12235>
- Oberski, D. (2016). Mixture models: Latent profile and latent class analysis. In J. Robertson & M. Kaptein (Eds.), *Modern statistical methods for HCI* (pp. 275–287). New York, NY: Springer.
- Paulus, M. (2014). The emergence of prosocial behavior: Why do infants and toddlers help, comfort, and share? *Child Development Perspectives*, 8(2), 77–81. <https://doi.org/10.1111/cdep.12066>
- Paulus, M., Kuhn-Popp, N., Licata, M., Sodian, B., & Meinhardt, J. (2013). Neural correlates of prosocial behavior in infancy: Different neurophysiological mechanisms support the emergence of helping and comforting. *NeuroImage*, 66, 522–530. <https://doi.org/10.1016/j.neuroimage.2012.10.041>
- Peall, K. J., Smith, D. J., Kurian, M. A., Wardle, M., Waite, A. J., Hedderly, T., ... Morris, H. R. (2013). SGCE mutations cause psychiatric disorders: Clinical and genetic characterization. *Brain*, 136(1), 294–303. <https://doi.org/10.1093/brain/aws308>
- Preckel, K., Kanske, P., & Singer, T. (2018). On the interaction of social affect and cognition: Empathy, compassion and theory of mind. *Current Opinion in Behavioral Sciences*, 19, 1–6. <https://doi.org/10.1016/j.cobeha.2017.07.010>
- Qi, T., Wu, Y., Zeng, J., Zhang, F. T., Xue, A. L., Jiang, L. D., ... Yang, J. (2018). Identifying gene targets for brain-related traits using transcriptomic and methylomic data from blood. *Nature Communications*, 9, 2282. <https://doi.org/10.1038/s41467-018-04558-1>
- Rakyan, V. K., Down, T. A., Balding, D. J., & Beck, S. (2011). Epigenome-wide association studies for common human diseases. *Nature Reviews Genetics*, 12(8), 529–541. <https://doi.org/10.1038/nrg3000>
- Rijlaarsdam, J., Cecil, C. A., Relton, C. L., & Barker, E. D. (2021). Epigenetic profiling of social communication trajectories and co-occurring mental health problems: A prospective, methylome-wide association study. *Development and Psychopathology*, 1–10. <https://doi.org/10.1017/S0954579420001662>
- Rijlaarsdam, J., Cecil, C. A. M., Walton, E., Mesirow, M. S. C., Relton, C. L., Gaunt, T. R., ... Barker, E. D. (2017). Prenatal unhealthy diet, insulin-like growth factor 2 gene (IGF2) methylation, and attention deficit hyperactivity disorder symptoms in youth with early-onset conduct problems. *Journal of Child Psychology and Psychiatry*, 58(1), 19–27. <https://doi.org/10.1111/jcpp.12589>
- Rijlaarsdam, J., Pappa, I., Walton, E., Bakermans-Kranenburg, M. J., Mileva-Seitz, V. R., Rippe, R. C. A., ... van IJzendoorn, M. H. (2016). An epigenome-wide association meta-analysis of prenatal maternal stress in neonates: A model approach for replication. *Epigenetics*, 11(2), 140–149. <https://doi.org/10.1080/15592294.2016.1145329>

- Saffari, A., Silver, M. J., Zavattari, P., Moi, L., Columbano, A., Meaburn, E. L., & Dudbridge, F. (2018). Estimation of a significance threshold for epigenome-wide association studies. *Genetic Epidemiology*, 42(1), 20–33. <https://doi.org/10.1002/gepi.22086>
- Safra, L., Tecu, T., Lambert, S., Sheskin, M., Baumard, N., & Chevallier, C. (2016). Neighborhood deprivation negatively impacts children's prosocial behavior. *Frontiers in Psychology*, 7, 1760. <https://doi.org/10.3389/fpsyg.2016.01760>
- Salvatore, J. E., & Dick, D. M. (2018). Genetic influences on conduct disorder. *Neuroscience & Biobehavioral Reviews*, 91, 91–101. <https://doi.org/10.1016/j.neubiorev.2016.06.034>
- Sasaki, J. Y., Kim, H. S., Mojaverian, T., Kelley, L. D. S., Park, I. Y., & Janusonis, S. (2013). Religion priming differentially increases prosocial behavior among variants of the dopamine D4 receptor (DRD4) gene. *Social Cognitive and Affective Neuroscience*, 8(2), 209–215. <https://doi.org/10.1093/scan/nsr089>
- Seno, M. M. G., Hu, P., Gwadry, F. G., Pinto, D., Marshall, C. R., Casallo, G., & Scherer, S. W. (2011). Gene and miRNA expression profiles in autism spectrum disorders. *Brain Research*, 1380, 85–97. <https://doi.org/10.1016/j.brainres.2010.09.046>
- Suderman, M., Staley, J. R., French, R., Arathimos, R., Simpkin, A., & Tilling, K. (2018). dmrff: Identifying differentially methylated regions efficiently with power and control. *bioRxiv*, 508556. <https://doi.org/10.1101/508556>
- Thapar, A., Langley, K., O'Donovan, M., & Owen, M. (2006). Refining the attention deficit hyperactivity disorder phenotype for molecular genetic studies. *Molecular Psychiatry*, 11(8), 714–720. <https://doi.org/10.1038/sj.mp.4001831>
- Van der Graaff, J., Carlo, G., Crocetti, E., Koot, H. M., & Branje, S. (2018). Prosocial behavior in adolescence: Gender differences in development and links with empathy. *Journal of Youth and Adolescence*, 47(5), 1086–1099. <https://doi.org/10.1007/s10964-017-0786-1>
- Van IJzendoorn, M. H., & Bakermans-Kranenburg, M. J. (2014). Prosocial development and situational morality: Neurobiological, parental, and contextual factors. In J. F. Leckman, C. Panter-Brick, & R. Salah (Eds.), *Pathways to peace: The transformative power of children and families* (Vol. 15, pp. 161–184). Cambridge, MA: MIT Press.
- Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36(3), 48. <https://doi.org/10.18637/jss.v036.i03>
- Walton, E., Pingault, J. B., Cecil, C. A. M., Gaunt, T. R., Relton, C. L., Mill, J., & Barker, E. D. (2017). Epigenetic profiling of ADHD symptoms trajectories: A prospective, methylome-wide study. *Molecular Psychiatry*, 22(2), 250–256. <https://doi.org/10.1038/mp.2016.85>
- Wu, Y., Cao, H., Baranova, A., Huang, H., Li, S., Cai, L., ... Wang, Q. (2020). Multi-trait analysis for genome-wide association study of five psychiatric disorders. *Translational Psychiatry*, 10(209), 1–11. <https://doi.org/10.1038/s41398-020-00902-6>
- Yang, B. Z., Zhang, H. P., Ge, W. J., Weder, N., Douglas-Palumberi, H., Perepletchikova, F., ... Kaufman, J. (2013). Child abuse and epigenetic mechanisms of disease risk. *American Journal of Preventive Medicine*, 44(2), 101–107. <https://doi.org/10.1016/j.amepre.2012.10.012>
- Zhang, T. Y., & Meaney, M. J. (2010). Epigenetics and the environmental regulation of the genome and its function. *Annual Review of Psychology*, 61, 439–466. <https://doi.org/10.1146/annurev.psych.60.110707.163625>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Luo, M., Meehan, A. J., Walton, E., Röder, S., Herberth, G., Zenclussen, A. C., Cosín-Tomás, M., Sunyer, J., Mulder, R. H., Cortes Hidalgo, A. P., Bakermans-Kranenburg, M. J., Felix, J. F., Relton, C., Suderman, M., Pappa, I., Kok, R., Tiemeier, H., van IJzendoorn, M. H., Barker, E. D., & Cecil, C. A. M. (2021). Neonatal DNA methylation and childhood low prosocial behavior: An epigenome-wide association meta-analysis. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics Part B*, 1–14. <https://doi.org/10.1002/ajmg.b.32862>