



King's Research Portal

DOI: 10.1123/ijsnem.2021-0139

Document Version Peer reviewed version

Link to publication record in King's Research Portal

Citation for published version (APA):

Witard, O., Bannock, L., & Tipton, K. D. (2021). Making Sense of Muscle Protein Synthesis: A Focus on Muscle Growth During Resistance Training. *International Journal of Sport Nutrition and Exercise Metabolism*. Advance online publication. https://doi.org/10.1123/ijsnem.2021-0139

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Making Sense of Muscle Protein Synthesis: A Focus on Muscle Growth During Resistance Training

Oliver C. Witard,¹ Laurent Bannock,² and Kevin D. Tipton^{2,3}

¹Centre for Human & Applied Physiological Sciences, King's College London, London, United Kingdom; ²The Institute of Performance Nutrition, Edinburgh, United Kingdom; ³Liverpool John Moores University, Liverpool, United Kingdom

The acute response of muscle protein synthesis (MPS) to resistance exercise and nutrition is often used to inform recommendations for exercise programming and dietary interventions, particularly protein nutrition, to support and enhance muscle growth with training. Those recommendations are worthwhile only if there is a predictive relationship between the acute response of MPS and subsequent muscle hypertrophy during resistance exercise training. The metabolic basis for muscle hypertrophy is the dynamic balance between the synthesis and degradation of myofibrillar proteins in muscle. There is ample evidence that the process of MPS is much more responsive to exercise and nutrition interventions than muscle protein breakdown. Thus, it is intuitively satisfying to translate the acute changes in MPS to muscle hypertrophy with training over a longer time frame. Our aim is to examine and critically evaluate the strength and nature of this relationship. Moreover, we examine the methodological and physiological factors related to measurement of MPS and changes in muscle hypertrophy that contribute to uncertainty regarding this relationship. Finally, we attempt to offer recommendations for practical and contextually relevant application of the information available from studies of the acute response of MPS to optimize muscle hypertrophy with training.

Keywords: myofibrillar protein synthesis, muscle hypertrophy, muscle remodeling, translational efficiency, ribosomal biogenesis, stable isotopes

Muscle protein synthesis (MPS) is the metabolic process that describes the incorporation of amino acids into bound skeletal muscle proteins. Muscle proteins can be crudely classified into the contractile myofibrillar proteins (i.e., myosin, actin, tropomyosin, troponin) and the energy producing mitochondrial proteins. The synthesis of myofibrillar proteins is primarily responsible for changes in skeletal muscle mass following resistance training; whereas, mitochondrial proteins are primarily synthesized in response to endurance type training (Wilkinson et al., 2008). The MPS is most commonly expressed as a rate of amino acid incorporation into bound muscle protein over a given time period, typically a single hour or a single day. Conversely, the metabolic process of muscle protein breakdown describes the degradation of bound muscle proteins into their amino acid precursors that occurs continuously and concurrently with MPS. As such, the aggregate difference in rates of MPS and muscle protein breakdown determines whether muscle protein is gained (MPS exceeds muscle protein breakdown) or muscle protein is lost (muscle protein breakdown exceeds MPS). Of the two metabolic processes, MPS is more responsive to exercise and nutritional stimuli (Tipton et al., 2018), at least in healthy individuals, and thus has garnered most scientific attention in the context of muscle adaptations to exercise training.

Q1

The assessment of the acute response of MPS to combined exercise and nutrition interventions is commonly used as the scientific basis to inform sport and exercise nutrition, in particular protein nutrition, recommendations for the training and performance of athletes and other exercisers. This longstanding assumption is predicated on a direct relationship existing between the acute response of MPS to a single bout of resistance exercise combined with a nutritional intervention (REx) and chronic phenotypic adaptations (i.e., muscle hypertrophy) to resistance exercise training and repeated dietary manipulation (RET). An early report by Balagopal et al. (1997) supported the idea that acute measurements of MPS are predictive of chronic muscle adaptation, correlating basal rates of MPS with measurements of muscle mass, strength, and muscle mass per unit muscle mass (indirect marker of muscle quality), albeit in untrained individuals across a mix of young, middle-aged, and older adult cohorts (Balagopal et al., 1997). In addition, more recent studies have demonstrated that the acute response of MPS to REx (Tang et al., 2009; Wilkinson et al., 2007) may predict longer term muscle growth with RET (Hartman et al., 2007; Volek et al., 2013). However, this relationship has been challenged (Tipton & Wolfe, 2001) with experimental evidence that a disconnect exists between acute measurements of MPS in response to REx and chronic changes in skeletal muscle mass following RET in previously untrained young men (Mayhew et al., 2009; Mitchell et al., 2014). As such, these data have cast into doubt the predictive value of acute measurements of MPS to inform evidence-based nutrition interventions, with clear implications for practitioners across various disciplines related to sport and exercise nutrition.

Viewed through the lens of the applied sport and exercise nutrition practitioner, it is crucial to understand the real-world significance of the stated and/or perceived superiority of one nutritional strategy over another. Clearly, there are many potential dietary, exercise training, and performance interventions available. An accurate translation of scientific evidence for these interventions to applied practice is going to be most relevant to the exerciser, and especially for competitive athletes. Indeed, this translation applies to whether the focus is on, for instance, the

Tipton (kevin@theiopn.com) is corresponding author.

total intake, type, or timing of a nutritional intervention. Nutritional strategies to enhance muscle hypertrophy are commonly determined on the basis of controlled laboratory studies that report MPS as the primary outcome measurement. Thus, it is crucial that the real-world significance of acute MPS measurements that are used to determine the superiority of nutritional interventions for muscle hypertrophy is understood within the context and limitations of these methods.

Therefore, the main purpose of this narrative review is to critically evaluate the relationship between acute measurements of MPS and chronic measurements of muscle adaptation, with specific reference to RET and muscle hypertrophy. We comprehensively discuss a range of physiological and methodological variables that, in our view and others (Mitchell et al., 2015a), underpin the complex relationship between the acute response of MPS to exercise and nutrition and chronic changes in muscle mass. Physiological variables relate to inherent variability in the response of MPS to exercise and nutrition, the modulation of muscle protein metabolism with changing training status, the influence of a multitude of training paradigms, and genetics. Methodological variables relate to subtle, yet important, technical considerations with regard to measurements of MPS and muscle hypertrophy. As such, our aim is to "make sense of muscle protein synthesis" by providing a balanced and contextually relevant interpretation of the relationship between the acute response of MPS and chronic changes in muscle mass through a lens of translating the science of MPS into real-world practice for the end user practitioner (physiologist or nutritionist), coach, athlete, and/or researcher.

Metabolic Basis of Muscle Hypertrophy

Muscle hypertrophy represents the primary phenotypic adaptation to RET (Goldberg et al., 1975; McGlory et al., 2017) owing, in large part, to the plasticity of skeletal muscle tissue in response to REx and nutrition. The precise definition of skeletal muscle hypertrophy is a topic of current debate among the scientific community (Damas et al., 2015; Figueiredo, 2019; Haun et al., 2019; Joanisse et al., 2020; Roberts et al., 2020). Traditionally, muscle hypertrophy is defined as an increase in skeletal muscle mass and cross-sectional area (CSA) at the whole tissue and cellular levels (Haun et al., 2019; Russell et al., 2000). This definition is underpinned by the notion that an accretion of contractile (i.e., myofibrillar) proteins occurs due to an increased abundance of sarcomeres within the preexisting myofibrils of muscle fibers, and leads to an increase in muscle fiber CSA (Russell et al., 2000).

The plasticity of skeletal muscle is mediated, at least in part, by the constant turnover or remodeling of muscle proteins. In this regard, two metabolic processes, MPS and muscle protein breakdown, act concurrently in response to various stimuli to repair, replace, and generate new muscle proteins leading to phenotypic adaptations. There is evidence that the fold change in MPS in response to REx and/or protein feeding is greater (as much as 2.5-fold) than muscle protein breakdown (Biolo et al., 1995, 1997), suggesting that MPS is the primary metabolic driver of RETinduced muscle hypertrophy (Tipton & Wolfe, 1998). Accordingly, it has been proposed that muscle hypertrophy following RET stems from a cumulative accretion of muscle proteins resulting from the repeated increase in response of myofibrillar-MPS to successive bouts of REx (Hawley et al., 2006). Hence, according to this traditional definition of muscle hypertrophy, it may seem intuitively satisfying that assessment of the acute response of MPS to REx provides an informative tool when devising RET and nutritional interventions to maximize muscle hypertrophy in athletes and other exercisers.

We acknowledge that an alternative definition of muscle hypertrophy relates to an increase in skeletal muscle size accompanied by an increase in mineral, protein, or substrate abundance (e.g., glycogen and intramuscular triglyceride) (Haun et al., 2019). This contemporary, and arguably more comprehensive, model of muscle hypertrophy also accounts for the growth of nonmyofibril components. Accordingly, three different types of muscle hypertrophy have been proposed, namely myofibrillar hypertrophy, sarcoplasmic hypertrophy, and connective tissue hypertrophy, each with their own biological definition (Haun et al., 2019). Myofibrillar hypertrophy is defined as an increase in the size and/or number of myofibrils accompanied by an increase in sarcomere number or sarcomere protein abundance directly related to the structure or contractile force generation of the muscle, that is, directly related to the more traditional definition of hypertrophy described above. Sarcoplasmic hypertrophy relates to a chronic increase in volume of the sarcolemma and/or sarcoplasm accompanied by an increase in the volume of mitochondria, sarcoplasmic reticulum, t-tubules, and/or sarcoplasmic enzyme or substrate content. Finally, connective tissue hypertrophy is defined as an increase in volume of the extracellular matrix of skeletal muscle accompanied by an increase in mineral or protein content. A critical evaluation of skeletal muscle hypertrophy as a biological construct is beyond the scope of this text, and the reader is referred to several recent reviews on this topic (Damas et al., 2018; Haun et al., 2019; Roberts et al., 2020). Nonetheless, we suggest that all three types of hypertrophy likely contribute to measured changes in muscle mass with RET, almost certainly to varying degrees depending on the type of training, as well as the type and timing of the measurement. Moreover, these factors potentially add to variability in the measurement of muscle hypertrophy with RET, leading to potential confusion for informing practice.

Muscle Protein Synthesis

The acute measurement of in vivo human rates of MPS in response to REx dates back to the 1990s (Tipton & Wolfe, 1998). A seminal study by Chesley et al. (1992) demonstrated that REx, performed in the fasted state, stimulated MPS (Chesley et al., 1992). Subsequently, it was shown that this response persisted for up to 48 hr postexercise (Phillips et al., 1997). Biolo et al. (1997) first demonstrated that hyperaminoacidemia (elevated arterial amino acid concentrations) following exercise further stimulated MPS (Biolo et al., 1997). Next, the first studies were published that demonstrated ingestion of essential amino acids immediately following REx increased the postexercise stimulation of MPS, resulting in a net accretion of muscle protein (Rasmussen et al., 2000; Tipton et al., 1999, 2001). Collectively, these data provided a platform for studies over the next 20 years (~2000 to present) to systematically investigate the interaction of exercise and nutrition for stimulation of MPS, with direct application to sport and exercise nutrition and exercise science.

Several methods have been used to measure the acute response of MPS to exercise and nutrition in humans. A comprehensive discussion of the methods used to measure MPS is beyond the scope of this review, but the interested reader is directed to a number of excellent recent reviews (Brook & Wilkinson, 2020; Millward & Smith, 2019; Wilkinson et al., 2017). The most common approach is the precursor–product method that allows

for the determination of muscle protein fractional synthesis rate (FSR). In practice, this method utilizes stable isotope labeled amino acids (i.e., ${}^{13}C_6$ phenylalanine, $1-{}^{13}C$ leucine), usually administered by intravenous infusion under controlled laboratory conditions, to directly trace the incorporation of free amino acids into newly synthesized bound muscle proteins, typically over an acute 3-12 hr time period following a single exercise and/or nutrition stimulus. Traditionally, FSR was calculated for mixed muscle proteins, that is, all muscle protein fractions combined. Methodological advances during the 1990s allowed for the separation of muscle protein fractions (Hasten et al., 1998; Rooyackers et al., 1996) and thus acute measurements of muscle myofibrillar FSR or muscle mitochondrial FSR were possible in an exercise science setting, dependent on the mode (resistance or endurance-based) of exercise stimulus (Wilkinson et al., 2008). Another recent advancement in the field is centered around the re-emergence of the orally administered deuterium oxide (D₂O) tracer method to measure free-living integrative rates of MPS. Rather than providing a single snapshot of the acute MPS response in just a few hours under tightly controlled laboratory conditions, the D₂O technique quantifies multiple acute MPS responses to exercise and/or nutritional stimuli integrated over hours, days (Wilkinson et al., 2014), weeks, or even months (Brook et al., 2015) providing greater real-world application to the athlete. Today, separation techniques have evolved further to measure FSR at the individual muscle protein level using D_2O . The focus of the review is on studies that directly determined MPS using the measurement of FSR.

The Controversy

The controversy surrounding the value of acute (i.e., 3–6 hr) measurements of MPS for predicting chronic (i.e., 10–16 weeks) changes in muscle mass with RET has been evident essentially since the measurement of MPS has been used to assess the metabolic response of muscle to REx (Tipton & Wolfe, 2001). More recently, an elegant study by Mitchell et al. (2014) cast doubt on the relationship. This study was novel in examining the withinparticipant (i.e., muscle mass of the same participants was measured pre and post RET) relationship between the acute response of myofibrillar-MPS to REx and protein feeding (administered as a single 30 g milk protein bolus), and the muscle hypertrophic response to 16 weeks of progressive RET in previously untrained young men. No measurement of muscle protein breakdown was conducted in this study. As such, this study design offered insight into whether any heterogeneity in the muscle hypertrophic response to RET could be explained by differences in the acute response of MPS to REx between the 23 participants that conducted the study. The muscle hypertrophic response was determined by measurement of pre-post RET changes in quadriceps volume and lean body mass using magnetic resonance imaging and dual-energy X-ray absorptiometry, respectively. The acute response of MPS was measured over a 6-hr recovery period following the first (of 64) bout of REx.

Perhaps surprisingly to many at the time, and refuting their original hypothesis, the study by Mitchell et al. (2014) revealed no association between the rate of myofibrillar–MPS measured over 6 hr following the initial bout of REx and protein ingestion and the change in muscle volume or lean body mass following 16 weeks of RET. Moreover, no correlation of the change in MPS from rest with the change in muscle volume was reported (Mitchell et al., 2014). Indeed, this observation is consistent with the results of a comparable 16 weeks RET study by Mayhew et al. (2009) that was

conducted in previously untrained young and older adult men (Mayhew et al., 2009). In this study, no relationship was observed between the acute response of mixed MPS measured in the fasted state 24 hr after the initial bout of REx and muscle hypertrophy as determined by measurement of muscle fiber cross-sectional area. Taken together, these data suggest that acute measurements of MPS offer limited quantitative value for predicting individualized chronic changes in muscle mass following progressive RET, at least when the acute response of MPS is measured following the initial exercise session of the RET period. These studies have contributed to some confusion—particularly for practitioners, students, and others without specialist knowledge of the strengths and limitation of stable isotope methodology—and controversy over the interpretation of data from the measurement of MPS in response to exercise and nutrition (Mitchell et al., 2015b).

In contrast, multiple lines of evidence support the notion that the acute response of MPS to REx, with or without nutritional intervention, is predictive of chronic changes in muscle mass with RET when repeatedly exposed to a comparable exercise or nutritional intervention, at least when studied on an averaged, group basis. First, ingesting an 18 g bolus of milk protein immediately after REx stimulated a greater acute response of MPS than a dosematched soy protein beverage in young men (Wilkinson et al., 2007). This finding was consistent with a longitudinal training study that reported a greater change in muscle hypertrophy when a milk protein beverage was consumed immediately after each REx session of a 12-week RET program versus a soy protein beverage in young men (Hartman et al., 2007). Similarly, the greater acute response of MPS to ingesting 20 g of whey protein versus casein immediately post REx (Tang et al., 2009) translated to greater muscle hypertrophy following 10 weeks of RET (Volek et al., 2013). Second, the acute response of MPS to REx when manipulating exercise workload (low vs. high) (Burd, West, et al., 2010) and exercise volume (i.e., 1 vs. 3 sets of REx) (Burd, Holwerda, et al., 2010) corresponded with the muscle hypertrophic response to RET protocols that manipulated these same training variables (Mitchell et al., 2012). Finally, REx-induced increases in putative anabolic hormones were not shown to increase the acute response of MPS (West et al., 2009) or enhance RET-induced muscle hypertrophy (West et al., 2010) in young men. When combined with data generated by Mitchell et al. (2014), these data highlight the complexity of the relationship between the acute response of MPS to REx and nutrition and subsequent changes in muscle mass with RET. In our view, and that of others (Damas et al., 2018; Mitchell et al., 2015), a series of physiological and methodological factors mediate this complex relationship between acute measurements of MPS and chronic changes in muscle hypertrophy, as detailed below.

Physiological Factors

Several physiological factors, related to both the acute response of MPS to REx and the muscle hypertrophic response to RET, appear to contribute to the observed discrepancy between measured rates of MPS and muscle hypertrophy. Muscle hypertrophy is a complex physiological process that is altered as training progresses. For the initial response of MPS to predict subsequent muscle hypertrophy during RET, it must be assumed that the measured response of MPS to REx is uniform throughout the training period. However, it is clear that the response of MPS is modified from the initiation of RET and as training progresses (Kim et al., 2005; Phillips et al., 2002; Tang et al., 2008). This modification takes place on a number

of levels that include the timecourse (amplitude and duration) and nature (directed to anabolic or nonanabolic processes) of the MPS response, as discussed below.

There is considerable evidence from both cross-sectional (Phillips et al., 1999) and longitudinal (Kim et al., 2005; Phillips et al., 2002; Tang et al., 2008) studies that the training status of an individual modifies the amplitude and duration of the acute response of MPS to REx. In the untrained state, the acute response of MPS has been shown to peak later, but remain elevated for longer, after REx compared with the trained state (Phillips et al., 2002; Tang et al., 2008). Conversely, in the trained state, the acute response of MPS to REx is more rapid but shorter lived than the untrained state (Phillips et al., 1999). As a result, the overall acute stimulation of MPS after REx is generally considered to be greater in untrained versus trained individuals, at least when the absolute workload of REx is matched between training states (Damas et al., 2015). Given that training status clearly modulates the acute response of MPS to REx, it follows that the relationship between the acute MPS response to REx and chronic muscle growth response to RET may be altered over the time course of the training process.

To date, the most comprehensive study to examine the influence of training status on the relationship between the acute MPS response to REx and the muscle growth response to RET was conducted by Damas et al. (2016). In this study, 10 untrained young (~27 years) men performed 10 weeks of RET consisting of two sessions of REx per week. The RET program was divided into three phases, namely the initial (i.e., at baseline), early (after 3 weeks of RET), and late (after 10 weeks of RET) phase of RET. Measurements of the acute MPS response to REx and muscle mass were obtained at each phase of RET. This elegant study design offered unique insight into the temporal relationship between acute measurements of MPS in response to REx, assessed in both the trained and untrained state, and the subsequent muscle growth response during RET.

The study by Damas, Phillips, Libardi, et al. (2016) presents data that reveals a time course-dependent relationship between the acute response of MPS to REx and chronic changes in muscle mass during RET. In this regard, no relationship was observed between the acute response of myofibrillar-MPS to the initial REx bout of the RET period and the change in muscle mass following 10 weeks of RET. As detailed above, this observation is consistent with previous studies that reported no association between the acute response of MPS to the initial REx bout and the change in muscle volume (Mitchell et al., 2014) and fiber cross-sectional area (Mayhew et al., 2009) following 16 weeks of RET in previously untrained men. In contrast, the acute response of MPS to REx measured at Weeks 3 and 10 were associated with chronic changes in muscle mass over the 10-week RET period (Damas, Phillips, Libardi, et al., 2016). These data are consistent with recent studies that reported associations between acute measurements of MPS and muscle hypertrophy over 3 (Brook et al., 2015) and 12 weeks (Reidy et al., 2017) of RET. Taken together, these data indicate the relationship between acute measurements of MPS and chronic changes in the muscle growth response becomes apparent as the training status of the individual progresses (Table 1). The predictive value of the acute response of MPS to nutrition and exercise interventions seems to be greater in trained than untrained individuals, who are not accustomed to muscle loading during REx (Damas et al., 2018). Thus, the researcher or practitioner may wish to consider the relative value of acute measurements of MPS for predicting chronic changes in muscle growth when formulating

training and nutrition recommendations, at least for trained individuals.

One physiological mechanism proposed to explain the temporal relationship between acute measurements of MPS in response to REx and chronic changes in the muscle growth response to RET relates to the nature of the response of MPS to REx (Damas et al., 2018). Damas et al. (2016) reported a greater acute response of MPS to the initial REx bout compared with the MPS response to REx performed during the early (Week 3) and later (Week 10) phase of RET. This trend aligned with the acute (48 hr) muscle damage response to REx that was highest after the initial unaccustomed REx bout, but was attenuated by the early (Week 3) phase of RET. The authors reasoned that during the early phase of a training program, the increased response of MPS to REx and protein ingestion is related more to the repair and remodeling of existing older, perhaps damaged, proteins (Damas, Phillips, Lixandrao, et al., 2016) than to hypertrophy. These early, more global, metabolic responses not only lead to the repair and remodeling of proteins, but also set the stage for future deposition of muscle proteins and muscle_growth (Brook, Wilkinson, Smith, & Atherton, 2016; Burd & De Lisio, 2017; Joanisse et al., 2020). Consistent with this notion, the greater muscle damage response to unaccustomed eccentric-based exercise versus a work-matched bout of concentric exercise has been shown to correspond with a greater acute response of MPS to eccentric REx (Moore et al., 2005; Pavis et al., 2021). As RET progresses, the responses of MPS to REx and nutrition become more refined toward muscle hypertrophy. This notion is supported by data showing that both mitochondrial and myofibrillar-MPS are increased following a REx in the untrained state (Wilkinson et al., 2008). However, following 10 weeks of RET, only myofibrillar FSR is increased. Taken together, these data suggest that with the progression of RET, and as the degree of exercise-induced muscle damage starts to diminish, the acute stimulation of MPS is directed almost exclusively to the accretion of new muscle proteins, thus explaining the correlation between acute rates of MPS and the muscle growth response during the later phase of RET (Trommelen et al., 2019).

The inherent variability in the response of MPS to REx and nutrition, as well as the response of muscle hypertrophy to RET, also contributes to our inability to utilize acute metabolic data to predict an individual response to RET (Figure 1). Individual responses to REx and nutrition may vary by as much as 100%, even within groups subjected to identical nutrition and exercise conditions. This variability in response to exercise and nutrition is reported consistently (Jackman et al., 2017; Macnaughton et al., 2016; McGlory et al., 2016) and is considered to represent normal physiological variability (Smith et al., 2011). While the source of this individual variability is not fully understood at this time, genetic variability must be a contributing factor (Clarkson et al., 2005; Pescatello et al., 2006; Riechman et al., 2004). Attempts to control prestudy activity and diet are common in these studies, yet the variability is evident. Moreover, in many studies, the population from which participants are selected is kept fairly tight. Yet, even when the range of muscle mass is restricted, there is considerable variation in the response of MPS (Macnaughton et al., 2016). The methodological conditions under which MPS is determined that may influence the measured response will be discussed below. However, in the examples illustrated in Figure 1, the method used to determine MPS, as well as the conditions under which it was measured, in each individual were identical within studies. Hence, methodological issues alone do not account for all the observed

Q6

Table 1 Rels	ationship Betw	veen Acute Measuremen	ts of MPS and Chronic	Changes in Muscle	Mass in Response to RET
Reference	Participants	Study design	Measurement of MPS	Measurement of muscle mass	Relationship
Balagopal et al. (1997)	24 healthy UT males and fe- males and fe- males Three age groups: Young (23 ± 1) years) Middle aged (52 ± 1) years) Older (77 ± 2) vears)	Cross-sectional No exercise training All measurements collected in basal state	Mixed muscle protein FSR MHC FSR Sarcoplasmic protein FSR L-[1- ¹³ C] leucine infusion 3-hr tracer incorporation period in laboratory setting	Estimated from daily urinary creatinine	Mixed muscle protein FSR correlated with muscle mass ($r = .30$, $p = .220$) MHC FSR correlated with muscle mass ($r = .48$, $p = .020$) No correlation between sarcoplasmic protein FSR and muscle mass
Mayhew et al. (2009)	years) 36 healthy UT males and fe- males Two age groups: Young $(28 \pm 1$ years) Older $(64 \pm 1$ years)	Longitudinal 16 weeks of progressive RET (3 days/week) MPS measured following first REx session of RET period Muscle mass measured immediately pre and post RET	Mixed muscle protein FSR L-[<i>ring</i> - ² H ₅] phenylalanine infusion 3-hr tracer incorporation period in laboratory setting	fCSA by immunofluo- rescence microscopy Thigh lean mass by DXA Total lean (fat and bone free) mass by DXA	No correlation between mixed muscle FSR and changes in muscle mass (all measurements) following RET
Mitchell et al. (2014)	23 UT young (24±1 year) males	Longitudinal 16 weeks of progressive RET (4 day/week) m·s measured following 1 st REx session of RET period Muscle mass measured immediately pre and post RET	Muscle myofibrillar protein FSR $L^{-}[ring^{-13}C_{6}]$ phenylalanine infusion 6-hr tracer incorporation period in laboratory setting	Quadriceps volume by MRI Total lean (fat and bone free) mass by DXA	No correlation between muscle myofibrillar protein FSR and changes in quadriceps volume following RET ($r = .10$, $p > .050$) No correlation between muscle myofibrillar protein FSR and changes in total lean mass following RET ($r = .13$, $p > .050$)
Damas, Phillips, Libardi, et al. (2016)	10 UT young (27±1 years) males	Longitudinal 10 weeks of progressive RET (2 days/week) MPS measured during Weeks 1, 3, and 10 of RET period Muscle mass measured at pre, Weeks 3 and 10 of RET	Integrated muscle myofi- brillar protein FSR Oral deuterium oxide tracer 48-hr tracer incorporation period under free-living conditions	fCSA by microscopy vCSA	No correlation between integrated muscle myofibrillar protein FSR at Week 1 and changes in fCSA following RET Integrated muscle myofibrillar protein FSR at Week 3 correlated with changes in ν CSA following RET ($r = .9$, $p = .002$) Integrated muscle myofibrillar protein FSR at Week 10 correlated with changes in fCSA ($r = .9$, $p = .032$) and ν CSA following RET ($r = .9$, $p = .000$)
					(continued)

0 2 ¢ No 0 Ż \$ ĉ 2 \$ Ido Pac NDC N ÷ ē Tahle 1 Relationshin Retu

Table 1 (co	ntinued)				
Reference	Participants	Study design	Measurement of MPS	Measurement of muscle mass	Relationship
Brook et al. (2015)	10 UT young (23 ± 1 years) males	Longitudinal 6 weeks of progressive uni- lateral lower limb RET (3 days/week) MPS measured over 3 and 6 weeks of RET period Muscle mass measured at pre, Weeks 3 and 6 of RET	Integrated muscle myofi- brillar protein FSR Oral deuterium oxide tracer Tracer incorporation over 6 weeks under free-living conditions	Thigh muscle thick- ness by ultrasound Thigh muscle mass by DXA	Integrated muscle myofibrillar protein FSR at Week 3 correlated with changes in thigh muscle thickness ($r_2 = .52$, $p = .010$)
Reidy et al. (2017)	31 UT young $(25 \pm 2 \text{ years})$ males	Longitudinal 12 weeks of progressive whole-body RET (3 days/ week) MPS measured pre and post RET	Mixed muscle protein FSR Muscle myofibrillar protein FSR L- <i>[ring</i> – ¹³ C ₆] phenylalanine infusion 6-hr tracer incorporation period in laboratory setting	Total lean (fat and bone free) mass by DXA Vastus lateralis muscle thickness by ultra- sound fCSA by immunohistochemistry	The pre-post RET change in mixed muscle FSR was correlated with the change in <i>vastus lateralis</i> muscle thickness (r =.22, p =.003)
<i>Note.</i> DXA = dual- exercise training; 1	energy X-ray absorp REx = resistance exer	tiometry, fCSA = muscle fiber cross-sciee; vCSA = vastus lateralis cross-s	s-sectional area; FSR = fractional s ectional area; MPS = muscle protei	ynthesis rates; MHC = myos in synthesis.	In heavy chain; MRI = magnetic resonance spectroscopy; RET = resistance

Q5



Figure 1 — Individual FSR responses to a combination of REx followed by protein ingestion in two previous studies. (a) Individual fasted FSR at rest (REST) and with ingestion of 30 g protein following resistance exercise (FEDEX) in two groups of trained young weightlifters (adapted from McGlory et al., 2016) and (b) individual FSR in response to ingestion of 20 and 40 g whey protein following REx in trained young weightlifters (adapted from Macnaughton et al., 2016). MPS = muscle protein synthesis; REx = resistance exercise; FSR = fractional synthetic rate.

Q7

variability. Inherent variability in the metabolic response to REx and nutrition contributes to uncertainty in predicting muscle growth based on measured rates of MPS in individuals.

One potential contributing factor to the variability of the response of MPS to identical REx and protein feeding conditions (Figure 1) might be differences in translational capacity, that is, the total number of ribosomes capable of producing peptide chains (Wen et al., 2016). The MPS is the metabolic process from which functional proteins are produced from polypeptide chains created by ribosomes. The measurement of FSR essentially represents translational efficiency, that is, the rate of translation for a given number of ribosomes. It is clear that ribosome number, that is, translational capacity, does not change acutely following REx (Brook, Wilkinson, Mitchell, et al., 2016; Chesley et al., 1992). However, differences in translational capacity between individuals would result in differences in FSR in response to a given REx and/ or nutrition stimulus. Thus, translational capacity may help explain the individual variability in response of MPS to anabolic stimuli.

Methodological Factors

The lack of ability to predict long-term muscle hypertrophic responses to RET with the acute measurement of MPS does not necessarily reflect the overall worth, or lack thereof, of information obtained from acute metabolic studies. Contributing factors to the uncertain relationship between the acute MPS response to REx and nutrition, and the muscle hypertrophic response to RET, include a lack of consistency in methods utilized, as well inherent variability resulting from the methods used (Mitchell et al., 2015). There also is heterogeneity in the response of muscle mass to RET that contributes to this disconnect. Accordingly, there are numerous reasons to suggest that the study design and methods chosen to determine hypertrophy in RET studies contributes to this quite heterogeneous response. A full evaluation of these methods is beyond the scope of this review, so interested readers are referred to an excellent presentation of the methodology by Haun et al. (2019).

Several factors related to study design and methods used to assess MPS must be considered when interpreting the relationship between the acute response of MPS- and RET-induced changes in muscle mass. Over the past 25-30 years, the vast majority of studies investigating the response of MPS have utilized the precursor-product method with direct incorporation of the stable isotopically labeled amino acids into muscle protein to determine FSR. Accurate prediction of muscle hypertrophy during RET by determining FSR in response to REx and nutrition requires certain assumptions to be made and met. First, we must assume that the initial measurement of FSR is representative of every subsequent stimulation of MPS for the remainder of the RET period, that is, the responses remain unchanged throughout RET (see discussion above). Next, the measured FSR captures the true response of MPS to REx and protein ingestion. Thus, methodological choices will be critical for determining the true response of MPS.

Methodological considerations influence the ability to capture the true response of MPS with measurement of the FSR in response to exercise and nutrition. Until recently, the majority of studies measuring FSR included an infusion of a labeled amino acid and multiple muscle biopsy samples. The FSR is reported as an hourly rate of synthesis in the time between the muscle samples. An important issue for any infusion study to determine FSR is the limited time period for incorporation of the labeled amino acid. One critical assumption is that the time between biopsies captures the true period of stimulation of MPS. Thus, regardless of the maximal magnitude of the response, if the second muscle sample is taken before the response of MPS returns to baseline, a portion of the true response of MPS may be missed and the determined FSR would be an underestimation (Figure 2). Of course, the converse would be true if the biopsy is taken too late to capture the true response. We must assume that the duration of the true MPS response is captured in the time between muscle biopsies and that this duration is not different between trials assessing the response to different nutrition and/or exercise interventions.

Another factor that contributes to a mismatch between the true response of MPS to REx is the prolonged enhancement of the utilization of amino acids from protein ingestion for MPS following a REx bout (Figure 3). The REx sensitizes the muscle to the anabolic stimulation of elevated amino acid levels from protein feeding (Biolo et al., 1997; Witard et al., 2014). It is clear that the sensitivity of muscle to amino acids remains enhanced for at least 24 hr following the exercise (Burd et al., 2011). Thus, any protein containing meal consumed within this 24-hr time period will result in a MPS response that is greater than that in response to a meal not preceded by REx. An acute measurement of MPS based on an infusion of labeled amino acids and biopsies for only a few hours after exercise would not be capable of capturing the contribution to muscle hypertrophy resulting from all of these enhanced postprandial elevations of MPS (Figure 3a). Thus, an acute measurement limited to only a few hours after REx would not reflect the entire influence of the exercise on MPS and subsequent muscle hypertrophy further contributing to the observed mismatch



Figure 2 — The response of MPS to a bout of REx and protein ingestion. (a) Infusion of $[^{13}C_6]$ phenylalanine and muscle samples taken at timepoints that capture the entire true response of MPS and (b) infusion ends and muscle samples are taken at 0 and 4 hr, but the true response of MPS remains elevated above baseline for 6 hr, so the response is underestimated. MPS = muscle protein synthesis; REx = resistance exercise.

between measurement of MPS and changes in muscle mass with training.

Over the past 15 years, another method has been revisited to determine an integrated FSR in free-living participants over a time period that is not limited by an infusion, that is, the D_2O method (Figure 3). Thus, MPS in various situations and in response to various exercise and nutrition interventions can be determined over the time course of days to weeks. The determined rate of MPS integrates the response to all physical activity and nutrient consumption during that time, including the prolonged response of MPS to subsequent meals following REx (Figure 3b). Thus, the D_2O method could be argued to provide a more holistic assessment

of MPS without the limitations inherent with the requirement for infusion of stable isotopes for measurement of MPS. It is perhaps not particularly surprising that integrated rates of MPS over longer time periods than are possible with isotope infusion studies, as well as inclusion of habitual physical activity and enhanced periods of postprandial MPS in response to exercise hours to days earlier, are better correlated with subsequent muscle hypertrophy. Several studies utilizing the D₂O measurement of FSR have reported correlations of MPS with subsequent muscle hypertrophy (Brook et al., 2015; Damas, Phillips, Libardi, et al., 2016; Franchi et al., 2015). Therefore, this method for assessing MPS seems to be more suitable for predicting muscle hypertrophy with RET.



Figure 3 — Comparison of measurement of MPS with (a) an infusion of labeled amino acids (${}^{13}C_6$ Phe) or (b) ingestion of deuterated water (D₂O). The response of MPS is enhanced following REx and this is captured by D₂O measurement of MPS. MPS = muscle protein synthesis; REx = resistance exercise.

The disconnect between the initial measurement of MPS and subsequent muscle hypertrophy during RET may be due to methodological choices made for measurement of changes in muscle mass in addition to MPS. Differences in study design and methods chosen to determine changes in muscle mass, in addition to inherent individual variability in the response of muscle to training (Mobley et al., 2018), contribute to variable results among RET studies. Factors including training duration, sleep quality, nontraining physical activity, nutrition, and other lifestyle variables may impact the training response (Haun et al., 2019; Mitchell et al., 2014). Proper control of many of these factors is virtually impossible in most RET study situations. This variability is further complicated by the various permutations possible with various combinations of these factors (Haun et al., 2019).

Perhaps a more prosaic factor contributing to the disconnect between the acute response of MPS and subsequent muscle hypertrophy with RET relates to the inherent limitations of methods used to measure changes in muscle mass in humans. Reported changes in muscle mass with RET are heavily dependent on the method chosen to assess those changes. Hence, the critical reader should consider the limitations of these methods when evaluating any particular training study. Changes in muscle mass may be measured on one or more of several levels, that is, biochemical, ultrastructural, histological, and gross anatomical levels. When multiple methods from these levels of hypertrophy are used, the agreement between methods is often poor (Haun et al., 2019). Moreover, as detailed above, there are different types of hypertrophy that must be considered in combination with the method chosen to assess changes in muscle mass. Strict control of methodological conditions, both at the time of measurement and/or laboratory conditions, is necessary (Haun et al., 2019). Three types of hypertrophy have been proposed: connective tissue, sarcoplasmic, and myofibrillar. Contributions of each type of hypertrophy to measured hypertrophy may vary with training status and/or the method used to assess hypertrophy. For example, there is evidence that hypertrophy measured at the early stage of a RET program may result from edema-induced, that is, muscle swelling and sarcoplasmic hypertrophy (Damas, Phillips, Libardi, et al., 2016). This means that if muscle hypertrophy is based on dual-energy X-ray absorptiometry or other methods without consideration of changes in intramuscular fluid, overestimations of true hypertrophy will be made. Clearly, changes in muscle mass with fluid infiltration are not related to MPS. These methodological factors should be considered when assessing the relationship between the acute response of MPS to changes in muscle mass with RET.

Practical Implications

Translating the science behind this complex relationship between the acute response of MPS to exercise and/or nutrition into clear, contextually relevant and practical messages is a priority for practitioners, coaches, athletes, and researchers. Based on our critical evaluation of existing evidence, we can make three practical implications.

(a) The acute response of MPS to REx plus protein ingestion will translate to chronic adaptations in muscle mass only in trained individuals. The predictive value of acute measurements of MPS for chronic adaptations in muscle mass in individuals at the beginning of a period of RET is limited due to the multiple regenerative roles of MPS beyond the accumulation of new muscle protein during the early stages of the training process. Nevertheless, greater rates of MPS in untrained individuals still may be considered beneficial since they are indicative of greater rates of protein turnover and muscle remodeling following exercise.

- (b) The predictive value of the acute response of MPS in distinguishing between the anabolic capacity of an exercise training or nutritional intervention warrants consideration when offering practical recommendations at a group level. In this regard, the practitioner may use this information as a general starting point to trial the effectiveness of an exercise or nutritional stimulus. However, the practitioner should remain open minded that a "one-size-fits-all" approach almost certainly does not apply, and there will likely be some athletes that do not respond to the intervention.
- (c) Finally, any recommendations made based on information, such as is described in (b), should not be based on quantitative differences between interventions that stimulate MPS. Despite one intervention being X% better than another according to acute metabolic data, this will not translate directly, at least quantitatively, to the magnitude of change for the parameter (i.e., muscle hypertrophy) of interest. Hence, the practitioner should manage expectations when explaining the potential gains afforded to the intervention of interest.

Conclusions

In this review, we have attempted to provide an evidence-based critical evaluation for the use of results from acute metabolic studies to predict changes in muscle mass with RET. It is clear that the measured acute response of MPS to an exercise/nutrition intervention is not predictive of muscle hypertrophy for any individual participating in a RET and nutrition program based on that particular combination of exercise parameters and nutrition. This lack of predictive power is especially true if the individual is beginning an unaccustomed exercise program. Nevertheless, this discrepancy should not be used to determine the value of studies measuring MPS in response to REx and protein nutrition. There are multiple examples of studies in which the acute response of MPS does predict the average hypertrophy on a group level (Hartman et al., 2007; Tang et al., 2009; Volek et al., 2013; West et al., 2009, 2010; Wilkinson et al., 2007). Moreover, measurement of the acute response of MPS to REx and nutrition interventions can provide valuable information. Regardless of training status, the acute response of MPS is indicative of protein turnover and muscle remodeling critical for recovery from exercise and adaptation to training.

The measurement of integrated MPS that includes the enhanced postprandial response of MPS to protein ingestion in free-living individuals certainly may provide predictive information about subsequent muscle growth, albeit not in individuals undergoing unaccustomed exercise. Moreover, the acute measurement of MPS also provides more sensitivity than chronic training studies over a much shorter time frame and can thus be viewed as a good starting point for determining nutritional recommendations. Given the nature of measurement of FSR, if a difference is detected in an acute study, for example, between different protein sources, then we can conclude with high confidence that the measured difference is physiologically relevant, at least qualitatively. In this regard, the protein source that engenders the greater FSR may be considered the higher quality protein source irrespective of whether chronic studies are able to detect differences in muscle hypertrophy under comparable conditions of protein source manipulation. Thus, we can use that information to inform subsequent RET studies.

Finally, the acute measurement of MPS in response to exercise and nutrition offers valuable mechanistic information. In fact, delineation of mechanisms of muscle protein metabolism was the aim of many of the seminal studies that are now used to contribute to the development of recommendations (Biolo et al., 1997; Phillips et al., 1997; Tipton et al., 1999, 2001). Thus, whereas practitioners should be aware of the potential pitfalls with reliance on acute metabolic studies for making nutritional recommendations for athletes and exercisers, with proper interpretation a great deal of valuable information may be gleaned from these studies. Acute measurement of MPS in response to various nutrition and exercise interventions should be viewed as yet another tool in the toolbox for use by practitioners and others.

References

- Balagopal, P., Ljungqvist, O., & Nair, K.S. (1997). Skeletal muscle myosin heavy-chain synthesis rate in healthy humans. *American Journal* of Physiology, 272(1), 45. https://doi.org/10.1152/ajpendo.1997.272. 1.E45
- Biolo, G., Maggi, S.P., Williams, B.D., Tipton, K.D., & Wolfe, R.R. (1995). Increased rates of muscle protein turnover and amino acid transport after resistance exercise in humans. *American Journal of Physiology*, 268, E514–E520. https://doi.org/10.1152/ajpendo.1995. 268.3.E514
- Biolo, G., Tipton, K.D., Klein, S., & Wolfe, R.R. (1997). An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *American Journal of Physiology*, 273, E122–E129. https://doi.org/10.1152/ajpendo.1997.273.1.E122
- Brook, M.S., & Wilkinson, D.J. (2020). Contemporary stable isotope tracer approaches: Insights into skeletal muscle metabolism in health and disease. *Experimental Physiology*, 105(7), 1081–1089. https:// doi.org/10.1113/EP087492
- Brook, M.S., Wilkinson, D.J., Mitchell, W.K., Lund, J.N., Phillips, B.E., Szewczyk, N.J., Greenhaff, P.L., Smith, K., & Atherton, P.J. (2016). Synchronous deficits in cumulative muscle protein synthesis and ribosomal biogenesis underlie age-related anabolic resistance to exercise in humans. *The Journal of Physiology*, 594(24), 7399– 7417. https://doi.org/10.1113/JP272857
- Brook, M.S., Wilkinson, D.J., Mitchell, W.K., Lund, J.N., Szewczyk, N.J., Greenhaff, P.L., Smith, K., & Atherton, P.J. (2015). Skeletal muscle hypertrophy adaptations predominate in the early stages of resistance exercise training, matching deuterium oxide-derived measures of muscle protein synthesis and mechanistic target of rapamycin complex 1 signaling. *The FASEB Journal*, 29(11), 4485–4496. https://doi. org/10.1096/fj.15-273755
- Brook, M.S., Wilkinson, D.J., Smith, K., & Atherton, P.J. (2016). The metabolic and temporal basis of muscle hypertrophy in response to resistance exercise. *European Journal of Sport Science*, 16(6), 633– 644. https://doi.org/10.1080/17461391.2015.1073362
- Burd, N.A., & De Lisio, M. (2017). Skeletal muscle remodeling: Interconnections between stem cells and protein turnover. *Exercise and Sport Sciences Reviews*, 45(3), 187–191. https://doi.org/10.1249/ JES.000000000000117
- Burd, N.A., Holwerda, A.M., Selby, K.C., West, D.W., Staples, A.W., Cain, N.E., Cashaback, J.G., Potvin, J.R., Baker, S.K., & Phillips, S.M. (2010). Resistance exercise volume affects myofibrillar protein synthesis and anabolic signalling molecule phosphorylation in young

men. The Journal of Physiology, 588(16), 3119–3130. https://doi.org/ 10.1113/jphysiol.2010.192856

- Burd, N.A., West, D.W., Moore, D.R., Atherton, P.J., Staples, A.W., Prior, T., Tang, J.E., Rennie, M.J., Baker, S.K., & Phillips, S.M. (2011). Enhanced amino acid sensitivity of myofibrillar protein synthesis persists for up to 24 h after resistance exercise in young men. *The Journal of Nutrition*, 141(4), 568–573. https://doi.org/10.3945/jn. 110.135038
- Burd, N.A., West, D.W., Staples, A.W., Atherton, P.J., Baker, J.M., Moore, D.R., Holwerda, A.M., Parise, G., Rennie, M.J., Baker, S.K., & Phillips, S.M. (2010). Low-load high volume resistance exercise stimulates muscle protein synthesis more than high-load low volume resistance exercise in young men. *PLoS One, 5*(8), Article e12033. https://doi.org/10.1371/journal.pone.0012033
- Chesley, A., MacDougall, J.D., Tarnopolsky, M.A., Atkinson, S.A., & Smith, K. (1992). Changes in human muscle protein synthesis after resistance exercise. *Journal of Applied Physiology*, 73(4), 1383– 1388. https://doi.org/10.1152/jappl.1992.73.4.1383
- Clarkson, P.M., Devaney, J.M., Gordish-Dressman, H., Thompson, P.D., Hubal, M.J., Urso, M., Price, T.B., Angelopoulos, T.J., Gordon, P.M., Moyna, N.M., Pescatello, L.S., Visich, P.S., Zoeller, R.F., Seip, R.L., & Hoffman, E.P. (2005). ACTN3 genotype is associated with increases in muscle strength in response to resistance training in women. *Journal of Applied Physiology*, 99(1), 154–163. https:// doi.org/10.1152/japplphysiol.01139.2004
- Damas, F., Libardi, C.A., & Ugrinowitsch, C. (2018). The development of skeletal muscle hypertrophy through resistance training: The role of muscle damage and muscle protein synthesis. *European Journal of Applied Physiology*, 118(3), 485–500. https://doi.org/10.1007/ s00421-017-3792-9
- Damas, F., Phillips, S., Vechin, F.C., & Ugrinowitsch, C. (2015). A review of resistance training-induced changes in skeletal muscle protein synthesis and their contribution to hypertrophy. *Sports Medicine*, 45(6), 801–807. https://doi.org/10.1007/s40279-015-0320-0
- Damas, F., Phillips, S.M., Libardi, C.A., Vechin, F.C., Lixandrao, M.E., Jannig, P.R., Costa, L.A., Bacurau, A.V., Snijders, T., Parise, G., Tricoli, V., Roschel, H., & Ugrinowitsch, C. (2016). Resistance training-induced changes in integrated myofibrillar protein synthesis are related to hypertrophy only after attenuation of muscle damage. *The Journal of Physiology*, 594(18), 5209–5222. https://doi.org/10. 1113/JP272472
- Damas, F., Phillips, S.M., Lixandrao, M.E., Vechin, F.C., Libardi, C.A., Roschel, H., Tricoli, V., & Ugrinowitsch, C. (2016). Early resistance training-induced increases in muscle cross-sectional area are concomitant with edema-induced muscle swelling. *European Journal of Applied Physiology*, 116(1), 49–56. https://doi.org/10.1007/s00421-015-3243-4
- Figueiredo, V.C. (2019). Revisiting the roles of protein synthesis during skeletal muscle hypertrophy induced by exercise. *American Journal of Physiology—Regulatory, Integrative and Comparative Physiology,* 317(5), R709–R718. https://doi.org/10.1152/ajpregu.00162.2019
- Franchi, M.V., Wilkinson, D.J., Quinlan, J.I., Mitchell, W.K., Lund, J.N., Williams, J.P., Reeves, N.D., Smith, K., Atherton, P.J., & Narici, M.V. (2015). Early structural remodeling and deuterium oxidederived protein metabolic responses to eccentric and concentric loading in human skeletal muscle. *Physiological Reports*, 3(11), Article e12593. https://doi.org/10.14814/phy2.12593
- Goldberg, A.L., Etlinger, J.D., Goldspink, D.F., & Jablecki, C. (1975). Mechanism of work-induced hypertrophy of skeletal muscle. *Medicine & Science in Sports & Exercise*, 7(3), 185–198.
- Hartman, J.W., Tang, J.E., Wilkinson, S.B., Tarnopolsky, M.A., Lawrence, R.L., Fullerton, A.V., & Phillips, S.M. (2007). Consumption of

fat-free fluid milk after resistance exercise promotes greater lean mass accretion than does consumption of soy or carbohydrate in young, novice, male weightlifters. *The American Journal of Clinical Nutrition*, *86*(2), 373–381. https://doi.org/10.1093/ajcn/86.2.373

- Hasten, D.L., Morris, G.S., Ramanadham, S., & Yarasheski, K.E. (1998). Isolation of human skeletal muscle myosin heavy chain and actin for measurement of fractional synthesis rates. *American Journal of Physiology*, 275(6), Article E1092. https://doi.org/10.1152/ajpendo. 1998.275.6.E1092
- Haun, C.T., Vann, C.G., Roberts, B.M., Vigotsky, A.D., Schoenfeld, B.J., & Roberts, M.D. (2019). A critical evaluation of the biological construct skeletal muscle hypertrophy: Size matters but so does the measurement. *Frontiers in Physiology*, *10*, 247. https://doi.org/ 10.3389/fphys.2019.00247
- Hawley, J.A., Tipton, K.D., & Millard-Stafford, M. (2006). Promoting training adaptations through nutritional interventions. *Journal of Sports Science*, 24(7), 709–721. https://doi.org/10.1080/02640410 500482727
- Jackman, S.R., Witard, O.C., Philp, A., Wallis, G.A., Baar, K., & Tipton, K.D. (2017). Branched-chain amino acid ingestion stimulates muscle myofibrillar protein synthesis following resistance exercise in humans. *Frontiers in Physiology*, *8*, 390. https://doi.org/10.3389/fphys. 2017.00390
- Joanisse, S., Lim, C., McKendry, J., Mcleod, J.C., Stokes, T., & Phillips, S.M. (2020). Recent advances in understanding resistance exercise training-induced skeletal muscle hypertrophy in humans. *F1000 Research*, 9. https://doi.org/10.12688/f1000research.21588.1
- Kim, P.L., Staron, R.S., & Phillips, S.M. (2005). Fasted-state skeletal muscle protein synthesis after resistance exercise is altered with training. *Journal of Physiology*, 568(1), 283–290. https://doi.org/ 10.1113/jphysiol.2005.093708
- Macnaughton, L.S., Wardle, S.L., Witard, O.C., McGlory, C., Hamilton, D.L., Jeromson, S., Lawrence, C.E., Wallis, G.A., & Tipton, K.D. (2016). The response of muscle protein synthesis following whole-body resistance exercise is greater following 40 g than 20 g of ingested whey protein. *Physiological Reports*, 4(15), Article e12893. https://doi.org/10.14814/phy2.12893
- Mayhew, D.L., Kim, J.S., Cross, J.M., Ferrando, A.A., & Bamman, M.M. (2009). Translational signaling responses preceding resistance training-mediated myofiber hypertrophy in young and old humans. *Journal of Applied Physiology*, 107(5), 1655–1662. https://doi.org/10. 1152/japplphysiol.91234.2008
- McGlory, C., Devries, M.C., & Phillips, S.M. (2017). Skeletal muscle and resistance exercise training; the role of protein synthesis in recovery and remodeling. *Journal of Applied Physiology*, 122(3), 541–548. https://doi.org/10.1152/japplphysiol.00613.2016
- McGlory, C., Wardle, S.L., Macnaughton, L.S., Witard, O.C., Scott, F., Dick, J., Bell, J.G., Phillips, S.M., Galloway, S.D., Hamilton, D.L., & Tipton, K.D. (2016). Fish oil supplementation suppresses resistance exercise and feeding-induced increases in anabolic signaling without affecting myofibrillar protein synthesis in young men. *Physiological Reports*, 4(6), Article e12715. https://doi.org/10. 14814/phy2.12715
- Millward, D.J., & Smith, K. (2019). The application of stable-isotope tracers to study human musculoskeletal protein turnover: A tale of bag filling and bag enlargement. *The Journal of Physiology*, 597(5), 1235–1249. https://doi.org/10.1113/JP275430
- Mitchell, C.J., Churchward-Venne, T.A., Cameron-Smith, D., & Phillips, S.M. (2015a). What is the relationship between the acute muscle protein synthesis response and changes in muscle mass? *Journal of Applied Physiology*, *118*(4), 495–497. https://doi.org/10.1152/ japplphysiol.00609.2014

- Mitchell, C.J., Churchward-Venne, T.A., Cameron-Smith, D., & Phillips, S.M. (2015b). Last word on viewpoint: What is the relationship between the acute muscle protein synthetic response and changes in muscle mass? *Journal of Applied Physiology*, *118*(4), 503. https://doi. org/10.1152/japplphysiol.01056.2014
- Mitchell, C.J., Churchward-Venne, T., Parise, G., Bellamy, L., Baker, S.K., Smith, K., Atherton, P.J., & Phillips, S.M. (2014). Acute postexercise myofibrillar protein synthesis is not correlated with resistance training-induced muscle hypertrophy in young men. *PLoS One*, 9(2), Article e89431. https://doi.org/10.1371/journal.pone.0089431
- Mitchell, C.J., Churchward-Venne, T., West, D.W., Burd, N.A., Breen, L., Baker, S.K., & Phillips, S.M. (2012). Resistance exercise load does not determine training-mediated hypertrophic gains in young men. *Journal of Applied Physiology*, 113(1), 71–77. https://doi.org/10. 1152/japplphysiol.00307.2012
- Mobley, C.B., Haun, C.T., Roberson, P.A., Mumford, P.W., Kephart, W.C., Romero, M.A., Osburn, S.C., Vann, C.G., Young, K.C., Beck, D.T., Martin, J.S., Lockwood, C.M., & Roberts, M.D. (2018). Biomarkers associated with low, moderate, and high vastus lateralis muscle hypertrophy following 12 weeks of resistance training. *PLoS One*, *13*(4), Article e0195203. https://doi.org/10.1371/journal.pone. 0195203
- Moore, D.R., Phillips, S.M., Babraj, J.A., Smith, K., & Rennie, M.J. (2005). Myofibrillar and collagen protein synthesis in human skeletal muscle in young men after maximal shortening and lengthening contractions. *American Journal of Physiology—Endocrinology and Metabolism*, 288(6), E1153–E1159. https://doi.org/10.1152/ajpendo. 00387.2004
- Pavis, G.F., Jameson, T.S.O., Dirks, M.L., Lee, B.P., Abdelrahman, D.R., Murton, A.J., Porter, C., Alamdari, N., Mikus, C.R., Wall, B.T., & Stephens, F.B. (2021). Improved recovery from skeletal muscle damage is largely unexplained by myofibrillar protein synthesis or inflammatory and regenerative gene expression pathways. *American Journal of Physiology—Endocrinology and Metabolism, 320*(2), E291–E305. https://doi.org/10.1152/ajpendo.00454.2020
- Pescatello, L.S., Kostek, M.A., Gordish-Dressman, H., Thompson, P.D., Seip, R.L., Price, T.B., Angelopoulos, T.J., Clarkson, P.M., Gordon, P.M., Moyna, N.M., Visich, P.S., Zoeller, R.F., Devaney, J.M., & Hoffman, E.P. (2006). ACE ID genotype and the muscle strength and size response to unilateral resistance training. *Medicine & Science in Sports & Exercise*, 38(6), 1074–1081. https://doi.org/10.1249/01. mss.0000222835.28273.80
- Phillips, S.M., Parise, G., Roy, B.D., Tipton, K.D., Wolfe, R.R., & Tamopolsky, M.A. (2002). Resistance-training-induced adaptations in skeletal muscle protein turnover in the fed state. *Canadian Journal* of Physiology and Pharmacology, 80(11), 1045–1053. https://doi. org/10.1139/y02-134
- Phillips, S.M., Tipton, K.D., Aarsland, A., Wolf, S.E., & Wolfe, R.R. (1997). Mixed muscle protein synthesis and breakdown after resistance exercise in humans. *American Journal of Physiology*, 273, 99. https://doi.org/10.1152/ajpendo.1997.273.1.E99
- Phillips, S.M., Tipton, K.D., Ferrando, A.A., & Wolfe, R.R. (1999). Resistance training reduces the acute exercise-induced increase in muscle protein turnover. *American Journal of Physiology*, 276, E118–E124. https://doi.org/10.1152/ajpendo.1999.276.1.E118
- Rasmussen, B.B., Tipton, K.D., Miller, S.L., Wolf, S.E., & Wolfe, R.R. (2000). An oral essential amino acid-carbohydrate supplement enhances muscle protein anabolism after resistance exercise. *Journal of Applied Physiology*, 88(2), 386–392. https://doi.org/10.1152/jappl. 2000.88.2.386
- Reidy, P.T., Borack, M.S., Markofski, M.M., Dickinson, J.M., Fry, C.S., Deer, R.R., Volpi, E., & Rasmussen, B.B. (2017). Post-absorptive

muscle protein turnover affects resistance training hypertrophy. *European Journal of Applied Physiology*, *117*(5), 853–866. https://doi.org/10.1007/s00421-017-3566-4

- Riechman, S.E., Balasekaran, G., Roth, S.M., & Ferrell, R.E. (2004). Association of interleukin-15 protein and interleukin-15 receptor genetic variation with resistance exercise training responses. *Journal* of Applied Physiology, 97(6), 2214–2219. https://doi.org/10.1152/ japplphysiol.00491.2004
- Roberts, M.D., Haun, C.T., Vann, C.G., Osburn, S.C., & Young, K.C. (2020). Sarcoplasmic hypertrophy in skeletal muscle: A scientific "unicorn" or resistance training adaptation? *Frontiers in Physiology*, *11*, 816. https://doi.org/10.3389/fphys.2020.00816
- Rooyackers, O.E., Adey, D.B., Ades, P.A., & Nair, K.S. (1996). Effect of age on in vivo rates of mitochondrial protein synthesis in human skeletal muscle. *Proceedings of the National Academy of Sciences of the United States of America*, 93(26), 15364–15369. https://doi.org/ 10.1073/pnas.93.26.15364
- Russell, B., Motlagh, D., & Ashley, W.W. (2000). Form follows function: How muscle shape is regulated by work. *Journal of Applied Physiology*, 88(3), 1127–1132. https://doi.org/10.1152/jappl.2000.88.3. 1127
- Smith, G.I., Patterson, B.W., & Mittendorfer, B. (2011). Human muscle protein turnover—Why is it so variable? *Journal of Applied Physiology*, *110*(2), 480–491. https://doi.org/10.1152/japplphysiol.00125. 2010
- Tang, J.E., Moore, D.R., Kujbida, G.W., Tarnopolsky, M.A., & Phillips, S.M. (2009). Ingestion of whey hydrolysate, casein, or soy protein isolate: Effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *Journal of Applied Physiology*, 107(3), 987–992. https://doi.org/10.1152/jappl physiol.00076.2009
- Tang, J.E., Perco, J.G., Moore, D.R., Wilkinson, S.B., & Phillips, S.M. (2008). Resistance training alters the response of fed state mixed muscle protein synthesis in young men. *American Journal of Physi*ology—Regulatory, Integrative and Comparative Physiology, 294(1), R172–R178. https://doi.org/10.1152/ajpregu.00636.2007
- Tipton, K.D., Ferrando, A.A., Phillips, S.M., Doyle, D.J., & Wolfe, R.R. (1999). Postexercise net protein synthesis in human muscle from orally administered amino acids. *American Journal of Physiology*, 276, E628–E634. https://doi.org/10.1152/ajpendo.1999.276.4.E628
- Tipton, K.D., Hamilton, D.L., & Gallagher, I.J. (2018). Assessing the role of muscle protein breakdown in response to nutrition and exercise in humans. *Sports Medicine*, 48, 53–64. https://doi.org/10.1007/ s40279-017-0845-5
- Tipton, K.D., Rasmussen, B.B., Miller, S.L., Wolf, S.E., Owens-Stovall, S., Petrini, B.E., & Wolfe, R.R. (2001). Timing of amino acidcarbohydrate ingestion alters anabolic response of muscle to resistance exercise. *American Journal of Physiology—Endocrinology and Metabolism, 281*(2), E197–E206. https://doi.org/10.1152/ajpendo. 2001.281.2.E197
- Tipton, K.D., & Wolfe, R.R. (1998). Exercise-induced changes in protein metabolism. Acta Physiologica Scandinavica, 162(3), 377–387. https://doi.org/10.1046/j.1365-201X.1998.00306.x
- Tipton, K.D., & Wolfe, R.R. (2001). Exercise, protein metabolism, and muscle growth. *International Journal of Sport Nutrition and Exercise*

Metabolism, 11(1), 109–132. https://doi.org/10.1123/ijsnem.11. 1.109

- Volek, J.S., Volk, B.M., Gomez, A.L., Kunces, L.J., Kupchak, B.R., Freidenreich, D.J., Aristizabal, J.C., Saenz, C., Dunn-Lewis, C., Ballard, K.D., Quann, E.E., Kawiecki, D.L., Flanagan, S.D., Comstock, B.A., Fragala, M.S., Earp, J.E., Fernandez, M.L., Bruno, R.S., Ptolemy, A.S., ... Kraemer, W.J. (2013). Whey protein supplementation during resistance training augments lean body mass. *Journal of the American College of Nutrition*, 32(2), 122–135. https://doi.org/ 10.1080/07315724.2013.793580
- Wen, Y., Alimov, A.P., & McCarthy, J.J. (2016). Ribosome biogenesis is necessary for skeletal muscle hypertrophy. *Exercise and Sport Sciences Reviews*, 44(3), 110–115. https://doi.org/10.1249/JES. 0000000000000082
- West, D.W., Burd, N.A., Tang, J.E., Moore, D.R., Staples, A.W., Holwerda, A.M., Baker, S.K., & Phillips, S.M. (2010). Elevations in ostensibly anabolic hormones with resistance exercise enhance neither training—Induced muscle hypertrophy nor strength of the elbow flexors. *Journal of Applied Physiology*, 108(1), 60–67. https://doi. org/10.1152/japplphysiol.01147.2009
- West, D.W., Kujbida, G.W., Moore, D.R., Atherton, P., Burd, N.A., Padzik, J.P., De Lisio, M., Tang, J.E., Parise, G., Rennie, M.J., Baker, S.K., & Phillips, S.M. (2009). Resistance exercise-induced increases in putative anabolic hormones do not enhance muscle protein synthesis or intracellular signalling in young men. *The Journal of Physiology*, 587(21), 5239–5247. https://doi.org/10. 1113/jphysiol.2009.177220
- Wilkinson, D.J., Brook, M.S., Smith, K., & Atherton, P.J. (2017). Stable isotope tracers and exercise physiology: Past, present and future. *The Journal of Physiology*, 595(9), 2873–2882. https://doi.org/10.1113/ JP272277
- Wilkinson, D.J., Franchi, M.V., Brook, M.S., Narici, M.V., Williams, J.P., Mitchell, W.K., Szewczyk, N.J., Greenhaff, P.L., Atherton, P.J., & Smith, K. (2014). A validation of the application of D(2)O stable isotope tracer techniques for monitoring day-to-day changes in muscle protein subfraction synthesis in humans. *American Journal* of Physiology—Endocrinology and Metabolism, 306(5), E571–E579. https://doi.org/10.1152/ajpendo.00650.2013
- Wilkinson, S.B., Phillips, S.M., Atherton, P.J., Patel, R., Yarasheski, K.E., Tarnopolsky, M.A., & Rennie, M.J. (2008). Differential effects of resistance and endurance exercise in the fed state on signalling molecule phosphorylation and protein synthesis in human muscle. *Journal of Physiology*, 586(15), 3701–3717. https://doi.org/10.1113/ jphysiol.2008.153916
- Wilkinson, S.B., Tarnopolsky, M.A., Macdonald, M.J., MacDonald, J.R., Armstrong, D., & Phillips, S.M. (2007). Consumption of fluid skim milk promotes greater muscle protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soy-protein beverage. *The American Journal of Clinical Nutrition*, 85(4), 1031–1040. https://doi.org/10.1093/ajcn/85.4.1031
- Witard, O.C., Jackman, S.R., Breen, L., Smith, K., Selby, A., & Tipton, K.D. (2014). Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *The American Journal of Clinical Nutrition*, 99(1), 86–95. https://doi.org/10.3945/ajcn.112.055517

Queries

- Q1. Please verify the inserted running title.
- Q2. Please provide ORCid for Corresponding author "Kevin D. Tipton."
- Q3. Please ensure author information is listed correctly here and within the byline.
- Q4. Please confirm whether the reference "Mitchell et al., 2015" cited in the text refers to "Mitchell et al., 2015a" or "Mitchell et al., 2015b."
- Q5. Please provide the expansion for "UT."
- Q6. Please provide the complete reference details for "Trommelen et al., 2019" to be included in the reference list.
- Q7. Please provide descriptions for "*" and "#" used in Figure 1.
- Q8. Please provide page number if available for reference "Joanisse et al. (2020)."