



## King's Research Portal

DOI:

[10.1093/ckj/sfab239](https://doi.org/10.1093/ckj/sfab239)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Mayes, J., Billany, R. E., Vadaszy, N., Young, H. M. L., Castle, E. M., Bishop, N. C., Bramham, K., Nixon, A. C., Wilkinson, T. J., Hamilton, A., Saynor, Z. L., Chilcot, J., Picariello, F., Macdonald, J., & Greenwood, S. A. (2022). The rapid development of a novel kidney-specific digital intervention for self-management of physical activity and emotional wellbeing during the COVID-19 pandemic and beyond: Kidney Beam. *Clinical Kidney Journal*, 15(3), 571-573. <https://doi.org/10.1093/ckj/sfab239>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.




- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## LETTER TO THE EDITOR

# The rapid development of a novel kidney-specific digital intervention for self-management of physical activity and emotional well-being during the COVID-19 pandemic and beyond: Kidney Beam

Juliet Mayes<sup>1</sup>, Roseanne E. Billany<sup>2</sup>, Noemi Vadaszy<sup>2</sup>,  
 Hannah M.L. Young <sup>2,3</sup>, Ellen M. Castle<sup>1,4</sup>, Nicolette C. Bishop<sup>5</sup>,  
 Kate Bramham<sup>1,4</sup>, Andrew C. Nixon<sup>6</sup>, Thomas J. Wilkinson<sup>2</sup>,  
 Alexander J. Hamilton<sup>7</sup>, Zoe L. Saynor<sup>8</sup>, Joseph Chilcot <sup>4</sup>,  
 Federica Picariello <sup>4</sup>, Jamie Macdonald<sup>9</sup> and Sharlene A. Greenwood<sup>1,4</sup>

<sup>1</sup>Therapies Department, King's College Hospital NHS Trust, London, UK, <sup>2</sup>Leicester Kidney Lifestyle Team, University of Leicester, Leicester, UK, <sup>3</sup>Leicester Diabetes Centre, University Hospitals of Leicester NHS Trust, Leicester, UK, <sup>4</sup>Renal Sciences, King's College London, London, UK, <sup>5</sup>School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, UK, <sup>6</sup>Department of Renal Medicine, Lancashire Teaching Hospitals NHS Foundation Trust, Preston, UK, <sup>7</sup>Population Health Sciences, University of Bristol, Bristol, UK, <sup>8</sup>School of Sport, Health and Exercise Science, University of Portsmouth, Portsmouth, UK and <sup>9</sup>School of Human and Behavioural Sciences, Bangor University, Wales, Bangor, UK

Correspondence to: Juliet Mayes; E-mail: [juliet.mayes@nhs.net](mailto:juliet.mayes@nhs.net)

Individuals living with chronic kidney disease (CKD) report high levels of physical inactivity and poor emotional well-being [1]. Despite disease-specific guidelines promoting physical activity (PA) participation [2], people living with CKD do not routinely receive PA or emotional well-being support as part of their clinical care [3].

The challenge to engage people living with CKD with PA and emotional well-being interventions was exacerbated by the coronavirus disease 2019 (COVID-19) global pandemic. Specifically, CKD was identified as a significant risk factor for more severe COVID-19 infection [4, 5], and people with CKD were advised to shield at home. The resultant impact on the physical and emotional well-being of these individuals was captured in a Kidney Care UK patient survey, which revealed that 41%

of respondents felt anxious, lonely or isolated during the pandemic [6].

The COVID-19 pandemic led to an increase in the use of digital technology and 'eHealth' interventions that could deliver aspects of usual and enhanced CKD care. This context led to the current authors rapidly developing and rolling out a kidney-specific PA and emotional well-being platform, Kidney Beam (the promotion was by social media only; no additional technical or clinical support was offered).

The Kidney Beam programme (<https://beamfeelgood.com/home>) is a contextually appropriate, theory-informed intervention that was designed to promote digitally delivered PA and emotional well-being self-management in people living with CKD, as guided by the Behaviour Change Wheel [7]. The

Received: 1.10.2021; Editorial decision: 25.11.2021

© The Author(s) 2021. Published by Oxford University Press on behalf of the ERA. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

**Table 1. Demographic data of participants who signed up to use the Kidney Beam platform**

	Total users (n)	Percentage (%)
Age (years)		
0–18	0	0
18–25	35	4
25–35	155	16
35–45	205	21
45–55	231	24
55–65	209	22
65–75	86	9
75+	35	4
Total	956	100
User type		
Living with condition	736	77
Exercise or healthcare practitioner	127	13
Caregiver	34	4
Other	50	5
Unknown	12	1
Gender		
Male	249	26
Female	707	74
Other	0	0
Kidney Care Clinic		
General Practitioner	121	25
General Nephrology Clinic	122	25
Low Clearance Clinic	15	3
Peritoneal Dialysis Clinic	23	5
Haemodialysis Clinic	34	7
Kidney Transplant Clinic	153	32
Other	14	3
Total	482	100

multifaceted digital intervention package was composed of on-line live and on-demand exercise training videos, careful selection of healthcare professionals and patient champions, use of email feedback and action planning, and online educational videos and blogs.

A 6-month prospective feasibility project was approved by King's College Hospital Research and Innovation team to allow rapid design, delivery and evaluation of Kidney Beam during the COVID-19 pandemic. The project was funded by Kidney Research UK (KRUK). All people in the UK living with CKD aged  $\geq 18$  years were eligible to sign up to the platform. Participants consented to complete a voluntary electronic survey on sign up to the platform and at 6 months, to establish whether participants were meeting current PA guidelines, investigate perceptions of health, and collect usability and acceptability data about the platform. Data were collected between 1 June 2020 and 30 November 2020.

**Table 2. Pre- and post-6-month participation voluntary survey results**

	Baseline, n (%)	6 months, n (%)	Percentage change baseline to 6 months ( $\Delta$ %)
Achieving 150 min of moderate-intensity physical activity (per week)	83 (30)	43 (60)	100
Achieving 75 min of vigorous-intensity physical activity (per week)	97 (35)	36 (42)	20
Achieving twice-weekly strength training	86 (31)	46 (54)	74
Perceived energy levels to be good or very good	55 (20)	26 (30)	50

In total, 959 adults signed up to use the platform within the 6-month time period (see Table 1). There was a 43% activation rate (people who sign-up to the platform completing one or more classes), which is higher than the UK national average for digital healthcare interventions. More than 1900 movement classes were completed during the 6-month time period. A total of 276 participants completed the voluntary survey at baseline, with 85 completing the follow-up survey at 6 months. Results suggest an increase in those participants achieving recommended PA guidelines and in those participants who perceived their energy levels to be good or very good by the end of the 6-month project (see Table 2). Promisingly, 96% of participants surveyed would recommend Kidney Beam, with the biggest reported benefit being that the intervention was kidney-specific and delivered by specialist kidney healthcare professionals.

Digital health interventions that are co-designed with people living with a long-term health condition may result in better clinical and person-centred outcomes [8, 9]. The Kidney Beam programme was rapidly co-developed and delivered with minimal resource during the COVID-19 pandemic. Despite known challenges around digital and health literacy, this digital self-management programme attracted 736 participants living with CKD to sign up to the platform. A randomized controlled wait-list trial is underway to evaluate the feasibility, clinical value and cost-effectiveness of the Kidney Beam programme delivered as part of clinical care (NCT04872933).

## ACKNOWLEDGEMENTS

We thank Kidney Research UK for rapidly funding this feasibility project and enabling the platform to be free at the point of access for all individuals living with CKD.

## FUNDING

The feasibility project was a collaboration between King's College Hospital, Kidney Research UK and Beam, a health technology platform supporting people with health conditions to stay physically active. Kidney Research UK did not have any role in study design; collection, analysis and interpretation of data; writing the report; or the decision to submit the report for publication. Juliet Mayes, Pre-Doctoral Clinical Academic Fellow, (NIHR301893), Hannah Young, Development and Skills Enhancement Fellow (NIHR301593), and Sharlene Greenwood, NIHR Advanced Research Fellow (ICA-CL-2017-03-020), are funded by National Institute for Health Research (NIHR) for this research project. The views

expressed in this publication are those of the authors and not necessarily those of the NIHR, NHS or the UK Department of Health and Social Care.

### AUTHORS' CONTRIBUTIONS

S.A.G. and J. Mayes contributed to acquisition, analysis and interpretation of data. S.A.G., J.Mayes, R.E.B., N.V., H.M.L.Y., E.M.C., N.C.B., A.H., Z.L.S., J.C., F.P., J. Macdonald, K.B., A.C.N. and T.J.W. contributed important intellectual content during manuscript drafting or revision, and accept accountability for the overall work by ensuring that questions pertaining to the accuracy or integrity of any portion of the work are appropriately investigated and resolved.

### CONFLICT OF INTEREST STATEMENT

The results presented in this article have not been published previously in whole or part.

### REFERENCES

1. Wilkinson TJ, Clarke AL, Nixon DGD et al. Prevalence and correlates of physical activity across kidney disease stages: an observational multicentre study. *Nephrol Dial Transplant* 2021; 36: 641–649
2. The Renal Association. *UKKA guidelines, Clinical Practice Guideline Exercise and Lifestyle in Chronic Kidney Disease*. Published April 2021. [https://ukkidney.org/sites/renal.org/files/Exercise%20and%20Lifestyle%20in%20CKD%20clinical%20practice%20guideline33\\_v4\\_FINAL\\_0.pdf](https://ukkidney.org/sites/renal.org/files/Exercise%20and%20Lifestyle%20in%20CKD%20clinical%20practice%20guideline33_v4_FINAL_0.pdf) (15 September 2021, date last accessed)
3. Koufaki P, Greenwood S, Painter P et al. The BASES expert statement on exercise therapy for people with chronic kidney disease. *J Sports Sci* 2015; 33: 1902–1907
4. ERA-EDTA Council, ERACODA Working Group. Chronic kidney disease is a key risk factor for severe COVID-19: a call to action by the ERA-EDTA. *Nephrol Dial Transplant* 2021; 36: 87–94
5. Clark A, Jit M, Warren-Gash C et al. Global, regional, and national estimates of the population at increased risk of severe COVID-19 due to underlying health conditions in 2020: a modelling study. *Lancet Glob Health* 2020; 8: e1003–e1017
6. KidneyCareUK. *KCUK Patient Survey Report 2020*. <https://www.kidneycareuk.org/news-and-campaigns/news/fears-kidney-patients-government-coronavirus-advice-leaves-thousands-dark/> (21 June 2021, date last accessed)
7. Michie S, Atkins L, West R. *The Behaviour Change Wheel: A Guide to Designing Interventions*. London: Silverback Publishing, 2014
8. Kitsiou S, Pare G, Jaana M et al. Effectiveness of mHealth interventions for patients with diabetes: an overview of systematic reviews. *PLoS One* 2017; 12: e0173160
9. Widmer RJ, Collins NM, Collins CS et al. Digital health interventions for the prevention of cardiovascular disease: a systematic review and meta-analysis. *Mayo Clin Proc* 2015; 90: 469–480