



King's Research Portal

DOI: 10.3390/metabo12070601

Document Version Publisher's PDF, also known as Version of record

Link to publication record in King's Research Portal

Citation for published version (APA):

Louca, B., Nogal MacHo, A., Moskal, A., Goudling, N., Shipley, M., Alkis, T., Lindbohm, J., Hu, J., Kifer, D., Wang, N., Chawes, B., Rexrode, K., Ben-Shlomo, Y., Kivimaki, M., Murphy, R., Yu, B., Gunter, M. J., Suhre, K., Lawlor, D. A., ... Menni, C. (2022). Cross-Sectional Blood Metabolite Markers of Hypertension: A Multicohort Analysis of 44,306 Individuals from the COnsortium of METabolomics Studies. *Metabolites*, *12*(7), Article 601. https://doi.org/10.3390/metabo12070601

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
You may not further distribute the material or use it for any profit-making activity or commercial gain
You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





Cross-Sectional Blood Metabolite Markers of Hypertension: A Multicohort Analysis of 44,306 Individuals from the **COnsortium of METabolomics Studies**

Panayiotis Louca ¹, Ana Nogal ¹, Aurélie Moskal ², Neil J. Goulding ^{3,4}, Martin J. Shipley ⁵, Taryn Alkis ⁶, Joni V. Lindbohm ^{5,7}, Jie Hu ⁸, Domagoj Kifer ⁹, Ni Wang ^{10,11}, Bo Chawes ¹⁰, Kathryn M. Rexrode ⁸, Yoav Ben-Shlomo^{3,12}, Mika Kivimaki⁵, Rachel A. Murphy^{13,14}, Bing Yu⁶, Marc J. Gunter², Karsten Suhre¹⁵, Deborah A. Lawlor^{3,4,16}, Massimo Mangino^{1,17}, and Cristina Menni^{1,*}

- 1 Department of Twin Research, King's College London, London SE1 7EH, UK;
- panayiotis.louca@kcl.ac.uk (P.L.); ana.nogal_macho@kcl.ac.uk (A.N.); massimo.mangino@kcl.ac.uk (M.M.) 2 Nutrition and Metabolism Section, International Agency for Research on Cancer, 69372 Lyon, France;
- aurelie.moskal@inserm.fr (A.M.); gunterm@iarc.fr (M.J.G.) 3 Population Health Sciences, Bristol Medical School, University of Bristol, Bristol BS8 2BN, UK;
- n.goulding@bristol.ac.uk (N.J.G.); y.ben-shlomo@bristol.ac.uk (Y.B.-S.); d.a.lawlor@bristol.ac.uk (D.A.L.)
- MRC Integrative Epidemiology Unit, University of Bristol, Bristol BS8 2BN, UK
- 5 Department Epidemiology and Public Health, University College London, London WC1E 7HB, UK; martin.shipley@ucl.ac.uk (M.J.S.); joni.lindbohm@helsinki.fi (J.V.L.); m.kivimaki@ucl.ac.uk (M.K.)
- Department of Epidemiology, Human Genetics and Environmental Sciences, School of Public Health, The University of Texas Health Science Center, Houston, TX 77030, USA; taryn.alkis@uth.tmc.edu (T.A.); bing.yu@uth.tmc.edu (B.Y.)
- 7 Clinicum, Department of Public Health, University of Helsinki, P.O. Box 20 Helsinki, Finland
- 8 Division of Women's Health, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02115, USA; jhu13@bwh.harvard.edu (J.H.); krexrode@bwh.harvard.edu (K.M.R.)
- Faculty of Pharmacy and Biochemistry, University of Zagreb, 10000 Zagreb, Croatia; dkifer@pharma.hr 10 Copenhagen Prospective Studies on Asthma in Childhood, Herlev and Gentofte Hospital, University of
- Copenhagen, 2820 Gentofte, Denmark; ni.wang@dbac.dk (N.W.); chawes@copsac.com (B.C.) 11 Department of Biotechnology and Biomedicine, Technical University of Denmark,
- 2800 Kongens Lyngby, Denmark 12
- NIHR Applied Research Collaboration West, University Hospitals Bristol and Weston National Health Service Foundation Trust, Bristol BS1 2NT, UK
- 13 School of Population and Public Health, University of British Columbia, Vancouver, BC V6T 1Z3, Canada; rachel.murphy@ubc.ca
- 14 Cancer Control Research, BC Cancer, Vancouver, BC V5Z 1G1, Canada
- 15 Department of Biophysics and Physiology, Weill Cornell Medicine-Qatar, Doha 24144, Qatar; karsten@suhre.fr
- 16 Bristol NIHR Biomedical Research Centre, University of Bristol, Bristol BS1 2NT, UK
- 17 NIHR Biomedical Research Centre at Guy's and St Thomas' Foundation Trust, London SE1 9RT, UK
 - Correspondence: cristina.menni@kcl.ac.uk; Tel.: +44-(0)-207-188-7188 (ext. 52594)

Abstract: Hypertension is the main modifiable risk factor for cardiovascular morbidity and mortality but discovering molecular mechanisms for targeted treatment has been challenging. Here we investigate associations of blood metabolite markers with hypertension by integrating data from nine intercontinental cohorts from the COnsortium of METabolomics Studies. We included 44,306 individuals with circulating metabolites (up to 813). Metabolites were aligned and inverse normalised to allow intraplatform comparison. Logistic models adjusting for covariates were performed in each cohort and results were combined using random-effect inverse-variance meta-analyses adjusting for multiple testing. We further conducted canonical pathway analysis to investigate the pathways underlying the hypertension-associated metabolites. In 12,479 hypertensive cases and 31,827 controls without renal impairment, we identified 38 metabolites, associated with hypertension after adjusting for age, sex, body mass index, ethnicity, and multiple testing. Of these, 32 metabolite associations, predominantly lipid (steroids and fatty acyls) and organic acids (amino-, hydroxy-, and keto-acids) remained after further adjusting for comorbidities and dietary intake. Among the identified metabolites, 5 were novel, including 2 bile acids, 2 glycerophospholipids, and ketoleucine. Pathway analysis further implicates the role of the amino-acids, serine/glycine, and bile acids in hypertension regulation.



Citation: Louca, P.; Nogal, A.; Moskal, A.; Goulding, N.J.; Shipley, M.J.; Alkis, T.; Lindbohm, J.V.; Hu, J.; Kifer, D.; Wang, N.; et al. Cross-Sectional Blood Metabolite Markers of Hypertension: A Multicohort Analysis of 44,306 Individuals from the COnsortium of METabolomics Studies. Metabolites 2022, 12, 601. https://doi.org/ 10.3390/metabo12070601

4

9

Academic Editor: Pollen K. Yeung

Received: 16 May 2022 Accepted: 9 June 2022 Published: 28 June 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).

In the largest cross-sectional hypertension-metabolomics study to date, we identify 32 circulating metabolites (of which 5 novel and 27 confirmed) that are potentially actionable targets for intervention. Further in-vivo studies are needed to identify their specific role in the aetiology or progression of hypertension.

Keywords: metabolomics; hypertension

1. Introduction

High blood pressure (BP) is the leading modifiable risk factor for cardiovascular disease (CVD), global morbidity, and mortality, and recognised as the greatest single risk factor contributing to the global burden of disease—responsible for 8.5 million deaths per year [1,2]. BP is complex and multifactorial by nature, influenced by genomic, lifestyle, and environmental factors and a multitude of physiological pathways [3,4]. Reducing the burden of hypertension is therefore important, and can be achieved via improving the coverage of treatment and primary prevention, which has become an objective of many global and national health initiatives [2]. However, the pace of progress in this regard has stalled [5], partly due to the deceleration of research investigating novel targets, which could then be used for drug/treatment development [6].

Circulating metabolites comprise the intermediate and end-products of metabolic pathways that reflect physiological processes [7]. Therefore, metabolites are well suited to characterise the influence of key measures of disease risk brought about by environmental, nutritional, or lifestyle factors, thus leading to the discovery of novel metabolic aspects of complex diseases [8,9].

Previous work has identified numerous metabolites linked with hypertension and blood pressure [4,10–13]. The majority of which falling into the lipid and amino acid classes [4]. This includes the dicarboxylic fatty acid hexadecanedioate that is associated with increased BP mortality [11] and heart failure [14], and the amino acid phenylacetylglutamine [15].

Though metabolomics has been promising in identifying disease biomarkers, assays are expensive, typically performed on <1000 subjects, and no single metabolomics platform covers the full range of detectable metabolites [16], with a modest overlap between platforms [16]. Accordingly, studies on the metabolic mechanisms underlying hypertension are limited, and lack sufficient sample size [16]. The COnsortium of METabolomics Studies (COMETS) is a partnership between many well-characterised prospective cohort studies to provide a framework for collaborative metabolomics research, specifically initiated to advance our understanding of the metabolomes' involvement in disease aetiology, diagnosis, and treatment, by overcoming some of the abovementioned challenges [16].

The aim of this study is to identify blood metabolite markers of essential hypertension, which may provide novel actionable targets and pathways for hypertension treatments. To achieve this, we will conduct a metabolome-wide association study (MWAS) of hypertension by integrating data from nine intercontinental cohorts, including 44,306 individuals of different ethnicities, sex, age, and metabolomic platforms, from COMETS. We then investigate canonical pathways underlying the hypertension-associated metabolites.

2. Results

2.1. Study Demographics

We included 12,479 essential hypertensive cases and 31,827 non-hypertensive controls from 9 cohorts (n = 44,306) and analysed a total of 813 serum/plasma metabolites. The demographic characteristics of the study sample are presented in Table 1. Average age ranged from 27.95 (± 5.69) to 74.67 (± 2.84) years, and on average subjects were overweight with an average body mass index (BMI) in the range of 26.1 (± 4.81) to 28.93 (± 5.93) kg/m². In total we included 24,864 female and 19,442 male individuals, of which 19,448 were of

Cohort ALSPAC ARIC BIB CaPS EPIC HealthABC QBB TwinsUK Whitehall II 9396 3293 16,418 1795 989 232 2906 4427 4850 n mean (SD) 74.7 (2.8) 27.1 (4.4) 138.1 (23.3) 56.2 (6) 26.3 (3.9) 125.2 (18.3) 40.9 (12.5) 52.9 (5.5) 28.9 (5.9) 28 (5.7) 26.8 (5.9) 61.4 (4.4) 26.8 (3.7) 54 (13.3) 56.1 (8) 39 (12) Age BMI 26.7 (4.1) 138.3 (22.5) 28.9 (5.9) 26.1 (5) 120.3 (13.8) 26.1 (4.8) 127.4 (18.6) 115.9 (17) SBF 129.8 (23.3) 110.6 (11.8) 148.6 (24.4) DBP 71.1 (9.1) 77.9 (14.3) 78.5 (10.8) 79.9 (13.5) 65.6 (8.5) 84.1 (13.2) 84.8 (12.6) 74 (11.2) 78.6 (11.3) n (%) n (%) n (%) n (%) n (%) n(%)n (%) n (%) n (%) 1007 (10.7) 1322 (29.9) HTN cases 1501 (45.6) 631 (63.8) 6897 (42) 48 (20.7) 409 (14.1) 646 (13.3) 18(1) Non-HTN 4204 (86.7) 8389 (89.3) 1792 (54.4) 1777 (99) 9521 (58) 184 (79.3) 2497 (85.9) 3105 (70.1) 358 (36.2) controls Sex 989 (100) 3062 (33) 1375 (41.8) 8659 (52.7) 232 (100) 1457 (50.1) 339 (7.7) 3329 (68.6) Males Females 6334 (67) 1918 (58.2) 1795 (100) Ò 7759 (47.3) 1449 (49.9) 4088 (92.3) 1521 (31.4) Ancestry 989 (100) White 8436 (89.8) 1053 (32) 0 3653 (82.5) 4475 (92.3) 842 (46.9) NA 2240 (68) 40 (0.4) NA 232 (100) 0 16 (0.4) 103 (2.1) Black Ò 43 (0.5) Asian/Hispanic 0 953 (53.1) 0 NA 2906 (100) 27 (0.6) 234 (4.8) Other 37 (0.4) 0 Ò 0 NA 0 0 Ò 37(0.7)

 Table 1. Descriptive characteristics of included cohorts.

Abbreviations: SD, standard deviation; ALSPAC, Avon Longitudinal Study of Parents and Children; ARIC, Atherosclerosis Risk in Communities Study; BIB, Born in Bradford; CaPS, Caerphilly Prospective Study; EPIC, European Prospective Investigation into Cancer and Nutrition; HealthABC, Health, Aging and Body Composition Study; QBB, Qatar Biobank; BMI, body mass index; SBP, systolic blood pressure; DBP diastolic blood pressure.

White/European ancestry, 2631 of black ancestry, 4163 of Asian or Hispanic ancestry, and

2.2. Discovery Analysis

74 of other ancestries.

To identify circulating metabolites associated with essential hypertension in the overall sample, we meta-analysed the age, age², sex, BMI, and ethnicity adjusted results from each of the nine contributing cohorts. We identified 38 metabolites associated with hypertension cross-sectionally, after adjusting for multiple testing (Figure 1 and Supplementary Table S2). To the best of our knowledge, 6 of these were novel hypertension associations. These 38 hypertension-associated metabolites were comprised of 14 lipids (36.8%) (6 fatty acyls, 5 steroids, and 3 glycerophospholipids), 13 organic acids (34.2%) (9 amino acids and peptides, 2 short-chain-keto acids, and 2 hydroxy acid), 5 organic oxygen compounds (13.2%) (3 carbohydrates and 2 alcohols), 5 organoheterocyclic compounds (13.2%) (3 purines, 1 indole, and 1 pyridoxine), and 1 (2.6%) phenylpropanoid (hydrocinnamic acid) (Figure 1).



Figure 1. Hypertension-associated metabolites after adjusting for traditional risk factors. Colours and groupings represent each metabolite super class, while a black border signifies novel-associations with hypertension/BP.

For the 38 metabolites that passed our multiple testing threshold, heterogeneity statistics ranged from 0% to 65% (median = 26.6%) for I^2 , and from 0 to 14 (median = 1.1) for Cochrane's Q.

Of the 14 lipids, all were positively associated with hypertension, and the strongest association was Glycerylphosphorylethanolamine (odds ratio (OR) [95%CI] = 1.31 [1.16, 1.48]). The strongest positive and negative associations of the 13 organic acids, was Homocitrulline (OR [95%CI] = 1.25 [1.17, 1.34]) and Serine (OR [95%CI] = 0.76 [0.72, 0.8]), respectively. Of 5 positively associated organic oxygen compounds, erythritol elicited the strongest association (OR [95%CI] = 1.2 [1.11, 1.31]). Similarly, all 5 organoheterocyclic were positively associated with hypertension, with pyridoxine (OR [95%CI] = 1.31 [1.24, 1.37]) having the highest OR. While the only phenylpropanoid to be identified, hydrocinnamic acid, was negatively associated with hypertension (OR [95%CI] = 0.88 [0.82, 0.94]) (Figure 2).



Figure 2. Odds ratio's and 95% confidence intervals of hypertension associated metabolites adjusting for traditional risk factors (TRF) (green) or TRF + diet + comorbidities (red). Semi-transparent bars illustrate analyses not passing the pre-defined alpha threshold. Base line colours represent the super class of metabolites, while sub-base labels indicate the metabolite sub class.

2.3. Sensitivity Analysis

To investigate the robustness of these associations, using inverse-variance metaanalysis we further pooled the fully adjusted model (further adjusting for dietary salt and alcohol intakes and comorbidities) estimates for the 38 hypertension-associated metabolites.

Out of the 38 metabolites, 32 remained significant after further adjustments and multiple testing ($p < 0.05/38 = 1.3 \times 10^{-3}$).

The more robust hypertension-associated metabolites were still predominantly lipids (12 of 14, 37.5%) and organic acids (11 of 13, 34.4%), but also 4 of the 5 organic oxygen com-

pounds and 4 of the 5 organoheterocyclic compounds remained, as did hydrocinnamic acid. Of the 32 metabolites, 5 of the novel hypertension-associations also remained associated (Figure 2 and Supplementary Table S2).

While further accounting for dietary influences and comorbidities, the organoheterocyclic metabolite, pyridoxine still elicited the strongest positive effect on hypertension (OR [95%CI] = 1.31 [1.24, 1.38]), while the amino acid serine also was the strongest negative association (OR [95%CI] = 0.78 [0.74, 0.83]) (Figure 2).

2.4. Stratified Analyses

To investigate if key demographic factors may be driving metabolite associations, we conducted a sub-analysis by stratifying our samples by (i) sex and (ii) race/ethnicity. We then pooled the estimates for both the traditional-risk factor adjusted and fully adjusted models for the 32 hypertension associated metabolites.

For the traditional risk factor adjusted estimates, when stratifying by sex, out of the 32 metabolites, 20 metabolites were associated in the male only sample, and 27 were associated in the female only sample. When we stratified by race/ethnicity 14 of the 32 metabolites were associated with hypertension in those of White/European ancestry, and 10 in those of Black ancestry (Figure 3).



Figure 3. Forest plot of traditional risk factor adjusted analyses stratified by sex, and ethnicity. Metabolites are grouped by class. Points represents odds ratio (OR), and tails show the 95% confidence interval (CI). For an improved scale, associations with an upper CI > 3 have been truncated, reducing the standard error by 2.5 times, and are identified by a * above the OR point. For true values see Supplementary Figure S2. Metabolites not passing an α level of 0.05 are shown with a white point. Metabolites where there were fewer than 2 cohorts with the metabolite detected in >80% of the stratified sample are missing.

Results when further adjusting for comorbidities and alcohol and salt intakes remained consistent. For the fully adjusted model estimates, 18 of the 20 metabolites remained associated in males, while all 27 hypertension-associated metabolites remained significant in females (at p < 0.05). For race/ethnicity, 15 metabolites were associated in White/European individuals and all 10 previously identified metabolites were associated in those of Black ancestry (at p < 0.05) (Supplementary Figure S2).

2.5. Pathway Analysis

We further conducted canonical pathway analysis of the 32 hypertension-associated metabolites using the IPA database to explore the metabolic footprint of those metabolites. Canonical pathway analysis suggests that the probe set of metabolites were likely to be involved in tRNA charging, glycine, leucine and serine pathways, folate pathways, and bile acid biosynthesis (Supplementary Figure S3).

3. Discussion

In the largest MWAS of essential hypertension study to date, including over 44,000 individuals from COMETS, we identify 32 circulating metabolites associated with hypertension, after adjusting for traditional risk factors, diet, comorbidities, and multiple testing. Five of these metabolites were novel and had not yet been previously shown to associate with hypertension or blood pressure. These include 4 lipids (2 Glycerophospholipids, and 2 bile acids) and 1 organic acid (the short-chain keto acid, ketoleucine). The majority of the identified metabolites were lipids, including steroids (2) and bile acids (2), and organic acids, including amino acids (7). Organic oxygen compounds (4), organoheterocyclic compounds (4), and phenylpropanoids (1) were also identified. Pathway analysis in the IPA database support these findings, identifying tRNA charging, as the pathway in which the metabolite probe set was most likely to be involved, and additionally showing significant overlap with glycine and serine pathways (Supplementary Figure S3). Pathway analysis also suggests a renal role via bile acids and folate.

Results in women and men were mostly statistically consistent with each other, as were those comparing White/European to African origin ethnicity. However, for malic acid the confidence intervals between males and females did not overlap, with an OR [95%CI] in men of 1.41 [1.26, 1.57] and 1.15 [1.09, 1.23] in women, despite these estimates being imprecise.

We find novel associations between circulating levels of the secondary bile acids glycocholic acid and chenodeoxycholic acid glycine conjugate (GCDCA), and hypertension, both of which were positively associated. Results were consistent when stratifying by sex and by race/ethnicity. Bile acids are recognised to facilitate transport of dietary lipids and fat-soluble vitamins; regulate glucose homeostasis, lipids and lipoprotein metabolism, energy expenditure; and influence inflammation [17]. In addition to these, the vasoactive properties of bile acids have been recognised for decades, although the mechanisms by which they act has not been defined [18]. The expression of bile acid receptors in endothelium has conjured hypotheses relating to nitric oxide production, a potent vasodilator [18]. In murine models of hypertension the infusion of bile acids (deoxycholic acid) significantly reduced arterial BP by 12 mmHg [19]. Although glycocholic acid has not previously been associated with hypertension or blood pressure, previous evidence suggests links with other cardiovascular factors, such as atrial fibrillation [20], and its precursor, cholic acid, has been causally linked with the development of hypertension within murine models [21]. GCDCA on the other hand has been linked with toxicity in cardiac mitochondria in vitro [22].

Amino acids, including serine, glycine, histidine, and alanine, have previously and consistently been implicated in a variety of cardiometabolic traits, including hypertension and CVD [7,11,23], and amino-acid metabolism is considered one of the critical pathways involved in hypertension regulation [24]. Indeed, amino acids have been implicated in intracellular signalling, whereby dysregulation of amino acid metabolism may result in inflammation, oxidative stress, and insulin resistance, states involved in the aetiology

of hypertension [25]. Essential amino acids cannot be synthesised de novo and must be acquired from dietary sources. Here we report negative associations between the essential amino acid histidine and hypertension, and positive associations for the essential amino acids isoleucine, and leucine, in accordance with previous studies on coronary heart disease [26]. For the non-essential amino acids, both glycine, and serine were negatively associated with hypertension. Abundance of non-essential amino acids are symbolic of protein catabolism and links have been made to their involvement in pulsatile haemodynamics [23]. These findings are in line with previous reports [7,11,23]. We have previously reported inverse associations between pulse wave velocity and various amino acids in a sample of 1797 normotensive White/European women [7]. Urinary levels of amino acids have been repeatedly correlated with blood pressure, hypertension, and vascular factors. Several amino acids, including, serine, glycine, and histidine, were inversely related to SBP and pulse pressure in individuals of Black ancestry [23]. On the other hand, of the seven hypertension-associated amino acids we find here, only isoleucine, remains significant when stratifying by Black ancestry. These findings are supported by our canonical pathway analysis in the IPA database, which also matches the findings of Zhao and colleagues [24].

There is a long-standing relationship between various lipids and CVD. Here, we identify 14 lipids positively associated with hypertension (Figure 3). Within these, 4 were fatty acids, including tetradecanedioic acid, stearidonic acid, palmitoylcarnitine, and acetylcarnitine.

Fatty acids are a major energy source for myocardial tissue, and long-chain fatty acids in particular have been implicated in cardiovascular events [27]. These reports are in line with previous evidence. In murine models of hypertension stearidonic acid was found to be significantly higher compared to normotensive controls [28]. Moreover, the alpha, omega-dicarboxylic acid, tetradecanedioic acid, is a close relative to hexadecanedioic acid, another long-chain dicarboxylic acid with causal evidence linked to increased BP [11].

Within the 14 positively associated lipids were also 2 hydroxy- and 1 androstanesteroids, namely, cortisol, cortisone and 5-androstenediol. Elevated levels of steroid hormones have been implicated with an increased risk for the development of hypertension [24]. Cushing's syndrome, a disorder of over secretion of cortisol, presents with high BP [29], and this is believed to be a direct consequence of the effect of cortisol on adrenocorticotrophic hormone. Moreover, an infusion of cortisol over a 5-day period has been shown to increase SBP by 21 mmHg [30].

The current study has several strengths. By leveraging the COMETS our study is to-date the largest MWAS of hypertension, with a substantial sample size to detect any rare metabolites. This also facilitates the inclusion of cohorts spanning multiple continents with diverse ethnic demographics. The MWAS approach also facilitated the inclusion of a range of metabolomics platforms, increasing our scope for potential hypertension-associated metabolites.

However, despite these strengths our results must be appreciated in the presence of several limitations. Firstly, although we account for potential confounders including, dietary intake and comorbidities, in large prospective-observational cohorts, food frequency questionnaires are the typical method for dietary data collection, while comorbidities are recorded by health questionnaires, rather than derived from medical records. The selfreported nature of this data may introduce reporting bias, which includes social desirability and selective recall, resulting in misclassification. Such misclassification of confounders could result in residual confounding. Furthermore, cohorts were included in the sensitivity analyses even if they did not have data on all potential confounders, and this would also introduce residual confounding. Moreover, non-hypertensive controls, may have had other comorbidities, which could contribute to changes in metabolism and therefor a potential source of confounding. Second, given the study designs of the contributing cohorts data included may not have been recorded/reported on the same day. Third, despite leveraging the COMETS, our sample was not represented equally by all race/ethnicities, and was mostly comprised of White/European individuals, consequently our stratified analysis in those of Black ethnicity may not have been sufficiently powered, and we were unable to have an appreciable overlap of metabolites to stratify our sample by Asian/Hispanic or other ancestries. Moreover, in our stratified analysis not all metabolites were present in >1 cohort and >80% of the stratified sample, therefore comparisons between sexes and ethnicities for some metabolites cannot be made. Additionally, regional disparities in the metabolome have previously been reported [31], but we were unable to account for it, as specific regional data was not available. Fourth, to incorporate all cohorts, here we have used cross-sectional metabolomics. This lack of longitudinal metabolomics prevents us from investigating causality or validating the predictive power of these metabolites in relation to hypertension. Moreover, hypertension was defined from a single timepoint, and we were unable to describe the potential duration of hypertension, as longitudinal BP data was unavailable in all cohorts. Lastly, although we include over 44,000 individuals and multiple metabolomic platforms, metabolites are not measured by each platform, thus the limited overlap between platforms reduces statistical power.

4. Methods

A flowchart of the study design is presented in Supplementary Figure S1.

4.1. Study Populations

This study is drawn from COMETS, which is described in detail elsewhere [16].

Here we included cohorts from COMETS that had concurrent systolic blood pressure (SBP), diastolic blood pressure (DBP), and metabolomics on participants without predetermined renal impairment. No a priori exclusion was applied to cohorts for the number of hypertensive cases (mean cases = 1387). Additional requested covariates included: age, sex, BMI, ethnicity, use of antihypertensive medication, co-morbidities (prevalence of cancer, diabetes, and heart disease), serum creatinine, and dietary information (salt and alcohol intake). We also included data from the Qatar Biobank cohort, a non-COMETS member, but with data matching the COMETS recruitment criteria (Supplementary Figure S1).

A total of nine prospective cohorts were recruited spanning three continents, and are described in Supplementary Table S1. They include Asia: Qatar Biobank (QBB), a population-based cohort study in Qatar, the first of its kind in the Gulf region [32]; Europe: Avon Longitudinal Study of Parents and Children (ALSPAC), a multi-generation birth cohort study, recruited from the former county of Avon in South-West England [33–36], Born in Bradford (BIB), a multi-ethnic pregnancy and birth cohort recruited from the north of England [37], Caerphilly Prospective Study (CaPS), an epidemiological cohort with a sample representative of a small town in South Wales, UK [38], European Prospective Investigation into Cancer and Nutrition (EPIC), a multi-centre cohort comprised of 23 centres in 10 European countries [39], TwinsUK, the largest cohort of community-dwelling adult twins in the UK [40], and Whitehall II (WII), a longitudinal cohort of British civil servants recruited from London, England [41]; North America: Atherosclerosis Risk in Communities Study (ARIC), a large prospective study recruited from four US communities of Minnesota, North Carolina, Maryland, and Mississippi [42], and Health, Aging and Body Composition Study (HealthABC), a mixed ethnicity cohort recruited from four US states, including Memphis, Tennessee or Pittsburgh, and Pennsylvania [43] (Supplementary Figure S1).

4.2. Phenotypes

4.2.1. Metabolomics

Metabolites were quantified from blood in each study using targeted or untargeted assays (Supplementary Table S1), with most studies utilising mass-spectrometry from Metabolon, Inc. (Morrisville, NC, USA). Metabolites were aligned between studies and across platforms using the universal ID developed by the COMETS harmonisation working group [44], and where not available by the Human Metabolome Database [45] (HMDB) identifier directly from each cohort.

4.2.2. Blood Pressure

Both SBP and DBP were measured within each study (mmHg) from a single timepoint, at the closest visit to the metabolomics blood sample collection. Essential hypertension was then defined following the European Society of Hypertension 2021 guidelines [1]. If participants' SBP \geq 140 or their DBP \geq 90, or they were using anti-hypertensive medication at the time of the BP measurement, they were classified as hypertensive cases, otherwise they were considered as non-hypertensive controls.

4.2.3. Covariates

Within each study, demographics, lifestyle factors (diet), anthropometrics, and medical data were collected at the closest visit to the metabolomics blood sample. Variables included in this study were age; sex; ethnicity; BMI (kg/m²); use of antihypertensive medication; presence of cancer, diabetes, or heart disease; serum creatinine (μ mol/L); dietary salt (g/day) and alcohol intake (g/day).

4.3. Statistical Analysis

Standardised pipelines were followed by each cohort to analyse data using the R programming language [46].

Ethnicity was defined using 4 levels: White/European, Black, Asian or Hispanic, or other ancestry. Salt intake was defined by a binary boundary, based on current recommendations [47], ≥ 6 g/day, or ≤ 6 g/day. Alcohol intake was categorised using 4 cut-offs, 0 g per day, >0 g and <15 g per day, ≥ 15 g per day and <30 g per day, or ≥ 30 g day.

For the current analyses, cohorts excluded all individuals predefined as renally impaired. To further mitigate potential confounding effects brought about by renal dysfunction, where serum creatinine was available, we estimated glomerular filtration rate (eGFR) using the Modification of Diet in Renal Disease equation [48], and individuals with an eGFR <60 were excluded. To account for inter-study differences and improve normality all metabolites were quantile normalised using a rank-based inverse normal transformation [49], and to mitigate spurious associations metabolites were excluded if present in <80% of the sample.

Logistic regressions were used to estimate the associations between metabolites and hypertension status. Covariates were determined a priori to the analysis based on previous literature. Basic models were adjusted for traditional risk factors (age, age², sex, BMI, and ethnicity) and, where necessary, batch correction was included. As a sensitivity analysis we ran a second multivariable-adjusted model further adjusted for dietary salt and alcohol intakes and the prevalence of comorbidities (dichotomous variables for cancer, diabetes, and heart disease,). As a sub-analysis, we also ran both models stratified by (i) sex and (ii) ethnicity.

As we wanted to include fully annotated and therefore actionable metabolites, here we only included metabolites that were identifiable by a HMDB id and were analysed in more than 1 cohort. First, we ran random-effects inverse variance meta-analyses using the R package "meta" to pool estimates from the basic model. *p* values were corrected for multiple testing using a Bonferroni correction ($0.05 \times 813 = 6.15 \times 10^{-5}$). For all hypertension-associated metabolites (after multiple testing) we tested the robustness of associations by further pooling the estimates for the multivariable adjusted model using the same methods (Supplementary Figure S1). Heterogeneity in study-specific estimates were examined using the Cochrane Q-value and I² statistics.

We then conducted a canonical pathway analysis using the Ingenuity pathway analysis (IPA) database [50] to investigate metabolic pathways underlying the hypertension-associated metabolites (Supplementary Figure S1). A right-tailed Fisher's exact test was used to calculate a *p*-value determining the probability ($\alpha = 0.05$) that the association between the hypertension-associated metabolites and the canonical pathway was not explained by chance alone.

5. Conclusions

In the largest metabolome wide association study of hypertension to date, including multi-ethnic cohorts, we report 5 novel hypertension-associated metabolites and confirm 27 previous hypertension-associations, highlighting the influence of amino acids, and lipids—including the novel positive-relationship with bile acids. Canonical pathway analysis supports these findings. The clinical implications of these metabolites lie in a series of follow-up studies investigating the molecular pathways related to hypertension, as this may lead to the identification of molecular mechanisms involved in cardiovascular diseases, particularly those linked to glycine, serine, and bile acids, that act through other pathways. The identification of key metabolites related to hypertension should encourage further research into this field.

Supplementary Materials: The following supporting information can be downloaded at: https: //www.mdpi.com/article/10.3390/metabo12070601/s1, Supplementary Figure S1: Flowchart of study analytical pipeline.; Supplementary Figure S2: Heatmap of overall and stratified analyses grouped by metabolite class; Supplementary Table S1: Descriptive table of contributing COHORTS; Supplementary Table S2: Hypertension-associated metabolites. References [51–62] are cited in the Supplementary Materials.

Author Contributions: C.M. conceived and designed the experiment; P.L., A.M., N.J.G., M.J.S., T.A., J.V.L., R.A.M. and K.S., ran the analysis; C.M. and P.L. verified the underlying data; P.L. and C.M. wrote the original manuscript. A.N., J.H., D.K., N.W., B.C., K.M.R., Y.B.-S., M.K., B.Y., M.J.G., D.A.L. and M.M., contributed methods/materials/analysis tools. All authors have read and agreed to the published version of the manuscript.

Funding: This work was funded by the Chronic Disease Research Foundation (CDRF) and the MRC AIM-HY project grant TwinsUK is funded by the Wellcome Trust, Medical Research Council, Versus Arthritis, European Union Horizon 2020, Chronic Disease Research Foundation (CDRF), Zoe Global Ltd. and the National Institute for Health Research (NIHR) Clinical Research Network (CRN) and Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in partnership with King's College London). The Health Aging and Body Composition Study was supported in part by the Intramural Research Program of the National Institutes of Health, National Institute on Aging (NIA) and National Cancer Institute; the NIA Contracts N01-AG-6-2101, N01-AG-6-2103, and N01-AG-6-2106; NIA Grant R01-AG028050, National Institute of Nursing Research Grant R01-NR-012459, the Wake Forest University Claude D. Pepper Older Americans for Independence Center (1P30AG21332); and the Pittsburgh Claude D. Pepper Center (P30 AG024827). Qatar Biobank is supported by the Qatar Foundation. The Whitehall II study is supported by the Wellcome Trust (221854/Z/20/Z), the UK Medical Research Council (MR/S011676/1), and the National Institute on Aging (NIH), US (R01AG056477). The Atherosclerosis Risk in Communities study has been funded in whole or in part with Federal funds from the National Heart, Lung, and Blood Institute, National Institutes of Health, Department of Health and Human Services, under Contract nos. (75N92022D00001, 75N92022D00002, 75N92022D00003, 75N92022D00004, 75N92022D00005). The Caerphlliy Propsective Study (CaPS) was undertaken by the former MRC Epidemiology Unit (South Wales) and was funded by the Medical Research Council of the United Kingdom. The University of Bristol Medical School act as the data custodians (https://www.bristol.ac.uk/population-health-sciences/projects/ caerphilly/, accessed on 8 June 2022) The UK Medical Research Council and Wellcome (Grant ref: 217065/Z/19/Z) and the University of Bristol provide core support for ALSPAC. Deborah Lawlor, and Neil Goulding contributed to this work and will serve as guarantors for the ALSPAC contents of this paper. A comprehensive list of grants funding is available on the ALSPAC website (http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf, accessed on 8 June 2022). Work from ALSPAC authors was specifically funded by the British Heart Foundation (Grant ref: SP/07/008/24066 and CS/15/6/31468), Wellcome Trust (Grant ref: WT092830/Z/10/ Z), Wellcome Trust and MRC (Grant ref: 092731) and NIHR (Grant ref: NF-SI-0611-10196). C.M. is funded by the Chronic Disease Research Foundation and by the MRC AIM-HY project grant. P.L. is funded by the Chronic Disease Research Foundation (CDRF-15/2018). A.N is funded by the Chronic Disease Research Foundation. M.M. is funded by the National Institute for Health Research (NIHR)-funded BioResource, Clinical Research Facility and Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in partnership with King's College London. R.A.M is

funded by the Michael Smith Foundation for Health Research (grant 17644). NG and DAL work in a Unit that receives support from the University of Bristol and the MRC (MC_UU_00011/6). K.S. is supported by the Biomedical Research Program at Weill Cornell Medicine in Qatar, a program funded by the Qatar Foundation, and by Qatar National Research Fund (QNRF) grant NPRP11C-0115-180010. The statements made herein are solely the responsibility of the authors. Bo Chawes has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No. 946228). M.K. is supported by the Wellcome Trust (221854/Z/20/Z), the UK Medical Research Council (MR/S011676/1), Academy of Finland (350426), and the National Institute on Aging (NIH), US (R01AG056477). JVL was supported by Academy of Finland (339568).

Institutional Review Board Statement: In accordance with the declaration of Helsinki ethical approval for each study was obtained by the ethical research boards pertaining to each study. TwinsUK: ethical approval was granted by the formerly known St. Thomas' Hospital Research Ethics Committee (REC). Following restructure and merging of REC, subsequent amendments were approved by the NRES Committee London—Westminster (REC ref: EC04/015, 1 November 2011); EPIC: The EPIC study, in particular the eight case-control studies nested within EPIC which provide data for this analysis, were approved by the ethics committee at the International Agency for Research on Cancer (IARC) (IEC 08-06, 10 April 2008), (liver cancer study: IEC 16-06, 11 February 2016), (breast cancer study: IEC 14-07, 7 April 2014), (colon cancer studies: IEC 14-08, 7 April 2014), (prostate cancer study: IEC 14-09, 7 April 2014), (kidney cancer study: IEC 15-06, 25 February 2015), (endometrial cancer study, IEC 16-20, 28 April 2016), (gallbladder cancer study: IEC 16-23, 28 June 2016); ALSPAC: Ethical approval was granted by the Bristol and Weston Health Authority (E1808, 28 November 1989), Southmead Health Authority (49/89, 5 April 1990), Frenchay Health Authority (90/8, 28 June 1990), Informed consent for the use of data collected via questionnaires and clinics was obtained from participants following the recommendations of the ALSPAC Ethics and Law Committee at the time and the local Research Ethics Committees. Consent for biological samples has been collected in accordance with the Human Tissue Act (2004); ARIC: Ethical approval was approved, and the protocol vetted by the institutional review board; BIB: Ethical approval was granted by the Bradford National Health Service Research Ethics Committee (ref 06/Q1202/48, 1 April 2008); Ethical approval granted by Gwent Research Ethics Committee (Ref JW/ST/01/69, 18 April 2002); HealthABC: AGES-Reykjavik was approved by the National Bioethics Committee in Iceland that acts as the Institutional Review Board for the Icelandic Heart Association (approval number: VSN-00-063), and by the National Institute on Aging Intramural Institutional Review Board; QBB: Ethical approval for the Qatar Biobank Study was granted by the Hamad Medical Corporation Ethics Committee; Whitehall II: Ethical approval for the Whitehall II study was obtained from the University College London Medical School committee on the ethics of human research.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The data used in this study are held by the department of Twin Research at King's College London. The data relating to TwinsUK can be released to bona fide researchers using our normal procedures overseen by the Wellcome Trust and its guidelines as part of our core funding (https://twinsuk.ac.uk/resources-for-researchers/access-our-data/, accessed on 8 June 2022). However, data relating to the Qatar Biobank, Avon Longitudinal Study of Parents and Children, Born in Bradford, Caerphilly Prospective Study, Whitehall II, Atherosclerosis Risk in Communities Study, Health, Aging and Body Composition Study, will require additional approval steps.

Acknowledgments: We thank all the participants and staff of the contributing cohorts for their time and effort and supporting our research. We also acknowledge Tamara Harris, and Eric Shiroma, of the HealthABC cohort for their support and facilitating NIA/NIH approval. The authors thank the staff and participants of the ARIC study for their important contributions. We are extremely grateful to all the families who took part in the ALSPAC study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses.

Conflicts of Interest: R.A.M. is a consultant for Pharmavite. All other authors declare no competing financial interests. Where authors are identified as personnel of the International Agency for Research on Cancer/World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy, or views of the International Agency for Research on Cancer/World Health Organization.

References

- Stergiou, G.S.; Palatini, P.; Parati, G.; O'Brien, E.; Januszewicz, A.; Lurbe, E.; Persu, A.; Mancia, G.; Kreutz, R.; on behalf of the European Society of Hypertension Council and the European Society of Hypertension Working Group on Blood Pressure Monitoring and Cardiovascular Variability; et al. 2021 European Society of Hypertension practice guidelines for office and out-of-office blood pressure measurement. *J. Hypertens.* 2021, *39*, 1293–1302. [CrossRef] [PubMed]
- 2. Zhou, B.; Perel, P.; Mensah, G.A.; Ezzati, M. Global epidemiology, health burden and effective interventions for elevated blood pressure and hypertension. *Nat. Rev. Cardiol.* **2021**, *18*, 785–802. [CrossRef] [PubMed]
- 3. Padmanabhan, S.; Dominiczak, A.F. Genomics of hypertension: The road to precision medicine. *Nat. Rev. Cardiol.* 2021, *18*, 235–250. [CrossRef] [PubMed]
- 4. Louca, P.; Menni, C.; Padmanabhan, S. Genomic Determinants of Hypertension With a Focus on Metabolomics and the Gut Microbiome. *Am. J. Hypertens.* 2020, *33*, 473–481. [CrossRef] [PubMed]
- Padmanabhan, S.; Caulfield, M.; Dominiczak, A.F. Genetic and Molecular Aspects of Hypertension. *Circ. Res.* 2015, 116, 937–959. [CrossRef]
- 6. Dzau, V.J.; Balatbat, C.A. Future of Hypertension. *Hypertension* 2019, 74, 450–457. [CrossRef]
- Menni, C.; Mangino, M.; Cecelja, M.; Psatha, M.; Brosnan, M.J.; Trimmer, J.; Mohney, R.P.; Chowienczyk, P.; Padmanabhan, S.; Spector, T.D.; et al. Metabolomic study of carotid-femoral pulse-wave velocity in women. *J. Hypertens.* 2015, 33, 791–796. [CrossRef]
- Menni, C.; Fauman, E.; Erte, I.; Perry, J.R.; Kastenmüller, G.; Shin, S.Y.; Petersen, A.K.; Hyde, C.; Psatha, M.; Ward, K.J.; et al. Biomarkers for type 2 diabetes and impaired fasting glucose using a nontargeted metabolomics approach. *Diabetes* 2013, 62, 4270–4276. [CrossRef]
- 9. Tang, W.H.; Wang, Z.; Levison, B.S.; Koeth, R.A.; Britt, E.B.; Fu, X.; Wu, Y.; Hazen, S.L. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *N. Engl. J. Med.* **2013**, *368*, 1575–1584. [CrossRef]
- Dietrich, S.; Floegel, A.; Weikert, C.; Prehn, C.; Adamski, J.; Pischon, T.; Boeing, H.; Drogan, D. Identification of Serum Metabolites Associated With Incident Hypertension in the European Prospective Investigation Into Cancer and Nutrition-Potsdam Study. *Hypertension* 2016, *68*, 471–477. [CrossRef]
- Menni, C.; Graham, D.; Kastenmüller, G.; Alharbi, N.H.; Alsanosi, S.M.; McBride, M.; Mangino, M.; Titcombe, P.; Shin, S.Y.; Psatha, M.; et al. Metabolomic identification of a novel pathway of blood pressure regulation involving hexadecanedioate. *Hypertension* 2015, *66*, 422–429. [CrossRef] [PubMed]
- 12. Hughes, G.S.; Mathur, R.S.; Margolius, H.S. Sex steroid hormones are altered in essential hypertension. *J. Hypertens.* **1989**, *7*, 181–187. [CrossRef] [PubMed]
- 13. Zheng, Y.; Yu, B.; Alexander, D.; Mosley, T.H.; Heiss, G.; Nettleton, J.A.; Boerwinkle, E. Metabolomics and Incident Hypertension Among Blacks. *Hypertension* **2013**, *62*, 398–403. [CrossRef] [PubMed]
- 14. Yu, B.; Li, A.H.; Metcalf, G.A.; Muzny, D.M.; Morrison, A.C.; White, S.; Mosley, T.H.; Gibbs, R.A.; Boerwinkle, E. Loss-of-function variants influence the human serum metabolome. *Sci. Adv.* **2016**, *2*, e1600800. [CrossRef] [PubMed]
- Zheng, Y.; Yu, B.; Alexander, D.; Manolio, T.A.; Aguilar, D.; Coresh, J.; Heiss, G.; Boerwinkle, E.; Nettleton, J.A. Associations between Metabolomic Compounds and Incident Heart Failure Among African Americans: The ARIC Study. *Am. J. Epidemiol.* 2013, 178, 534–542. [CrossRef]
- Yu, B.; Zanetti, K.A.; Temprosa, M.; Albanes, D.; Appel, N.; Barrera, C.B.; Ben-Shlomo, Y.; Boerwinkle, E.; Casas, J.P.; Clish, C.; et al. The Consortium of Metabolomics Studies (COMETS): Metabolomics in 47 Prospective Cohort Studies. *Am. J. Epidemiol.* 2019, *188*, 991–1012. [CrossRef]
- 17. Bujak, R.; Mateo, J.; Blanco, I.; Izquierdo-García, J.L.; Dudzik, D.; Markuszewski, M.J.; Peinado, V.I.; Laclaustra, M.; Barberá, J.A.; Barbas, C.; et al. New Biochemical Insights into the Mechanisms of Pulmonary Arterial Hypertension in Humans. *PLoS ONE* **2016**, *11*, e0160505. [CrossRef]
- 18. Pablo Arab, J.; Barrera, F.; Arrese, M. Bile Acids and Portal Hypertension. Ann. Hepatol. 2017, 16, S83–S86. [CrossRef]
- 19. Tominaga, T.; Suzuki, H.; Ogata, Y.; Imafuku, T.; Saruta, T. Bile acids are able to reduce blood pressure by attenuating the vascular reactivity in spontaneously hypertensive rats. *Life Sci.* **1988**, *42*, 1861–1868. [CrossRef]
- 20. Liu, L.; Su, J.; Li, R.; Luo, F. Changes in Intestinal Flora Structure and Metabolites Are Associated With Myocardial Fibrosis in Patients With Persistent Atrial Fibrillation. *Front. Nutr.* **2021**, *8*, 702085. [CrossRef]
- Wu, P.; Zhang, Y.; Liu, Y.; Wang, X.; Guo, Z.; Zhang, Y.; Liang, X.; Lai, W. Effects of cholic acid on blood pressure and production of vascular aldosterone and corticosterone. *Steroids* 1999, *64*, 291–295. [CrossRef]
- 22. Ferreira, M.; Coxito, P.M.; Sardão, V.A.; Palmeira, C.M.; Oliveira, P.J. Bile acids are toxic for isolated cardiac mitochondria. *Cardiovasc. Toxicol.* **2005**, *5*, 63–73. [CrossRef]
- Mels, C.M.; Delles, C.; Louw, R.; Schutte, A.E. Central systolic pressure and a nonessential amino acid metabolomics profile: The African Prospective study on the Early Detection and Identification of Cardiovascular disease and Hypertension. *J. Hypertens.* 2019, 37, 1157–1166. [CrossRef] [PubMed]

- Zhao, H.; Liu, Y.; Li, Z.; Song, Y.; Cai, X.; Liu, Y.; Zhang, T.; Yang, L.; Li, L.; Gao, S.; et al. Identification of essential hypertension biomarkers in human urine by non-targeted metabolomics based on UPLC-Q-TOF/MS. *Clin. Chim. Acta* 2018, 486, 192–198. [CrossRef] [PubMed]
- Flores-Guerrero, J.L.; Groothof, D.; Connelly, M.A.; Otvos, J.D.; Bakker, S.J.L.; Dullaart, R.P.F. Concentration of Branched-Chain Amino Acids Is a Strong Risk Marker for Incident Hypertension. *Hypertension* 2019, 74, 1428–1435. [CrossRef]
- Akbaraly, T.; Würtz, P.; Singh-Manoux, A.; Shipley, M.J.; Haapakoski, R.; Lehto, M.; Desrumaux, C.; Kähönen, M.; Lehtimäki, T.; Mikkilä, V.; et al. Association of circulating metabolites with healthy diet and risk of cardiovascular disease: Analysis of two cohort studies. *Sci. Rep.* 2018, *8*, 8620. [CrossRef]
- 27. Polak-Iwaniuk, A.; Harasim-Symbor, E.; Gołaszewska, K.; Chabowski, A. How Hypertension Affects Heart Metabolism. *Front. Physiol.* **2019**, *10*, 435. [CrossRef]
- Lu, Y.; Wang, G.; Hao, H.; Huang, Q.; Yan, B.; Zha, W.; Gu, S.; Ren, H.; Zhang, Y.; Fan, X.; et al. Gas chromatography/time-offlight mass spectrometry based metabonomic approach to differentiating hypertension- and age-related metabolic variation in spontaneously hypertensive rats. *Rapid Commun. Mass Spectrom.* 2008, 22, 2882–2888. [CrossRef]
- Whitworth, J.A.; Williamson, P.M.; Mangos, G.; Kelly, J.J. Cardiovascular consequences of cortisol excess. *Vasc. Health Risk Manag.* 2005, 1, 291–299. [CrossRef]
- 30. Whitworth, J.A.; Saines, D.; Scoggins, B.A. Blood pressure and metabolic effects of cortisol and deoxycorticosterone in man. *Clin. Exp. Hypertens. A* **1984**, *6*, 795–809. [CrossRef]
- Robinson, O.; Carter, A.R.; Ala-Korpela, M.; Casas, J.P.; Chaturvedi, N.; Engmann, J.; Howe, L.D.; Hughes, A.D.; Järvelin, M.-R.; Kähönen, M.; et al. Metabolic profiles of socio-economic position: A multi-cohort analysis. *Int. J. Epidemiol.* 2020, 50, 768–782. [CrossRef] [PubMed]
- 32. Al Kuwari, H.; Al Thani, A.; Al Marri, A.; Al Kaabi, A.; Abderrahim, H.; Afifi, N.; Qafoud, F.; Chan, Q.; Tzoulaki, I.; Downey, P.; et al. The Qatar Biobank: Background and methods. *BMC Public Health* **2015**, *15*, 1208. [CrossRef]
- Boyd, A.; Golding, J.; Macleod, J.; Lawlor, D.A.; Fraser, A.; Henderson, J.; Molloy, L.; Ness, A.; Ring, S.; Davey Smith, G. Cohort Profile: The 'Children of the 90s'—The index offspring of the Avon Longitudinal Study of Parents and Children. *Int. J. Epidemiol.* 2012, 42, 111–127. [CrossRef] [PubMed]
- Fraser, A.; Macdonald-Wallis, C.; Tilling, K.; Boyd, A.; Golding, J.; Davey Smith, G.; Henderson, J.; Macleod, J.; Molloy, L.; Ness, A.; et al. Cohort Profile: The Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *Int. J. Epidemiol.* 2012, 42, 97–110. [CrossRef] [PubMed]
- Northstone, K.; Lewcock, M.; Groom, A.; Boyd, A.; Macleod, J.; Timpson, N.; Wells, N. The Avon Longitudinal Study of Parents and Children (ALSPAC): An update on the enrolled sample of index children in 2019 [version 1; peer review: 2 approved]. Wellcome Open Res. 2019, 4, 51. [CrossRef] [PubMed]
- Harris, P.A.; Taylor, R.; Thielke, R.; Payne, J.; Gonzalez, N.; Conde, J.G. Research electronic data capture (REDCap)—A metadatadriven methodology and workflow process for providing translational research informatics support. *J. Biomed. Inform.* 2009, 42, 377–381. [CrossRef]
- Wright, J.; Small, N.; Raynor, P.; Tuffnell, D.; Bhopal, R.; Cameron, N.; Fairley, L.; Lawlor, D.A.; Parslow, R.; Petherick, E.S.; et al. Cohort Profile: The Born in Bradford multi-ethnic family cohort study. *Int. J. Epidemiol.* 2012, *42*, 978–991. [CrossRef]
- Garden, F.L.; Toelle, B.G.; Mihrshahi, S.; Webb, K.L.; Almqvist, C.; Tovey, E.R.; Brew, B.K.; Ayer, J.G.; Skilton, M.R.; Jones, G.; et al. Cohort profile: The Childhood Asthma Prevention Study (CAPS). *Int. J. Epidemiol.* 2018, 47, 1736–1736k. [CrossRef]
- Riboli, E.; Hunt, K.J.; Slimani, N.; Ferrari, P.; Norat, T.; Fahey, M.; Charrondière, U.R.; Hémon, B.; Casagrande, C.; Vignat, J.; et al. European Prospective Investigation into Cancer and Nutrition (EPIC): Study populations and data collection. *Public Health Nutr.* 2002, 5, 1113–1124. [CrossRef]
- 40. Verdi, S.; Abbasian, G.; Bowyer, R.C.E.; Lachance, G.; Yarand, D.; Christofidou, P.; Mangino, M.; Menni, C.; Bell, J.T.; Falchi, M.; et al. TwinsUK: The UK Adult Twin Registry Update. *Twin Res. Hum. Genet.* **2019**, *22*, 523–529. [CrossRef]
- 41. Marmot, M.; Brunner, E. Cohort Profile: The Whitehall II study. Int. J. Epidemiol. 2005, 34, 251–256. [CrossRef]
- Wright, J.D.; Folsom, A.R.; Coresh, J.; Sharrett, A.R.; Couper, D.; Wagenknecht, L.E.; Mosley, T.H.; Ballantyne, C.M.; Boerwinkle, E.A.; Rosamond, W.D.; et al. The ARIC (Atherosclerosis Risk In Communities) Study: JACC Focus Seminar 3/8. *J. Am. Coll. Cardiol.* 2021, 77, 2939–2959. [CrossRef] [PubMed]
- Santanasto, A.J.; Goodpaster, B.H.; Kritchevsky, S.B.; Miljkovic, I.; Satterfield, S.; Schwartz, A.V.; Cummings, S.R.; Boudreau, R.M.; Harris, T.B.; Newman, A.B. Body Composition Remodeling and Mortality: The Health Aging and Body Composition Study. J. Gerontol. Ser. A 2016, 72, 513–519. [CrossRef] [PubMed]
- Temprosa, M.; Moore, S.C.; Zanetti, K.A.; Appel, N.; Ruggieri, D.; Mazzilli, K.M.; Chen, K.L.; Kelly, R.S.; Lasky-Su, J.A.; Loftfield, E.; et al. COMETS Analytics: An online tool for analyzing and meta-analyzing metabolomics data in large research consortia. *Am. J. Epidemiol.* 2021, 191, 147–158. [CrossRef] [PubMed]
- 45. Wishart, D.S.; Feunang, Y.D.; Marcu, A.; Guo, A.C.; Liang, K.; Vázquez-Fresno, R.; Sajed, T.; Johnson, D.; Li, C.; Karu, N.; et al. HMDB 4.0: The human metabolome database for 2018. *Nucleic Acids Res.* **2017**, *46*, D608–D617. [CrossRef] [PubMed]
- 46. R Core Team. *R: A Language and Environment for Statistical Computing*, Version 3.6.3; R Foundation for Statistical Computing: Vienna, Austria, 2020.
- 47. Rust, P.; Ekmekcioglu, C. Impact of Salt Intake on the Pathogenesis and Treatment of Hypertension. *Adv. Exp. Med. Biol.* 2017, 956, 61–84. [CrossRef]

- Al-Maqbali, S.R.; Mula-Abed, W.A. Comparison between Three Different Equations for the Estimation of Glomerular Filtration Rate in Omani Patients with Type 2 Diabetes Mellitus. *Sultan Qaboos Univ. Med. J.* 2014, 14, e197–e203.
- Bliss, C.I.; Greenwood, M.L.; White, E.S. A Rankit Analysis of Paired Comparisons for Measuring the Effect of Sprays on Flavor. Biometrics 1956, 12, 381–403. [CrossRef]
- Krämer, A.; Green, J.; Pollard, J., Jr.; Tugendreich, S. Causal analysis approaches in Ingenuity Pathway Analysis. *Bioinformatics* 2013, 30, 523–530. [CrossRef]
- 51. Taylor, K.; McBride, N.; JGoulding, N.; Burrows, K.; Mason, D.; Pembrey, L.; Yang, T.; Azad, R.; Wright, J.; ALawlor, D. Metabolomics datasets in the Born in Bradford cohort [version 2; peer review: 1 approved, 1 approved with reservations]. Wellcome Open Res. 2021, 5, 264. [CrossRef]
- Goïta, Y.; Chao de la Barca, J.M.; Keïta, A.; Diarra, M.B.; Dembélé, K.C.; Chabrun, F.; Dramé, B.S.I.; Kassogué, Y.; Diakité, M.; Mirebeau-Prunier, D.; et al. Sexual Dimorphism of Metabolomic Profile in Arterial Hypertension. *Sci. Rep.* 2020, *10*, 7517. [CrossRef] [PubMed]
- 53. Li, L.; Zhong, S.; Hu, S.; Cheng, B.; Qiu, H.; Hu, Z. Changes of gut microbiome composition and metabolites associated with hypertensive heart failure rats. *BMC Microbiol.* **2021**, *21*, 141. [CrossRef] [PubMed]
- 54. Li, J.; Zhao, F.; Wang, Y.; Chen, J.; Tao, J.; Tian, G.; Wu, S.; Liu, W.; Cui, Q.; Geng, B.; et al. Gut microbiota dysbiosis contributes to the development of hypertension. *Microbiome* **2017**, *5*, 14. [CrossRef]
- Lu, J.; Tian, Y.; Gu, J.; Qiu, M.; Lu, Y.; Sun, W.; Kong, X. Comparative Study of Metabolite Changes after Antihypertensive Therapy with Calcium Channel Blockers or Angiotensin Type 1 Receptor Blockers. J. Cardiovasc. Pharmacol. 2021, 77, 228–237. [CrossRef]
- 56. Hao, Y.; Wang, Y.; Xi, L.; Li, G.; Zhao, F.; Qi, Y.; Liu, J.; Zhao, D. A Nested Case-Control Study of Association between Metabolome and Hypertension Risk. *BioMed Res. Int.* 2016, 2016, 7646979. [CrossRef]
- 57. Toba, H.; Nakamori, A.; Tanaka, Y.; Yukiya, R.; Tatsuoka, K.; Narutaki, M.; Tokitaka, M.; Hariu, H.; Kobara, M.; Nakata, T. Oral L-histidine exerts antihypertensive effects via central histamine H3 receptors and decreases nitric oxide content in the rostral ventrolateral medulla in spontaneously hypertensive rats. *Clin. Exp. Pharm. Physiol.* **2010**, *37*, 62–68. [CrossRef]
- Schwabedal, P.E.; Pietrzik, K.; Wittkowski, W. Pantothenic Acid Deficiency as a Factor Contributing to the Development of Hypertension. *Cardiology* 1985, 72 (Suppl. S1), 187–189. [CrossRef]
- 59. Ho, A.; Sinick, J.; Esko, T.; Fischer, K.; Menni, C.; Zierer, J.; Matey-Hernandez, M.; Fortney, K.; Morgen, E.K. Circulating glucuronic acid predicts healthspan and longevity in humans and mice. *Aging* **2019**, *11*, 7694–7706. [CrossRef]
- 60. Fregly, M.J.; Cade, J.R. Effect of Pyridoxine and Tryptophan, Alone and Combined, on the Development of Deoxycorticosterone Acetate-Induced Hypertension in Rats. *Pharmacology* **1995**, *50*, 298–306. [CrossRef]
- Dzúrik, R.; Fekovská, N.; Brimichová, G.; Tiso, P. Blood Pressure, 5-OH Indoleacetic Acid, and Vanilmandelic Acid Excretion and Blood Platelet Aggregation in Hypertensive Patients Treated with Ketanserin. J. Cardiovasc. Pharmacol. 1985, 7, 29–31. [CrossRef] [PubMed]
- 62. Alam, M.A. Anti-hypertensive Effect of Cereal Antioxidant Ferulic Acid and Its Mechanism of Action. *Front. Nutr.* **2019**, *6*, 121. [CrossRef] [PubMed]