



## King's Research Portal

DOI:

[10.1007/s00345-022-04171-4](https://doi.org/10.1007/s00345-022-04171-4)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Sharma, G., Shah, M., Ahluwalia, P., Dasgupta, P., Challacombe, B. J., Bhandari, M., Ahlawat, R., Rawal, S., Buffi, N. M., Sivaraman, A., Porter, J. R., Rogers, C., Motttrie, A., Abaza, R., Rha, K. H., Moon, D., Yuvaraja, T. B., Parekh, D. J., Capitanio, U., ... Gautam, G. (2022). Perioperative outcomes following robot-assisted partial nephrectomy in elderly patients. *World Journal of Urology*, 40(11), 2789-2798. <https://doi.org/10.1007/s00345-022-04171-4>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## **Perioperative outcomes following robot-assisted partial nephrectomy in elderly patients**

**Running title: Robot-assisted partial nephrectomy in elderly patients.**

### **Authors**

1. Gopal Sharma<sup>1</sup>, Vatikutti Fellow in Robotic Urologic Oncology Max Institute of Cancer Care, Saket, New Delhi, India. Email: [gopal.26669192@gmail.com](mailto:gopal.26669192@gmail.com).
2. Milap Shah<sup>1</sup>, Fellow in Robotic Urologic Oncology, Max Institute of Cancer Care, Saket, New Delhi, India. Email: [drmilapshah@gmail.com](mailto:drmilapshah@gmail.com)
3. Puneet Ahluwalia<sup>1</sup>, Principal Consultant, Department of Urologic Oncology, Max Institute of Cancer Care, Saket, New Delhi, India. Email: [puneet1923@gmail.com](mailto:puneet1923@gmail.com).
4. Prokar Dasgupta<sup>2</sup>, Faculty of Life Sciences and Medicine, King's College, London, King's Health Partners, UK, Email: [prokarurol@gmail.com](mailto:prokarurol@gmail.com).
5. Benjamin J Challacombe<sup>3</sup>, Consultant Urological Surgeon, Guy's and St. Thomas' NHS Foundation Trust, London, UK Email: [ben.challacombe@gstt.nhs.uk](mailto:ben.challacombe@gstt.nhs.uk)
6. Mahendra Bhandari<sup>4</sup>, Vattikuti foundation, Detroit, MI, USA, Email: [Mahendra@vattikutifoundation.com](mailto:Mahendra@vattikutifoundation.com).
7. Rajesh Ahlawat<sup>5</sup>, Medanta – The Medicity Hospital, New Delhi, India, Email: [rajesh.ahlawat@gmail.com](mailto:rajesh.ahlawat@gmail.com).
8. Sudhir Rawal<sup>6</sup>, Rajiv Gandhi Cancer Institute and Research Centre, New Delhi, India, Email: [sudirrawal85@gmail.com](mailto:sudirrawal85@gmail.com).
9. Nicolo M Buffi<sup>7</sup>, Humanitas Research Hospital, MI, Italy, Email: [buffi.nicolomaria@gmail.com](mailto:buffi.nicolomaria@gmail.com).
10. Ananthakrishanan Sivaraman<sup>8</sup>, Chennai Urology and Robotics Institute, , Chennai, India, Email: [ananthsiv@gmail.com](mailto:ananthsiv@gmail.com).
11. James R. Porter<sup>9</sup>, Swedish Medical Center, Seattle, WA, USA, Email: [porter@swedishurology.com](mailto:porter@swedishurology.com)
12. Craig Rogers<sup>10</sup>, Henry Ford Hospital, Detroit, MI, USA, Email: [crogers2@hfhs.org](mailto:crogers2@hfhs.org)
13. Alexandre Mottrie<sup>11</sup>, ORSI Academy, Melle, Belgium, Email: [a.mottrie@telenet.be](mailto:a.mottrie@telenet.be)
14. Ronney Abaza<sup>12</sup>, Central Ohio Urology Group and Mount Carmel Health System Prostate Cancer Program, Columbus, OH, USA,, Email: [ronneyabaza@hotmail.com](mailto:ronneyabaza@hotmail.com)

15. Khoon Ho Rha<sup>13</sup>, Yonsei University Health System, Seoul, South Korea, Email: khrha@yuhs.ac.kr
16. Daniel Moon<sup>14</sup>, Peter MacCallum Hospital, University of Melbourne, Royal Melbourne Clinical School, Melbourne, Australia, Email: drdanielmoon25@gmail.com.
17. Thyavihally B. Yuvaraja<sup>15</sup>, Kokilaben Dhirubhai Ambani Hospital, Mumbai, India, Email: yuvaraja.tb@relianceada.com.
18. Dipen J. Parekh<sup>16</sup>, University of Miami Health System, Miami, Florida, USA, Email: parekhd@med.miami.edu
19. Umberto Capitanio<sup>17</sup>, Unit of Urology, Division of Experimental Oncology, Urological Research Institute (URI), IRCCS Ospedale San Raffaele, Milan, Italy, Email: capitanio.umberto@hsr.it
20. Kris K. Maes<sup>18</sup>, Center for Robotic and Minimally Invasive Surgery, Hospital Da Luz, Luz Saúde, Portugal, Email: kris.k.maes@gmail.com.
21. Francesco Porpiglia<sup>19</sup>, San Luigi Gonzaga Hospital of Orbassano, Turin, Italy, Email: porpiglia@libero.it.
22. Levent Turkeri<sup>20</sup>, Acibadem M.A., Aydınlar University, Altuzinade Hospital, Department of Urology, Istanbul, Turkey, Email: turkeri@marmara.edu.tr.
23. Gagan Gautam<sup>1</sup>, Director, Department of Urologic Oncology, Max Institute of Cancer Care, Saket, New Delhi, India. Email: gagangg@gmail.com

Department(s) and institution(s) where the work was carried out:

<sup>1</sup>Department of Urologic Oncology, Max Institute of Cancer Care, New Delhi

<sup>2</sup>King's College, London, King's Health Partners, UK

<sup>3</sup>Guys and St Thomas' Hospital, London, UK.

<sup>4</sup>Vattikuti foundation, Detroit, MI, USA

<sup>5</sup>The Medicity Hospital, New Delhi, India

<sup>6</sup>Rajiv Gandhi Cancer Institute and Research Centre, New Delhi, India

<sup>7</sup>Humanitas Research Hospital, MI, Italy

<sup>8</sup>Chennai Urology and Robotics Institute, Chennai, India

<sup>9</sup>Swedish Medical Center, Seattle, WA, USA

<sup>10</sup>Henry Ford Hospital, Detroit, MI

<sup>11</sup>ORSI Academy, Melle, Belgium

<sup>12</sup> Central Ohio Urology Group and Mount Carmel Health System Prostate Cancer Program, Columbus, OH, USA, <sup>13</sup>Yonsei University Health System, Seoul, South Korea

<sup>14</sup>Peter MacCallum Cancer Centre, Royal Melbourne Clinical School, University of Melbourne Melbourne, Australia

<sup>15</sup>Kokilaben Dhirubhai Ambani Hospital, Mumbai, India

<sup>16</sup>University of Miami Health System, Miami, Florida, USA

<sup>17</sup>Urological Research Institute (URI), IRCCS Ospedale San Raffaele, Milan, Italy

<sup>18</sup>Center for Robotic and Minimally Invasive Surgery, Hospital Da Luz, Luz Saúde, Portugal

<sup>19</sup>San Luigi Gonzaga Hospital of Orbassano, Turin, Italy

<sup>20</sup>Acibadem M.A., Aydınlar University, Altuzinade Hospital, Department of Urology, Istanbul, Turkey

**Corresponding Author:**

**Dr. Gagan Gautam**

Director, Department of Urologic Oncology,  
Max Institute of Cancer Care, New Delhi, India.

Email: gagangg@gmail.com

Mobile number: (+91)9560966455

Total number of pages: 11

Total number of Tables/Graphs: 4

Total number of Images:0

Word counts

for abstract: 231

for the text: 1975

## **Abstract**

**Objective:** To compare perioperative outcomes following robot-assisted partial nephrectomy (RAPN) in patients with age  $\geq 70$  years to age  $< 70$  years.

**Methods:** Using Vattikuti Collective quality initiative (VCQI) database for RAPN we compared perioperative outcomes following RAPN between the two age groups. Primary outcome of the study was to compare trifecta outcomes between the two groups. Propensity matching using nearest neighbourhood method was performed with trifecta as primary outcome for sex, body mass index (BMI), solitary kidney, tumor size and Renal nephrometry score (RNS).

**Results:** Group A (age  $\geq 70$  years) included 461 patients whereas group B included 1932 patients. Before matching the two groups were statistically different for RNS and solitary kidney rates. After propensity matching, the two groups were comparable for baseline characteristics such as BMI, tumor size, clinical symptoms, tumor side, face of tumor, solitary kidney and tumor complexity. Among the perioperative outcome parameters there was no difference between two groups for operative time, blood loss, intraoperative transfusion, intraoperative complications, need for radical nephrectomy, positive margins and trifecta rates. Warm ischemia time was significantly longer in the younger age group (18.1 mins vs. 16.3 mins,  $p=0.003$ ). Perioperative complications were significantly higher in the older age group (11.8% vs. 7.7%,  $p=0.041$ ). However, there was no difference between the two groups for major complications.

**Conclusion:** RAPN in well-selected elderly patients is associated with comparable trifecta outcomes with acceptable perioperative morbidity.

**Keywords:** robotic; partial nephrectomy; elderly; propensity-matching.

## **Introduction**

Partial nephrectomy (PN) as a treatment option for small localized renal masses has become the standard of care<sup>1</sup>. Utilization rates of partial nephrectomy for renal masses have increased across all age groups<sup>2, 3</sup>. Superior functional and comparable oncological outcomes for PN compared to radical nephrectomy (RN) have translated into increased adoption of PN<sup>4, 5</sup>. In recent years, there has been a trend toward increased utilization of partial nephrectomy as a treatment option for managing localized small renal masses<sup>6, 7</sup>. However, its utilization in the elderly population remains poor<sup>2, 3</sup>. Reasons for this underutilization could be manifold. There is little doubt about the efficacy of partial nephrectomy in renal function preservation compared to RN<sup>8-10</sup>. However, elderly patients are at a significantly higher risk of death due to competing causes of mortality. Therefore, they may not extract long-term cardiovascular benefits from renal function

preservation. Thus, the lack of proven benefits in terms of overall and cancer-specific survival may be one of the deterrents for undergoing PN in the elderly population<sup>9-12</sup>.

Some studies have even reported superior overall survival due to a decrease in other causes of mortality such as cardiovascular events due to better renal function preservation with PN<sup>13, 14</sup>. However, this comes at the expense of increased perioperative complications associated with PN<sup>15-17</sup>. The apprehension of increased complications without any proven oncological benefit has led to a lower acceptance of PN for the elderly patient population. Population-based studies have shown lower rates of increased adoption of PN for elderly patients compared to their younger counterparts<sup>2, 3</sup>. Literature on perioperative outcomes following PN for the elderly population is limited. Furthermore, non-invasive treatment options as focal therapy has provided an alternative to PN in frail elderly patients with comorbidities<sup>1</sup>.

Few studies have reported outcomes following robot-assisted partial nephrectomy (RAPN) in the elderly population<sup>18-20</sup>. These studies have reported acceptable outcomes following RAPN in well-selected elderly patients. However, none of these studies have compared results with younger patients. Hence, with this Vattikuti collective quality initiative (VCQI) database study, we aimed to compare perioperative outcomes between patients aged  $\geq 70$  years and  $< 70$  years following RAPN.

## **Materials and Methods**

### **Vattikuti Collective Quality Initiative (VCQI) database**

VCQI is a prospective web-based multi-institutional collaborative database for various robotic surgical procedures<sup>21-24</sup>. Details of centers contributing to the database is provided in the supplementary file. Ethics clearance was obtained from each participating institution prior to data sharing. Due to the multi-institutional nature of the database, patients without adequate data had to be excluded from the study. For every patient, demographic, perioperative and postoperative data were collected as detailed in Table 1. Perioperative complications were graded as per Clavien-Dindo classification<sup>25</sup>. The primary objective of this study was the comparison of trifecta outcomes

between patients over the age of 70 years and those aged less than 70 years. Subgroup analysis of patients over the age of 80 years and those aged less than 80 years was also performed. Trifecta outcome was achieved without any complications, negative surgical margins, and warm ischemia time  $\leq 25$  minutes or zero ischemia<sup>24, 26</sup>.

### **Statistical analysis**

We checked the normality of continuous data using Kolmogorov-Smirnov and Shapiro tests of normality. An independent sample Student's t-test was used if data were normally distributed. Kruskal-Wallis test was used for non-normally distributed variables. For categorical variables, chi-square tests or Fisher's exact tests were used. Propensity matching using the nearest neighborhood method was performed with trifecta as the primary outcome for sex, BMI, solitary kidney, tumor size and RNS. All the statistical tests were two-sided and performed with a significance level of  $p < 0.05$ . The statistical tests were double-sided and conducted using SPSS version 23 (IBM corporation, New York, USA) and Stata (version 16; StataCorp, College Station, TX, USA). A p-value performed  $< 0.05$  was used to define significance.

### **Results**

From October 2014 to 2020, the participating centers contributed data of 3,801 patients who underwent RAPN. Of the 3,801 patients, 2,393 patients with complete data were included for the final analysis.

### **Prematching**

Group A (age  $\geq 70$  years) included 461 patients, whereas group B included 1932 patients. A comparison of two groups for baseline factors revealed that the two groups were comparable for sex, BMI and clinical tumor size. Mean age in A group was 75.1 years and 52.8 years ( $p=0.000$ ) in group B. There was no significant difference between the two groups for clinical symptoms, side of tumor, face of tumor, polar location of tumor and number of lesions operated. However, group A included a significantly higher number of patients with solitary kidneys (4.7% vs. 1.96%,  $p=0.000$ ). The mean RNS score was higher in group B patients (7.07 vs 6.88,  $p=0.03$ ). Group B



also had a significantly higher number of patients in the ‘high complexity’ stratification of RNS (10.4% vs. 6.3%). Preoperative hemoglobin (12.98 vs. 13.4,  $p=0.000$ ), and eGFR (68.4 vs. 83.5,  $p=0.000$ ) were significantly lower in group A, whereas creatinine was significantly higher in group A (1.08 vs. 0.94,  $p=0.000$ ) (Table 1).

In comparison of operative variables, two groups were comparable for surgical access (transperitoneal or retroperitoneal), duration of surgery, blood loss, positive margin intraoperative transfusion and intraoperative complications. Conversion to radical was significantly higher in the older age group (2.4% vs. 0.9%,  $p=0.010$ ) (Table 1). The postoperative complication rate was significantly higher in group older age group (12.5% vs. 7.8%,  $p=0.001$ ). However, the rate of major (grade III/IV) complications was similar in the two groups (2.6% vs. 2%,  $p=0.841$ ). Overall, major complications were noted in 52 of the patients. Among the patients with major complications, organ failure/ need for intensive care was required in 11 patients. Angioembolization was needed in 21 patients, 11 patients required Double J stenting for urine leak and reexploration was needed in 9 patients. The two groups showed no statistically significant difference for trifecta (71.5% vs. 71.1%,  $p=0.860$ ).

### **Postmatching**

Propensity matching was possible for 440 patients in either group. After propensity matching, the two groups were comparable for baseline characteristics such as BMI, tumor size, clinical symptoms, tumor side, face of tumor, solitary kidney and tumor complexity (Table 2). Among the perioperative outcome parameters there was no difference between the two groups for operative time, blood loss, intraoperative transfusion, intraoperative complications, need for radical nephrectomy, positive margins and trifecta rates. Warm ischemia time was significantly longer in the younger age group (18.1 mins vs. 16.3 mins,  $p=0.003$ ). Perioperative complications were significantly higher in the older age group (11.8% vs. 7.7%,  $p=0.041$ ). However, there was no difference between the two groups for major complications. On multivariate analysis, gender, tumor size and renal nephrometry score were identified as independent predictors of trifecta (Table 3). Standardized mean difference and variance ratios for the continuous covariates postmatching have been provided in the supplementary table.

### **Subgroup analysis**

Comparison of patients aged more than 80 years (n = 69) and less than 80 years (n= 2,324) showed that the two groups were comparable for certain baseline characteristics such as tumor size, sex, clinical symptoms, tumor side, face of tumor, tumor location, solitary kidney and tumor complexity (Table 4). There was no difference between the two groups for operative time, blood loss, intraoperative transfusion, intraoperative complications, need for radical nephrectomy, positive margins and trifecta rates. Warm ischemia time was significantly longer in the younger age group (18 mins vs. 14.2 mins,  $p=0.001$ ). Perioperative complications were significantly higher in the older age group (16% vs. 8.5%,  $p=0.031$ ). However, there was no difference between the two groups for major complications (2.9% vs. 2.1%,  $p=0.675$ ).

## Discussion

In the present study, before matching the two age groups were comparable in sex, BMI and clinical tumor size. However, the two groups differed significantly for Charlson comorbidity index, solitary kidney rates and renal nephrometry scores. There was no difference in the two groups for operative time, intraoperative complications, need for blood transfusion and blood loss. Mean WIT was significantly longer in the younger age group ( $18.3 \pm 9.26$  vs.  $16.2 \pm 8.7$ ,  $p=0.000$ ). Mean WIT remained longer in the younger age group even after matching. Furthermore, the conversion to radical nephrectomy was significantly higher in the elderly age group. However, there was no difference between the two groups for conversion to radical nephrectomy after matching (2% vs. 1.1%,  $p=0.281$ ). We noted significantly higher complications in group A (age  $\geq 70$  years) in the present study (before and after matching). However, this increased predilection was limited to the minor complications (grade 1 and 2), with rates of major complications being the same between the two groups. Similar results were noted when we compared for subgroup analysis for patients with age greater and lesser than 80 years. Literature is divided on the complication rates following PN compared to RN. Some studies have reported similar<sup>27-29</sup> and others have reported increased<sup>15, 17</sup> complications in patients undergoing PN compared to RN irrespective of the age group. A similar predicament related to complication rates for PN compared to RN is noted in studies reporting outcomes specifically in the elderly population. Two studies have reported (Lowrance et al<sup>30</sup> and Veccia et al<sup>12</sup>) significantly higher complication rates for elderly patients who underwent

PN compared to RN. However, An et al<sup>10</sup> and Antonelli et al<sup>8</sup> reported similar complications between RN and PN in their patient cohort.

Only a handful of studies have previously reported outcomes of PN in the elderly patient population<sup>12, 18-20</sup>. In their cohort of patients with a median age of 78 years, Ingels et al reported rates of blood transfusion, trifecta, intraoperative complications and major complications of 14.7%, 45%, 9% and 6.2%, respectively<sup>18</sup>. In contrast, patients above  $\geq 70$  years in the present study had much higher trifecta rates (71.5%) with lower perioperative morbidity. However, it is to be pointed out that in the study by Ingels et al different surgical modalities (open, laparoscopic and robotic) were employed and a robotic approach was predictive of lower complication rates<sup>18</sup>. In their cohort of elderly patients who underwent RAPN, Vartolomei et al reported perioperative outcomes similar to the present study<sup>20</sup>. Authors reported median operative time, blood loss, warm ischemia time and length of stay of 180 minutes, 100 ml, 14.5 minutes, and 5 days respectively. Positive surgical margins, overall complications and trifecta outcomes were reported in 1.9%, 15.4% and 71.2% respectively<sup>20</sup>. Similar results were reported by Bindayi et al. in their study for PN in their cohort of elderly patients<sup>19</sup>. Veccia et al compared RAPN to robotic RN in patients older than 65 years of age<sup>12</sup>. Authors reported positive surgical margin, overall complications, major complications and blood transfusion rates as 6%, 24%, 19% and 6%, respectively, in the RAPN group<sup>12</sup>. Superior rates of these perioperative outcomes were noted in the present study. Results of RAPN stated in the present study for the elderly group compare well for perioperative outcomes of the contemporary RAPN series<sup>31-34</sup>.

Our study is not without limitations. Firstly, due to the study's retrospective nature, the probability of a selection bias in patient inclusion is high. This is highlighted by the fact that elderly patients had lower complexity tumors in general, as compared to the matched cohort of younger patients. Propensity-matching between the two groups was performed to make two groups comparable for baseline factors. Furthermore, of the 3,801 patients, we included only 2,393 patients with complete data in this study. This could be one of the major limitations of this study. Secondly, the VCQI database also lacks surgeon experience or center caseload data. Lastly, Thirdly, there is heterogeneity in surgical techniques, learning curves, and perioperative management of patients due to the broader reach of the VCQI database. However, for precisely the same reasons, we believe

that our study is closer to the ‘real world scenario’ of the outcomes of RAPN in elderly patients and may provide unique insights regarding the same.

## **Conclusion**

Robot-assisted partial nephrectomy in well-selected elderly patients may be associated with comparable trifecta outcomes. However, the rates of overall perioperative complications were significantly higher in the elderly patient population.

## **DECLARATIONS**

### **Statements and Declarations**

#### **Authors contribution**

All the authors were involved in data acquisition and manuscript editing. G.S, M.S, P.A & G.G were involved in conception, design, analysis, manuscript writing and editing. G.G was involved in supervision of the project.

#### **Compliance with Ethical Standards**

Conflict of interests: Ronney Abaza is a speaker for Intuitive surgical, Conmed Inc and VTI. Benjamin J Challacombe, Kris K Maes, Rajesh Ahlawat and Gagan Gautam are proctors for Intuitive surgical. Other authors report no conflict of interests.

**Research involving Human Participants and/or Animals:** This study is retrospective analysis of VCQI database and ethics approval was obtained from all the participating centers prior to data collection.

**Informed consent:** Need for consent waived off by ethics committee.

**Data availability statement:** Corresponding author had full access to data and same can be provided on request to genuine authors.

**Acknowledgments:** Vattikuti Foundation for providing data. The authors would also like to thank Mrs. Tanvi Sharma (M Phil English) for English language editing

## **References**

1. Ljungberg B, Albiges L, Bedke J, Bex A, Capitanio U, Giles RH, et al. members of EAU guidelines on Renal Cell Carcinoma 2021 [accessed on 27.02.2022]. Available from: <https://uroweb.org/guideline/renal-cell-carcinoma/>.
2. Leppert JT, Mittakanti HR, Thomas IC, Lamberts RW, Sonn GA, Chung BI, et al. Contemporary Use of Partial Nephrectomy: Are Older Patients With Impaired Kidney Function Being Left Behind? *Urology*. 2017;100:65-71. Epub 2016/09/17.
3. Tan HJ, Daskivich TJ, Shirk JD, Filson CP, Litwin MS, Hu JC. Health status and use of partial nephrectomy in older adults with early-stage kidney cancer. *Urol Oncol*. 2017;35:153 e7-e14. Epub 2016/12/14.
4. Mir MC, Derweesh I, Porpiglia F, Zargar H, Mottrie A, Autorino R. Partial Nephrectomy Versus Radical Nephrectomy for Clinical T1b and T2 Renal Tumors: A Systematic Review and Meta-analysis of Comparative Studies. *Eur Urol*. 2017;71:606-17. Epub 2016/09/12.
5. Ljungberg B, Albiges L, Abu-Ghanem Y, Bensalah K, Dabestani S, Fernandez-Pello S, et al. European Association of Urology Guidelines on Renal Cell Carcinoma: The 2019 Update. *Eur Urol*. 2019;75:799-810. Epub 2019/02/26.
6. Woldrich JM, Palazzi K, Stroup SP, Sur RL, Parsons JK, Chang D, et al. Trends in the surgical management of localized renal masses: thermal ablation, partial and radical nephrectomy in the USA, 1998-2008. *BJU Int*. 2013;111:1261-8. Epub 2013/03/09.
7. Kim SP, Gross CP, Meropol N, Kutikov A, Smaldone MC, Shah ND, et al. National treatment trends among older patients with T1-localized renal cell carcinoma. *Urol Oncol*. 2017;35:113 e15- e21. Epub 2016/11/15.
8. Antonelli A, Veccia A, Pavan N, Mir C, Breda A, Takagi T, et al. Outcomes of Partial and Radical Nephrectomy in Octogenarians - A Multicenter International Study (Resurge). *Urology*. 2019;129:139-45. Epub 2019/03/28.
9. Chung JS, Son NH, Lee SE, Hong SK, Lee SC, Kwak C, et al. Overall survival and renal function after partial and radical nephrectomy among older patients with localised renal cell carcinoma: a propensity-matched multicentre study. *Eur J Cancer*. 2015;51:489-97. Epub 2015/01/13.
10. An JY, Ball MW, Gorin MA, Hong JJ, Johnson MH, Pavlovich CP, et al. Partial vs Radical Nephrectomy for T1-T2 Renal Masses in the Elderly: Comparison of Complications, Renal Function, and Oncologic Outcomes. *Urology*. 2017;100:151-7. Epub 2016/11/28.

11. Ristau BT, Handorf EA, Cahn DB, Kutikov A, Uzzo RG, Smaldone MC. Partial nephrectomy is not associated with an overall survival advantage over radical nephrectomy in elderly patients with stage Ib-II renal masses: An analysis of the national cancer data base. *Cancer*. 2018;124:3839-48. Epub 2018/09/13.
12. Veccia A, Dell'oglio P, Antonelli A, Minervini A, Simone G, Challacombe B, et al. Robotic partial nephrectomy versus radical nephrectomy in elderly patients with large renal masses. *Minerva Urol Nefrol*. 2020;72:99-108. Epub 2019/09/19.
13. Sun M, Trinh QD, Bianchi M, Hansen J, Hanna N, Abdollah F, et al. A non-cancer-related survival benefit is associated with partial nephrectomy. *Eur Urol*. 2012;61:725-31. Epub 2011/12/17.
14. Ficarra V, Bhayani S, Porter J, Buffi N, Lee R, Cestari A, et al. Predictors of warm ischemia time and perioperative complications in a multicenter, international series of robot-assisted partial nephrectomy. *Eur Urol*. 2012;61:395-402. Epub 2011/11/15.
15. Van Poppel H, Da Pozzo L, Albrecht W, Matveev V, Bono A, Borkowski A, et al. A prospective, randomised EORTC intergroup phase 3 study comparing the oncologic outcome of elective nephron-sparing surgery and radical nephrectomy for low-stage renal cell carcinoma. *Eur Urol*. 2011;59:543-52. Epub 2010/12/28.
16. Van Poppel H, Da Pozzo L, Albrecht W, Matveev V, Bono A, Borkowski A, et al. A prospective randomized EORTC intergroup phase 3 study comparing the complications of elective nephron-sparing surgery and radical nephrectomy for low-stage renal cell carcinoma. *Eur Urol*. 2007;51:1606-15. Epub 2006/12/05.
17. Lesage K, Joniau S, Fransis K, Van Poppel H. Comparison between open partial and radical nephrectomy for renal tumours: perioperative outcome and health-related quality of life. *Eur Urol*. 2007;51:614-20. Epub 2006/11/14.
18. Ingels A, Duc S, Bensalah K, Bigot P, Paparel P, Beauval JB, et al. Postoperative outcomes of elderly patients undergoing partial nephrectomy. *Sci Rep*. 2021;11:17201. Epub 2021/08/27.
19. Bindayi A, Autorino R, Capitanio U, Pavan N, Mir MC, Antonelli A, et al. Trifecta Outcomes of Partial Nephrectomy in Patients Over 75 Years Old: Analysis of the RENal SURGery in Elderly (RESURGE) Group. *Eur Urol Focus*. 2020;6:982-90. Epub 2019/02/26.

20. Vartolomei MD, Matei DV, Renne G, Tringali VM, Crisan N, Musi G, et al. Long-term oncologic and functional outcomes after robot-assisted partial nephrectomy in elderly patients. *Minerva Urol Nefrol.* 2019;71:31-7. Epub 2018/09/20.
21. Arora S, Abaza R, Adshead JM, Ahlawat RK, Challacombe BJ, Dasgupta P, et al. 'Trifecta' outcomes of robot-assisted partial nephrectomy in solitary kidney: a Vattikuti Collective Quality Initiative (VCQI) database analysis. *BJU Int.* 2018;121:119-23. Epub 2017/07/28.
22. Arora S, Bronkema C, Porter JR, Mottrie A, Dasgupta P, Challacombe B, et al. Omission of Cortical Renorrhaphy During Robotic Partial Nephrectomy: A Vattikuti Collective Quality Initiative Database Analysis. *Urology.* 2020;146:125-32. Epub 2020/09/18.
23. Arora S, Heulitt G, Menon M, Jeong W, Ahlawat RK, Capitanio U, et al. Retroperitoneal vs Transperitoneal Robot-assisted Partial Nephrectomy: Comparison in a Multi-institutional Setting. *Urology.* 2018;120:131-7. Epub 2018/07/28.
24. Sharma G, Shah M, Ahluwalia P, Dasgupta P, Challacombe BJ, Bhandari M, et al. Comparison of perioperative outcomes following transperitoneal versus retroperitoneal robot-assisted partial nephrectomy: a propensity-matched analysis of VCQI database. *World J Urol.* 2022. Epub 2022/07/23. eng.
25. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg.* 2004;240:205-13. Epub 2004/07/27.
26. Khalifeh A, Autorino R, Hillyer SP, Laydner H, Eyraud R, Panumatrassamee K, et al. Comparative outcomes and assessment of trifecta in 500 robotic and laparoscopic partial nephrectomy cases: a single surgeon experience. *J Urol.* 2013;189:1236-42. Epub 2012/10/20.
27. Lau WK, Blute ML, Weaver AL, Torres VE, Zincke H. Matched comparison of radical nephrectomy vs nephron-sparing surgery in patients with unilateral renal cell carcinoma and a normal contralateral kidney. *Mayo Clin Proc.* 2000;75:1236-42. Epub 2000/12/29.
28. Corman JM, Penson DF, Hur K, Khuri SF, Daley J, Henderson W, et al. Comparison of complications after radical and partial nephrectomy: results from the National Veterans Administration Surgical Quality Improvement Program. *BJU Int.* 2000;86:782-9. Epub 2000/11/09.
29. Stephenson AJ, Hakimi AA, Snyder ME, Russo P. Complications of radical and partial nephrectomy in a large contemporary cohort. *J Urol.* 2004;171:130-4. Epub 2003/12/11.

30. Lowrance WT, Yee DS, Savage C, Cronin AM, O'Brien MF, Donat SM, et al. Complications after radical and partial nephrectomy as a function of age. *J Urol*. 2010;183:1725-30. Epub 2010/03/20.
31. Simhan J, Smaldone MC, Tsai KJ, Li T, Reyes JM, Canter D, et al. Perioperative outcomes of robotic and open partial nephrectomy for moderately and highly complex renal lesions. *J Urol*. 2012;187:2000-4. Epub 2012/04/14.
32. Dulabon LM, Kaouk JH, Haber GP, Berkman DS, Rogers CG, Petros F, et al. Multi-institutional analysis of robotic partial nephrectomy for hilar versus nonhilar lesions in 446 consecutive cases. *Eur Urol*. 2011;59:325-30. Epub 2010/12/15.
33. Scoll BJ, Uzzo RG, Chen DY, Boorjian SA, Kutikov A, Manley BJ, et al. Robot-assisted partial nephrectomy: a large single-institutional experience. *Urology*. 2010;75:1328-34. Epub 2010/01/19.
34. Patel MN, Krane LS, Bhandari A, Laungani RG, Shrivastava A, Siddiqui SA, et al. Robotic partial nephrectomy for renal tumors larger than 4 cm. *Eur Urol*. 2010;57:310-6. Epub 2009/12/01.

## Table legends

**Table 1:** Comparison of baseline characteristics between the two age groups (Age  $\geq 70$  years and  $< 70$  years)

**Table 2:** Comparison of perioperative outcomes between the two age groups post matching (Age  $\geq 70$  years and  $< 70$  years).

**Table 3:** Multivariate analysis for the predictors of the trifecta outcomes.