



King's Research Portal

Link to publication record in King's Research Portal

Citation for published version (APA): Chen, Y., Murthy, S., & Turiaci, G. J. (2024). *Gravitational index of the heterotic string.*

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Gravitational index of the heterotic string

Yiming Chen¹, Sameer Murthy^{2,3}, Gustavo J. Turiaci⁴

¹ Stanford Institute for Theoretical Physics, Stanford University, Stanford, CA 94305, USA

² Department of Mathematics, King's College London, The Strand, London WC2R 2LS, UK

³ School of Natural Sciences, Institute for Advanced Study, Princeton, NJ, USA

⁴ Physics Department, University of Washington, Seattle, WA, USA

ABSTRACT: The fundamental heterotic string has a tower of BPS states whose supersymmetric index has an exponential growth in the charges. We construct the saddle-point of the gravitational path integral corresponding to this index. The saddle-point configuration is a supersymmetric rotating non-extremal Euclidean black hole. This configuration is singular in the two-derivative theory. We show that the addition of higher-derivative terms in four-dimensional $\mathcal{N} = 2$ supergravity resolves the singularity. In doing so, we extend the recently-developed "new attractor mechanism" to include the effect of higher-derivative terms. Remarkably, the one-loop, four-derivative F-term contribution to the prepotential leads to a precise match of the gravitational and microscopic index. We also comment, using the effective theory near the horizon, on the possibility of a string-size near-extremal black hole. Our results clarify the meaning of different descriptions of this system in the literature. The thermal state transitions to a winding condensate and a gas of strings without ever reaching a small black hole, while the index is captured by the rotating Euclidean black hole solution and is constant and thus smoothly connected to the microscopic ensemble.

Contents

1	Introduction		
2	 The two-charge black hole and its transition to free strings 2.1 Review of the two-charge black hole 2.2 Transition to free strings via the charged Horowitz-Polchinski solution 	4 5 10	
3	 Gravitational configurations that contribute to the index 3.1 Imposing supersymmetry 3.2 Explicit form of the solution 3.3 Summary of properties 	13 14 16 18	
4	The index and new attractor mechanism in 4d $\mathcal{N} = 2$ supergravity 4.1 $\frac{1}{2}$ -BPS solutions of $\mathcal{N} = 2$ supergravity 4.2 Euclidean black holes and near-horizon behavior 4.3 Two-charge solution	19 20 21 24	
5	Higher derivative corrections to the gravitational index	28	
6	Can the low-temperature physics be nearly conformal?	33	
7	Discussion	36	
A	Solution generation for rotating two-charge systems A.1 Generating α' corrections in various cases	37 40	

1 Introduction

Consider a fundamental heterotic string of energy M, with w units of winding and n units of momentum around a circle of finite size. In the weakly-coupled theory the string has a tower of excited states whose degeneracy grows exponentially in the excitation numbers. In the strongly-coupled theory, the effective description of the string is as a source for the metric and other low-energy fields of supergravity, and the solution of the low-energy theory is generically a black hole in the (d+1)-dimensional non-compact spacetime whose entropy corresponds to the exponentially growing degeneracy of states [1–3].

When all the right-movers are in their ground states, the heterotic string configuration preserves half the supersymmetry of the theory and obeys $M^2 = Q_R^2/2$ where Q_R is the right-moving momentum. The resulting tower of $\frac{1}{2}$ -BPS Dabholkar-Harvey states has a degeneracy which grows as $\exp(4\pi\sqrt{nw})$ [4]. What is the strongly-coupled description of this ensemble of supersymmetric states? One natural possibility is that they form a supersymmetric black hole. Such a black hole would be extremal with $AdS_2 \times S^{d-1}$ nearhorizon geometry. In this paper we mainly discuss four non-compact directions (d = 3), although we make a few comments on other dimensions as well.

As is well-known, this simple picture runs into various problems.¹ The zeroth-order problem, which was already discussed in the early days, is that the two-derivative supergravity solutions with a fundamental string source is singular, in that the horizon lies on top of the curvature singularity [6]. A more sophisticated analysis suggested that including higher-derivative terms in the low-energy supergravity stretches the horizon to string scale, thus resolving the singularity [7], [8–10].

However, there are still problems with this picture. Firstly, if we think of obtaining the extremal black hole solution by lowering the mass to its extremal value, there is good evidence that there is a transition at low temperatures to the so-called Horowitz-Polchinski solution [11] instead of obtaining an extremal black hole [12].² Secondly, there is no known $AdS_2 \times S^{d-1}$ solution in supergravity which admits 16 Killing spinors. This could be an issue of simply not having found such a solution but, in fact, we can sharpen this argument by analyzing possible superconformal groups in the near-horizon region. As we explain in Section 6, by studying the quantum super Schwarzian theory near the horizon, one can rule out the existence of a decoupled near-horizon AdS_2 region with 16 or more supercharges describing the heterotic string with all its symmetries.

In order to clarify the situation, it is useful to focus on particular observables that are amenable to precise calculation. In particular, the supersymmetric index [14] is independent of coupling, and can be precisely compared in the two descriptions. For the $\frac{1}{2}$ -BPS states that we are considering, the relevant index is a helicity supertrace in flat spacetime (see e.g. [15]). This quantity is protected against changes of coupling, for the usual reason of bosonic and fermionic supermultiplets appearing in pairs. In the free theory, the generating function of this index is the chiral partition function of 24 free bosonic fields i.e. $1/\eta(\tau)^{24}$ [4], whose coefficient at level nw grows as $\exp(4\pi\sqrt{nw})$ as mentioned above. The question we would like to address is: what is the saddle-point of the dual gravitational path integral of this index?

The aim of this paper is to present and analyze this gravitational saddle-point. Here, by saddle-point we mean a Euclidean solution of the low-energy supergravity, treating G_N in perturbation theory. The main subtlety—and difference from previous treatments—is the inclusion of $(-1)^F$ in the gravitational functional integral. We use the construction of the saddle-point for the gravitational index discussed in [16, 17]. The basic idea is that $(-1)^F$ can be included in the gravitational functional integral by adding an imaginary part to a background bosonic field that couples to an *R*-symmetry. This implies that the solution is necessarily complex. The Euclidean solution is non-extremal and has the topology of a cigar times S^{d-1} , with the horizon at the tip of the cigar.

In asymptotic flat space, the relevant background field that implements $(-1)^{\mathsf{F}}$ is the

¹Notably, this picture does hold in three- and four-charge systems, and leads to the famous agreement of statistical and thermodynamic entropy of black holes in string theory [5].

 $^{^{2}}$ See also [13] that studies the validity of the small black hole analysis.

angular velocity Ω which obeys $\beta\Omega = 2\pi i$. The solution has non-zero rotation along one axis, which defines a north and south pole at the horizon. Applying this idea to fourdimensional $\mathcal{N} = 2$ supergravity coupled to an arbitrary number of vector multiplets leads to the *new attractor mechanism* [18]. The metric of these solutions can be written in the Israel-Wilson-Perjés (IWP) form [19, 20], which is sourced by harmonic functions with two simple poles at the north and south poles, respectively. Near the poles, the values of the scalars of supergravity get fixed in terms of the charges of the black hole. The complex nature of the solution is reflected in that the holomorphic scalars are fixed at the north pole and vanish at the south pole. (The anti-holomorphic scalar has the opposite behavior.)

In this paper we construct the rotating Euclidean black hole as a solution to the effective four-dimensional supergravity describing the heterotic string theory on T^6 . We also write the solution in the variables of the ten-dimensional heterotic theory. As it turns out, the solution is singular at the two-derivative level. In the Einstein frame in four dimensions, the metric is smooth but the string coupling goes to zero and so the string-frame metric is singular. This opens the possibility that higher-derivative effects can resolve this singularity.

In order to resolve this singularity, we study the effect of higher-derivative corrections in the supergravity action on the Euclidean rotating black hole solution. We discuss a class of higher-derivative corrections that are summarized in the holomorphic prepotential of four-dimensional $\mathcal{N} = 2$ supergravity, and can be calculated using topological string methods in the dual Type II theory. In particular, the one-loop term in the prepotential leads to the supersymmetric completion of a particular four-derivative term proportional to the square of the Weyl tensor. This leads to an extension of the new attractor mechanism to include higher-derivative terms.

In the two-charge system, we find that the singularity is resolved in the higherderivative theory and, in particular, the string coupling at the tip is now non-zero. Note that the resolution involves a play-off between 2-derivative and 4-derivative effects. This would lead to an uncontrollable approximation for generic observables. However, the calculation for supersymmetric quantities such as the index has a better chance of leading to a precise result, along the lines of the non-renormalization theorem discussed in [21, 22]. In particular, if *D*-terms vanish on our supersymmetric configurations, the fact that the topological string expansion for Type II on $K3 \times T^2$ is one-loop exact implies that our result would be rigorous. Quite remarkably we find that, with these assumptions, the gravitational free-energy of the index agrees precisely with that of the microscopic index to give the entropy $4\pi\sqrt{nw}$ including the numerical coefficient³.

Now we briefly summarize the overall picture that we propose. First consider the thermal partition function and states in the Lorentzian theory. Away from extremality the fundamental heterotic string is described by the two-charge black hole solution with

³To evaluate the action we assume it is temperature independent and compute it in the large β limit. At two derivative level we have proved this earlier [18]. The calculation of the on-shell action (the Gibbons-Hawking free energy of the system) at finite β including the effect of 4-derivative terms is complicated and we postpone it to future work.

a macroscopic horizon. As we lower the mass towards extremality there is a transition to a gas of free strings, without ever reaching the extremal black hole solution [23]. Now consider the gravitational path integral for the supersymmetric index as in the present paper. The saddle-point is a supersymmetric non-extremal Euclidean geometry valid at all temperatures. The saddle-point is really like an instanton rather than describing Lorentzian states. At the two-derivative level the solution has a large horizon area, but contains curvature singularities near the poles of the horizon. These singularities are resolved by the inclusion of string-scale higher-derivative corrections in the effective action coming from string theory. The index calculated in this manner is independent of coupling, and agrees with the index of a gas of strings at weak-coupling.⁴ In this respect, the index behaves in a manner similar to the supersymmetric indices corresponding to big black holes⁵ [16–18]. They do not depend on coupling, and have continuously connected dual pictures in the gravitational and microscopic regimes, rather than a sharp transition.

The plan of paper is as follows. In Section 2 we review the family of two-charge black hole solutions in heterotic string theory, their thermodynamics, and the transition to the gas of free strings via the Horowitz-Polchinski solution. In Section 3 we discuss the gravitational saddle-point for the index, i.e. the Euclidean rotating black hole, in the heterotic string variables. In Section 4 we switch to the four-dimensional effective supergravity description at the two-derivative level. We review the new attractor mechanism and the embedding of the Euclidean rotating black hole in this description. In Section 5 we include a fourderivative term in the 4d supergravity coming from the string compactification. We show how it desingularizes the Euclidean rotating black hole, and analyze its effect on the entropy. In Section 6 we analyze the superconformal groups that can possible arise in the nearhorizon geometry and the quantum effects of the respective super-Schwarzians. We show that AdS₂ cannot arise at the quantum level in the two-charge system. In Section 7 we discuss further possible directions. In the appendix we discuss how the solution generating technique [24, 25] relates the rotating charged and uncharged solutions.

Note added While this paper was in preparation we received [26], which contains overlap on the two-derivative solutions that calculate the index.

2 The two-charge black hole and its transition to free strings

In this section we review the properties of the two-charge black hole in the heterotic string theory on T^6 . In particular, we discuss some peculiar features of the naive extremal limit of this solution. Then we review the construction of the charged Horowitz-Polchinski solution and how it modifies the approach to extremality.

⁴There may even exist a Horowitz-Polchinski-type supersymmetric solution at intermediate coupling. The details of this would be interesting and may involve a perhaps suitably complexified sigma-model. We comment further on this possibility in Section 3.1.

⁵Big black holes refer to black holes with with horizon curvatures parameterically smaller than the string and Planck scale.

2.1 Review of the two-charge black hole

We begin by reviewing the two-charge black hole, constructed by Sen in [27], as a solution to the four dimensional supergravity theory arising from a toroidal compactification of heterotic string theory.

2.1.1 Heterotic string on T^6

The four dimensional theory has $\mathcal{N} = 4$ supersymmetry and contains the supergravity multiplet, and 22 $\mathcal{N} = 4$ vector multiplets. The supergravity multiplet has six graviphotons, resulting in a total gauge group U(1)²⁸. We collectively denote the potentials by A^a with $a = 1, \ldots, 28$. The moduli space consists of the dilaton Φ and 132 scalar fields organized in a symmetric 28×28 dimensional matrix M that satisfies $MLM^T = L$. L is a diagonal matrix $L_{aa} = -1$ for $a = 1, \ldots, 22$ and $L_{aa} = +1$ for $a = 23, \ldots, 28$. The scalars M encode deformations of the torus, of the background B-field, and of the Wilson lines. Besides these fields, we also have a four dimensional B-field $B_{\mu\nu}$, which is dual to a scalar.

The action, in Lorentzian signature, is given by

$$I = \frac{1}{16\pi} \int d^4x \sqrt{-g} e^{-\Phi} \left[R + (\partial \Phi)^2 + \frac{1}{8} \text{Tr} \left[\partial_{\mu} M L \partial^{\mu} M L \right] - \frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} - F^a_{\mu\nu} (LML)_{ab} F^{b\mu\nu} \right] + (\text{fermions}) + (\text{bdy terms}), \quad (2.1)$$

where we denote the U(1)²⁸ field strength by $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a$ and the *B*-field field strength by $H_{\mu\nu\rho} = \partial_\mu B_{\nu\rho} + 2A_\mu^a L_{ab} F_{\nu\rho}^b + (\text{cyclic})$. The action is invariant under an $O(6, 22, \mathbb{R})$ transformation which acts on a, b indices only, so it does not act on the dilaton, metric or *B* field. Consistency with quantization of electromagnetic charge restricts it to $O(6, 22; \mathbb{Z})$.

The solutions we discuss in the following have boundary conditions at infinity such that locally

$$M_{ab}|_{\infty} = \delta_{ab}, \qquad \Phi|_{\infty} = 0, \qquad g_{\mu\nu}|_{\infty} = \eta_{\mu\nu}.$$
 (2.2)

For the gauge fields, both the electromagnetic ones and the B-field, we can choose to fix their holonomies or their charges at infinity.

2.1.2 Sen's black hole solution

The procedure to find the electrically charged black holes described in [27] is the following. First, one realizes that for metrics with a U(1) isometry (corresponding to Euclidean time), the symmetry of the action gets enhanced to O(7, 23). One can think about this in terms of a $T^7 = T^6 \times S^1$ compactification of the heterotic string, but [27] verifies it by explicit construction of the O(7, 23) action. We discuss such transformations more generally in Appendix A, not necessarily restricting our attention to black hole solutions. Let us present the final results. The metric in the Einstein frame $g^E_{\mu\nu} = e^{-\Phi}g_{\mu\nu}$ and Lorentzian signature is given by

$$e^{-\Phi} ds^{2} = \Delta^{1/2} \left[-\frac{r^{2} + a^{2} \cos^{2} \theta - 2mr}{\Delta} dt^{2} + \frac{dr^{2}}{r^{2} + a^{2} - 2mr} + d\theta^{2} + \frac{\sin^{2} \theta}{\Delta} [\Delta + a^{2} \sin^{2} \theta (r^{2} + a^{2} \cos^{2} \theta + 2mr \cosh \alpha \cosh \beta)] d\phi^{2} - \frac{2mra \sin^{2} \theta (\cosh \alpha + \cosh \beta)}{\Delta} dt d\phi \right], \qquad (2.3)$$

where we define the function

 $\Delta = (r^2 + a^2 \cos^2 \theta)^2 + 2mr(r^2 + a^2 \cos^2 \theta)(\cosh \alpha \cosh \beta - 1) + m^2 r^2 (\cosh \alpha - \cosh \beta)^2.$ (2.4)

The dilaton is given by

$$\Phi = \frac{1}{2} \log \frac{(r^2 + a^2 \cos^2 \theta)^2}{\Delta}.$$
 (2.5)

We can decompose the vector fields into a 22-dimensional vector of one-forms \vec{A}_L , made up of the first 22 components of A, and a 6-dimensional vector of one-forms \vec{A}_R made of the last 6 components of A. These gauge potentials and the two-form are given by

$$\vec{A}_{L} = -\frac{\vec{n}}{\sqrt{2}} \frac{mr \sinh \alpha ((r^{2} + a^{2} \cos^{2} \theta) \cosh \beta + mr (\cosh \alpha - \cosh \beta))}{\Delta} dt + \frac{\vec{n}}{\sqrt{2}} \frac{mra \sinh \alpha \sin^{2} \theta (r^{2} + a^{2} \cos^{2} \theta + mr \cosh \beta (\cosh \alpha - \cosh \beta))}{\Delta} d\phi, \quad (2.6)$$
$$\vec{A}_{R} = -\frac{\vec{p}}{\sqrt{2}} \frac{mr \sinh \beta ((r^{2} + a^{2} \cos^{2} \theta) \cosh \alpha + mr (\cosh \beta - \cosh \alpha))}{\Delta} dt + \frac{\vec{p}}{\sqrt{2}} \frac{mra \sinh \beta \sin^{2} \theta (r^{2} + a^{2} \cos^{2} \theta + mr \cosh \alpha (\cosh \beta - \cosh \alpha))}{\Delta} d\phi, \quad (2.7)$$
$$B = \frac{mra \sin^{2} \theta (\cosh \alpha - \cosh \beta)}{\Delta} (r^{2} + a^{2} \cos^{2} \theta + mr (\cosh \alpha \cosh \beta - 1)) dt d\phi. (2.8)$$

The scalar moduli are given by

$$M = I_{28} + \begin{pmatrix} \mathsf{P}nn^T \ \mathsf{Q}np^T \\ \mathsf{Q}pn^T \ \mathsf{P}pp^T \end{pmatrix}, \qquad (2.9)$$

where

$$\mathsf{P} = \frac{2m^2 r^2 \sinh^2 \alpha \sinh^2 \beta}{\Delta},\tag{2.10}$$

$$\mathsf{Q} = -\frac{2mr\sinh\alpha\sinh\beta}{\Delta}(r^2 + a^2\cos^2\theta + mr(\cosh\alpha\cosh\beta - 1)).$$
(2.11)

Here \vec{n} and \vec{p} are arbitrary 22 and 6 dimensional unit vectors. It will be important to know the charges and chemical potentials as functions of the parameters appearing in the solution, which are $(m, a, \alpha, \beta, \vec{n}, \vec{p})$. Define \vec{Q}_L to be a 22-dimensional vector made out of the first 22 components of Q^a . Similarly define \vec{Q}_R to be the last 6 components. To compute the charges we can evaluate the large r limit of the gauge potentials and read them off from $A \sim Q/r \, dt$. Using that $\Delta \sim r^4$ as $r \to \infty$ we obtain

$$\vec{Q}_L = \frac{m}{\sqrt{2}} \sinh \alpha \cosh \beta \, \vec{n} \,, \qquad \vec{Q}_R = \frac{m}{\sqrt{2}} \sinh \beta \cosh \alpha \, \vec{p} \,.$$
 (2.12)

We will also use Q_L, Q_R to denote the modulus of the respective charge vectors. Since the black holes are electrically charged and rotating there will also be a non-trivial magnetic dipole moment, but it will not be important for the discussion here.

Let us now move on to the other charges encoded in the metric. The mass and angular momentum of the black hole can be read off from the large r behavior of g_{tt} and $g_{t\phi}$ and are given by

$$M = \frac{m(1 + \cosh \alpha \cosh \beta)}{2}, \qquad J = \frac{ma(\cosh \alpha + \cosh \beta)}{2}. \tag{2.13}$$

The metric has an outer horizon r_+ at the locations where g_{rr}^{-1} vanishes

$$r^2 - 2mr + a^2 = 0 \quad \Rightarrow \quad r_{\pm} = m \pm \sqrt{m^2 - a^2}.$$
 (2.14)

The temperature and angular velocities are

$$\beta = \frac{2\pi m (\cosh \alpha + \cosh \beta) (m + \sqrt{m^2 - a^2})}{\sqrt{m^2 - a^2}}, \qquad (2.15)$$

$$\Omega = \frac{a}{m(\cosh \alpha + \cosh \beta)(m + \sqrt{m^2 - a^2})}.$$
(2.16)

We see that for the Lorentzian solution to make sense we need $m \ge a$ and the limit $m \to a$ is an extremal limit (in the sense that inner and outer horizons have the same area) although the temperature *does not* vanish. We will expand on this later. When $a \to 0$ and $\alpha, \beta \to 0$ then the solution becomes the Schwarszchild black hole. We can check this by looking at the potentials, for example $\beta \to 8\pi m$.

Having found the location of the horizon and the angular velocity, we can evaluate the electric chemical potentials. These quantities are given by $\mu = i_V A|_{r\to\infty} - i_V A|_{hor}$, where $V = \partial_t + \Omega \partial_\phi$. This is a gauge invariant statement, but to be more rigorous we should apply a large gauge transformation such that $i_V A|_{hor} = 0$. The electric potentials are

$$\vec{\mu}_L = \frac{\vec{n}}{\sqrt{2}} \frac{\sinh \alpha}{\cosh \alpha + \cosh \beta}, \qquad \vec{\mu}_R = \frac{\vec{p}}{\sqrt{2}} \frac{\sinh \beta}{\cosh \alpha + \cosh \beta}.$$
(2.17)

Finally we can compute the area of the horizon, with respect to the Einstein frame metric,

$$A = 4\pi m (\cosh \alpha + \cosh \beta) (m + \sqrt{m^2 - a^2}).$$
(2.18)

The on-shell action of the black hole is, in the grand canonical ensemble where we set Dirichlet boundary conditions for all fields, given by

$$\log Z = -I_{\text{on-shell}} = \frac{A}{4} - \beta M + \beta \vec{\mu}_L \cdot \vec{Q}_L + \beta \vec{\mu}_R \cdot \vec{Q}_R + \beta \Omega J. \qquad (2.19)$$

We can compare the supersymmetric limit with the extremal limit. We already mentioned the extremal limit is m = a. The solution is supersymmetric when

SUSY:
$$M^2 = \frac{\bar{Q}_R^2}{2}.$$
 (2.20)

Combining with (2.13) and (2.12) we see that it implies

$$m^2 (1 + \cosh \alpha \cosh \beta)^2 = m^2 \cosh^2 \alpha \sinh^2 \beta. \qquad (2.21)$$

If all parameters are finite and real we cannot satisfy (2.21) as we need either (i) m = 0, implying M = 0 and $\vec{Q}_R = \vec{0}$ and, moreover, A = 0—implying a singularity at the horizon; or (ii) we have $m \neq 0$ but the solution is complex, for example we can see that if we solve for $\cosh \alpha$ as a function of $\cosh \beta$ then $\cosh \alpha < 1$ for all $\cosh \beta \in (+1, \infty)$. As explained by Sen the supersymmetric solutions have $m \to 0$ and either β going to infinity or α, β going to infinity together, but it is not always extremal in the sense of being zero temperature.

2.1.3 Thermodynamics without $(-1)^{\mathsf{F}}$ insertion

We want to describe the termodynamics of non-rotating electric black holes in this setup, without any insertion of $(-1)^{\mathsf{F}}$ (which we will discuss in later sections). We then set a = 0 and the expressions in the previous section simplify. The mass and entropy of the black hole are given by

$$M = \frac{m(1 + \cosh \alpha \cosh \beta)}{2}, \quad S = 2\pi m^2 (\cosh \alpha + \cosh \beta), \quad (2.22)$$

and the temperature

$$\beta = 4\pi m (\cosh \alpha + \cosh \beta). \tag{2.23}$$

The expressions for the charges and electric potentials are unmodified.

There are several distinct limits one can consider. In this section we assume the validity of (2.22) and (2.23), and discuss the thermodynamics implied by them. However, we should keep in mind that these relations have α' corrections which can become important when the black hole becomes small and, in fact, as we will discuss in Section 2.2, the correct description in certain regime is in fact given by horizonless geometries.

Low temperature limit The simplest limit one can consider is the low temperature limit, i.e. $\beta \to \infty$, while holding the charges fixed. In this limit we have

$$\sinh \alpha \sim \sqrt{2} \frac{8\pi Q_L}{\beta} \sim 0, \qquad \cosh \alpha \sim 1,$$
 (2.24)

$$\sinh\beta \sim \sqrt{2} \frac{8\pi Q_R}{\beta} \sim 0, \qquad \cosh\beta \sim 1.$$
 (2.25)

For this reason, if we compute the mass and entropy as functions of temperature we get

$$M \sim \frac{\beta}{8\pi}, \qquad S \sim \frac{\beta^2}{16\pi}.$$
 (2.26)

We simply recover the behavior of a regular Schwarszchild black hole, the bigger the black hole the bigger the mass and the colder it gets. Since the black hole becomes large and weakly curved, the black hole description is trustworthy even though the ensemble is thermodynamically unstable. The naive extremal limit Since we are considering here the special case a = 0, what we mean by the extremal limit is $m \to 0$. However, to keep the charges (2.12) fixed, the limit is not so straightforward. The only solution to this problem is to take either $|\alpha| \to \infty$ or $|\beta| \to \infty$. For simplicity, here we will focus on the cases where $\alpha, \beta > 0$. The generalization of the discussion to the cases where one or both of them are negative is straightforward.

We will consider first the case of $\beta \to \infty$, which as it turns out corresponds to the case of $Q_R > Q_L$. To keep the charges fixed, we take $\beta \to \infty$ and $m \to 0$ while keeping

$$m\cosh\beta \to m_0$$
 (2.27)

finite. The first surprise, under the assumption that we can trust the black hole solution, is that this extremal limit does not imply vanishing temperature. Instead

$$\beta_{\text{ext}} = 4\pi m_0, \qquad Q_L = \frac{m_0}{\sqrt{2}} \sinh \alpha, \qquad Q_R = \frac{m_0}{\sqrt{2}} \cosh \alpha.$$
 (2.28)

From the last two equations of (2.28) we see that $\tanh \alpha = Q_L/Q_R$ and $m_0 = \sqrt{2}\sqrt{Q_R^2 - Q_L^2}$. Therefore the extremal limit has a finite, charge dependent, temperature

$$T_{\max} = \frac{1}{4\sqrt{2}\pi\sqrt{Q_R^2 - Q_L^2}}.$$
(2.29)

This solution is precisely Sen's small black hole since the area of the horizon now vanishes

$$A \sim m^2 \cosh\beta \sim m_0 m \to 0. \tag{2.30}$$

We saw earlier that at low temperature the black hole is very large since it behaves as the Schwarszchild solution. As the temperature increases the mass decreases until it reaches extremality at β_{ext} . This is the situation if $Q_R > Q_L$. At this point the entropy vanishes since the area of the horizon vanishes. If we further extrapolate the expressions to temperature higher than T_{max} , the mass parameter becomes negative m < 0, leaving a naked singularity. Moreover, the parameter β goes to the complex plane. At the temperature T_{max} the value of the mass is given by $M_{\text{ext}}^2 = Q_R^2/2$ which is precisely the BPS mass which is the lowest possible from the dual quantum mechanical point of view. In Figure 1, we show how the mass and entropy behave as functions of temperature. We should however stress that we should not trust the curves when the temperature becomes too close to T_{max} and the black hole becomes string scale. We will discuss the correct behavior in Section 2.2.

Above we considered the case of $\beta \to \infty$ and therefore $Q_R > Q_L$. The discussion of the case $Q_L > Q_R$ essentially mirrors the previous case, but the interpretation is slightly different. In this case as the black hole temperature rises, the solution eventually has $\alpha \to \infty$ instead. The expressions above are still true, but now the lowest mass black hole has $M = Q_L/\sqrt{2}$ which is not supersymmetric. Since $Q_L > Q_R$ there is now a range of masses, namely $Q_L^2/2 \ge M^2 \ge Q_R^2/2$ which is allowed by unitarity in the microscopic theory, but where black holes do not exist. Therefore depending on the sign of $Q_R - Q_L$ the ground state might be BPS or not.



Figure 1: (a) An illustration of the black hole mass (divided by its extremal value M_{ext}) as a function of temperature (divided by $T_{\text{max}} = 1/(4\pi\sqrt{2}\sqrt{|Q_R^2 - Q_L^2|}))$. The mass decreases until the temperature reaches maximum. The dashed part of the blue line means that the black hole is of string scale and the curve is just an extrapolation. (b) An illustration of the black hole entropy (divided by the extremal entropy S_{ext} from microstate counting) as a function of temperature. The naive extrapolation of the entropy simply vanishes when we reach the maximum temperature.

Finally, if $Q_L = Q_R$ then $\alpha = \beta$. We will not go into the details of this special case, but just note that we need to take both of them to infinity, and the extrapolation of (2.23) suggests that the extremal limit corresponds to infinite temperature $\beta_{\text{ext}} = 0$.

We note that the black hole solution in the two derivative theory (2.1) is symmetric with respect to turning on the left or right moving charges. However, from the microscopic theory point of view, we wouldn't expect such symmetry. For one, the solution should be supersymmetric only when $M = Q_R/\sqrt{2}$ but not when $M = Q_L/\sqrt{2}$. As it turns out, this symmetry is broken once one considers the leading α' corrections to the black hole [28].

2.2 Transition to free strings via the charged Horowitz-Polchinski solution

As we saw in the previous section, the extrapolation of the gravity solution (2.3) to the extremal limit has some peculiar features. The black hole solution becomes singular as the horizon shrinks to zero size. Of course, once the size of the horizon becomes comparable to string scale, the higher curvature corrections would become important and there is no reason to trust the solution (2.3). In this section, we will review the result in [12], which states that the approach to extremality of the two-charge system is in fact described by a charged analogue of the Horowitz-Polchinski solution [11].

Here we will only sketch the idea behind the construction of the solution as details can be found in [12]. The original uncharged Horowitz-Polchinski solution can be constructed explicitly by considering the effective action of the thermal winding modes coupled to gravity and dilaton. The uncharged solution is trustworthy when the temperature is below but very close to the Hagedorn temperature, $(\tilde{\beta} - \beta_H)/\beta_H \ll 1$,⁶ where we can focus on only the winding modes with winding number ±1 and the fluctuation of the g_{tt} component of the metric. The solution carries a classical entropy, which agrees with the Hagedorn density of states for a gas of strings close to the Hagedorn temperature. This suggests that the physical interpretation of the solution is describing a gas of highly excited strings, weakly interacting via gravity (see also [29]). It is further suggested in [12] that in Heterotic string theory, this classical solution could potentially be smoothly connected to the Schwarzschild black hole solution as we lower the temperature.⁷

An important property of the solution which differs from the Schwarzschild black hole is that as we increase the temperature, the physical size ℓ_{size} of the solution expands as $1/\sqrt{\tilde{\beta} - \beta_H}$. Physically, this means the classical solution is describing a more and more dilute string gas. When the temperature gets so close to the Hagedorn temperature such that the size of the classical solution approaches the size of a free string gas with the same mass, the classical solution breaks down and the correct description is simply the quantum description of free string gas. In four dimensions, the solution is only valid when

$$\tilde{g}_s^{\frac{4}{3}} \lesssim (\tilde{\beta} - \beta_H) / \beta_H \ll 1.$$
(2.31)

Given the uncharged solution, [12] performed a solution generating transformation to find the charged solution. The explicit form of the solution can be found in [12], while here we will only mention a few main properties. It turns out that the $\tilde{\beta} \to \beta_H$ limit of the "seed" solution is mapped to the extremal limits $M \sim Q_R/\sqrt{2}$ or $M \sim Q_L/\sqrt{2}$ of the generated solutions. Differing from the naive extrapolation of the black hole results in Section 2.1.3, where the temperature rises and reaches a maximum, the temperature of the generated Horowitz-Polchinski solution goes to zero, i.e. $\beta \to \infty$ in the extremal limit. Furthermore, the solution correctly reproduces the entropy as expected from counting microstates of long strings carrying momentum and winding in the extra directions

$$S \sim \sqrt{2\pi\ell_s} \sqrt{Q_R^2 - Q_L^2}, \qquad M \sim \frac{Q_R}{\sqrt{2}},$$

$$S \sim \pi\ell_s \sqrt{Q_L^2 - Q_R^2}, \qquad M \sim \frac{Q_L}{\sqrt{2}}.$$
(2.32)

The thermodynamics of the charged solution can be determined purely from the thermodynamics of the seed solution plus the form of the solution generating transformation at spatial infinity. We will discuss the derivation of such relations in Appendix A, extending the analysis in [12] to the cases where the seed solution has rotation. Specifying (A.29)– (A.33) to the case without rotation, and expanding the temperature of the seed solution

⁶We will use tilded quantities to denote the quantities in the uncharged solution, which are distinct from those of the charged solution but related via the solution generating procedure, see Appendix. A for a more detailed discussion.

⁷While in the Type II theories, a worldsheet index argument suggests that the two solutions cannot be smoothly interpolated.

around $\beta_H = 2\pi R_H = 2\pi (1 + 1/\sqrt{2})\ell_s$, we get

$$M = \frac{\tilde{S}'}{2\pi R_H} \cosh \alpha \cosh \beta ,$$

$$S = \frac{\tilde{S}'}{2} \left[\left(1 - \frac{\alpha'}{2R_H^2} \right) \cosh \alpha + \left(1 + \frac{\alpha'}{2R_H^2} \right) \cosh \beta \right] ,$$

$$\beta = \frac{\beta_H}{2} \left[\left(1 + \frac{\alpha'}{2R_H^2} \right) \cosh \alpha + \left(1 - \frac{\alpha'}{2R_H^2} \right) \cosh \beta \right] ,$$

$$Q_L = \frac{\tilde{S}'}{\sqrt{2\pi R_H}} \cosh \beta \sinh \alpha ,$$

$$Q_R = \frac{\tilde{S}'}{\sqrt{2\pi R_H}} \cosh \alpha \sinh \beta ,$$

(2.33)

where \tilde{S}' is the entropy of the seed solution after stripping off the dependence on the string coupling, and is given by

$$\tilde{S}' \approx c_0 \sqrt{\ell_s^3(\tilde{\beta} - \beta_H)}.$$
 (2.34)

Here $c_0 \approx 2.41$ is an order one number coming from the properties of the seed solution.

In summary, (2.33) and (2.34) describe the thermodynamic quantities of the charged Horowitz-Polchinski solution, parameterized by three parameters $\{\tilde{\beta}, \alpha, \beta\}$. They are only valid when we satisfy (2.31). Note that the lower end of the inequility in (2.31) is expressed in terms of the string coupling constant in the seed solution, rather than the generated solution.

With (2.33), we can reexamine how the naive approach to extremality in section 2.1.3 gets modified. As we mentioned, now the extremal limit corresponds to $\tilde{\beta} \to \beta_H$, and to fix the charges, we need to combine it with the limits $\alpha \to \infty$ or $\beta \to \infty$. For the case of $\alpha \to \infty$, we have $M \to Q_L/\sqrt{2}$, and for $\beta \to \infty$, we have $M \to Q_R/\sqrt{2}$. From the expression of β in (2.33), we clearly see that both would correspond to the limit where $\beta \to \infty$, i.e. the temperature goes to zero in the extremal limit.

We can consider some special cases for which we can write down simpler formulas for how the energy and entropy depends on the temperature. Consider the case $Q_L = 0$ $(\alpha = 0, \beta \rightarrow \infty)$, at low temperature we have

$$\frac{M - M_{\text{ext}}}{M_{\text{ext}}} \sim \frac{S - S_{\text{ext}}}{S_{\text{ext}}} \sim 2e^{-2\beta} \sim (\pi \ell_s T)^2, \qquad (2.35)$$

where $M_{\text{ext}} = Q_R/\sqrt{2}$ and $S_{\text{ext}} = \sqrt{2}\pi \ell_s Q_R$. Translating (2.31) into the parameters of the generated solution, we find that the solution is trustworthy when

$$\left(\frac{\ell_s}{Q_R}\right)^{\frac{1}{2}} \ll Q_R T \ll 1.$$
(2.36)

On the other hand, for the case of $Q_R = 0$ ($\beta = 0, \alpha \to \infty$), at low temperature we have

$$\frac{M - M_{\text{ext}}}{M_{\text{ext}}} \sim \frac{S - S_{\text{ext}}}{S_{\text{ext}}} \sim 2e^{-2\alpha} \sim 2(\pi \ell_s T)^2, \qquad (2.37)$$



Figure 2: (a) The blue curves are from Figure 1 for comparison. The orange curve is the mass of the charged Horowitz-Polchinski solution as a function of the temperature. It is trustworthy when $\sqrt{\ell_s/|Q|} \ll T/T_{\max} \ll 1$, where |Q| can be taken to be the greater of $|Q_L|, |Q_R|$. The dashed lines are extrapolations. We see that the temperature goes to zero in the extremal limit. (b) The orange curve is the entropy of the Horowitz-Polchinski as a function of the temperature. We see that S/S_{ext} goes to one in the extremal limit. The precise shape of the curves depends on the values of the charges and ℓ_s in Planck units. In the microcanonical ensemble, where we vary the mass, we expect the two curves to be joined together near where their extrapolations cross.

where $M_{\text{ext}} = Q_L/\sqrt{2}$ and $S_{\text{ext}} = \pi \ell_s Q_L$. The region of validity of the solution is similar to (2.36), but with Q_R replaced by Q_L . Note that we have $\delta S \propto \delta M$, which is expected from the microscopic entropy formula

$$S = 2\pi \ell_s \left[\sqrt{M^2 - \frac{Q_L^2}{2}} + \frac{1}{\sqrt{2}} \sqrt{M^2 - \frac{Q_R^2}{2}} \right]$$
(2.38)

when we expand around extremality.

We note that the low temperature thermodynamics is qualitatively different from what one would expect if there were an AdS_2 region in the geometry. In Section 6 we will argue that the fact that the low energy physics is not given by AdS_2 could be argued from more general principle. In Figure 2 we illustrate that after considering the charged Horowitz-Polchinski solution, the naive extremal limit in Figure 1 is modified significantly.

We close this section by mentioning that other than the thermodynamic quantities, one can also look at the physical size of the Horowitz-Polchinski solution, and it matches with the size of the BPS string expected from a random walk picture [12],

$$\ell_{\rm phys} \sim (nw)^{\frac{1}{4}} \ell_s \,. \tag{2.39}$$

3 Gravitational configurations that contribute to the index

We now initiate our study of the gravitational path integral that evaluates the index of two-charge states. We first construct supersymmetric solutions by taking a non-trivial limit of the two-charge black hole analyzed in the previous section. The situation is more subtle than its 'big black hole' counterpart and we emphasize some puzzling features. Those features will go away later when we include higher derivative corrections.

3.1 Imposing supersymmetry

We want to construct supersymmetric finite temperature black hole solutions that contribute to the gravitational path integral for the index. We actually need to consider the helicity supertrace, but one can simply absorb the fermion zero modes and we will not worry about this here. Imposing supersymmetry at the boundary (i.e. the asymptotic region of flat space) requires fermions to be periodic. As explained in [17] this constrains the angular velocity Ω . Supersymmetry in the bulk requires the existence of a globally defined Killing spinor. For the black hole geometries presented in Section 2 this constrains the mass parameter M. Explicitly these two conditions are given by

> Boundary Supersymmetry : $\beta \Omega = 2\pi i \mod 4\pi \mathbb{Z}$, Bulk Supersymmetry : $M = Q_R/\sqrt{2}$.

The first condition fixes the fermion periodicity conditions because this choice of angular velocity amounts to an insertion of $e^{\beta\Omega J} \rightarrow e^{2\pi i J} = (-1)^{\mathsf{F}}$. Since all excitations have integral or half-integral angular momentum this condition defines the angular velocity only up to a shift multiple of $4\pi/\beta$. Bulk and boundary supersymmetry are in principle independent. For example, in the case of Reissner-Nordström, if $\beta\Omega = 2\pi i$ then this implies that M = Q and a globally defined Killing spinor exists, but if $\beta\Omega = 6\pi i$ or any other odd multiple of $2\pi i$, then $M \neq Q$ and supersymmetry is not preserved in the bulk. One unexpected feature of the two-charge system is that even for $\beta\Omega = 2\pi i$ bulk supersymmetry is not guaranteed.

The discussion here will assume that in the presence of $(-1)^{\mathsf{F}}$ the gravitational path integral is dominated by a black hole-like saddle. Can there be another saddle similar to the Horowitz-Polchinski one that contributes to the index? One option, since the thermal circle is non-contractible, is to take the Horowitz-Polchinski solution of the previous section and impose fermions to be periodic. The problem with this proposal is that the solution is not supersymmetric and therefore does not contribute to the index due to fermion zeromodes.⁸ Note also that the Horowitz-Polchinski solution also breaks down before we reach exact extremality due to quantum corrections. Even if this had worked, the problem is that the equations of motion that Horowitz-Polchinski solves are not correct in the presence of $(-1)^{\mathsf{F}}$. At least in flat space, periodic fermions lead to the standard GSO projection and therefore the light winding mode that condenses in the Horowitz-Polchinski solution is not part of the low energy spectrum [30]. It is therefore not clear which equations we are supposed to solve to find such a solution. It is possible that by turning on an imaginary angular potential, similar to the black hole case, one can evade the flat space GSO projection argument and arrive at a supersymmetric solution, but we are not sure

 $^{{}^{8}}$ Even if it did, it produces a contribution that depends on temperature in the free energy, inconsistent with the behavior of an index.

how to construct such a solution explicitly (see related discussion in [31]). We should emphasize that even if such a solution was found and produces the correct index, it would not invalidate our discussion below. On the contrary, it would be in harmony with the conjecture that there is no phase transition between the Horowitz-Polchinski solution and the black hole in the heterotic theory. Since the supergravity sector is part of the low energy spectrum when fermions are periodic, and since there are black hole solutions in this sector, this will be our focus in the rest of the paper.

After these introductory remarks let us impose $\beta \Omega = 2\pi i$ and see what the implications are for the two-charge black hole. Since the angular velocity is imaginary it is convenient to introduce a = ia, so a will then be real. The condition becomes then

$$\frac{\beta\Omega}{2\pi i} = \frac{\mathsf{a}}{\sqrt{m^2 + \mathsf{a}^2}} = 1.$$
(3.1)

It is convenient sometimes to replace the variables (m, \mathbf{a}) by $(m, x = \mathbf{a}/m)$. Then the boundary supersymmetry condition becomes $x = \sqrt{1 + x^2}$, and the only solution is to take $x \to \infty$. We are working in the thermal ensemble and the black hole temperature, in the limit $x = \mathbf{a}/m \to \infty$, becomes

$$\beta = \frac{2\pi m (\cosh \alpha + \cosh \beta) (m + \sqrt{m^2 + \mathbf{a}^2})}{\sqrt{m^2 + \mathbf{a}^2}} \Big|_{\mathbf{a}/m \to \infty} = 2\pi m (\cosh \alpha + \cosh \beta). \quad (3.2)$$

Since the temperature is a free parameter set by our choice of boundary conditions at asymptotic flat space, the right hand side of this expression should remain finite. We analyze below two ways to implement this condition.

Setting $\mathbf{a} \to \infty$ and *m* finite As a first possibility we can keep *m*, α and β finite, with one combination of them set by the temperature, and send $\mathbf{a} \to \infty$ to implement boundary supersymmetry. We will see now this option is ruled out after imposing bulk supersymmetry. For *m* finite, combining (2.13) with (2.12) one can show that setting $M^2 = Q_R^2/2$ implies the following relation between the parameters α and β

$$(1 + \cosh \alpha \cosh \beta)^2 = \cosh^2 \alpha \sinh^2 \beta .$$
(3.3)

There is no real solution of this equation for finite α and β , since the constraint implies $e^{\alpha+\beta} < 0$. It is not clear that the black hole solves the correct equations of motion since for complex values of α and β the transformation used to generate the solution is not in O(7,23) any longer. There is one more problem with this scenario that is worth mentioning. We evaluate the area of the horizon, given in (2.18), in the large a limit

$$A \to 4\pi m (\cosh \alpha + \cosh \beta) m x \quad \Rightarrow \quad A = 2\beta m x \,.$$
 (3.4)

We see that at fixed temperature and keeping m finite, the area is $A \propto \mathbf{a} \to \infty$. Now instead of vanishing, the area actually diverges! It is not obvious that this feature is necessarily problematic, as explained in [23], but we would like to avoid it if possible since the interpretation of such a saddle is unclear. For these reasons we will not consider this direction any further.

Setting $m \to 0$ and a finite Another way of implementing the bulk supersymmetry condition is to keep a finite while taking the mass parameter m to zero. This is the regime considered in [27], in the absence of the $(-1)^{\mathsf{F}}$ insertion. Looking at the expression for the temperature given in (3.2) we see that this limit requires taking $(\cosh \alpha + \cosh \beta) \to \infty$. How this limit is taken is fixed by bulk supersymmetry. For this purpose let us take the $m \to 0$ limit of the charges

$$\vec{Q}_L = \frac{\beta}{2\pi\sqrt{2}} \frac{\sinh\alpha\cosh\beta}{\cosh\alpha + \cosh\beta} \vec{n}, \qquad \vec{Q}_R = \frac{\beta}{2\pi\sqrt{2}} \frac{\sinh\beta\cosh\alpha}{\cosh\alpha + \cosh\beta} \vec{p}.$$
(3.5)

and similarly the mass

$$M = \frac{\beta}{4\pi} \frac{1 + \cosh \alpha \cosh \beta}{\cosh \alpha + \cosh \beta} \,. \tag{3.6}$$

One can check that $\beta \to \infty$ while keeping α finite does provide a solution of $M^2 = Q_R^2/2$, as pointed out in [27]. To guarantee both boundary and bulk supersymmetry, as well as a finite temperature, we then take

$$m \to 0, \quad \beta \to \infty, \qquad \text{such that} \qquad m \cosh \beta \to m_0,$$
 (3.7)

where we introduced the finite parameter m_0 . In terms only of parameters that are finite in the limit, the temperature and charges are

$$\beta = 2\pi m_0, \qquad Q_L = \frac{m_0}{\sqrt{2}} \sinh \alpha, \qquad Q_R = \frac{m_0}{\sqrt{2}} \cosh \alpha. \tag{3.8}$$

We can use the second and third equations to solve for m_0 and α in terms of the charges. Explicitly, we obtain

$$m_0 = \sqrt{2(Q_R^2 - Q_L^2)}, \quad \tanh \alpha = \frac{Q_L}{Q_R}.$$
 (3.9)

Using these values, we see that the temperature is fixed in terms of the charges! Moreover, the parameter **a** is undetermined. We will comment more on this later. This solution only exists if $Q_R > Q_L$. No supersymmetric solution of this kind exists if $Q_L > Q_R$ instead.⁹ This is consistent with the spectrum of heterotic strings compactified on a torus, and consistent with the spectrum of Lorentzian black holes in the absence of the $(-1)^{\mathsf{F}}$ insertion discussed in Section 2. Finally, the chemical potentials of the left and right charges have magnitudes $\mu_L = 0$ and $\mu_R = 1/\sqrt{2}$, independent of the parameters m_0 and α .

3.2 Explicit form of the solution

Now that we have determined the scaling of all parameters in the supersymmetric limit we can write down explicitly how the solution looks like. The same solution was written down

⁹What happens with the supergravity solution when $Q_L = Q_R$? In this case $\alpha = \beta$ and bulk supersymmetry demands that $\alpha = \beta \to \infty$. But it is evident from (3.5) that in this case keeping the temperature finite leads to $Q_L, Q_R \to \infty$, therefore no solution exists with finite $Q_L = Q_R$. Another option is to take $\beta \to 0$ such that βe^{β} is finite. This implies now that $m \sim e^{-2\beta}$, vanishing faster than before. To keep the black hole area finite we need to send $\mathbf{a} \to \infty$. Since its not clear whether this is a physical solution or not we will not consider it further and restrict ourselves to $Q_R > Q_L$.

in [6, 32] although it was interpreted differently. In the supersymmetric limit the metric simplifies slightly,

$$e^{-\Phi} ds^{2} = \Delta^{\frac{1}{2}} \left\{ \frac{r^{2} - \mathsf{a}^{2} \cos^{2} \theta}{\Delta} dt_{E}^{2} + \frac{dr^{2}}{r^{2} - \mathsf{a}^{2}} + d\theta^{2} + \frac{\sin^{2} \theta}{\Delta} [\Delta - \mathsf{a}^{2} \sin^{2} \theta (r^{2} - \mathsf{a}^{2} \cos^{2} \theta + 2m_{0}r \cosh \alpha)] d\phi^{2} - \frac{2m_{0}r \mathsf{a} \sin^{2} \theta}{\Delta} dt_{E} d\phi \right\},$$
(3.10)

where the supersymmetric limit of the function Δ becomes

$$\Delta = (r^2 - \mathsf{a}^2 \cos^2 \theta)^2 + 2m_0 r (r^2 - \mathsf{a}^2 \cos^2 \theta) \cosh \alpha + m_0^2 r^2 \,. \tag{3.11}$$

Here, we remind the reader that ds^2 is defined in the string frame. The dilaton is still given by $\Phi = \log(r^2 - a^2 \cos^2 \theta) / \Delta^{1/2}$. The solution is real in Euclidean space. From the second term in the first line we see that the horizon is located at

$$r_{+} = a,$$
 (3.12)

giving a geometric interpretation to **a**. One can verify directly with the supersymmetric solution that smoothness of the metric at the horizon imposes values of the temperature and angular velocity consistent with the previous analysis, namely $\beta = 2\pi m_0$ and $\Omega = i/m_0$, such that $\beta \Omega = 2\pi i$. In fact, the solution is smooth everywhere on the sphere except at the poles, which we comment on later. The area of the horizon in Einstein frame is

$$A = \int d\theta d\phi \sqrt{g_{\theta\theta}^E g_{\phi\phi}^E} = \int d\theta d\phi m_0 \mathbf{a} \sin\theta = 4\pi m_0 \mathbf{a}, \qquad (3.13)$$

which is finite and non-vanishing! The rotation necessary to implement the $(-1)^{\mathsf{F}}$ has turned the small black hole (at extremality) into a macroscopic one at finite temperature for the index. This seems to suggest we are now able to use the gravitational path integral to evaluate the index of the two-charge state avoiding small black holes, but this is too naive. Even though the solution has a macroscopic horizon, the black hole geometry is actually singular at the poles of the sphere $\theta = 0, \pi$ so the rotation alone has not fixed all issues. This is clear when looking in the Einstein frame metric which is given by the string frame one multiplied by e^{Φ} . Near the horizon

$$e^{\Phi}\big|_{r=\mathsf{a}} = \Delta^{-\frac{1}{2}} \mathsf{a}^2 \sin^2 \theta, \qquad (3.14)$$

and $\Delta|_{r=a}$ is a non-vanishing finite function of θ . Therefore we see that e^{Φ} vanishes at the two poles of the sphere. A more detailed analysis shows that the string frame metric is singular, while the vanishing of e^{Φ} at the poles makes the Einstein frame metric smooth, although higher derivative corrections in the Einstein frame can be large due to dilaton diverging.

The solution is specified by the gauge potentials besides the metric. These also have a finite limit in the supersymmetric limit given by

$$\vec{A}_{L} = -\frac{\vec{n}}{\sqrt{2}} \frac{m_{0}r \sinh \alpha (r^{2} - a^{2} \cos^{2} \theta)}{\Delta} dt - i\frac{\vec{n}}{\sqrt{2}} \frac{m_{0}^{2}r^{2}a \sinh 2\alpha \sin^{2} \theta}{2\Delta} d\phi,$$

$$\vec{A}_{R} = -\frac{\vec{p}}{\sqrt{2}} \frac{m_{0}r((r^{2} - a^{2} \cos^{2} \theta) \cosh \alpha + m_{0}r)}{\Delta} dt$$

$$+ i\frac{\vec{p}}{\sqrt{2}} \frac{m_{0}ra \sin^{2} \theta (r^{2} - a^{2} \cos^{2} \theta + m_{0}r \cosh \alpha)}{\Delta} d\phi,$$

$$B = -i\frac{m_{0}ra \sin^{2} \theta}{\Delta} (r^{2} - a^{2} \cos^{2} \theta + m_{0}r \cosh \alpha) dt d\phi.$$
(3.15)

These gauge potentials are smooth and one can extract from them the chemical potentials, verifying that $\mu_L = 0$ and $\mu_R = 1/\sqrt{2}$. Finally, the functions appearing in the moduli scalar profiles M also simplify

$$\mathsf{P} = \frac{2m_0^2 r^2 \sinh^2 \alpha}{\Delta}, \qquad \mathsf{Q} = -\frac{2m_0 r \sinh \alpha}{\Delta} (r^2 - \mathsf{a}^2 \cos^2 \theta + m_0 r \cosh \alpha). \tag{3.16}$$

These functions are smooth and finite at the horizon.

In general, the Euclidean action of the two-charge heterotic black hole is given by $I_{\text{on-shell}} = \beta M - A/4 - \beta \Omega J$ and comes solely from the boundary terms in the action. This is the result in an ensemble of fixed electric charges. If we have a supersymmetric solution then the first term is $\beta Q_R^2/2$. In the rest of the paper we will always include an appropriate boundary term that has the effect of shifting the zero-point energy such that $I \rightarrow I - \beta Q_R^2/2$. After this modification, $-I_{\text{on-shell}}$ simplifies to $A/4 + \beta \Omega J$. We can show now that whenever a supersymmetric saddle exists, then these two terms cancel each other and the classical two-derivative contribution to the on-shell action, and therefore to $\log Z_{\text{index}}$ vanishes. The area and the angular momentum can be written as

$$\frac{A}{4} = \frac{\beta \mathsf{a}}{2}, \qquad J = i\frac{\beta \mathsf{a}}{4\pi}. \tag{3.17}$$

We already discussed the area. The angular momentum can be read off directly from the supersymmetric solution or taking the appropriate limit of (2.13). Using that $\beta \Omega = 2\pi i$ we get the desired cancellation

$$\frac{A}{4} + \beta \Omega J = 0, \qquad \Rightarrow \qquad \log Z_{\text{index}} = -I_{\text{on-shell}} = 0. \tag{3.18}$$

We found a black hole with a macroscopic area that nevertheless gives a vanishing contribution to the index to leading order. This is the desired conclusion, since a non-zero action at the two-derivative level would, by dimensional analysis, grow too fast with the charges and would not reproduce the microscopic index.

3.3 Summary of properties

To recap, we have found a supersymmetric finite temperature black hole with two-charges that evaluates the index avoiding the small black hole regime. Even after including the rotation necessary to implement the $(-1)^{\mathsf{F}}$ some puzzles remain to be resolved. We finish this section by summarizing and expanding on some of them. In the next sections we will resolve them by including higher derivative corrections.

- 1. The solution we found has a curvature singularity at the poles of the S^2 horizon $r = r_+ = a$ at $\theta = 0$ and π . This forces us to include higher derivative corrections even though the horizon is macroscopic.
- 2. For a given set of charges, the solution only exists for a unique value of the temperature. This is surpring since charges and temperatures can be independently determined by boundary conditions at infinity. After taking the supersymmetric limit we found

$$\beta = 2\pi \sqrt{2(Q_R^2 - Q_L^2)}.$$

For comparison, in the Reissner-Nordström case analyzed in [17] the inverse temperature is $\beta = \pi Q(Q + 2\mathbf{a})/\mathbf{a}$, while $r_+ = Q + \mathbf{a}$. A solution therefore exists for any temperature and the horizon radius depends on its value. For the case considered here, the singular feature is that the dependence on \mathbf{a} disappeared completely from β , making it a function only of the charge. What solution contributes to the gravitational path integral when $\beta \neq 2\pi \sqrt{2(Q_R^2 - Q_L^2)}$?

3. In the Reissner-Nordström case quoted in the previous item, not only is the temperature a tunable parameter but it also determines a unique value of \mathbf{a} . In the two-charge case another consequence of the fact that \mathbf{a} drops out of the temperature¹⁰ is that there is a moduli space of solutions labeled by \mathbf{a} from 0 to ∞ . All values of \mathbf{a} contribute equally and have to be integrated over. This is a disturbing state of affairs and we will see that higher derivative corrections will determine a unique value of \mathbf{a} .

Before addressing the effect of higher derivative corrections, we will show in the next section that these features are common to a class of two-charge black holes in $\mathcal{N} \geq 2$ supergravity.

4 The index and new attractor mechanism in 4d $\mathcal{N} = 2$ supergravity

In this section we move to the $\mathcal{N} = 4$ four-dimensional effective theory that describes the compactification of the heterotic string on T^6 . From the string-string duality, this is equivalent to Type II theory on $K3 \times T^2$. More generally, we can consider four-dimensional $\mathcal{N} = 2$ theories that arise from the compactification of Type II on a Calabi-Yau 3-fold. These theories are described by $\mathcal{N} = 2$ supergravity coupled to vector and hyper multiplets. The fields that participate in the black hole solutions live in the graviton multiplet and the

¹⁰The fact that **a** drops out, and the temperature is fixed in terms of the charges, is special to compactifications to four dimensions. Similar black hole solutions in toroidal compactifications to D > 4 dimensions do not have this problem [26]. The reason we insist on working in D = 4 is the fact that supersymmetric higher derivative solutions are known and we use them in Section 5.

vector multiplet, and we focus on this sector. In this section we discuss the theory with a two-derivative action, and in the next section we consider higher-derivative corrections.

The conformal supergravity formalism is an elegant and powerful formalism to describe the theory of $\mathcal{N} = 2$ supergravity coupled to a number of vector multiplets [33, 34]. In this formalism, local conformal invariance is a gauge symmetry of the theory. Accordingly, each field is assigned a conformal or Weyl weight. Eventually, the conformal symmetry is gauge-fixed by assigning a dimensionful number like $m_{\rm Pl}$ to a field with non-zero Weyl weight. This gauge-fixing leads to Poincaré supergravity.

The bosonic physical field content of each vector multiplet consists of a vector field A^I and a complex scalar X^I , $I = 0, 1, ..., n_v$. The scalars have Weyl weight 1. Solutions of the theory carry electric and magnetic charges (Q_I, P^I) under these gauge fields. The two-derivative action is completely specified by the prepotential F(X), which is a homogeneous function of degree two and has Weyl weight 2. An important role is played by the first derivatives $F_I = \partial F / \partial X^I$ which have Weyl weight 1. At the two-derivative level the equations of motion of the theory have $Sp(2n_v + 2; \mathbb{R})$ symmetry, under which the charge vector $\Gamma = (P^I, Q_I)$ and the period vector $\Omega_{\text{hol}} = (X^I, F_I)$ transform as $2(n_v + 1)$ dimensional vectors.

Two important invariants of the symplectic symmetry are^{11} the generalized Kähler potential (of Weyl weight 2)

$$e^{-\mathcal{K}(X,\overline{X})} = i \left(X^I \overline{F}_I - \overline{X}^I F_I \right) = i \left\langle \Omega_{\text{hol}}, \overline{\Omega}_{\text{hol}} \right\rangle, \qquad (4.1)$$

and the central charge function (of Weyl weight 0),

$$Z(\Gamma;\Omega) = \langle \Gamma, \Omega \rangle = e^{\frac{1}{2}\mathcal{K}(X,\overline{X})} \left(P^I F_I - Q_I X^I \right).$$
(4.2)

The conjugate central function $\overline{Z}(\Gamma; \Omega) = Z(\Gamma; \overline{\Omega})$ is also used below.

4.1 $\frac{1}{2}$ -BPS solutions of $\mathcal{N} = 2$ supergravity

The general $\frac{1}{2}$ -BPS stationary solution of this supergravity theory is described as follows [35, 36]. The input to the solution the vector $H(\mathbf{x}) = (\widetilde{H}^{I}(\mathbf{x}), H_{I}(\mathbf{x}))$ of harmonic functions on base space. These functions are sources of electric and magnetic fields.

The metric takes the form

$$ds^{2} = e^{2U}(dt_{E} + i\omega)^{2} + e^{-2U}dx^{m}dx^{m}, \qquad (4.3)$$

where $U(\mathbf{x})$ is a function of the base space coordinates x^m , m = 1, 2, 3, and $\omega = \omega_m dx^m$ is a one-form on the base space that governs the angular rotation of the solution. In order to have asymptotically flat space, one has $e^{-2U(\infty)} = 1$. Note that we have directly described the Euclidean solution with compact time coordinate t_E . The warping of the base space and the connection ω in (4.3) are given by

$$e^{-2U} = i \left(\mathcal{Y}^I \,\overline{\mathcal{G}}_I - \overline{\mathcal{Y}}^I \,\mathcal{G}_I \right), \qquad \star \,\mathrm{d}\,\omega = i \langle \mathrm{d}H, H \rangle, \tag{4.4}$$

¹¹Here, and below, the symplectic product of two vectors $A = (\tilde{A}^I, A_I)$ and $B = (\tilde{B}^I, B_I)$ is given by $\langle A, B \rangle = \tilde{A}^I B_I - \tilde{B}^I A_I$.

where the weight-0 normalized fields $(\mathcal{Y}^I, \mathcal{G}_I)$ are given by

$$\left(\mathcal{Y}^{I}, \mathcal{G}_{I}\right) = e^{\frac{1}{2}\mathcal{K}(X,\overline{X})} \overline{Z}(H;\Omega) \left(X^{I}, F_{I}\right)$$

$$(4.5)$$

and similarly $\overline{\mathcal{Y}}^I$, $\overline{\mathcal{G}}_I$ by the complex conjugate equations. The scalar fields obey the generalized stabilization equations¹²

$$\mathcal{Y}^{I} - \overline{\mathcal{Y}}^{I} = i\widetilde{H}^{I}, \qquad \mathcal{G}_{I} - \overline{\mathcal{G}}_{I} = iH_{I}, \qquad (4.6)$$

The above equations of supersymmetry for the $(\mathcal{Y}^I, \mathcal{G}_I)$ fields can be presented in terms of a prepotential function $\mathcal{G}(\mathcal{Y})$ with $\mathcal{G}_I = \partial \mathcal{G}/\partial \mathcal{Y}^I$, where \mathcal{G} is the same mathematical function as $F.^{13}$

Besides the metric and the scalars, the solution is specified by the n_v+1 vector multiplet U(1) gauge fields. These are also determined by the harmonic functions in a specific way, presented for example in equation (3.21) of [18].

4.2 Euclidean black holes and near-horizon behavior

In this subsection we review the "new attractor mechanism" that was found in [18]. Here we present all the equations in coordinate language. We refer the reader to [18] for details of the coordinate-invariant equations.

Extremal black holes Supersymmetric extremal solutions are obtained by choosing the harmonic functions in the above discussion to have electric and magnetic sources at one point, say the origin, i.e.,

$$H(\mathbf{x}) = h + \frac{\Gamma}{r}, \qquad r = |\mathbf{x}|.$$
(4.7)

The equations (4.4) show that the solution is spherically symmetric and static, i.e. $\omega = 0$ in (4.3). The vector $h \equiv H(\infty)$ is the asymptotic value of sources. The generalized stabilization equations (4.6) imply that the asymptotic behavior of the scalars as $r \to 0$ is

$$\mathcal{Y}^I \sim \frac{Y^I_*}{r}, \qquad \mathcal{G}_I \sim \frac{G_{I*}}{r}, \qquad (4.8)$$

where the constant values Y_*^I, G_{I*} (and therefore Z_*) are determined by the *extremal* attractor equations

$$Y_*^I - \overline{Y}_*^I = iP^I, \qquad G_{I*} - \overline{G}_{I*} = iQ_I.$$

$$(4.9)$$

The behavior of the metric as $r \to 0$ is

$$e^{-2U} \sim \frac{i(Y_*^I \overline{G}_{I*} - \overline{Y}_*^I G_{I*})}{r^2}.$$
 (4.10)

From the form of the metric (4.3) we see that the Bekenstein-Hawking entropy of the black hole is given by $S_{\rm BH} = \pi i (Y_*^I \overline{G}_{I*} - \overline{Y}_*^I G_{I*}).$

 $^{^{12}}$ Note that the signs in these equations are opposite to the one in [18].

¹³One should note that the rescaling between X^{I} and \mathcal{Y}^{I} is not holomorphic, but one can still apply the concept of holomorphic and anti-holomorphic to the \mathcal{Y} variables directly. Indeed, we encounter holomorphic (and anti-holomorphic) functions of \mathcal{Y}^{I} in the discussion of the entropy.

Non-extremal supersymmetric solutions In order to obtain non-extremal solutions relevant for the index, we replace the double pole behavior (4.10) of the extremal metric by two single poles. Accordingly, we split the sources into a pair of sources at two different points called the north and south pole. The total charge is split as

$$\gamma_{N} = \frac{1}{2} \left(P^{I} + in^{I}, Q_{I} + im_{I} \right), \qquad \gamma_{S} = \frac{1}{2} \left(P^{I} - in^{I}, Q_{I} - im_{I} \right), \qquad \Gamma = \gamma_{N} + \gamma_{S},$$
(4.11)

and the corresponding harmonic functions are given by

$$H(\mathbf{x}) = h + \frac{\gamma_N}{|\mathbf{x} - \mathbf{x}_N|} + \frac{\gamma_S}{|\mathbf{x} - \mathbf{x}_S|}.$$
 (4.12)

Compared to the extremal solution we have introduced electric and magnetic dipole charges (m_I, n^I) and the distance $|\mathbf{x}_N - \mathbf{x}_S|$ between the north and south poles. The equation for ω in (4.4) shows that the metric necessarily has angular momentum. All these parameters of the solution are determined by the *new attractor mechanism* [18] that we now recall.

The main point is that the spinning non-extremal solution near one of the poles has a simple scaling behavior, and the values of the scalars at the poles are therefore determined completely by the charges, similar to the extremal solution. There is, however, an important difference in that now we only have simple poles. As we briefly review below, one of the consequences is that the solution is necessarily complex.

The full solution has a symmetry which exchanges the north with the south pole and the scalars with their complex conjugates. Therefore we focus on the north pole and let the parameter $\rho = |\mathbf{x} - \mathbf{x}_N|$ be very small. From (4.12) we have, as $\rho \to 0$,

$$H(\mathbf{x}) \sim \frac{\gamma_N}{\rho} + H^{(0)} + O(\rho), \qquad H^{(0)} = h + \frac{\gamma_S}{|\mathbf{x}_N - \mathbf{x}_S|}.$$
 (4.13)

The generalized stabilization equations (4.6) then imply that

$$\mathcal{Y}^{I} \sim \frac{Y_{N}^{I}}{\rho} + Y^{I(0)} + O(\rho), \qquad \mathcal{G}_{I} \sim \frac{G_{NI}}{\rho} + G_{I}^{(0)} + O(\rho), \qquad (4.14)$$

and similarly for the conjugate scalars, with

$$(Y_N^I - \overline{Y}_N^I, G_{IN} - \overline{G}_{IN}) = i\gamma_N, \qquad (Y^{I(0)} - \overline{Y}^{I(0)}, G_I^{(0)} - \overline{G}_I^{(0)}) = iH^{(0)}.$$
 (4.15)

Note the Laurent expansion (4.14) combined with the definition of the scalar fields (4.5) and the homogeneity property of the prepotential leads to the following identity that we use below,

$$G_I^{(0)} = \frac{\partial \mathcal{G}_I}{\partial \mathcal{Y}^J}(\mathbf{x}_N) Y^{J(0)} \quad \Rightarrow \quad Y_N^I G_I^{(0)} = G_{JN} Y^{J(0)}.$$
(4.16)

Now, the equation (4.4) for the metric implies that

$$e^{-2U} \sim i\left(\frac{Y_N^I}{\rho} + Y^{I(0)}\right) \left(\frac{\overline{G}_{IN}}{\rho} + \overline{G}_I^{(0)}\right) - i\left(\frac{\overline{Y}_N^I}{\rho} + \overline{Y}^{I(0)}\right) \left(\frac{\overline{G}_{IN}}{\rho} + \overline{G}_I^{(0)}\right) + \mathcal{O}(\rho) \,. \tag{4.17}$$

Demanding a simple pole at \mathbf{x}_N implies that

$$\left(\overline{Y}_{N}^{I}, \overline{G}_{IN}\right) = 0 \quad \Rightarrow \quad \left(Y_{N}^{I}, G_{IN}\right) = i\gamma_{N}, \qquad (4.18)$$

where we use the first equation of (4.15) to obtain the second equality. The real part of these equations,

$$Y_N^I - (Y_N^I)^* = iP^I, \qquad G_{IN} - (G_{IN})^* = iQ_I,$$
(4.19)

are recognized as the extremal attractor equations (4.9). Assuming a unique solution, we obtain $(Y_N^I, G_{IN}) = (Y_*^I, G_{I*})$. The scalars at the north pole are thus fixed by the solution to the extremal attractor equations. Further, the dipole charges can then be read off to be

$$-n^{I} = Y_{N}^{I} + (Y_{N}^{I})^{*}, \qquad -m_{I} = G_{IN} + (G_{IN})^{*}.$$
(4.20)

The metric now only has a simple pole,

$$e^{-2U} \sim \frac{i}{\rho} \left(Y_N^I \overline{G}_I^{(0)} - G_{IN} \overline{Y}^{I(0)} \right) + O(1) ,$$

= $-\frac{1}{\rho} \left(Y_N^I H_I^{(0)} - G_{IN} \widetilde{H}^{I(0)} \right) + \frac{i}{\rho} \left(Y_N^I G_I^{(0)} - G_{IN} Y^{I(0)} \right) + O(1) ,$ (4.21)
= $-\frac{1}{\rho} \left(Y_N^I H_I^{(0)} - G_{IN} \widetilde{H}^{I(0)} \right) + O(1) .$

Here, we use the second equation in (4.15) to reach the second line, and the identity (4.16) in order to obtain the final equality.

Finally, the behavior of the connection ω near the poles is given by expanding the equation (4.4). Using (4.14) we obtain

$$\star d\omega = \frac{\widehat{n}}{\rho^2} \langle \gamma_N, H^{(0)} \rangle + \mathcal{O}(1/\rho), \qquad (4.22)$$

where $\hat{n} = (\mathbf{x} - \mathbf{x}_N)/|\mathbf{x} - \mathbf{x}_N|$, the three-dimensional unit vector with the north pole as origin. This equation implies that there is a three-dimensional Dirac string associated with ω emanating from the north pole. There is a similar condition at the south pole. The condition of smoothness implies that this Dirac string should end smoothly at the poles. This regularity condition leads to (see the discussion in [37, 38] for more details)

$$\frac{i\langle\gamma_N,\gamma_S\rangle}{|\mathbf{x}_N-\mathbf{x}_S|} + i\langle\gamma_N,h\rangle = \frac{\beta}{4\pi}.$$
(4.23)

This equation fixes the distance between the north and south poles in terms of the temperature, charges at the poles, and the asymptotic values of the moduli. As $\beta \to \infty$, we obtain the attractor solution with $\Omega \to 0$, $|\mathbf{x}_N - \mathbf{x}_S| \to 0$, and an infinite throat.

To summarize, we obtain a Euclidean metric with a cigar-like topology. The consistency of the periods of angular and time coordinates, leads to the condition $\beta\Omega = 2\pi i$. The values of the scalars at the tip of the cigar at the north and south poles are fixed by the charges in a manner similar to the extremal attractor mechanism, but now all the equations are split into a holomorphic piece at the north pole and an anti-holomorphic piece at the south pole. The holomorphic scalar fields take their attractor values at the north pole, and the anti-holomorphic scalars vanish. At the south pole we have the conjugate equations, and the anti-holomorphic scalars take their attractor values and the holomorphic scalars vanish.

4.3 Two-charge solution

With this set-up, we study the behavior of these $\frac{1}{2}$ -BPS rotating Euclidean solutions with two-charges. We begin by making some general comments that apply to any theory of $\mathcal{N} = 2$ supergravity. We will make contact with the black hole studied in sections 2 and 3 at the end. The classical on-shell action of the non-extremal supersymmetric solution reviewed above is given, after removing a factor of $\beta M_{\rm BPS}$ by a boundary term, by

$$-I_{\text{on-shell}} = \pi(q_I Y_N^I + 2i\mathcal{G}(Y_N)) + \pi(q_I \overline{Y}_S^I - 2i\overline{\mathcal{G}(Y_S)}), \qquad (4.24)$$

$$= \pi i \langle \gamma_N, \gamma_S \rangle, \qquad (4.25)$$

where \mathcal{G} is the same function as F, as discussed below (4.6). The result in the first line comes from showing that the on-shell action is a total derivative up to delta functions localized at the poles [18]. Using the new attractor equations it can be put in the form of the second line¹⁴. This quantity $\pi i \langle \gamma_N, \gamma_S \rangle$ is *not* the area of the horizon of the nonextremal solution, which is instead equal to

$$A = \beta |\mathbf{x}_N - \mathbf{x}_S|. \tag{4.26}$$

This can be easily derived (see [18]) from the metric (4.3) together with the equation for ω in (4.4) and we will not repeat it here. The on-shell action and the area only coincide at zero temperatures, such that $\log Z_{\text{index}}$ is proportional to the extremal area. This is due to the fact that these black holes satisfy the quantum statistical relation, i.e. the on-shell action is equal to $A/4 + 2\pi i J$ (since $\beta \Omega = 2\pi i$) and at low temperatures the angular momentum vanishes.

Given the discussion in the previous paragraph, in a general $\mathcal{N} = 2$ theory of supergravity, solutions that have vanishing area at extremality are characterized by

$$\langle \gamma_N, \gamma_S \rangle = 0. \tag{4.27}$$

This condition guarantees that the extremal area of the black hole is zero at the twoderivative level, and the index of the black hole is small. Since $\gamma_{N/S}$ are determined in terms of the solution to the extremal attractor equations, this can be rephrased as a condition on the choice of charges that the black hole carries. Finally, as emphasized above, the condition (4.27) still allows for the finite temperature supersymmetric solution of [18] to have a finite size horizon with area (4.26).

For the choice of charges that satisfy $\langle \gamma_N, \gamma_S \rangle = 0$, this might suggest that we have succeeded in finding a smooth solution that contributes to the gravitational path integral for the index. However, this is too quick. Let us recall the regularity condition at the horizon (4.23). When charges are chosen to satisfy $\langle \gamma_N, \gamma_S \rangle = 0$, the condition degenerates to:

$$\beta = 4\pi i \langle \gamma_N, h \rangle. \tag{4.28}$$

This is problematic for the following reasons

¹⁴The rewriting in the second line is only valid at the two-derivative level.

- 1. First, the temperature (4.28) is fixed in terms of the black hole charges and asymptotic moduli, while we should be free to set them independently.
- 2. The second problem is that $|\mathbf{x}_N \mathbf{x}_S|$ has disappeared completely from the equation. There is nothing fixing the separation between the two poles and therefore there is a moduli space of solutions, characterized for example by the area of the horizon. In particular $|\mathbf{x}_N - \mathbf{x}_S| = 0$ is part of the moduli space which gives rise to the original singular small black hole we were trying to avoid.
- 3. The extremal area is also proportional to $Z_*(\Gamma)\overline{Z}_*(\Gamma)$, which implies $Z_*(\Gamma) = 0$ for the small black hole. In the non-extremal solution Z = 0 and $\overline{Z} = 0$ at the north and south poles. The solution is then singular whenever $\langle \gamma_N, \gamma_S \rangle = 0$.

These are precisely the shortcomings enumerated in Section 3.3 for the two-charge black hole in the heterotic string. We see here all those features are universal to $\mathcal{N} = 2$ supergravity with vanishing extremal area at the two-derivative level.

To provide some more intuition we will consider some special cases, and connect to the black hole in Section 3. To leading order, compactifications of type II or heterotic string theory to four dimensions can be described by the following prepotential

$$F(X) = -\frac{1}{6} \frac{C_{IJK} X^{I} X^{J} X^{K}}{X^{0}}.$$
(4.29)

The value of C_{IJK} depends on the model. For example for type IIA compactified on $K3 \times T^2$ the coefficients C_{IJK} are the intersection numbers of a basis of 4-cycles. This case is related by duality to the toroidally compactified heterotic string theory in Sections 2 and 3. Instead of studying this theory we will analyze a simpler version where only X^1, X^2, X^3 are non-zero and moreover $X^2 = X^3$. Therefore consider a theory with $n_v = 2$ and a prepotential of the form

$$F(X) = -\frac{X^1 (X^2)^2}{X^0}.$$
(4.30)

This theory is equivalent to a sector of (2.1), as explained in [8]. We make more comments on this below. A concrete solution with vanishing extremal area involves turning on only the following two charges

$$Q_I = (Q_0, 0, 0), \qquad P^I = (0, P^1, 0).$$
 (4.31)

We take $Q_0 < 0$ and $P^1 > 0$. In the context of the toroidally compactified heterotic string these charges have the interpretation of momentum n and winding w along a circle $S^1 \subset T^6$, namely

$$Q_0 = -n, \qquad P^1 = w. (4.32)$$

Moreover the interpretation of the moduli is $X^1/X^0 = ie^{-\Phi}$ while X^2/X^0 parametrizes the volume of a cycle inside T^6 . The solution to the extremal attractor equations (4.9) is given by

$$Y_*^I = \left(0, \frac{iP^1}{2}, 0\right), \qquad G_{I*} = \left(\frac{iQ_0}{2}, 0, -\sqrt{P^1|Q_0|}\right).$$
(4.33)

This result should be surprising. Since $G_{2*} = -2Y_*^1Y_*^2/Y_*^0$, how come $X^0 = X^2 = 0$ and G_{2*} is finite? What is the physical value of the moduli X^2/X^0 that describes the geometry of T^6 ? To see how this comes about let us turn a small magnetic charge P^2 . (Physically the way this is regulated is by turning on $\tilde{H}^2 = h^2 \neq 0$ and going slightly away from the horizon, keeping $P^2 = 0$. In practice it is easier to turn on a small P^2 at the attractor point, both lead to the same answer.) Then the attractor equation becomes

$$Y_*^I = \left(\frac{P^1 P^2}{2\sqrt{P^1 |Q_0|}}, \frac{iP^1}{2}, \frac{iP^2}{2}\right).$$
(4.34)

We see that as $P^2 \to 0$ the ratio between Y^2_* and Y^0_* is finite

$$\frac{Y_*^2}{Y_*^0} \to i\sqrt{\frac{|Q_0|}{P^1}} = i\sqrt{\frac{n}{w}}.$$
(4.35)

In the application to the heterotic string this guarantees that near the horizon (or near \mathbf{x}_N and \mathbf{x}_S for the non-extremal metric) the volume of T^6 is finite, and also explains why G_{2*} is non-vanishing. The other moduli controlling the string coupling does diverge

$$\frac{Y_*^1}{Y_*^0} = i\sqrt{P^1|Q_0|} \frac{1}{P^2} \to i\sqrt{nw} \times \infty.$$
(4.36)

This divergence will be regulated by higher derivative corrections. In string theory this implies that the string coupling vanishes at the horizon (extremal) or at the poles $\mathbf{x}_{N/S}$ (non-extremal).

The new attractor equations imply that (Y_*^I, G_{I*}) and $(\bar{Y}_*^I, \bar{G}_{I*})$ are precisely the local charge at the north and south poles respectively

$$\gamma_N = \left(0, \frac{P^1}{2}, 0; \frac{Q_0}{2}, 0, i\sqrt{P^1|Q_0|}\right), \quad \gamma_S = (\gamma_N)^*.$$
 (4.37)

We can replace these local charges into equation (4.12). Once these harmonic functions are known, the full non-extremal solution can be constructed. Let us assume for simplicity that the moduli at infinity are chosen such that $H_1 = \tilde{H}^0 = 0$ and $\tilde{H}^2 = h^2$. We also choose $h_2 = 0$, although this does not imply $H_2 = 0$ since $\gamma_{N/S}$ include a dipole in this direction. For concreteness we also pick $X^1/X^0|_{\infty} = i2$ and $X^2/X^0|_{\infty} = i$, implying $h^1 = -h_0 = 1$ and $h^2 = 1/2$.¹⁵ The emblackening factor is

$$e^{-2U} = 2\tilde{H}^2 \sqrt{-\tilde{H}^1 H_0} \tag{4.38}$$

$$= \left\{ \left(1 + \frac{\frac{1}{2}P^{1}}{|\mathbf{x} - \mathbf{x}_{N}|} + \frac{\frac{1}{2}P^{1}}{|\mathbf{x} - \mathbf{x}_{S}|} \right) \left(1 + \frac{\frac{1}{2}|Q_{0}|}{|\mathbf{x} - \mathbf{x}_{N}|} + \frac{\frac{1}{2}|Q_{0}|}{|\mathbf{x} - \mathbf{x}_{S}|} \right) \right\}^{\frac{1}{2}}.$$
 (4.39)

Note that $e^{-2U} = -i\mathcal{Y}^1/\mathcal{Y}^0 = e^{-\Phi}$. Curiously this is independent of H_2 , which carries the information of the dipole. (Of course the dipole does appear in the scalars and gauge fields.) One can readily verify that the charge vector at the poles satisfies $\langle \gamma_N, \gamma_S \rangle = 0$.

¹⁵This specific choice of asymptotic moduli is done mainly to connect later to the solution in Section 3. The solution can be easily generalied to arbitrary $X^{1,2}/X^0|_{\infty}$.

All the previous considerations then apply to this black hole. A solution only exist with a unique temperature, it has a moduli space of solutions labeled by the horizon area, and the scalars are singular at the poles of the horizon.

Let us now show that this solution is precisely the same as the one analyzed in Section 3 earlier¹⁶. This has to be true since it was proven in [8] that the action (2.1) can be obtained from $\mathcal{N} = 2$ supergravity with prepotential (4.30), so we will be brief. Consider the metric given in (3.10) and change variables (t_E, r, θ, ϕ) to (t_E, x^1, x^2, x^3) according to

$$x^1 = \sqrt{r^2 - \mathsf{a}^2} \sin \theta \cos \phi \,, \tag{4.40}$$

$$x^2 = \sqrt{r^2 - \mathbf{a}^2} \sin \theta \sin \phi, \qquad (4.41)$$

$$x^3 = r\cos\theta. \tag{4.42}$$

In this coordinates the metric (3.10) has the form [26]

$$ds^{2} = e^{2\Phi} (dt_{E} + i\omega)^{2} + dx^{m} dx^{m}.$$
(4.43)

where $i\omega = am_0(r^2 - a^2\cos^2\theta)^{-1}r\sin^2\theta d\phi$ and the dilaton becomes

$$e^{-2\Phi} = \left(1 + \frac{m_0 e^{\alpha}}{2|\mathbf{x} - \mathbf{x}_N|} + \frac{m_0 e^{\alpha}}{2|\mathbf{x} - \mathbf{x}_S|}\right) \left(1 + \frac{m_0 e^{-\alpha}}{2|\mathbf{x} - \mathbf{x}_N|} + \frac{m_0 e^{-\alpha}}{2|\mathbf{x} - \mathbf{x}_S|}\right),$$
(4.44)

with $\mathbf{x}_N = (0, 0, \mathbf{a})$ and $\mathbf{x}_S = (0, 0, -\mathbf{a})$. This metric looks different than the one in (4.3). The solution is to recall (3.10) is given in string frame while the formalism used here for $\mathcal{N} = 2$ supergravity generates an action with canonical Einstein-Hilbert term. Therefore we should translate (3.10) into Einstein frame, which gives

$$e^{-\Phi} ds^2 = e^{\Phi} (dt_E + i\omega)^2 + e^{-\Phi} dx^m dx^m.$$
 (4.45)

It is possible now to put this metric in the form (4.3) with $2U = \Phi$ for a suitable choice of the constants h. This also implies that $P^1 = m_0 e^{\alpha}$ while $Q_0 = -m_0 e^{-\alpha}$. This is consistent with the interpretation of P^1 as winding and Q_0 as momentum. More importantly, the separation between the poles is identified with

$$|\mathbf{x}_N - \mathbf{x}_S| = 2a. \tag{4.46}$$

For example, (4.26) implies that the horizon metric in IWP form is $A = 2\beta a$ which matches the result in Section 3. Moreover, for our choice of h and γ_N we obtain $i\langle\gamma_N,h\rangle = \sqrt{P^1|Q_0|}/2 = m_0/2$ which implies from (4.28) a value of the temperature $\beta = 2\pi m_0$, matching Section 3 as well.

¹⁶The reader might wonder why we used two different formalism to describe the solution. The presentation in Section 3 is useful since it can be extended outside of the supersymmetric regime as in Section 2. The advantage of the formalism in this section is that, within supersymmetric solutions, it will allow us to incorporate higher derivative corrections, as in Section 5.

5 Higher derivative corrections to the gravitational index

In this section we present the rotating black hole solution including higher-derivative corrections in the $\mathcal{N} = 2$ supergravity coming from string theory. The higher-derivative corrections in $\mathcal{N} = 2$ supergravity are of two types: chiral superspace integrals of the holomorphic prepotential function, and full superspace integral, sometimes called F-terms and D-terms, respectively, in the $\mathcal{N} = 2$ context. We only consider the first type of terms, which are summarized by corrections to the tree-level holomorphic prepotential function $\mathcal{G}(\mathcal{Y}^I)$ discussed in the previous section. We then make comments on D-terms and the controllability of the solution.

The treatment begins by introducing another chiral field \hat{A} of weight 2, whose top component is proportional to the Weyl-squared tensor in four-dimensions. The prepotential is extended to a function $F(X^I, \hat{A})$ and, in addition to the first derivatives $F_I = \partial F / \partial X^I$, we have $F_{\hat{A}} = \partial F / \partial \hat{A}$. The particular form of the prepotential that arises in Calabi-Yau three-fold compactifications of Type II string theory is

$$F(X^{I}, \widehat{A}) = -\frac{1}{6} \frac{C_{IJK} X^{I} X^{J} X^{K}}{X^{0}} - c \frac{\widehat{A}}{64} \frac{X^{1}}{X^{0}} + \dots$$
(5.1)

Here \widehat{A} effectively counts the loops in the topological string expansion. As \widehat{A} contains four derivatives, we can also think of the series as an expansion in the number of derivatives.

The cubic or tree-level term was discussed in the previous section, and we have shown the one-loop term. Here $c = c_2/24$ where c_2 is the second Chern class of the Calabi-Yau. In addition there are typically an infinite series of terms given by the higher loops of the topological string expansion. In Type II theory on $K3 \times T^2$, the term X^1/X^0 above is completed by worldsheet instanton effects to a modular function of X^1/X^0 . Importantly, the expansion terminates at this order due to the fermion zero modes on T^2 .

In our case of interest here, i.e. the two-charge black hole, the two-derivative result is singular, so that in the full solutions the correction terms are as important as the twoderivative term. This means that, a priori, one does not have a controllable expansion. Nevertheless it is useful to see explicitly that including higher-derivative terms can lead to a smooth solution, as we show below.

A proof that D-terms vanish on configurations that preserve supersymmetry would justify the approach where we only keep F-terms.¹⁷ In some situations, like the attractor black hole entropy, one can indeed show that D-terms do not affect the quantity of interest [21, 22], and therefore one obtains an exact formula [39–42]. In our present context, as we see below, the entropy arising from the prepotential (5.1) agrees with the microscopic string theory index including the numerical coefficient, pointing to a non-renormalization result.

The analysis of supersymmetry equations and solutions of the higher-derivative theory was performed in the impressive set of papers [36, 43, 44]. Just as we define the rescaled

¹⁷As opposed to previous work [8–10], we do not expect the generic expansion to be controllable for the thermal partition function which does not include an insertion of $(-1)^{\mathsf{F}}$.

fields $(\mathcal{Y}^I, \mathcal{G}_I)$ in (4.5), we also define a rescaled field $\Upsilon = e^{\mathcal{K}(X,\overline{X})} \overline{Z}^2 \widehat{A}$. The higherderivative equations are, generally speaking, more complicated than the two-derivative theory. The exception is the equations governing the scalar fields, which are still given by the generalized stabilization equations (4.6), although they now depend on Υ .

New attractor with higher-derivative corrections

The solution is specified by a vector of harmonic functions $H = (\tilde{H}^I, H_I)$. The generalized stabilization equations are now

$$\mathcal{Y}^{I} - \overline{\mathcal{Y}}^{I} = i\widetilde{H}^{I}, \qquad \mathcal{G}_{I}(\mathcal{Y}, \Upsilon) - \overline{\mathcal{G}}_{I}(\mathcal{Y}, \Upsilon) = iH_{I}, \qquad (5.2)$$

and they determine \mathcal{Y}^I in terms of H and Υ . The metric is of the IWP form as in the two-derivative theory

$$ds^{2} = e^{2U} (dt_{E} + i\omega_{m} dx^{m})^{2} + e^{-2U} dx^{m} dx^{m}.$$
(5.3)

The functions U and ω are now governed by

$$e^{-2U} = i \left(\mathcal{Y}^{I} \overline{\mathcal{G}}_{I} - \mathcal{G}_{I} \overline{\mathcal{Y}}^{I} \right) + 128i e^{U} \nabla^{p} \left((\nabla_{p} e^{-U}) (\mathcal{G}_{\Upsilon} - \overline{\mathcal{G}}_{\Upsilon}) \right) - 32i e^{4U} (R(\omega)_{p})^{2} (\mathcal{G}_{\Upsilon} - \overline{\mathcal{G}}_{\Upsilon}) - 64e^{2U} R(\omega)_{p} \nabla^{p} (\mathcal{G}_{\Upsilon} + \overline{\mathcal{G}}_{\Upsilon}) .$$
(5.4)

$$-iR(\omega)_p = H_I \overleftrightarrow{\nabla}_p H^I - 128i \nabla^q \left[\nabla_{[p} (e^{2U} R(\omega)_{q]} (G_{\Upsilon} - \overline{G}_{\Upsilon})) \right] - 128 \nabla^q \left(2\nabla_{[p} U \nabla_{q]} (G_{\Upsilon} + \overline{G}_{\Upsilon}) \right).$$
(5.5)

In the two-derivative theory $\mathcal{G}_{\Upsilon} = 0$ and the above equations reduce to the simpler equations (4.4). Finally, the value of Υ is

$$\Upsilon = -64 \left(\nabla_p U - \frac{i}{2} e^{2U} R(\omega)_p \right)^2.$$
(5.6)

Here we follow the notation of [43] with ∇_a the derivative in the three-dimensional flat base-space, and $R(\omega)^p = \varepsilon^{pqs} \nabla_q \omega_s$. For the non-extremal rotating saddle we set, as in Section 4

$$H(\mathbf{x}) = h + \frac{\gamma_N}{|\mathbf{x} - \mathbf{x}_N|} + \frac{\gamma_S}{|\mathbf{x} - \mathbf{x}_S|}, \qquad \gamma_N + \gamma_S = \Gamma.$$
(5.7)

Therefore part of the problem is to determine γ_N , γ_S and $|\mathbf{x}_N - \mathbf{x}_S|$ using smoothness, just like we did at the two derivative level [18].

As an exercise we can study the behavior of all the fields in an extremal attractor background as in the discussion near (4.7), as $r \to 0$,

Extremal:
$$R(\omega) = 0$$
, $\mathcal{Y}^I \sim \frac{Y_*}{r} + \dots$, $e^{-2U} \sim \frac{a}{r^2} + \dots$, $\Upsilon \sim \frac{\Upsilon_*}{r^2} + \dots$ (5.8)

The fact that $U \sim \log r$ together with (5.6) implies that $\Upsilon_* = -64$. The scalars Y_*^I are determined by $Y_*^I - \bar{Y}_*^I = iP^I$ and $G_{I*} - \bar{G}_{I*} = iQ_I$. The scalar fields therefore behave as in the two-derivative theory. The scaling behavior of the other fields can be read off from

their conformal weights—this can be explained either by using a conformal compensator, or by thinking of the other fields as functions of the scalar fields. The only thing to be careful of is that coefficients can be zero, as happens for $R(\omega)$ because of the spherical symmetry. The value $\Upsilon_* = -64$ is precisely the one required for the Wald entropy of the extremal black hole to equal the logarithm of the growth of microscopic states of the fundamental heterotic string [45].

Now we repeat the analysis for the rotating non-extremal solution, and determine the value of $\gamma_{N/S}$. We focus on the behavior of the solution near the north pole, since the behavior at the south pole is related by complex conjugation. Recall that the idea in the rotating non-extremal solution is to demand that the metric factor e^{-2U} has only a single pole, leading to a finite temperature horizon. Accordingly, we assume the following behavior for the non-extremal solution, as $\rho = |\mathbf{x} - \mathbf{x}_N| \to 0$,

$$\mathcal{Y}^{I} \sim \frac{Y_{N}}{\rho} + \dots, \qquad e^{-2U} \sim \frac{a_{1}}{\rho} + \dots, \qquad R(\omega)_{p} = \frac{a_{2}}{\rho^{2}} \,\widehat{n} + \dots, \qquad \Upsilon \sim \frac{\Upsilon_{*}}{\rho^{2}} + \dots,$$
(5.9)

where $\hat{n} = (\mathbf{x} - \mathbf{x}_N)/|\mathbf{x} - \mathbf{x}_N|$. Now we analyze the equations (5.4)–(5.6). The upshot is that the solution to the extremal attractor (Y_*^I, G_{I*}) together with $\Upsilon_* = -64$ determines γ_N . Together with the smoothness condition, this completely determines the sources H. Now the full solution can be determined in terms of the boundary data (Γ and β)¹⁸.

We begin with (5.5). \mathcal{G}_{Υ} has Weyl weight 0 and so $\mathcal{G}_{\Upsilon} \sim O(1)$ as $\rho \to 0$. The leading behavior of the second term on the right hand side is therefore proportional to $\nabla_{[p}(e^{2U}R(\omega)_{q})$. Now, $e^{2U}R(\omega)_q \sim \frac{a_2}{\rho}\hat{n}_q + \ldots$, and therefore the anti-symmetry of the action of the derivatives annihilates this term so as to give no contribution at $O(1/\rho^2)$. A similar analysis applies to the last term in (5.5). This shows that the value of $R(\omega)$ at the pole in the higher-derivative theory is the same as its value given in (4.22) in the two-derivative theory. We conclude that $a_2 = i\langle \gamma_N, H^{(0)} \rangle$. In particular, this means that the condition of there being no Dirac-string singularity implies the same relation (4.23) as in the two-derivative theory, which we repeat here for convenience

$$\frac{i\langle\gamma_N,\gamma_S\rangle}{|\mathbf{x}_N-\mathbf{x}_S|} + i\langle\gamma_N,h\rangle = \frac{\beta}{4\pi}.$$
(5.10)

Now we move to (5.4). Let us start by considering the first term on the right hand side. This is the value of e^{-2U} in the two-derivative theory. We consider a solution that cancels the double pole coming from this term by imposing $\bar{Y}_N^I = \bar{G}_{IN} = 0$ (and $Y_S^I = G_{IS} = 0$ for the south pole). The stabilization equation, together with the old attractor, would then imply

$$i\gamma_N = (Y^I_*, G_{I*}).$$
 (5.11)

This solution is not yet complete, as we have not yet determined Υ_* which appears implicitly in G_I . We will come back to this point shortly. First, we explain how the other

¹⁸We assume that, given H, a solution to the partial differential equations (5.4) and (5.5) exists and is unique. This has been verified in the low temperature limit numerically.

terms in (5.4) lead to a vanishing double pole. As $\rho \to 0$, calling the constant $\mathcal{G}_{\Upsilon} = C_1$, the leading behavior of the second and third terms is governed by

$$C_1 \left(128i \, e^U \nabla^2 e^{-U} - 32i \, e^{4U} (R(\omega)_p)^2 \right) \sim -32i \, C_1 \left(1 + \frac{a_2^2}{a_1^2} \right) \frac{1}{\rho^2} + \dots$$
 (5.12)

We see that in order to have a vanishing double pole we must have $a_2 = \pm ia_1$. In the two-derivative theory we have, from (4.21) and (4.18), the value $a_1 = -\langle \gamma_N, H^{(0)} \rangle$ so that $a_2 = ia_1$. We choose this solution also in the higher-derivative theory. This implies a condition of vanishing of other O(1/ ρ) terms of the right-hand side of (5.4) involving higher derivative terms. From the equation (5.6), we deduce that

$$\Upsilon \sim -\frac{64}{\rho^2} \left(\frac{1}{2} - \frac{i}{2} \frac{a_2}{a_1} \right) = -\frac{64}{\rho^2}.$$
(5.13)

This completes the derivation since now we determined $\Upsilon_* = -64$. As in the two-derivative solution, we see that the holomorphic (anti-holomorphic) scalar fields take the attractor values at the north (south) pole. Furthermore they determine the dipole charges γ_N (γ_S) through the relation (5.11).

So far we described the metric and scalar profiles. The gauge fields are also determined by the harmonic sources through the equations $F_{0p}^{I} = -\nabla_{p}[e^{2U}(\mathcal{Y}^{I} + \bar{\mathcal{Y}}^{I})]$ and $G_{0p}^{I} = -\nabla_{p}[e^{2U}(\mathcal{G}_{I} + \bar{\mathcal{G}}_{I})]$ where $F_{\mu\nu}^{I}$ and $G_{\mu\nu}^{I}$ determine the field strength and its dual. These equations are the same as in the two-derivative theory.

To conclude the characterization of the solution, we propose a formula for the onshell action of such configurations, including boundary terms that remove the zero-point energy $\beta M_{\rm BPS}$, to be

$$-I_{\text{on-shell}} = \pi(q_I Y_N^I + 2i\mathcal{G}(Y_N)) + \pi(q_I \overline{Y}_S^I - 2i\overline{\mathcal{G}(Y_S)}).$$
(5.14)

We have two reasons that justify this expression. First, it reduces to the two-derivative action (4.24) when \mathcal{G} is independent of Υ , for any temperature. (Note that when \mathcal{G} depends on Υ (4.25) is different from (4.24).) Second, one can prove this formula in the presence of arbitrary higher-derivative corrections, under the assumption that the on-shell action is temperature independent. The action in the zero-temperature limit was explicitly evaluated in [39] and matches (5.14) (as well as the Wald entropy derived in [43]).

The reason we have not attempted an explicit evaluation of the action is that it requires solving the differential equations (5.4) and (5.5). Nevertheless, there might be a more elegant way to obtain (5.14). The split form of the action in (5.14) is similar to observations of the factorization of the partition function in [46, 47]. An efficient way to prove this could be to apply the techniques of [48], we leave this for future work.

The corrected two-charge solution

We now come back to the specific theory relevant to the two-charge black hole in heterotic string theory, with only $X^1, X^2 = X^3$, with prepotential

$$F(X^{I}, \widehat{A}) = -\frac{X^{1}(X^{2})^{2}}{X^{0}} - c\frac{\widehat{A}}{64}\frac{X^{1}}{X^{0}}.$$
(5.15)

For the heterotic string c = 1 but we find it useful to keep it general to keep track of the effect of higher derivative corrections. The non-zero monopole charges in the solution are taken to be Q_0, P^1 and recall that $Q_0 < 0$ and $P^1 > 0$. According to the discussion in Sections 4.2, 4.3, we choose sources $(\tilde{H}^0, \tilde{H}^1, \tilde{H}^2, H_0, H_1, H_2)$. Let us begin by solving the new attractor equations which determine $\gamma_{N/S}$. The starting point is the extremal attractor solution

$$Y_*^I = \left(\sqrt{\frac{cP^1}{|Q_0|}}, \frac{iP^1}{2}, 0\right), \qquad G_{I*} = \left(\frac{iQ_0}{2}, -\sqrt{\frac{cP^1}{|Q_0|}}, 0\right), \qquad \Upsilon_* = -64, \quad (5.16)$$

which is equal to Y_N at the north pole while $\bar{Y}_N = 0$. The divergence in the dilaton (near the north and south pole) at the horizon is now regulated since Y^0_* is finite. The fact that now $G_{2*} = 0$ and $G_{1*} \neq 0$ might seem like a major modification compared to the twoderivative level discussion. To see how this comes about one can turn a small charge P^2 (or, equivalently, go slightly away from the pole, as explained in Section 4.3). For example, this produces a term $G_{2*} \sim \sqrt{P^1 |Q_0|} \frac{P^2}{4c+(P^2)^2}$. We see explicitly that the limits $c \to 0$ and $P^2 \to 0$ do not commute and this is the source of discrepancy. This is clearly due to the singular nature of the two-derivative solution and would not happen for a big black hole.

According to the new attractor solution (5.11) the charge vectors associated to the north and south pole contributions are

$$\gamma_N = \left(i\sqrt{\frac{cP^1}{|Q_0|}}, \frac{P^1}{2}, 0, \frac{Q_0}{2}, -i\sqrt{\frac{cP^1}{|Q_0|}}, 0\right)$$
(5.17)

and $\gamma_S = (\gamma_N)^*$. Notice again that this is not a small correction to the two-derivative solution for γ_N ! One can check that now we have

$$i\langle\gamma_N,\gamma_S\rangle = 2\sqrt{cP^1|Q_0|}.$$
(5.18)

The final ingredient to determine the sources (5.7) and therefore the full solution is to choose h. At infinity we have $\Upsilon \to 0$ and therefore we can repeat the same analysis as in the two-derivative theory to establish the values $h^I = \tilde{H}^I|_{\infty}$ and $h_I = H_I|_{\infty}$ that allow us to compare with the solution in Sections 2 and 3. The answer found in Section 4.3 is $h_I =$ (-1, 0, 0) and $h^I = (0, 1, \frac{1}{2})$. The solution to the generalized stabilization equations (4.6) with prepotential (5.1) becomes then an algebraic equation that relates $(\mathcal{Y}^I, \mathcal{G}_I)$ to (\tilde{H}^I, H_I) and Υ , which can be solved easily¹⁹.

The relation (4.23) fixes the parameter $2\mathbf{a} = |\mathbf{x}_N - \mathbf{x}_S|$ in terms of β , as anticipated earlier in Section 3. Since, in the language of that section, $i\langle\gamma_N,\gamma_S\rangle = 2\sqrt{cP^1|Q_0|} = 2\sqrt{cm_0}$ and $i\langle\gamma_N,h\rangle = \frac{\sqrt{c(P^1-Q_0)}}{\sqrt{P^1|Q_0|}} = 2\sqrt{c}\cosh\alpha$, the relation (4.23) gives

$$\frac{2\sqrt{c}m_0}{2\mathsf{a}} + 2\sqrt{c}\cosh\alpha = \frac{\beta}{4\pi}, \qquad \Rightarrow \quad \mathsf{a} = \frac{4\pi m_0}{\frac{\beta}{\sqrt{c}} - 8\pi\cosh\alpha}. \tag{5.19}$$

¹⁹It is important to emphasize that the stabilization equations are not enough to determine \mathcal{Y} in terms of H in the higher-derivative theory. The reason is that they involve Υ , which depends on U and ω . Therefore the stabilization equations are not decoupled from (5.4) and (5.5).

This relation resolves the problem appearing in the two-derivative expression (4.28), namely that the temperature was apparently fixed in terms of the charges and asymptotic moduli. Now we see that the temperature and the moduli are independent quantities. The relation (5.19) shows the solution is only valid as long as the denominator is positive. Curiously, the denominator changes sign at a finite value of β . (The same phenomenon is found even for the big black hole where $|\mathbf{x}_N - \mathbf{x}_S|$ becomes negative above a charge-dependent temperature. Explaining this remains an open question.)

In summary, we find a solution that is smooth with the dilaton finite everywhere, all the parameters of the solution are determined in terms of boundary data which are independent values of the temperature and moduli.

To conclude this section we evaluate the on-shell action of the solution we constructed, using (5.14). Since $Z_{\text{index}} \sim e^{-I_{\text{on-shell}}}$ we find that it makes the following contribution to the gravitational index

$$\log Z_{\text{index}} = \pi (q_I Y_N^I + 2i\mathcal{G}(Y_N)) + \pi (q_I \overline{Y}_S^I - 2i\overline{\mathcal{G}(Y_S)}), \qquad (5.20)$$

$$= 4\pi \sqrt{cP^1 |Q_0|} \,. \tag{5.21}$$

This is now finite. To compare with the heterotic string action we recall that c = 1. Therefore we find a gravitational index given, in terms of the charges $n = -Q_0$ and $w = P^1$, by

$$\log Z_{\text{index}} = -I_{\text{on-shell}} = 4\pi\sqrt{nw}.$$
(5.22)

(In terms of the parameter m_0 of Section 3 this is $4\pi m_0$.) This result precisely matches the microscopic evaluation of the index counting BPS fundamental heterotic strings, as reviewed in the introduction.

Note that the same phenomenon is known to occur if one assumes an attractor AdS_2 geometry [45]. What is remarkable here is that one obtains the same result even though the rotating geometry for the index preserves much less symmetry, and is compatible with the discussion in Section 2 on the partition function.

6 Can the low-temperature physics be nearly conformal?

In Section 2 we studied two-charge black holes in string theory. At low temperatures the black hole becomes small, i.e. it has a horizon area that vanishes at the two-derivative level. Following [12], we explained how a transition takes place to a winding condensate without an event horizon, before we reach the small black hole, and ultimately the ground state is described by a gas of strings and not a black hole.

Is the behavior described above universal? Could there be a string theory solution that does not transition to a gas of strings or other stringy objects without a horizon, such that the ground state is really described by a small black hole? In this section we use effective theory arguments to comment on the possibility of a string-size near-extremal black hole. Our arguments also apply for big black holes with 16 or more supercharges.

By a small black hole we mean a gravitational solution with a finite (string scale) area that exists at very low temperatures, which develops an AdS_2 throat. This assumption can be justified in some cases where there is a smooth limit to the two-derivative theory [49, 50]. We also assume that the index is a function that grows exponentially with the charges. The latter requirement is satisfied by most examples in string theory (an exception being e.g. the D0 brane black hole). Finally, we assume that the extremal limit of the small black hole preserves at least 16 supercharges.

The near-extremal AdS_2 phase of such black holes implies the presence of a nearlyconformal symmetry of the quantum system describing the black hole microstates. Its low energy description is given by the Schwarzian theory. Since we assume the system preserves a certain number of supercharges, we have a supersymmetric generalization of the Schwarzian theory.

It is possible to label the different supersymmetric extensions of the Schwarzian theory by the superconformal group of symmetries broken by the near-AdS₂ solution. For example the bosonic Schwarzian theory would be labeled by $PSL(2, \mathbb{R})$, the one describing the near-BPS limit of the 1/16-BPS black holes in AdS₅ would be labeled by SU(1, 1|1) and the one describing the near-BPS limit of large black holes in flat space by PSU(1, 1|2). Therefore we can address this problem by analyzing the possible superconformal groups. To perform our analysis we use that the supersymmetric Schwarzian action acts as a U(1) generator in the space of super-reparametrization, such that we can apply the localization principle of [51]. We assume that the R-symmetry of the superconformal group is realized in a geometric manner in spacetime.²⁰ We now show that no Schwarzian theory exists with these properties.

Given the choice of a superconformal group we can derive the partition function of such a theory including quantum corrections. These corrections are one-loop exact and have the following form. Denote by $\vec{\mu}$ the vector of chemical potentials that couple to the $r = \operatorname{rank}(G)$ Cartan generators of the *R*-symmetry group *G*, with the periodicity $\mu \sim \mu + 1$. The partition function then takes the form

$$Z = \sum_{\vec{n} \in \mathbb{Z}^r} \beta^{\frac{N_F - N_B}{2}} Z_{\text{one-loop}}(\vec{\mu} + \vec{n}) e^{S_0 + \frac{1}{\beta}I(\vec{\mu} + \vec{n})}, \qquad (6.1)$$

where $Z_{\text{one-loop}}(\vec{\mu} + \vec{n})$, and $I(\vec{\mu} + \vec{n})$, are functions of chemical potentials only arising from the one loop determinant, and on-shell action, respectively. The sum over integers \vec{n} is a sum over saddles related by large gauge transformations. S_0 is the extremal entropy while N_F and N_B correspond to the number of fermionic generator and bosonic generators in the superconformal group. The overall prefactor of $\beta^{\frac{N_F - N_B}{2}}$ in the one-loop determinant comes from subtracting a contribution from zero-modes.

When $N_F \ge N_B$ the theory can predict the presence of a large number of ground states of order e^{S_0} . For example for SU(1,1|1), $N_B - N_F = 0$ so that the one-loop determinant is temperature independent and in the zero-temperature limit the action becomes temperature independent, resulting in $Z(\beta \to \infty) \sim e^{S_0} + O(e^{-\beta})$. For PSU(1,1|2), instead, $N_F = 8 > N_B = 6$ and the one-loop determinant diverges as $\beta \to \infty$. In this case the

²⁰These assumptions do not rule out the possibility of a sigma model description. See [52–56] for a related discussion on the AdS_3 near-horizon configurations of a fundamental string.

sum over saddles is important in order to regulate this divergence and once again we find $Z(\beta \to \infty) \sim e^{S_0}$.

When $N_B > N_F$ the one-loop determinant vanishes in the $\beta \to \infty$ limit and its unavoidable to obtain $Z(\beta \to \infty) = 0$. This implies that the quantum corrections due to the Schwarzian mode would destroy the classical degeneracy of ground states. In other words $\log Z \sim S_0 - \frac{N_F - N_B}{2} \log \beta \to -\infty$ in the large β limit. Since the index gives a lower bound on the number of ground states, we can use this criterion to rule out the existence of a small black hole horizon with an AdS₂ factor. In other words, there is no mass-gap and therefore no decoupled AdS₂ geometry in such theories.

By the arguments in the previous paragraph we reduced the problem to an analysis of the number of bosonic vs fermionic generators of superconformal groups [57]. The list of such groups can be found e.g. in Table 1 of the review [58], see [59] for an extensive list.

Let us begin with superconformal groups with more than 16 supercharges. There are three possible families. The first family are SU(1, 1|n) with n > 4, which has a SU(n)×U(1) *R*-symmetry. This group has $N_B = n^2 + 3$ and $N_F = 4n$. For all n > 1 one has $N_B > N_F$ and therefore the ground states do not survive quantum corrections. The same is true for OSp(n|2) for n > 8 since $N_B = \frac{1}{2}(n^2 - n) + 3$ and $N_F = 2n$ and for $OSp(4^*|2n)$ with n > 2which has $N_B = 2n^2 + n + 6$ and $N_F = 8n$. Therefore, if an index predicts a large ground state degeneracy, those states cannot describe small black holes preserving more than 16 supercharges.

Next, we consider the case of systems with 16 supercharges. In this case there are four superconformal groups, which are

Superconformal Group	N_B/N_F	<i>R</i> -symmetry	
$OSp(4^* 4)$	16/16	$SU(2) \times Spin(5)$	
$\mathrm{SU}(1,1 4)$	19/16	$\mathrm{SU}(4) \times \mathrm{U}(1)$	(6.2)
F(4)	24/16	SO(7)	
OSp(8 2)	31/16	SO(8)	

The last three of these groups have $N_B > N_F$ and therefore do not have a ground state degeneracy. The case $OSp(4^*|4)$ is special since $N_B = N_F$. Here, the Spin(5) R-symmetry does not act as a geometric rotation of any subspace of $T^{6,21}$

Thus, there is no superconformal group, and therefore no Schwarzian theory, which could describe the low energy phase of such small black holes. This argument can be extended to theories with 14 supercharges. Below 14 there are multiple possibilities of nearlyconformal theories with a large number of ground states such as SU(1, 1|1) or PSU(1, 1|2), and other superconformal groups which haven't found an application yet. Another theory that deserves further study along these lines is $D(2, 1; \alpha)$.

²¹It is possible that the supercharges transform as a spinor of the tangent space Spin(5). This possibility was discussed in related AdS_3 near-horizon configurations in [53].

7 Discussion

We would like to conclude with some comments and future directions.

We have evaluated the gravitational path integral that corresponds to the index of the two-charge states in heterotic string theory. This calculation involved constructing a solution that combines two-derivative with four-derivative interactions. An obvious future direction is to make this construction more precise. The advantage of the present paper compared with previous attempts is that we turned this question into a concrete program, starting with proving that D-terms vanish due to supersymmetry. Besides this, it would be necessary to prove that the action coming from F-terms is temperature independent and reproduces (5.14) for any prepotential.

Another natural direction is to compute quantum corrections around the black hole solution evaluating the index. We hope that one can use the recent techniques in [60] and [61] to reproduce the logarithmic correction to the index in the large $N = \sqrt{nw}$ limit. An even more ambitious goal can be to use supergravity localization to reproduce the exact generating function $1/\eta^{24}(\tau)$.

Finally, it is interesting to consider whether this approach can be extended to black holes with one charge. An example is a 10 dimensional type IIA string theory black hole with D0-charge N. This leads to a small black hole in the extremal limit and its nearextremal excitations are dual to the BFSS matrix model. In this case even a microscopic calculation of the index is hard and has only been completed for the N = 2 theory [62], although there is a simple M-theory prediction for it for all N. To attack this problem using the gravitational path integral, one can easily find a rotating D0-brane black hole starting with the uncharged Myers-Perry solution in 10 dimensions (t, x^1, \ldots, x^9) , uplifting to 11 dimensions by adding an extra circle x^{10} , performing a boost in the $t - x^{10}$ coordinates, and finally dimensionally reducing back to 10 dimensions. Imposing the boundary conditions relevant for the index one can find, at the two-derivative level, a supersymmetric rotating non-extremal black hole with a macroscopic area and zero on-shell action, similar to its two-charge counterpart of Section 3 [63]. The solution is singular at the poles of the horizon and the string coupling becomes large at those locations. An evaluation of the index of the BFSS model using the gravitational path integral for type IIA seems to require a new idea and cannot be computed by a simple extension of the techniques used here.

Acknowledgements

It is a pleasure to thank Pietro Benetti-Genolini, Jan Boruch, Matt Heydeman, Luca Iliesiu, Gabriel Lopes Cardoso, Juan Maldacena, Thomas Mohaupt, Ashoke Sen, and Edward Witten for interesting and useful discussions. Y.C. acknowledges the KITP program "What is String Theory? Weaving Perspectives Together", during which this work is completed. S.M. acknowledges the support of the J. Robert Oppenheimer Visiting Professorship at the Institute for Advanced Study, Princeton and the STFC grants ST/T000759/1, ST/X000753/1. The work of GJT is supported by the University of Washington and the DOE award DE-SC0024363.

A Solution generation for rotating two-charge systems

In this appendix, we discuss how one can use the solution generating procedure to work out the properties of the rotating two-charge system from the uncharged (but rotating) solution. We will derive the thermodynamic quantities of the charged solution with the information of the thermodynamic quantities of the uncharged solution as well as the transformation at infinity. The derivation applies to general classical solutions in string theory, including black hole solutions as well as potential rotating Horowitz-Polchinski solutions that have not yet been constructed. One application of our derivation, which we discuss in appendix. A.1, is that one can work out the α' corrections to the rotating two-charge black hole in [27], without needing to find the corrected geometry explicitly. The discussion here mostly follows the derivation in [12], with the extra ingredient that the seed solution can carry rotation.

The starting point is a seed solution in D dimensions which has an asymptotic time circle with length $\tilde{\beta} = 2\pi \tilde{R}$. For the purpose of applying to the discussion in the main text, we have D = 4, though the discussion is more general. We will assume the existence of a U(1) isometry, time translation, everywhere in the solution. For simplicity, we consider the transformations that generate charges along one extra circle, and the relevant symmetry group is O(2, 2) which acts on fields in d = D - 1 dimensional space once we dimensional reduce on both the time circle and the extra circle. This is a subgroup of the O(7, 23)transformation of heterotic string theory compactified on T^6 we considered in the main text, but it also applies to the Bosonic/Type II theories compactified on a torus. We parametrize the transformation by the parameters (α, β).

The main equation, which we refer to [12] for justification, is the invariance of the action under the solution generating transformation.

$$\log Z(R,\Omega,r,\mu_n,\mu_w,\Phi_D) = \log Z(R,\Omega,\Phi_D).$$
(A.1)

Let's unpack this equality by explaining the notations. All the tilded quantities refer to the seed solution, which has an angular potential $\tilde{\Omega}$ other than the inverse temperature (over 2π) \tilde{R} and asymptotic value of the dilaton $\tilde{\Phi}_D$. The quantities without tildes are those for the generated solution, which also has r being the size of the extra circle, and μ_n, μ_w being the chemical potentials for the momentum and winding charges Q_n and Q_w . We will fix r = 1 in the following discussion.²² We can factor out the dependence of the dilaton and define the primed quantities²³

$$e^{-\Phi_D}\log Z'(R,\Omega,\mu_n,\mu_w) \equiv \log Z(R,\Omega,\mu_n,\mu_w,\phi_D), \quad e^{-\bar{\Phi}_D}\log \tilde{Z}'(\tilde{R},\tilde{\Omega}) \equiv \log \tilde{Z}(\tilde{R},\tilde{\Omega},\tilde{\Phi}_D)$$
(A.2)

Using the invariance of the d = D - 1 dimensional dilaton, the asymptotic value of the dilaton transforms as

$$e^{-\Phi_D}\tilde{R} = e^{-\Phi_D}R\tag{A.3}$$

 $^{^{22}}$ This is only valid since we are using r to denote the asymptotic size of the circle. Inside the spacetime it will be varying.

 $^{^{23}}$ For the dilaton, here we follow the convention used in the main text, see (2.1).

and therefore from (A.1) we have

$$\log Z' = \frac{R}{\tilde{R}} \log \tilde{Z}'. \tag{A.4}$$

For the generated solution, we have the general statistical expression for the action

$$\log Z = e^{-\Phi_D} \log Z' = S - 2\pi R M + 2\pi R \Omega J + 2\pi R \mu_n Q_n + 2\pi R \mu_w Q_w$$
(A.5)

where various charges and entropy can be computed as

$$2\pi RQ_n = e^{-\Phi_D} \partial_{\mu_n} \log Z', \quad 2\pi RQ_w = e^{-\Phi_D} \partial_{\mu_w} \log Z', \quad 2\pi RJ = e^{-\Phi_D} \partial_\Omega \log Z',$$

$$2\pi M = e^{-\Phi_D} \left(-\partial_R + \frac{\mu_n}{R} \partial_{\mu_n} + \frac{\mu_w}{R} \partial_{\mu_w} + \frac{\Omega}{R} \partial_\Omega \right) \log Z', \quad S = e^{-\Phi_D} (1 - R\partial_R) \log Z'.$$
(A.6)

On the other hand, for the seed solution, we have

$$\log \tilde{Z}' = \tilde{S}' - 2\pi \tilde{R}\tilde{M}' + 2\pi \tilde{R}\tilde{\Omega}\tilde{J}',$$

$$2\pi \tilde{R}\tilde{J}' = \partial_{\tilde{\Omega}}\log \tilde{Z}', \quad 2\pi \tilde{M}' = \left(-\partial_{\tilde{R}} + \frac{\tilde{\Omega}}{\tilde{R}}\partial_{\tilde{\Omega}}\right)\log \tilde{Z}', \quad \tilde{S}' = \left(1 - \tilde{R}\partial_{\tilde{R}}\right)\log \tilde{Z}'.$$
(A.7)

Note that here similar to \tilde{Z}' we have also defined primed thermodynamic quantities which are simply themselves but without the dilaton dependence (for example, $\tilde{S} = e^{-\tilde{\Phi}_D}\tilde{S}'$). Now, combining the expression for Q_n in (A.6) with (A.7), we get

$$2\pi R Q_n = e^{-\Phi_D} \partial_{\mu_n} \left(\frac{R}{\tilde{R}} \log \tilde{Z}' \right) = e^{-\Phi_D} R \left[\frac{\partial \tilde{R}}{\partial \mu_n} \partial_{\tilde{R}} \left(\frac{1}{\tilde{R}} \log \tilde{Z}' \right) + \frac{1}{\tilde{R}} \frac{\partial \tilde{\Omega}}{\partial \mu_n} \partial_{\tilde{\Omega}} \left(\log \tilde{Z}' \right) \right]$$
$$= e^{-\Phi_D} R \left[-\frac{1}{\tilde{R}^2} \frac{\partial \tilde{R}}{\partial \mu_n} \tilde{S}' + 2\pi \frac{\partial \tilde{\Omega}}{\partial \mu_n} \tilde{J}' \right].$$
(A.8)

Similar derivations lead to

$$2\pi Q_w = e^{-\Phi_D} \left[-\frac{1}{\tilde{R}^2} \frac{\partial \tilde{R}}{\partial \mu_w} \tilde{S}' + 2\pi \frac{\partial \tilde{\Omega}}{\partial \mu_w} \tilde{J}' \right], \tag{A.9}$$

$$2\pi J = e^{-\Phi_D} \left[-\frac{1}{\tilde{R}^2} \frac{\partial \tilde{R}}{\partial \Omega} \tilde{S}' + 2\pi \frac{\partial \tilde{\Omega}}{\partial \Omega} \tilde{J}' \right], \qquad (A.10)$$

$$S = e^{-\Phi_D} \left[\frac{R^2}{\tilde{R}^2} \frac{\partial \tilde{R}}{\partial R} \tilde{S}' - 2\pi R^2 \frac{\partial \tilde{\Omega}}{\partial R} \tilde{J}' \right], \tag{A.11}$$

and finally, for the energy, we have

$$2\pi M = 2\pi e^{-\Phi_D} \tilde{M}' + e^{-\Phi_D} \left[-1 + \frac{R}{\tilde{R}} \frac{\partial \tilde{R}}{\partial R} - \frac{\mu_n}{\tilde{R}} \frac{\partial \tilde{R}}{\partial \mu_n} - \frac{\mu_w}{\tilde{R}} \frac{\partial \tilde{R}}{\partial \mu_w} - \frac{\Omega}{\tilde{R}} \frac{\partial \tilde{R}}{\partial \Omega} \right] \frac{\tilde{S}'}{\tilde{R}} + 2\pi e^{-\Phi_D} \left[-1 - \frac{R}{\tilde{\Omega}} \frac{\partial \tilde{\Omega}}{\partial R} + \frac{\Omega}{\tilde{\Omega}} \frac{\partial \tilde{\Omega}}{\partial \Omega} + \frac{\mu_n}{\tilde{\Omega}} \frac{\partial \tilde{\Omega}}{\partial \mu_n} + \frac{\mu_w}{\tilde{\Omega}} \frac{\partial \tilde{\Omega}}{\partial \mu_w} \right] \tilde{\Omega} \tilde{J}'.$$
(A.12)

(A.8) - (A.11) constitute as generalizations of the formulas in [12]. These relations are general and we didn't use the specific form of the solution generating transformation yet. All the information of the specific solution generation is contained in the map between

$$\{\tilde{R}, \tilde{\Omega}\} \leftrightarrow \{R, \Omega, \mu_n, \mu_w\}.$$
 (A.13)

To find these relations, only the asymptotic form of the solution generating transformation is needed. To zeroth order in α' , the transformation is the same in the bosonic/Type II and heterotic case, while they are different at order α' , which we defer to A.1. Following the derivation of [12] (which uses the formalism of [64]), we have²⁴

$$\tilde{R} = R\sqrt{1-\mu_n^2}\sqrt{1-\mu_w^2}, \qquad \tilde{\Omega} = \frac{\Omega}{\sqrt{1-\mu_n^2}\sqrt{1-\mu_w^2}}.$$
 (A.14)

An easy way to understand the relation between Ω and $\tilde{\Omega}$ is the following. Let's say that the rotation is in φ direction, then in the seed solution we have periodic identification $(t_E, \varphi) \sim (t_E + 1, \varphi + 2\pi i \tilde{R} \tilde{\Omega})$, where we've chosen the Euclidean time coordinate such that its periodicity is one. This identification should remain invariant under the transformation, meaning that $\tilde{R} \tilde{\Omega}$ should be invariant, which explains (A.14). To connect with the discussion in the main text, we can further parameterize μ_n and μ_w in terms of α, β

$$\mu_n = \tanh \frac{\alpha + \beta}{2}, \qquad \mu_w = \tanh \frac{\alpha - \beta}{2}.$$
(A.15)

Using (A.14) and (A.15), as well as $Q_{L,R} = Q_n \pm Q_w$, we can simplify (A.8) - (A.11) and get

$$2\pi Q_L = e^{-\Phi_D} \cosh\beta \sinh\alpha \left[\frac{\tilde{S}'}{\tilde{R}} + 2\pi\tilde{\Omega}\tilde{J}'\right], \qquad (A.16)$$

$$2\pi Q_R = e^{-\Phi_D} \cosh \alpha \sinh \beta \left[\frac{\tilde{S}'}{\tilde{R}} + 2\pi \tilde{\Omega} \tilde{J}' \right], \qquad (A.17)$$

$$2\pi J = 2\pi e^{-\Phi_D} \frac{\cosh \alpha + \cosh \beta}{2} \tilde{J}', \qquad (A.18)$$

$$S = e^{-\Phi_D} \tilde{S}' \frac{\cosh \alpha + \cosh \beta}{2}, \tag{A.19}$$

$$2\pi M = 2\pi e^{-\Phi_D} \tilde{M}' + e^{-\Phi_D} (\cosh \alpha \cosh \beta - 1) \left(\frac{\tilde{S}'}{\tilde{R}} + 2\pi \tilde{\Omega} \tilde{J}'\right).$$
(A.20)

The point of (A.16) - (A.20) is that once we know $\{\tilde{M}', \tilde{S}', \tilde{J}'\}$ of the uncharged solution, the thermodynamic quantities of the charged solution will be completely determined. For example, a consistency check of these expressions is that if we plug in the $\{\tilde{M}', \tilde{S}', \tilde{J}'\}$ for a four dimensional Kerr black hole, we correctly reproduce the thermodynamics of the two-charge black hole in [27].

As an aside, it is interesting to note from the expressions that the combination of $\frac{\tilde{S}'}{\tilde{R}} + 2\pi \tilde{\Omega} \tilde{J}'$ is the "charge" being transformed.

²⁴Here we follow the convention of [12] for the normalization of the gauge fields and charges, which leads to a $\sqrt{2}$ factor difference compared to the main text, i.e. $Q_{\text{appendix}} = Q_{\text{main text}}/\sqrt{2}$.

A.1 Generating α' corrections in various cases

A practical application of (A.16) - (A.20) is that one could use them to work out the α' correction to the rotating two-charge system using the knowledge of the α' correction to the seed solution, without needing to find the explicit corrected solution. Of course, this is assuming that we don't encounter any singular solutions along the solution generating transformation, so perturbative α' corrections are under control. For this reason, one cannot apply the results here directly to the solution we studied in the main text, which is singular at the two derivative level. Nonetheless, the formulas would apply when we are far away from the singular limit $\Omega = 2\pi i/\beta$.

The starting point of the derivation is the α' correction to the thermodynamics of the Kerr black hole (focusing on D = 4). This can be derived by simply evaluating the leading α' correction to the action on the uncorrected black hole background. The end results, in the bosonic/heterotic cases, are

$$\tilde{\beta} = \frac{4\pi r_+(a^2 + r_+^2)}{r_+^2 - a^2}, \quad \tilde{\Omega} = \frac{a}{r_+^2 + a^2}, \quad (A.21)$$

$$\tilde{M}' = \frac{r_+^2 + a^2}{2r_+}, \quad \tilde{S}' = \pi (r_+^2 + a^2) + 2\pi\lambda, \quad \tilde{J}' = a\tilde{M}'.$$
 (A.22)

Here $\lambda = \alpha'/2, \alpha'/4$ for bosonic and heterotic string theory, respectively. λ vanishes for Type II theories and the leading order correction happens at α'^3 , therefore we won't discuss it here. We emphasize that r_+, a in (A.21) and (A.22) are only parameters to parametrize $\tilde{\beta}$ and $\tilde{\Omega}$ and they differ from the corresponding geometric quantities by α' corrections.

For bosonic string theory, we can then plug (A.21), (A.22) into (A.16) - (A.20) can get the leading α' corrections to the rotating two charge system:

$$\frac{1}{T} = \frac{4\pi r_+ (r_+^2 + a^2)}{r_+^2 - a^2} \frac{\cosh \alpha + \cosh \beta}{2}, \quad \Omega = \frac{a}{r_+^2 + a^2} \frac{2}{\cosh \alpha + \cosh \beta}, \tag{A.23}$$

$$Q_L = \sinh \alpha \cosh \beta \left[\frac{\pi (r_+^2 + a^2)}{2r_+} + \lambda \frac{\pi (r_+^2 - a^2)}{r_+ (r_+^2 + a^2)} \right], \tag{A.24}$$

$$Q_R = \cosh \alpha \sinh \beta \left[\frac{\pi (r_+^2 + a^2)}{2r_+} + \lambda \frac{\pi (r_+^2 - a^2)}{r_+ (r_+^2 + a^2)} \right], \tag{A.25}$$

$$J = \frac{\cosh \alpha + \cosh \beta}{2} a \frac{r_{+}^{2} + a^{2}}{2r_{+}},$$
 (A.26)

$$2\pi M = \frac{\pi (r_+^2 + a^2)}{2r_+} (1 + \cosh \alpha \cosh \beta) + \lambda \frac{\pi (r_+^2 - a^2)}{r_+ (r_+^2 + a^2)} (\cosh \alpha \cosh \beta - 1), \qquad (A.27)$$

$$S = \frac{\cosh \alpha + \cosh \beta}{2} (\pi (r_+^2 + a^2) + 2\pi\lambda), \qquad (A.28)$$

where we've set $G_N = 1$ in the above expressions.

For the case of the heterotic string theory, the correct solution generating transformation in fact differs from the bosonic and type II theory by terms proportional to α' . The correct transformation was motivated by studying the Horowitz-Polchinski solution in [12], but was later independently verified by studying the α' corrections to the non-rotating twocharge system [28]. The derivation here follows that of [12], with the main new ingredient being that the angular potential Ω is related to $\tilde{\Omega}$ of the seed by $R\Omega = \tilde{R}\tilde{\Omega}$. We omit the intermediate steps and present the results analogous to (A.16) to (A.20).

$$R = \frac{\tilde{R}}{2} \left[\left(1 + \frac{\alpha'}{2\tilde{R}^2} \right) \cosh \alpha + \left(1 - \frac{\alpha'}{2\tilde{R}^2} \right) \cosh \beta \right], \quad \Omega = \frac{\tilde{R}}{R} \tilde{\Omega}, \tag{A.29}$$

$$2\pi Q_L = e^{-\Phi_D} \cosh\beta \sinh\alpha \left[\frac{\tilde{S}'}{\tilde{R}} + 2\pi\tilde{\Omega}\tilde{J}'\right], \quad 2\pi Q_R = e^{-\Phi_D} \cosh\alpha \sinh\beta \left[\frac{\tilde{S}'}{\tilde{R}} + 2\pi\tilde{\Omega}\tilde{J}'\right],$$
(A.30)

$$J = e^{-\Phi_D} \frac{\tilde{J}'}{2} \left[\left(1 + \frac{\alpha'}{2\tilde{R}^2} \right) \cosh \alpha + \left(1 - \frac{\alpha'}{2\tilde{R}^2} \right) \cosh \beta \right], \tag{A.31}$$

$$S = e^{-\Phi_D} \left[\frac{\cosh \alpha + \cosh \beta}{2} \tilde{S}' - \frac{\alpha' (\cosh \alpha - \cosh \beta)}{4\tilde{R}^2} (\tilde{S}' + 2\pi \tilde{R} \tilde{\Omega} \tilde{J}') \right],$$
(A.32)

$$2\pi M = 2\pi e^{-\Phi_D} \tilde{M}' + e^{-\Phi_D} (\cosh \alpha \cosh \beta - 1) \left(\frac{\tilde{S}'}{\tilde{R}} + 2\pi \tilde{\Omega} \tilde{J}' \right).$$
(A.33)

We can further plug (A.21) and (A.22) and get the explicit formulas for the α' corrected thermodynamic quantities for rotating two-charge solutions in heterotic string theory.

References

- M. J. Bowick, L. Smolin and L. C. R. Wijewardhana, Role of String Excitations in the Last Stages of Black Hole Evaporation, Phys. Rev. Lett. 56 (1986) 424.
- [2] L. Susskind, Some speculations about black hole entropy in string theory, hep-th/9309145.
- G. T. Horowitz and J. Polchinski, A Correspondence principle for black holes and strings, Phys. Rev. D 55 (1997) 6189-6197, [hep-th/9612146].
- [4] A. Dabholkar and J. A. Harvey, Nonrenormalization of the Superstring Tension, Phys. Rev. Lett. 63 (1989) 478.
- [5] A. Strominger and C. Vafa, Microscopic origin of the Bekenstein-Hawking entropy, Phys. Lett. B 379 (1996) 99-104, [hep-th/9601029].
- [6] A. Dabholkar, J. P. Gauntlett, J. A. Harvey and D. Waldram, Strings as solitons and black holes as strings, Nucl. Phys. B 474 (1996) 85-121, [hep-th/9511053].
- [7] A. Sen, Extremal black holes and elementary string states, Mod. Phys. Lett. A 10 (1995) 2081-2094, [hep-th/9504147].
- [8] A. Sen, How does a fundamental string stretch its horizon?, JHEP 05 (2005) 059, [hep-th/0411255].
- [9] V. Hubeny, A. Maloney and M. Rangamani, String-corrected black holes, JHEP 05 (2005) 035, [hep-th/0411272].
- [10] A. Dabholkar, R. Kallosh and A. Maloney, A Stringy cloak for a classical singularity, JHEP 12 (2004) 059, [hep-th/0410076].

- G. T. Horowitz and J. Polchinski, Selfgravitating fundamental strings, Phys. Rev. D 57 (1998) 2557-2563, [hep-th/9707170].
- [12] Y. Chen, J. Maldacena and E. Witten, On the black hole/string transition, JHEP 01 (2023) 103, [2109.08563].
- [13] P. A. Cano, P. F. Ramirez and A. Ruipérez, The small black hole illusion, JHEP 03 (2020) 115, [1808.10449].
- [14] E. Witten, Dynamical Breaking of Supersymmetry, Nucl. Phys. B 188 (1981) 513.
- [15] E. Kiritsis, Introduction to nonperturbative string theory, AIP Conf. Proc. 419 (1998) 265-308, [hep-th/9708130].
- [16] A. Cabo-Bizet, D. Cassani, D. Martelli and S. Murthy, Microscopic origin of the Bekenstein-Hawking entropy of supersymmetric AdS₅ black holes, JHEP 10 (2019) 062, [1810.11442].
- [17] L. V. Iliesiu, M. Kologlu and G. J. Turiaci, Supersymmetric indices factorize, JHEP 05 (2023) 032, [2107.09062].
- [18] J. Boruch, L. V. Iliesiu, S. Murthy and G. J. Turiaci, New forms of attraction: Attractor saddles for the black hole index, 2310.07763.
- [19] Z. Perjes, Solutions of the coupled Einstein Maxwell equations representing the fields of spinning sources, Phys. Rev. Lett. 27 (1971) 1668.
- [20] W. Israel and G. A. Wilson, A class of stationary electromagnetic vacuum fields, J. Math. Phys. 13 (1972) 865–871.
- [21] D. Butter, B. de Wit and I. Lodato, Non-renormalization theorems and N=2 supersymmetric backgrounds, JHEP 03 (2014) 131, [1401.6591].
- [22] S. Murthy and V. Reys, Quantum black hole entropy and the holomorphic prepotential of N=2 supergravity, JHEP 10 (2013) 099, [1306.3796].
- [23] Y. Chen and G. J. Turiaci, Spin-Statistics for Black Hole Microstates, 2309.03478.
- [24] K. S. Narain, New Heterotic String Theories in Uncompactified Dimensions < 10, Phys. Lett. B 169 (1986) 41–46.
- [25] K. S. Narain, M. H. Sarmadi and E. Witten, A Note on Toroidal Compactification of Heterotic String Theory, Nucl. Phys. B 279 (1987) 369–379.
- [26] C. Chowdhury, A. Sen, P. Shanmugapriya and A. Virmani, Supersymmetric Index for Small Black Holes, 2401.13730.
- [27] A. Sen, Black hole solutions in heterotic string theory on a torus, Nucl. Phys. B 440 (1995) 421-440, [hep-th/9411187].
- [28] S. Massai, A. Ruipérez and M. Zatti, Revisiting α' corrections to heterotic two-charge black holes, 2311.03308.
- [29] T. Damour and G. Veneziano, Selfgravitating fundamental strings and black holes, Nucl. Phys. B 568 (2000) 93-119, [hep-th/9907030].
- [30] J. J. Atick and E. Witten, The Hagedorn Transition and the Number of Degrees of Freedom of String Theory, Nucl. Phys. B 310 (1988) 291–334.
- [31] N. Ceplak, R. Emparan, A. Puhm and M. Tomasevic, The correspondence between rotating black holes and fundamental strings, JHEP 11 (2023) 226, [2307.03573].

- [32] G. T. Horowitz and A. Sen, Rotating black holes which saturate a Bogomolny bound, Phys. Rev. D 53 (1996) 808-815, [hep-th/9509108].
- [33] B. de Wit, J. W. van Holten and A. Van Proeyen, Transformation Rules of N=2 Supergravity Multiplets, Nucl. Phys. B 167 (1980) 186.
- [34] B. de Wit, J. W. van Holten and A. Van Proeyen, Structure of N=2 Supergravity, Nucl. Phys. B 184 (1981) 77.
- [35] K. Behrndt, D. Lust and W. A. Sabra, Stationary solutions of N=2 supergravity, Nucl. Phys. B 510 (1998) 264-288, [hep-th/9705169].
- [36] G. Lopes Cardoso, B. de Wit, J. Kappeli and T. Mohaupt, Stationary BPS solutions in N=2 supergravity with R^{**2} interactions, JHEP 12 (2000) 019, [hep-th/0009234].
- [37] J. B. Hartle and S. W. Hawking, Solutions of the Einstein-Maxwell equations with many black holes, Commun. Math. Phys. 26 (1972) 87–101.
- [38] B. Whitt, ISRAEL-WILSON METRICS, Annals Phys. 161 (1985) 241.
- [39] A. Dabholkar, J. Gomes and S. Murthy, Quantum black holes, localization and the topological string, JHEP 06 (2011) 019, [1012.0265].
- [40] A. Dabholkar, J. Gomes and S. Murthy, Localization & Exact Holography, JHEP 04 (2013) 062, [1111.1161].
- [41] A. Dabholkar, J. Gomes and S. Murthy, Nonperturbative black hole entropy and Kloosterman sums, JHEP 03 (2015) 074, [1404.0033].
- [42] L. V. Iliesiu, S. Murthy and G. J. Turiaci, Black hole microstate counting from the gravitational path integral, 2209.13602.
- [43] G. Lopes Cardoso, B. de Wit, J. Kappeli and T. Mohaupt, Supersymmetric black hole solutions with R^{**}2 interactions, PoS tmr99 (1999) 011, [hep-th/0003157].
- [44] G. Lopes Cardoso, B. de Wit, J. Kappeli and T. Mohaupt, Examples of stationary BPS solutions in N=2 supergravity theories with R**2 interactions, Fortsch. Phys. 49 (2001) 557-563, [hep-th/0012232].
- [45] A. Dabholkar, Exact counting of black hole microstates, Phys. Rev. Lett. 94 (2005) 241301, [hep-th/0409148].
- [46] P. Benetti Genolini, J. M. Perez Ipiña and J. Sparks, Localization of the action in AdS/CFT, JHEP 10 (2019) 252, [1906.11249].
- [47] S. M. Hosseini, K. Hristov and A. Zaffaroni, Gluing gravitational blocks for AdS black holes, JHEP 12 (2019) 168, [1909.10550].
- [48] P. Benetti Genolini, J. P. Gauntlett and J. Sparks, Equivariant Localization in Supergravity, Phys. Rev. Lett. 131 (2023) 121602, [2306.03868].
- [49] H. K. Kunduri, J. Lucietti and H. S. Reall, Near-horizon symmetries of extremal black holes, Class. Quant. Grav. 24 (2007) 4169–4190, [0705.4214].
- [50] P. Figueras, H. K. Kunduri, J. Lucietti and M. Rangamani, Extremal vacuum black holes in higher dimensions, Phys. Rev. D 78 (2008) 044042, [0803.2998].
- [51] D. Stanford and E. Witten, Fermionic Localization of the Schwarzian Theory, JHEP 10 (2017) 008, [1703.04612].

- [52] A. Giveon and D. Kutasov, Fundamental strings and black holes, JHEP 01 (2007) 071, [hep-th/0611062].
- [53] A. Dabholkar and S. Murthy, Fundamental Superstrings as Holograms, JHEP 02 (2008) 034, [0707.3818].
- [54] C. V. Johnson, Heterotic Coset Models of Microscopic Strings and Black Holes, 0707.4303.
- [55] J. M. Lapan, A. Simons and A. Strominger, Nearing the Horizon of a Heterotic String, 0708.0016.
- [56] P. Kraus, F. Larsen and A. Shah, Fundamental Strings, Holography, and Nonlinear Superconformal Algebras, JHEP 11 (2007) 028, [0708.1001].
- [57] W. Nahm, Supersymmetries and their Representations, Nucl. Phys. B 135 (1978) 149.
- [58] R. Britto-Pacumio, J. Michelson, A. Strominger and A. Volovich, Lectures on Superconformal Quantum Mechanics and Multi-Black Hole Moduli Spaces, NATO Sci. Ser. C 556 (2000) 255-284, [hep-th/9911066].
- [59] L. Frappat, P. Sorba and A. Sciarrino, *Dictionary on Lie superalgebras*, hep-th/9607161.
- [60] A. A. H., P. V. Athira, C. Chowdhury and A. Sen, Logarithmic Correction to BPS Black Hole Entropy from Supersymmetric Index at Finite Temperature, 2306.07322.
- [61] A. H. Anupam, C. Chowdhury and A. Sen, Revisiting Logarithmic Correction to Five Dimensional BPS Black Hole Entropy, 2308.00038.
- [62] S. Sethi and M. Stern, D-brane bound states redux, Commun. Math. Phys. 194 (1998) 675-705, [hep-th/9705046].
- [63] G. Turiaci Unpublished notes .
- [64] J. Maharana and J. H. Schwarz, Noncompact symmetries in string theory, Nucl. Phys. B 390 (1993) 3-32, [hep-th/9207016].