



King's Research Portal

Document Version
Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Greenwood, S. A., Briggs, J., Walklin, C., Mangahis, E., Young, H. M. L., Castle, E. M., Billany, R. E., Asgari, E., Bhandari, S., Bishop, N., Bramham, K., Burton, J. O., Campbell, J., Chilcot, J., Cooper, N., Deelchand, V., Graham-Brown, M. P. M., Haggis, L., Hamilton, A., ... Macdonald, J. (2024). Kidney Beam - a cost-effective digital intervention to improve mental health. *Kidney International Reports*, 9(11), 3204-3217.

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Journal Pre-proof



Kidney Beam - a cost-effective digital intervention to improve mental health.

Sharlene A. Greenwood, Juliet Briggs, Christy Walklin, Emmanuel Mangahis, Hannah ML. Young, Ellen M. Castle, Roseanne E. Billany, Elham Asgari, Sunil Bhandari, Nicolette Bishop, Kate Bramham, James O. Burton, Jackie Campbell, Joseph Chilcot, Nicola Cooper, Vashist Deelchand, Matthew PM. Graham-Brown, Lynda Haggis, Alexander Hamilton, Mark Jesky, Philip A. Kalra, Pelagia Koufaki, Kieran McCafferty, Andrew C. Nixon, Helen Noble, Zoe L. Saynor, Maarten W. Taal, James Tollitt, David C. Wheeler, Thomas J. Wilkinson, Hannah Worboys, Jamie Macdonald

PII: S2468-0249(24)01915-6

DOI: <https://doi.org/10.1016/j.ekir.2024.08.030>

Reference: EKIR 2977

To appear in: *Kidney International Reports*

Received Date: 26 May 2024

Revised Date: 19 July 2024


Accepted Date: 27 August 2024

Please cite this article as: Greenwood SA, Briggs J, Walklin C, Mangahis E, Young HM, Castle EM, Billany RE, Asgari E, Bhandari S, Bishop N, Bramham K, Burton JO, Campbell J, Chilcot J, Cooper N, Deelchand V, Graham-Brown MP, Haggis L, Hamilton A, Jesky M, Kalra PA, Koufaki P, McCafferty K, Nixon AC, Noble H, Saynor ZL, Taal MW, Tollitt J, Wheeler DC, Wilkinson TJ, Worboys H, Macdonald J, Kidney Beam - a cost-effective digital intervention to improve mental health., *Kidney International Reports* (2024), doi: <https://doi.org/10.1016/j.ekir.2024.08.030>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 Published by Elsevier Inc. on behalf of the International Society of Nephrology.

Kidney Beam- a Cost-Effective Digital Intervention to Improve Mental Health



KI REPORTS
Kidney International Reports

Greenwood SA et al, 2024

Visual abstract by:
Abdul Qader, MD
✉ @md_abdulqader83

Conclusion The Kidney BEAM physical activity digital health intervention is clinically valuable and cost-effective means to improve mental health related quality of life in people with CKD.

Journal Pre-proof

1 **Kidney Beam - a cost-effective digital intervention to improve mental health.**

2 Sharlene A Greenwood^{1,2}, Juliet Briggs¹, Christy Walklin¹, Emmanuel Mangahis¹, Hannah ML
3 Young^{3,4}, Ellen M Castle⁵, Roseanne E Billany⁶, Elham Asgari⁷, Sunil Bhandari⁸, Nicolette
4 Bishop⁹, Kate Bramham², James O Burton⁶, Jackie Campbell¹⁰, Joseph Chilcot¹¹, Nicola
5 Cooper¹², Vashist Deelchand¹³, Matthew PM Graham-Brown⁶, Lynda Haggis², Alexander
6 Hamilton¹⁴, Mark Jesky¹⁵, Philip A Kalra¹⁶, Pelagia Koufaki¹⁷, Kieran McCafferty¹⁸, Andrew
7 C Nixon^{19, 20}, Helen Noble²¹, Zoe L Saynor²², Maarten W Taal²³, James Tollitt¹⁶, David C
8 Wheeler²⁴, Thomas J Wilkinson²⁵, Hannah Worboys¹², Jamie Macdonald²⁶.

9

10 ¹ Renal Therapies Department, King's College Hospital, London.

11 ² Centre for Nephrology, Urology and Transplantation, Faculty of Life Sciences, King's
12 College London, London.

13 ³ Leicester Diabetes Centre, University Hospitals of Leicester NHS Trust, Leicester.

14 ⁴ Diabetes Research Centre, University of Leicester, Leicester.

15 ⁵ School of Physiotherapy, Department of Health Sciences, Brunel University, London.

16 ⁶Department of Cardiovascular Sciences, University of Leicester, Leicester.

17 ⁷ Dept of Nephrology, Guys and St Thomas's Hospital, London.

18 ⁸ Dept of Nephrology, Hull University Teaching Hospitals NHS Trust, Hull.

19 ⁹ School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough.

20 ¹⁰ Faculty of Health, Education and Society, University of Northampton, Northampton.

21 ¹¹ Department of Psychology, Psychology & Neuroscience, King's College London, London.

- 1 ¹² Department of Population Health Science, University of Leicester, Leicester.
- 2 ¹³ Dept of Nephrology Royal Free Hospital, London.
- 3 ¹⁴ Dept of Nephrology, Royal Devon and Exeter NHS Foundation Trust, Exeter.
- 4 ¹⁵ Dept of Nephrology, Nottingham NHS Trust, Nottingham.
- 5 ¹⁶ Dept of Nephrology Royal Hospital, Northern Care Alliance NHS Foundation Trust, Salford.
- 6 ¹⁷ Dietetics, Nutrition & Biological Sciences, Physiotherapy, Podiatry & Radiography Division,
7 Queen Margaret University, Edinburgh.
- 8 ¹⁸ Dept of Nephrology, Barts Health NHS Trust, London.
- 9 ¹⁹ Dept of Renal Medicine, Lancashire Teaching Hospitals NHS Foundation Trust, Preston.
- 10 ²⁰ Division of Cardiovascular Sciences, The University of Manchester, Manchester, United
11 Kingdom.
- 12 ²¹ School of Nursing and Midwifery, Queen's University Belfast, Belfast.
- 13 ²² School of Sport, Health and Exercise Science, University of Southampton, Southampton.
- 14 ²³ Centre for Kidney Research and Innovation, School of Medicine, University of Nottingham,
15 Nottingham.
- 16 ²⁴ Department of Renal Medicine, University College London, London.
- 17 ²⁵ National Institute of Health Research Leicester Biomedical Research Centre, Leicester, UK.
- 18 ²⁶ Institute for Applied Human Physiology, Bangor University, Bangor.
- 19
- 20

1 **Correspondence to:**

2 Dr Sharlene Greenwood

3 Department of Renal Medicine

4 King's College Hospital NHS Trust

5 London SE5 9RS

6 Telephone: 020881947470

7 Email: sharlene.greenwood@nhs.net

8

9 **Running head:** Clinical value and cost effectiveness of Kidney BEAM

10

Journal Pre-proof

1 Abstract

2 **Background:** There is inequity in provision of physical rehabilitation services for people living
3 with chronic kidney disease (CKD). The Kidney BEAM trial evaluated the clinical value and
4 cost effectiveness of a physical activity digital health intervention in CKD.

5 **Methods:** In a single-blind, 11 centre, randomised controlled trial, 340 adult participants with
6 CKD were randomly assigned to either the Kidney BEAM physical activity digital health
7 intervention or a waitlist control. This study assesses the difference in the Kidney Disease
8 Quality of Life Short Form 1.3 Mental Component Summary (KDQoL-SF1.3 MCS) between
9 intervention and control groups at 6 months, and cost-effectiveness of the intervention.

10 **Results:** At 6 months there was a significant difference in mean adjusted change in KDQoL
11 MCS score between Kidney BEAM and waitlist control (intention-to-treat adjusted mean: 5.9
12 {95% confidence interval: 4.4 to 7.5} arbitrary units, $p < 0.0001$), and a 93% and 98% chance
13 of the intervention being cost-effective at a willingness to pay threshold of £20,000 and
14 £30,000 per quality-adjusted life year gained.

15 **Conclusion:** The Kidney BEAM physical activity digital health intervention is a clinically
16 valuable and cost-effective means to improve mental health related quality of life in people
17 with CKD (trial registration no. NCT04872933).

18

19 **Keywords:** Chronic kidney disease, cost-effectiveness, digital health intervention, Physical
20 activity, quality of life.

21

1 Introduction

2 Chronic kidney disease (CKD) affects more than 10% of the adult population worldwide,
3 amounting to in excess of 800 million individuals, and is predicted to be the fifth highest cause
4 of years of life lost worldwide by 2040. ¹ Physical inactivity is the fourth leading risk factor
5 for global mortality, is a major risk factor for multimorbidity in people with chronic disease
6 and has been associated with poor mental health-related quality of life (HRQoL). ^{2, 3}
7 Consequently, interventions to enhance physical activity, mental health and HRQoL are of
8 global interest and have been the focus of disease-specific guidelines, including those for
9 people living with CKD. ⁴⁻⁶

10

11 Whilst there may be benefits to in-person kidney rehabilitation, ⁷ this has not been provided
12 routinely in the United Kingdom, ⁸ and policy-related barriers restrict access to exercise
13 provision globally, leading to health inequality. ⁹ One of the barriers to implementation has
14 been a dearth of cost-effectiveness data to support the adoption of kidney-specific physical
15 rehabilitation programmes into already financially stretched healthcare systems. ¹⁰ Even where
16 there has been evidence published, such as the results from a UK study that reported the cost-
17 effectiveness of intra-dialytic cycling programmes, ¹¹ further complexities around availability
18 of exercise personnel, equipment and unit-level support have resulted in little meaningful
19 adoption to date. ¹⁰ Additionally, physical activity and exercise training trials in this patient
20 population often neglect to report on whether there are sustained benefits from structured
21 physical activity interventions, questioning the longer-term benefit and cost efficiency of these
22 interventions when considering commissioning. We have anticipated these requirements by
23 providing the 6-month patient outcome and healthcare utilisation analyses reported here within.

24

1 The importance of digital health interventions has been highlighted in the World Health
2 Organization (WHO) global strategy on digital health 2020–2025.¹² Furthermore, the
3 utilisation of digital health interventions can activate patients to engage in online lifestyle
4 interventions and education, which can promote self-management and improve health
5 outcomes for those with chronic disease.¹³

6

7 The 12-week Kidney BEAM physical activity digital health intervention (DHI) demonstrated
8 clinically meaningful and statistically significant improvements in mental HRQoL, physical
9 function, and patient activation (the ability to self-manage health behaviours) for people living
10 with CKD¹⁴, strongly supporting the efficacy of physical activity DHIs in the short-term.
11 However, the Transtheoretical Model suggests that maintenance of a behaviour can only be
12 assumed if sustained for at least 6 months.¹⁵ Therefore, we hypothesised that 6 months of a
13 physical activity DHI would reveal clinically meaningful improvements in mental HRQoL and
14 be a cost-effective solution to deliver physical activity interventions for people living with
15 CKD. The trial was co-designed with people with lived experience and targeted mental health-
16 related quality of life as this was the most important outcome to the patients who we consulted
17 with. Quality of life, and life participation, has been highlighted by the SONG initiative as
18 being important to people living with CKD across the disease trajectory.¹⁶

19

20 **Materials and methods**

21 **Study design**

22 The 6-month Kidney BEAM Trial was a multi-centre, randomised, single-blind, controlled
23 waitlist trial to assess the clinical value and cost-effectiveness of a physical activity digital
24 health intervention on health-related quality of life in people with CKD that was conducted at

1 eleven centres in the United Kingdom (UK). The trial design, protocol, and baseline
2 characteristics of the participants have been published previously,^{17, 18} as have the 12-week
3 results of the Kidney Beam Trial.¹⁴ The protocol was approved by the UK Bromley Research
4 Ethics Committee at King's College Hospital NHS Trust, London, UK. The trial was designed
5 and overseen by a trial steering committee and a data monitoring committee.

6

7 **Participants**

8 Adults with established CKD, including those who were pre-dialysis (CKD stages 2-4) and
9 those on kidney replacement therapy (dialysis and kidney transplantation), were eligible for a
10 digital health intervention if they had access to a digital device and WiFi connectivity.
11 Recruitment occurred at kidney centres across England, UK, intentionally chosen to represent
12 the geographical diversity of the UK CKD population. Potential participants underwent
13 screening, and their clinical records were reviewed to confirm eligibility. Trained research staff
14 approached suitable adults face-to-face during clinic visits or via telephone. Exclusions
15 included self-reported participation in a recent exercise program or use of a physical activity
16 digital health intervention within the last three months, persistent uncontrolled hypertension,
17 unstable angina, and conditions preventing engagement in a physical activity intervention, such
18 as peripheral vascular or musculoskeletal diseases. Decisions to exclude participants based on
19 the severity of peripheral vascular or musculoskeletal disease were adjudicated by the study
20 team to prevent risk to the patient rather than an exclusion based on chart diagnosis alone.
21 Informed written consent was obtained from all participants, and a detailed list of inclusion and
22 exclusion criteria can be found in the methods paper.¹⁷

23

24 **Randomisation and masking**

1 Participants were randomly assigned in a 1:1 ratio to the Kidney BEAM intervention group or
2 the waitlist control group. Randomisation was performed with the use of a Web-based system,
3 in randomly permuted blocks of six. Randomisation and treatment allocation was performed
4 by an independent member of the research team and the allocation list was stored in a password-
5 protected database. Given the nature of the intervention, it was not possible to blind the
6 healthcare professionals providing the programme or the participants. Outcome assessors were,
7 however, blinded to treatment allocation. The statistical analysis plan and the health economic
8 analysis plan¹⁷ were developed *a priori* by an independent statistician and health economist
9 and were approved by the trial steering committee. Data entry and quality assurance was
10 undertaken by data entry clerks unaware of treatment allocation. Data cleaning and analysis of
11 outcome data was conducted by the independent statistician and health economist unaware of
12 treatment allocation.

13

14 **Outcomes**

15 The primary objective for this 6-month trial was to evaluate the change in the Kidney Disease
16 Quality of Life Short Form 1.3 Mental Component Summary (KDQoL-SF1.3 MCS) between
17 baseline and 24 weeks and to assess cost effectiveness. The MCS is composed of all scales of
18 the SF-36 but is more heavily weighted to the vitality (energy/fatigue), social functioning, role
19 emotional and mental health subscales of the KDQoL questionnaire. Secondary objectives
20 included evaluating changes in the KDQoL-SF1.3 Physical Component Score (PCS) at 24
21 weeks (which is more heavily weighted to the physical functioning, role-physical, bodily pain,
22 general health sub-scales), other KDQoL sub-scales, the European Quality of Life 5-
23 dimension, 5-level (EQ-5D-5L) questionnaire (converted to EQ-5D-3L to allow comparison
24 with UK normative data) and healthcare utilisation data. All outcome measures were chosen as

1 valid and reliable tools to measure the primary and secondary outcomes in this patient
2 population.²⁴ All patient-reported outcome measures were completed via an online survey.
3 Health utilisation data was also obtained via video conference with participants. Safety
4 outcomes were based on adverse-event reporting. An independent data monitoring committee
5 had oversight of trial safety.

6

7 **Healthcare utilisation**

8 Data on associated hospital costs, primary care consultations, and social care usage were
9 collected via patient interview for the pre-trial and with-in trial period. Prescribed medication
10 costs were collected from hospital records. Intervention costs assume a cost of £15 per
11 participant per year and consisted of physiotherapy time, physiotherapy assistant time and
12 running costs for the Kidney BEAM platform. One experienced physiotherapy assistant at
13 whole time (1.0 whole time equivalent (WTE)), and 1 senior, experienced physiotherapist at
14 10% of their whole time (0.1 WTE) per 340 participants were costed in at current NHS staff
15 salary rates.¹⁹ This intervention cost reflects a proposed population-based contract assuming a
16 10% sign-up rate to the intervention across the CKD population of England. Resources were
17 valued using national tariffs.^{20, 21} All costs were expressed in 2021/2022 UK pounds (£). All
18 costs were expressed in 2021/22 UK (£) inflated to this base year where appropriate using the
19 UK Consumer Price Health Index.¹⁹

20

21 **Intervention**

22 The 12-week structured physical activity intervention has been described in detail elsewhere.
23 ^{24, 14} In brief, the 6-month Kidney BEAM intervention (<https://beamfeelgood.com/home>),
24 which included a rolling 12-week structured digitally delivered physical activity intervention,

1 was delivered by specialist kidney physiotherapists through ‘live’ sessions, which were
2 delivered in real-time via the digital platform, and a pre-recorded on-demand kidney
3 rehabilitation programme, followed by 12 weeks of self-managed physical activity accessed
4 through the Kidney BEAM platform. The structured 12-week sessions comprised a 10-minute
5 warm-up and cool-down involving general upper and lower limb mobility and stretching. The
6 core session included 20-30 minutes of moderate-intensity aerobic and resistance exercises,
7 delivered both in a standing and seated position. Additionally, participants received 15 minutes
8 of disease-specific education on topics related to managing kidney health, such as managing a
9 kidney diet and understanding diabetes, weekly. A physiotherapy assistant, trained in
10 motivational interviewing, provided ongoing general encouragement through weekly
11 telephone or email communication. Participants could review their progress via their
12 personalised dashboard on the platform. After completing the 12-week programme and
13 assessing outcomes, participants in the intervention group were advised by the physiotherapy
14 assistant to maintain self-management of their physical activity behaviour with ongoing access
15 to the Kidney BEAM platform. Participants who were allocated to the wait-list control group
16 did not participate in a 12-week structured exercise programme and were only sign-posted to
17 Kidney BEAM following the 12-week assessment.

18

19 **Statistical Analysis**

20 The trial was designed to detect a clinically meaningful 3 arbitrary unit (AU) difference in
21 HRQoL KDQoL-SF1.3 MCS score between groups at 12 weeks and 6 months. An estimated
22 sample size of 106 participants in each group (total n = 212) based on an MCS with a mean of
23 45 AU, SD 10 AU and correlation between repeated measures of 0.7, would allow a clinically
24 meaningful difference of 3 AU to be detected at 80% power and 5% alpha. Specifically, a 3-

1 point difference in MCS is associated with an odds ratio of 1.13 for being unable to work or an
2 odds ratio of 1.16 for 1-year job loss. The probability of using mental health services is
3 increased by approximately 30% (odds ratio = 1.31), and there is a 30% increased risk of
4 depression (odds ratio = 1.34). It is also associated with a 10% higher 1-year mortality risk
5 (odds ratio = 1.10). 340 patients were included to allow for a 30% drop-out and to ensure power
6 for secondary outcomes.²² The baseline characteristics were described using summary
7 statistics.¹⁷ Primary and secondary outcomes at 6 months were analysed with an analysis of
8 covariance model, with baseline data and age as covariates. Independence of covariates and
9 approximated normality of residuals were confirmed for all analyses. All analyses were
10 performed in the *intention-to-treat* (ITT) population using a last observation carried forward
11 (LOCF) approach to missing data as this gives the most conservative result. The results from
12 the LOCF analysis for the primary outcome were compared to those from a multiple imputation
13 sensitivity analysis using pooled results from 5 linear regression imputations. *Per protocol* (PP)
14 analyses in which only cases with observations at both baseline and week 24 were included,
15 were also completed to assess efficacy under ideal conditions. Two-sided p values of less than
16 0.05 were considered to indicate statistical significance. Analyses were performed with SPSS
17 (version 28, IBM, New York).

18

19 The reporting of the Health Economic Analysis adheres to the CHEERS 2022 Checklist.²³
20 The within-trial economic analyses were performed using individual patient level data
21 collected from the trial. The base case analysis included all participants completing the 12
22 week and 6-month follow-up with missing resource use items imputed using a last value
23 carried forward (LVCF) approach. Area under the curve methods were used to calculate the
24 QALYs accrued by each person during the intervention period based on EQ-5D-5L cost
25 utility data collected at baseline and at 3 and 6 months. The trial was conducted in the UK,

1 which has a National Health Service (NHS) providing publicly funded healthcare, primarily
2 free of charge at the point of use. The primary economic analysis was from the NHS and
3 personal social services perspective. The primary economic analysis compared the costs and
4 consequences of each arm over the 6 months following randomisation. For the analysis, we
5 adopted a bivariate model for estimating incremental costs and effects in WinBUGS using
6 Markov Chain Monte Carlo (MCMC) simulation methods²⁴ with costs and 1-QALYs
7 expressed as Gamma distributions. Bayesian methods require the specification of prior
8 distributions for parameters of the distributions. Here we used prior distributions intended to
9 be non-informative, as we wanted the resulting inferences to only depend on the data. For the
10 base-case analysis, the bivariate model incorporated adjustment for baseline costs (12 weeks
11 prior to intervention) and EQ-5D to allow for imbalance between the groups using the
12 methods proposed by Nixon and Thompson 2005.²⁵ Posterior distributions of the parameters
13 of interest for the inferences about cost-effectiveness were derived from 20,000 iterations of
14 the Markov chain, after an initial 20,000 iterations were discarded to ensure convergence.
15 Results were expressed in terms of cost per QALY gained (i.e., the incremental cost-
16 effectiveness ratio), which was estimated for the Kidney BEAM group compared with the
17 wait-list control group.

18

19 **Inclusion and ethics**

20 The trial was designed and overseen by a trial steering committee and a data monitoring
21 committee. The protocol and related documents were approved by Bromley NHS Research
22 Ethics Committee (REC) (21/LO/0243) and the Health Research Authority (HRA) and was
23 prospectively registered (NCT04872933) on 5th May 2021. All methods were carried out in
24 accordance with relevant guidelines and regulations. Informed consent was obtained from all
25 subjects and/or their legal guardian(s).

1

2 **Results**

3 **Participants**

4 From May 06, 2021, to October 30, 2022, 1102 people were assessed for eligibility (Fig. 1).
5 After excluding 721 (65%) people; 381 (35%) participants were consented and a total of 340
6 (31%) participants from eleven centres attended a baseline visit. The two main reasons for not
7 engaging with the trial were time constraints associated with the research trial and potential
8 participants that passed screening but were not able to be contacted to consent and participate
9 in the trial. 173 (51%) people were randomly assigned to the Kidney BEAM intervention
10 group, and 167 (49%) were assigned to the waitlist control group. Of these, 247 (73%)
11 participants completed the 6-month trial: 105 in the intervention group (61% of those
12 randomised) and 142 in the waitlist control group (85% of those randomised). All 340
13 participants were included in the *intention-to-treat* analysis. Overall, the two groups were
14 generally well balanced with respect to baseline characteristics (Table 1), albeit the mean EQ-
15 5D-3L utility scores were lower in the intervention group and there was more self-reported
16 burden of kidney disease, pain and sexual dysfunction in the intervention group (Table 2).

17

18 **Participant adherence**

19 A median of 15 (IQR 9-22) of the recommended 24 sessions of structured physical activity
20 were completed by participants in the Kidney BEAM intervention group during the structured
21 12-week physical activity component, representing a median adherence rate of 63 (IQR 38-92)
22 %. Participants completed a median of 529 (IQR 283-814) minutes of structured physical
23 activity (video/session length x number of sessions), the equivalent of 44 minutes per week. A
24 median of 6 (IQR 1-10) of the recommended 12 sessions of education were completed,

1 representing a median adherence rate of 50 (IQR 8-83) %. 65 out of 105 (62%) participants
2 from the Kidney BEAM intervention group continued to use the Kidney BEAM platform to
3 complete self-managed physical activity sessions after the 12-week assessment. Between 12
4 weeks and 6 months, participants in the Kidney BEAM group completed a median of 7 (IQR
5 3-41) sessions of self-managed physical activity sessions on the platform and completed a
6 median of 286 (IQR 103-1792) minutes of self-managed physical activity through the platform.
7 As per protocol, participants from the wait-list control group were informed at consent that
8 they could access the Kidney BEAM platform following the 12-week assessment. This was not
9 actively encouraged by the team and only 15 out of 142 (11%) participants from the waitlist
10 control group did choose to self-sign-up to the platform and complete self-managed physical
11 activity sessions on the Kidney BEAM platform between 12 weeks and 6 months. Participants
12 from the wait-list control group completed a median of 11 (IQR 5-46) sessions of self-managed
13 physical activity using the platform, and a median of 119 minutes (IQR 90.5-1822) minutes of
14 self-managed physical activity using the platform.

15

16 **Primary outcomes**

17 Using the most conservative last observation carried forward (LOCF) approach, there was a
18 clinically relevant and statistically significant improvement in the KDQoL SF 1.3 mental
19 component summary score after 6 months in the Kidney BEAM group compared to the control
20 group of 5.9 {95% confidence interval: 4.4 to 7.5} arbitrary units ($p < 0.0001$) (Table 2).
21 Sensitivity analysis confirmed this result, by using multiple imputation of the 6-month missing
22 values, and five iterations of linear regression imputation, revealing a pooled mean difference
23 of 5.8 {3.1 to 8.4} arbitrary units ($p < 0.0001$).

24

1 Regarding cost effectiveness, the adjusted *intention-to-treat* base case model, assuming a cost
2 per participant of £15 per year, showed a mean cost saving of £93 {95% confidence interval: -
3 £360 to £613} per participant in health care utilisation costs and a significant increment in
4 quality-adjusted life years (QALYs) of 0.027 {95% confidence interval: 0.013 to 0.040} years
5 per participant, resulting in a cost per QALY of £3,446 for the Kidney BEAM intervention
6 (Table 3 and Supplementary Table 1). This resulted in a 93% and 98% probability (indicated
7 by the proportion of the ellipses below the willingness to pay threshold line, Figure 2) of the
8 Kidney BEAM intervention being cost-effective, compared with wait-list control, at the
9 willingness to pay thresholds of £20,000 and £30,000 per QALY gained, respectively (Fig. 2
10 and Table 3). The adjusted *complete-case* model, assuming a cost per participant of £15 per
11 year, showed a mean cost saving of £273.60 {95% confidence interval: -£323 to £996.7} per
12 participant in health care utilisation costs and a significant increment in quality-adjusted life
13 years (QALYs) of 0.026 {95% confidence interval: 0.009 to 0.043} years per participant,
14 resulting in a cost per QALY of £10,523.08 for the Kidney BEAM intervention. This resulted
15 in a 75% and 87% probability of the Kidney BEAM intervention being cost-effective,
16 compared with wait-list control, at the willingness to pay thresholds of £20,000 and £30,000
17 per QALY gained (Fig. 2). The significant increase in KDQoL MCS in the Kidney BEAM
18 intervention group compared with wait-list control is associated with an incremental cost
19 effectiveness ratio (ICER) of £14.44 per one unit change in KDQoL MCS (Supplementary data
20 table 2). Exploratory analyses comparing the cost effectiveness of the Kidney BEAM digital
21 health intervention at varying costs per participant per year for the intervention (£30, £50 and
22 £100) did not result in any change to the ICER (Supplementary data table 3). Primary care,
23 medication, hospital-associated, and total costs are presented by group at 12 weeks pre-trial,
24 and at 12 weeks and 6 months during the trial (Supplementary data table 4, 5).

25

1 **Secondary outcomes**

2 The change in the KDQoL MCS was primarily due to mean between-group improvements in
3 the individual components of the KDQoL SF 1.3 questionnaire at the same time-point,
4 including: the social function, energy/fatigue, role emotional, and emotional wellbeing scales
5 (Table 2).

6

7 Analysis of secondary outcomes also revealed a significant improvement at 6 months in the
8 EQ-5D-3L utility score of 0.10 {95% confidence interval: 0.07 to 0.13} units ($p < 0.0001$) in
9 favour of the Kidney BEAM group (Table 2). The mean between-group difference in the
10 KDQoL PCS and the cognitive function sub-scale at 6 months were not significant ($p = 0.055$
11 and 0.082, respectively (Table 2)) but were significant on *per protocol* analysis (Supplementary
12 Data Table 6). All other sub-scales revealed significant mean between-group differences at 6
13 months in favour of the intervention group (Table 2).

14

15 There were nine unrelated Serious Adverse Events (SAE's) recorded in a total of 9 out of the
16 340 participants, with a similar incidence across both groups: 4 of the 9 (3%) in the Kidney
17 BEAM group and 5 of the 9 (3%) in the control group across the 6-month trial period. There
18 were no expected related or unrelated Serious Adverse Events recorded in either group during
19 the duration of the trial (Table 4).

20

21 **Participant dropouts and missing data**

22 There was no obvious difference in participant characteristics between participants that
23 completed the 6-month outcome assessment and participants that did not (Supplementary data

1 table 7). 47 (77%) of the 68 participants that did not complete the trial in the intervention group
2 withdrew within the first week post baseline assessment due to time constraints. As expected,
3 the number of missing data points for the cost effectiveness analyses increased as the trial
4 progressed, but at 6 months there were still 229 data points available for analysis (Kidney
5 BEAM intervention group: N=93; Control Group N = 136) (Supplementary data table 8).

6

7 **Discussion**

8 The results from this 6-month trial demonstrate that the Kidney BEAM physical activity digital
9 health intervention resulted in a clinically meaningful, sustained improvement in mental health
10 related quality of life in people with CKD and was cost-effective. Our data will support
11 commissioning of the Kidney BEAM innovation within the National Health System (NHS)
12 and inform commissioning of similar services in other healthcare systems.

13

14 Interventions that afford improvements in mental HRQOL are important for all people living
15 with CKD, and may be particularly important for those people receiving dialysis therapy where
16 lower levels of HRQOL have been associated with morbidity and mortality, and where every
17 1-point increase in MCS has been associated with a 2% reduction in the relative risk of death
18 and a 1% reduction in the relative risk of hospitalisation.²⁶ Specifically, a 3-point difference
19 in MCS is associated with an odds ratio of 1.13 for being unable to work or an odds ratio of
20 1.16 for 1-year job loss. The probability of using mental health services is increased by
21 approximately 30% (odds ratio = 1.31), and there is a 30% increased risk of depression (odds
22 ratio = 1.34). It is also associated with a 10% higher 1-year mortality risk (odds ratio – 1.10).

23 ²²

1

2 The continued improvements in mental health-related quality of life determinants resulting
3 from the 6-month Kidney BEAM intervention in the *intention-to-treat* analysis, were
4 accompanied by an increase in physical health-related quality of life determinants that weren't
5 observed at the 12-week assessment point. Mean KDQoL PCS scores in the intervention group
6 increased ($p=0.055$ in *intention-to-treat*; $p<0.0001$ in *per protocol* analysis) and were driven
7 by improvements in the sub-scales of the KDQoL questionnaire that make up the composite
8 score; including significant improvements in scores in the *intention-to-treat* population in role
9 physical, physical functioning, pain and general health. It is postulated that the perception of
10 being able to complete, participate, and be confident in undertaking physical tasks may require
11 an initial improved psychological perspective and the physiological gain in physical function
12 associated with an initial supervised programme, to achieve longer term gains in perception of
13 physical well-being. A structured physical activity programme as a 'kick-start' precursor to
14 physical health-related quality of life improvements, consolidated with a further 12 weeks of
15 self-managed physical activity behaviour appears to be essential to realise important physical
16 health-related quality of life gains in a patient population where high levels of sedentary
17 behaviour are common and the role of exercise counselling to improve both mental and
18 physical health outcomes is far from routine in kidney care management.¹⁰

19

20 This trial revealed that the Kidney BEAM 6-month physical activity digital health intervention,
21 specifically designed for people living with CKD, significantly improved mental HRQoL
22 compared with wait-list control with a 93% and 98% chance of the Kidney BEAM intervention
23 being cost-effective compared to wait-list control at a willingness to pay of £20,000 and
24 £30,000 per QALY gained. Every increment in QALYs resulting from a 6-month programme

1 of Kidney BEAM is associated with an ICER of £3,445.56, and every increment of 1AU in the
2 KDQoL MCS is associated with an ICER of £14.44. Assuming comparative effectiveness of
3 the kidney BEAM intervention compared with in-person kidney rehabilitation,^{27, 28} the
4 average cost implication is £708 per participant per year for in-person rehabilitation compared
5 to £15 per participant per year for delivery of the kidney BEAM intervention, a suggested cost
6 saving of £693 per participant.

7
8 Digital health interventions present a real opportunity for healthcare payers such as the NHS
9 to deliver essential services where fiscal resource and workforce are not available to deliver
10 face-to-face care. Furthermore, digital interventions offer convenience for patients who
11 participate from home and choose when to exercise. The Kidney BEAM digital health
12 intervention is the first virtual solution in the kidney rehabilitation space to be proven to be
13 cost-effective. Cost benefits of a similar magnitude have been realised with in-person and
14 home-based exercise interventions in other long-term condition populations, such as people
15 with cardiac and pulmonary conditions,²⁹⁻³¹ and a recent systematic review revealed that
16 cardiac rehabilitation digital health interventions were as cost effective as in-person cardiac
17 rehabilitation.³² Kidney Beam has now been rolled-out across all eight regions of England as
18 part of an implementation project in preparation for commissioning. Results from the Kidney
19 Beam Trial, together with practical experience gained through NHS implementation, will
20 ensure that there is a clear plan for long-term adoption by the NHS. Additionally, because the
21 Kidney BEAM programme is delivered online from a single centre, it is simple to establish in
22 a wide variety of health care systems and to offer to people across large geographical areas.

23

1 The Kidney BEAM physical activity digital health intervention was developed using the
2 Behaviour Change Wheel methodology,³³ a methodology based upon 19 frameworks of
3 behaviour change theory including the transtheoretical model of behaviour change.^{34,35} Careful
4 consideration and preparation of a logic model¹⁸ that incorporated key intervention functions
5 to facilitate a change in behaviour and overcome common barriers to engagement with physical
6 activity³⁶ was co-developed with people with lived experience and experts in the field. The
7 intention of the initial 12-week structured programme of physical activity was to support people
8 living with CKD to make important initial physiological and psychological gains in health
9 outcomes to promote and sustain self-managed physical activity behaviour following
10 completion of the programme. Evidence suggests that for meaningful behaviour change to be
11 achieved, there is a need for the 'active' behaviour to be maintained over a 6-month period.³⁷
12 The Kidney BEAM intervention was deliberately designed to meet this expectation, combining
13 the initial 12-week structured and supported physical activity digital health intervention with a
14 12-week self-managed digital health intervention component. This type of 'kick-start'
15 programme has been successfully utilised in in-person kidney-specific rehabilitation²⁷ as well
16 as in-person physical rehabilitation for other chronic conditions³⁸⁻⁴⁰ and has resulted in a
17 maintenance of health outcome gains and physical activity behaviour in the longer term²⁹⁻³¹.

18

19 The significant improvement we continue to report in the KDQoL mental component summary
20 at 6 months was likely driven by changes in the KDQoL subscales of emotional wellbeing, role
21 emotional, social function and vitality (energy/fatigue) scales, as these sub-scales are more
22 heavily weighted in the calculation of the MCS score. However, the improved physical
23 functioning, role physical, bodily pain and general health scores were also all improved, so
24 those subscales will also have contributed to the improvement in MCS score. It is noteworthy
25 that improvements in mental HRQoL, patient activation, and physical function were realised

1 at 12 weeks ¹⁴ suggesting the BEAM platform ‘kick-started’ improvements in health-related
2 quality of life during the initial 12-week structured component of the intervention. It is
3 encouraging to witness sustained and continued mental health-related quality of life gains with
4 the self-managed physical activity component of the intervention, particularly in a patient
5 population where lower patient activation levels have been recognised and are associated with
6 a lower HRQoL in people living with CKD. ⁴¹

7
8 The Kidney Beam Trial was inclusive of people living with CKD from across the disease
9 trajectory, including pre-dialysis and those people requiring dialysis treatment or living with a
10 kidney transplant. Whilst it is acknowledged that the mental burden of symptoms associated
11 with kidney disease, which vary along with disease stage and are highest among dialysis
12 recipients, ⁴² may be a challenge to treat with a one-size-fits-all physical activity DHI, the
13 inclusion of a seated option as well as a standing option for performing the activity did allow
14 for an inclusive approach and the health coaching provided by the physiotherapy assistant
15 encouraged a tailored approach to commencement and progression of the programme for all
16 participants. The baseline global physical activity questionnaire (GPAQ) revealed a mean score
17 of only 110 minutes per week. The mean additional physical activity minutes recorded on the
18 platform was 44 minutes at 12 weeks, and 22 minutes at 6 months, almost 50% and 25%
19 increases respectively. Additionally, as the GPAQ may over-estimate scores, the increase in
20 physical activity as a result of the Kidney BEAM intervention is important, especially as even
21 small increases in physical activity can have a major impact upon health outcomes for this
22 patient population. ⁴ An adherence rate of 63% with the 12-week ‘kick-start’ programme may
23 be considered as moderate, but compared favourably with physical activity DHIs for other
24 long-term conditions (55%) ⁴³ and face-to-face renal rehabilitation programmes (59%). ²⁸
25 Although we aimed to encourage participant engagement with behavioural change techniques

1 such as motivational interviewing, it is acknowledged that further work to personalise digital
2 health interventions may lead to better engagement with these physical activity interventions.

3

4 A limitation of the trial was the restriction of the trial sites to a single country and delivery of
5 the intervention in the English language only. Whilst the Kidney BEAM physical activity
6 platform was deliberately co-developed with people living with the condition, including people
7 with generally poor digital literacy, people from lower socio-economic backgrounds, minority
8 ethnic groups and elderly patients, there is acknowledgement that further work is required to
9 meet the needs of these populations who are expected to benefit the most from health promoting
10 strategies in the setting of CKD, including digital health interventions. Sub-studies are
11 underway to expand relevant content, translate the website into other languages and address
12 digital literacy and access. These limitations may partially explain the limited recruitment rate
13 observed in the Kidney BEAM trial and does mean that the generalisability of the trial findings
14 to CKD populations worldwide will require further evaluation.

15

16 The primary and secondary outcomes were self-reported and as participants were not blinded
17 to the allocated treatment, this method will have produced bias. We could also not mask the
18 supporting physiotherapy assistants. However, the health economist and statisticians were
19 masked. Healthcare utilisation for primary and social care were collected via patient interview,
20 which may have introduced recall bias. Concurrent medication usage and sleep quality were
21 not analysed as part of this current trial, and it is acknowledged that these may affect mental
22 health-related quality of life. There was a dropout rate of 39.8% from the intervention group at
23 6 months which required data to be imputed and may increase imprecision in estimates. There
24 was no obvious difference in participant characteristics between groups for complete and

1 incomplete cases and over 75% of the dropouts were within the first week of the trial. The last
2 observation carried forward approach to missing data generally offers a conservative estimate
3 of the patient's outcome trajectory in a study,⁴⁴ but can lead to an overestimation of the size of
4 the effect of the intervention. Per protocol analyses were conducted to confirm the results.

5

6 Recruitment for this trial was during the COVID-19 pandemic, a time where recruitment to
7 trials was particularly challenging, especially for more vulnerable patients (such as the elderly
8 and those with comorbidities). This contributed to the slightly younger and less comorbid
9 population we recruited. However, the study recruited a more diverse and representative
10 population than previous exercise interventions.⁴⁵ The inclusion of earlier CKD stages was a
11 strength of this current study, as most healthcare systems do not have capacity to support these
12 patients using traditional methods of face-to-face exercise intervention.

13

14 Participants from the waiting-list control group were offered access to the kidney beam
15 intervention at 12 weeks. We acknowledge that it would have been ideal to ask people from
16 the waitlist control group to wait until 6 months to access the platform but as this randomised
17 controlled trial was conducted during the COVID-19 pandemic, withholding access to a
18 potentially useful intervention for promoting mental HRQOL was deemed unethical. Only 11%
19 of people from this group chose to access the platform during this time, but it is acknowledged
20 that this may have led to an underestimation of the size of the effect between the Kidney BEAM
21 group and the wait-list control group.

22

23 Overall, this trial demonstrates that the Kidney BEAM physical activity platform is a clinically
24 beneficial and cost-effective digital health intervention to improve mental health related quality

1 of life in people with CKD. The results provide evidence to support commissioning within the
2 UK NHS.

3

4 **Data availability**

5 Data collected during the study, including deidentified participant data will be made available
6 on reasonable request, and following trial steering committee approval, by contacting
7 corresponding author on sharlene.greenwood@nhs.net. The study protocol, statistical
8 analysis plan, and other study forms can be obtained by visiting BMC Nephrology.²⁴ The
9 health economic analysis plan can be found in supplemental files.

10

11 **Acknowledgements**

12 This study was funded by a grant from Kidney Research UK: SP/BEAM/2020 HY, JB, RB,
13 NB, MGB, TW are supported by the National Institute for Health Research Leicester
14 Biomedical Research Centre. HW is funded by the National Institute for Health Research
15 (NIHR) Applied Research Collaboration East Midlands (ARC EM). The views expressed are
16 those of the authors and not necessarily those of the NHS, the National Institute for Health
17 Research, or the Department of Health.

18

19 **Funding**

20 The trial was funded by a research grant from Kidney Research UK; the funders of the trial had
21 no role in trial design, data collection, data analysis, data interpretation, or involvement in
22 writing of the report.

23

1 **Authors contributions**

2 Authorship followed ICMJE guidelines. S.G, H.Y, R.B, N.B, J.B, E.C, Z.S, H.N, A.H, K.B,
3 A.N, M.G.B, T.W and J.M were responsible for the inception and design of the project and
4 prepared the manuscript. S.G, H.Y, R.B, N.B, J.B, E.C, A.H, V.D, H.N, A.N, T.W, N.B, J.C,
5 N.C, H.W, S.B, J.B, P.K, P.A.K, D.W, J.T, M.J, M.T, J.B, E.A, K.M, Z.S, M.G.B, T.W, JM
6 contributed to the design of the study, provided methodological input, and wrote the manuscript
7 text and prepared tables 1-3. All authors reviewed the manuscript.

8

9 **Conflict of interest**

10 King's College Hospital NHS Trust and SG were involved in the conception and development
11 of Kidney BEAM. SG became a director of Kidney Beam Ltd in August 2023. SB was a
12 previous Trustee of Kidney Research UK. DW has an ongoing consultancy contract with
13 AstraZeneca and has received honoraria/consultancy fees from Astellas, Boehringer
14 Ingelheim, Bayer, Eledon, Galderma, GlaxoSmithKline, Gilead, Janssen, Mundipharma,
15 ProKidney, Tricida, Vifor and Zydus for activities related to education and clinical trials. JC
16 and NC were both independent contractors but were paid by the grant.

17

18 **Supplementary material**

19

20 Supplementary data table 1: Summary of QALYS (PDF)

21 Supplementary data table 2: Base-case model KDQoI MCS (PDF)

22 Supplementary data table 3: Sensitivity analysis reporting different values for cost of the
23 intervention per person (PDF)

24 Supplementary data table 4: Summary of observed costs by category, time-period and
25 intervention group (PDF)

26 Supplementary data table 5: Sources of resource use and unit costs (PDF)

27 Supplementary data table 6: Response of primary and secondary outcomes to the Kidney
28 BEAM intervention (per protocol analyses) (PDF)

29 Supplementary Data Table 7: Comparison of missing data between complete cases and
30 missing cases due to trial dropouts (PDF)

31 Supplementary Data Table 8: Table of missingness in the data

- 1 CONSORT 2010 Checklist of information to include when reporting a randomised trial
- 2
- 3
- 4
- 5 Supplementary information is available at KI Report's website.

Journal Pre-proof

References

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

1. Kovesdy, C.P., *Epidemiology of chronic kidney disease: an update 2022*. *Kidney International Supplements*, 2022. **12**(1): p. 7-11.
2. Zelle, D.M., et al., *Physical inactivity: a risk factor and target for intervention in renal care*. *Nature Reviews Nephrology*, 2017. **13**(3): p. 152-168.
3. Rampersad, C., et al., *Association of Physical Activity and Poor Health Outcomes in Patients With Advanced CKD*. *American Journal of Kidney Diseases*, 2021. **78**(3): p. 391-398.
4. Baker, L.A., et al., *Clinical practice guideline exercise and lifestyle in chronic kidney disease*. *BMC Nephrology*, 2022. **23**(1).
5. Group, K.D.I.G.O.K.C.W., . *KDIGO 2012 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease*. *Kidney international*, 2012: p. 1-150.
6. Bennett, P.N., et al., *Physical activity and exercise in peritoneal dialysis: International Society for Peritoneal Dialysis and the Global Renal Exercise Network practice recommendations*. *Peritoneal Dialysis International*, 2022. **42**(1): p. 8-24.
7. Greenwood, S.A., et al., *Mortality and morbidity following exercise-based renal rehabilitation in patients with chronic kidney disease: The effect of programme completion and change in exercise capacity*. *Nephrology Dialysis Transplantation*, 2019. **34**(4): p. 618-625.
8. Greenwood, S.A., et al., *Exercise counselling practices for patients with chronic kidney disease in the UK: a renal multidisciplinary team perspective*. *Nephron Clinical Practice*, 2014. **128**(1-2): p. 67-72.
9. Bennett, P.N., et al., *Global Policy Barriers and Enablers to Exercise and Physical Activity in Kidney Care*. *Journal of Renal Nutrition*, 2022. **32**(4): p. 441-449.
10. Coyne, E., et al., *Achieving consensus on psychosocial and physical rehabilitation management for people living with kidney disease*. *Clinical Kidney Journal*, 2023.
11. March, D.S., et al., *A Cost-Effective Analysis of the CYCLE-HD Randomized Controlled Trial*. *Kidney Int Rep*, 2021. **6**(6): p. 1548-1557.
12. Organization, W.H., *Global strategy on digital health 2020-2025*, W.H. Organization, Editor. 2021.
13. Association, A.D., *1. Promoting Health and Reducing Disparities in Populations*. *Diabetes Care*, 2016. **40**(Supplement_1): p. S6-S10.
14. Greenwood, S.A., et al., *Evaluating the effect of a digital health intervention to enhance physical activity in people with chronic kidney disease (Kidney BEAM): a multicentre, randomised controlled trial in the UK*. *The Lancet Digital Health*.
15. Abraham, C. and S. Denford, *Design, Implementation, and Evaluation of Behavior Change Interventions: A Ten-Task Guide*, in *The Handbook of Behavior Change*, M.S. Hagger, et al., Editors. 2020, Cambridge University Press: Cambridge. p. 269-284.
16. Tong, A., et al., *Implementing core outcomes in kidney disease: report of the Standardized Outcomes in Nephrology (SONG) implementation workshop*. *Kidney Int*, 2018. **94**(6): p. 1053-1068.
17. Walklin, C.G., et al., *The effect of a novel, digital physical activity and emotional well-being intervention on health-related quality of life in people with chronic kidney disease: trial design and baseline data from a multicentre prospective, wait-list randomised controlled trial (kidney BEAM)*. *BMC Nephrology*, 2023. **24**(1): p. 122.

- 1 18. Young, H.M.L., et al., *The development and internal pilot trial of a digital physical*
2 *activity and emotional well-being intervention (Kidney BEAM) for people with chronic*
3 *kidney disease*. Scientific Reports, 2024. **14**(1): p. 700.
- 4 19. Unit, P.S.S.R., *Unit costs of health and social care*
5 . 2022.
- 6 20. NHS England. Available from: [https://www.england.nhs.uk/costing-in-the-](https://www.england.nhs.uk/costing-in-the-nhs/national-cost-collection/#ncc1819)
7 [nhs/national-cost-collection/#ncc1819](https://www.england.nhs.uk/costing-in-the-nhs/national-cost-collection/#ncc1819).
- 8 21. Formulary, B.N., 2022.
- 9 22. Maruish ME, K.M., Bjorner JB, Gandek B, Turner-Bowker DM, Ware JE. , *Quality Metric*
10 *Incorporated; 2011. Chapter 10.*, in *User's manual for the SF-36v2 health survey*. 2011,
11 Lincoln (RI).
- 12 23. Husereau, D., et al., *Consolidated Health Economic Evaluation Reporting Standards*
13 *2022 (CHEERS 2022) statement: updated reporting guidance for health economic*
14 *evaluations*. *Bjog*, 2022. **129**(3): p. 336-344.
- 15 24. Lunn, D., et al., *WinBUGS - A Bayesian modeling framework: Concepts, structure and*
16 *extensibility*. *Statistics and Computing*, 2000. **10**: p. 325-337.
- 17 25. Nixon, R.M. and S.G. Thompson, *Methods for incorporating covariate adjustment,*
18 *subgroup analysis and between-centre differences into cost-effectiveness evaluations*.
19 *Health Econ*, 2005. **14**(12): p. 1217-29.
- 20 26. Lowrie, E.G., et al., *Medical outcomes study short form-36: a consistent and powerful*
21 *predictor of morbidity and mortality in dialysis patients*. *Am J Kidney Dis.*, 2003. **41**(6):
22 p. 1286-92. doi: 10.1016/s0272-6386(03)00361-5.
- 23 27. Greenwood, S.A., et al., *Mortality and morbidity following exercise-based renal*
24 *rehabilitation in patients with chronic kidney disease: the effect of programme*
25 *completion and change in exercise capacity*. *Nephrol Dial Transplant*, 2018.
- 26 28. Greenwood, S.A., et al., *Evaluation of a pragmatic exercise rehabilitation programme*
27 *in chronic kidney disease*. *Nephrology Dialysis Transplantation*, 2012. **27**(SUPPL. 3): p.
28 iii126-iii134.
- 29 29. Davidson, P.M., et al., *Can a heart failure-specific cardiac rehabilitation program*
30 *decrease hospitalizations and improve outcomes in high-risk patients?* *Eur J Cardiovasc*
31 *Prev Rehabil*, 2010. **17**(4): p. 393-402.
- 32 30. Hui, K.P. and A.B. Hewitt, *A simple pulmonary rehabilitation program improves health*
33 *outcomes and reduces hospital utilization in patients with COPD*. *Chest*, 2003. **124**(1):
34 p. 94-7.
- 35 31. Griffiths, T.L., et al., *Results at 1 year of outpatient multidisciplinary pulmonary*
36 *rehabilitation: a randomised controlled trial*. *Lancet*, 2000. **355**(9201): p. 362-8.
- 37 32. Batalik, L., et al., *The cost-effectiveness of exercise-based cardiac telerehabilitation*
38 *intervention: a systematic review*. *Eur J Phys Rehabil Med*, 2023. **59**(2): p. 248-258.
- 39 33. Michie, S., Van Stralen, M.M. and West, R., *The behaviour change wheel: a new*
40 *method for characterising and designing behaviour change interventions*.
41 *Implementation Science*, 2011. **6**(1): p. 42.
- 42 34. Prochaska, J.O. and C.C. DiClemente, *Stages and processes of self-change in smoking:*
43 *Towards an intergrated model of change*. *Journal of consulting and clinical Psychology*,
44 1983. **51**(3): p. 390-395.
- 45 35. Prochaska, J.O. and C.C. DiClemente, *Transtheoretical Therapy: Toards a more*
46 *integrative model of change*. *Psychotherapy: theory, research and practice*, 1982.
47 **19**(3): p. 276-288.

- 1 36. Mayes, J., et al., *Cultural influences on physical activity and exercise beliefs in patients*
2 *with chronic kidney disease: The Culture-CKD Study'-a qualitative study*. *BMJ Open*,
3 2022. **12**(1).
- 4 37. Jiménez-Zazo, F., et al., *Transtheoretical Model for Physical Activity in Older Adults:*
5 *Systematic Review*. *Int J Environ Res Public Health*, 2020. **17**(24).
- 6 38. Bolton, C.E., et al., *British Thoracic Society guideline on pulmonary rehabilitation in*
7 *adults: accredited by NICE*. *Thorax*, 2013. **68**(Suppl 2): p. ii1-ii30.
- 8 39. Silberman, A., et al., *The effectiveness and efficacy of an intensive cardiac*
9 *rehabilitation program in 24 sites*. *Am J Health Promot*, 2010. **24**(4): p. 260-6.
- 10 40. Clarkson, P., et al., *Digital tools to support the maintenance of physical activity in*
11 *people with long-term conditions: A scoping review*. *Digit Health*, 2022. **8**: p.
12 20552076221089778.
- 13 41. Magadi, W., et al., *Patient activation and its association with symptom burden and*
14 *quality of life across the spectrum of chronic kidney disease stages in England*. *BMC*
15 *Nephrology*, 2022. **23**(1): p. 45.
- 16 42. Fletcher, B.R., et al., *Symptom burden and health-related quality of life in chronic*
17 *kidney disease: A global systematic review and meta-analysis*. *PLoS Med*, 2022. **19**(4):
18 p. e1003954.
- 19 43. Jakob, R., et al., *Factors Influencing Adherence to mHealth Apps for Prevention or*
20 *Management of Noncommunicable Diseases: Systematic Review*. *J Med Internet Res*,
21 2022. **24**(5): p. e35371.
- 22 44. Liu, X., *Chapter 14 - Methods for handling missing data*, in *Methods and Applications*
23 *of Longitudinal Data Analysis*, X. Liu, Editor. 2016, Academic Press: Oxford. p. 441-473.
- 24 45. Heiwe, S. and S.H. Jacobson, *Exercise training for adults with chronic kidney disease*.
25 *Cochrane Database of Systematic Reviews*, 2011. **Oct 5**(10): p. CD003236.

1 **Fig. 1: Flowchart of participants through the trial.**

2

3 **Fig. 2: Cost-effectiveness plane with 95% confidence region.**

4

5 **Table 1.** Baseline demographic data.

Journal Pre-proof

	n	All	n	Kidney BEAM	n	Waitlist control
Age (years) (SD)	340	53.8 (13.5)	173	53.9 (13.6)	167	53.8 (13.5)
Sex (n) (%)	340		173		167	
Male		185 (54)		96 (55)		89 (53)
Female		155 (46)		77 (45)		78 (47)
Ethnicity (n) (%)	339		173		166	
Black		39 (11.5)		20 (11.6)		19 (11.4)
White		254 (74.9)		127 (73.4)		127 (76.5)
Asian		39 (11.5)		22 (12.7)		17 (10.2)
Biracial		7 (2.1)		4 (2.3)		3 (2.1)
Body Mass Index (kg/m²) (IQR)	327	28.4 (24.8, 33.3)	165	27.9 (24.7, 33.4)	162	28.8 (24.9, 33.0)
Smoking (n) (%)	339		172		167	
Current		16 (4.7)		5 (2.9)		11 (6.6)
Former		130 (38.3)		77 (44.8)		53 (31.7)

Never		193 (56.9)		90 (52.3)		103 (61.7)
Alcohol consumption (n) (%)	339		172		167	
More than recommended		26 (7.7)		14 (8.1)		12 (7.2)
Less than recommended		174 (51.3)		89 (51.7)		85 (50.9)
Non-drinker		139 (41.0)		69 (40.1)		70 (41.9)
Blood pressure (mm Hg) (SD)	307		154		153	
SBP		136.5 (18.4)		135.3 (19.3)		137.8 (17.5)
DBP		79.7 (10.7)		78.6 (11.1)		80.7 (10.2)
Resting heart rate (bpm) (SD)	207	77.6 (14.7)	103	77.8 (14.6)	104	77.3 (14.8)
Medical History (n) (%)	340		173		167	
CVA		8 (2.4)		4 (2.4)		4 (2.4)
MI		8 (2.4)		3 (1.7)		5 (3)
Diabetes		76 (22.4)		37 (21.4)		39 (23.4)
Hypertension		235 (69.1)		115 (68.9)		120 (69.4)
Cause of kidney disease (n) (%)	340		173		167	
Diabetic nephropathy		31 (9.1)		13 (7.5)		18 (10.8)

Hypertension		38 (11.2)		21 (12.1)		17 (10.2)
Nephrosclerosis		1 (0.3)		1 (0.6)		0 (0)
IgA nephropathy		39 (11.5)		18 (10.4)		21 (12.6)
Tubulointerstitial nephritis		5 (1.5)		2 (1.2)		3 (1.8)
PKD		60 (17.6)		31 (17.9)		29 (17.4)
Obstructive nephropathy		7 (2.1)		2 (1.2)		5 (3)
Medullary sponge kidney disease		0 (0)		0 (0)		0 (0)
Membranous nephropathy		5 (1.5)		5 (2.9)		0 (0)
Lupus nephritis		5 (1.5)		4 (2.3)		1 (0.6)
Unknown		65 (19.1)		33 (19.1)		32 (19.2)
Other		84 (24.7)		43 (24.9)		41 (24.6)
CKD stage (%)	339		172		167	
Stage 2		55 (16.2)		27 (15.7)		28 (16.8)
Stage 3A		62 (18.3)		29 (16.9)		33 (19.8)
Stage 3B		76 (22.4)		45 (26.2)		31 (18.6)
Stage 4		67 (19.8)		34 (19.8)		33 (19.8)

Stage 5		79 (23.3)		37 (21.5)		42 (25.1)
Treatment modality (n) (%)	340		173		167	
Non-dialysis dependent kidney disease		160 (47)		75 (43)		85 (51)
Kidney transplant recipient		118 (35)		65 (38)		53 (32)
Dialysis therapy		62 (18)		33 (19)		29 (17)
HbA1c (mmol/mol)	124	39 (35, 48)	64	39 (34, 50)	60	39 (36, 47)
Creatinine (micromol/L)	332	159 (106, 293)	170	159 (109, 279)	162	161 (106, 330)
CRP (mg/L)	169	4 (2, 9)	92	3.9 (2, 10)	77	4 (2, 9)

- 1 Data are mean (standard deviation), median (interquartile range), or number (%), as appropriate. Kidney BEAM, Kidney BEAM intervention group (physical activity training
- 2 and education plus usual care); Waitlist control, waitlist control group; n, total number of available data; CKD, chronic kidney disease; CVA, cerebrovascular accident; MI,
- 3 myocardial infarction; SBP, systolic blood pressure; DBP, diastolic blood pressure; HbA1C, glycated haemoglobin; CRP, C-reactive protein; IQR, inter-quartile range; PKD,
- 4 polycystic kidney disease

1 **Table 2:** Response of primary and secondary outcome measures to the Kidney BEAM intervention (intention to treat analysis)

Outcome measure	n	Baseline mean (SD)	6 months mean (SD)	Mean difference in change between groups (Kidney BEAM - waitlist control) mean {95% CI}	p value	Observed power
Primary outcome						
KDQoL MCS (AU)						
Kidney BEAM	171	44.6 (10.8)	48.7 (10.5)	5.9 (4.4-7.5)	<.0001	1.00
Waitlist control	167	48.1 (10.5)	43.5 (10.3)			
Secondary outcomes						
KDQOL PCS (AU)						
Kidney BEAM	171	40.0 (11.7)	42.9 (11.02)	1.5 (-0.03-2.9)	0.055	0.48
Waitlist control	167	41.3 (11.2)	42.5 (11.3)			
Symptom problem list						

Kidney BEAM	140	76.6 (18.2)	77.8 (17.9)	0.6 (-2.2-3.3)	0.67	0.07
Waitlist control	143	79.9 (16.8)	79.7 (18.7)			
Effects of Kidney Disease						
Kidney BEAM	166	69.1 (26.5)	72.3 (26.1)	1.0 (-2.8-4.9)	0.59	0.08
Waitlist control	161	75.6 (23.6)	76.3 (26.2)			
Burden of kidney disease						
Kidney BEAM	172	55.1 (31.2)	61.7 (30.7)	5.3 (2.0-8.6)	0.0017	0.88
Waitlist control	167	64.9 (30.5)	64.7 (29.9)			
Work status						
Kidney BEAM	84	61.8 (40.6)	61.2 (38.1)	-5.2 (-12.3-2.0)	0.15	0.29
Waitlist control	120	61.7 (41.4)	65.8 (37.8)			
Cognitive function						
Kidney BEAM	172	74.7 (19.3)	78.5 (17.9)	2.3 (-0.3-4.9)	0.082	0.41
Waitlist control	167	78.7 (19.5)	78.5 (17.9)			
Quality of social interaction						
Kidney BEAM	172	72.0 (18.9)	76.9 (17.7)	7.1 (4.1-10.0)	<.0001	1.00

Waitlist control	167	73.6 (18.2)	70.8 (18.7)			
Sexual function						
Kidney BEAM	102	42.3 (41.6)	41.5 (41.1)	-3.4 (-11.7-5.0)	0.427	0.124
Waitlist control	102	48.5 (41.7)	49.1 (43.4)			
Sleep						
Kidney BEAM	171	55.6 (19.5)	60.6 (18.7)	6.5 (3.5-9.5)	<.0001	0.99
Waitlist control	166	57.7 (20.3)	55.7 (21.0)			
Social support						
Kidney BEAM	158	72.7 (27.6)	77.0 (25.7)	4.0 (-1.0-9.0)	0.117	0.35
Waitlist control	150	75.7 (28.3)	74.7 (28.7)			
Dialysis staff encouragement						
Kidney BEAM	77	78.7 (24.3)	75.8 (26.3)	-6.1 (-12.2- -0.03)	0.049	0.51
Waitlist control	68	77.2 (27.4)	80.7 (27.3)			
Overall health						
Kidney BEAM	85	60.1 (19.9)	62.6 (18.0)	-1.3(-5.5-2.9)	0.55	0.09
Waitlist control	118	58.1 (18.1)	62.5 (20.1)			

Patient satisfaction						
Kidney BEAM	93	73.5 (22.8)	75.6 (21.2)	1.8 (-2.6-6.3)	0.417	0.128
Waitlist control	87	73.7 (24.3)	74.1 (22.4))			
Physical functioning						
Kidney BEAM	171	60.9 (30.1)	68.0 (28.2)	6.29 (2.9-9.7)	0.0003	0.95
Waitlist control	167	64.2 (30.7)	64.3 (30.5)			
Role physical						
Kidney BEAM	171	48.1 (41.8)	62.9 (42.6)	9.1 (1.8-16.3)	0.014	0.69
Waitlist control	167	51.0 (43.4)	55.4 (44.3)			
Pain						
Kidney BEAM	172	61.1 (26.4)	66.7 (26.0)	8.0 (3.8-12.2)	0.0002	0.96
Waitlist control	167	67.8 (27.7)	63.6 (29.8)			
General health						
Kidney BEAM	171	40.3 (21.6)	45.1 (22.2)	4.3 (1.6-7.0)	0.0018	0.88
Waitlist control	167	42.7 (21.6)	42.7 (22.0)			
Emotional wellbeing						

Kidney BEAM	171	67.0 (20.5)	74.3 (20.2)	4.0 (-1.0-9.0)	<0.0001	0.35
Waitlist control	167	70.3 (18.7)	65.9 (19.6)			
Role emotional						
Kidney BEAM	171	60.5 (42.5)	72.1 (39.4)	10.7 (3.1-18.4)	0.0058	0.79
Waitlist control	166	63.2 (42.3)	55.9 (43.7)			
Social function						
Kidney BEAM	172	61.6 (27.6)	69.4 (27.9)	10.1 (6.3-13.8)	<0.0001	1.00
Waitlist control	167	64.3 (30.2)	61.3 (28.9)			
Energy/fatigue						
Kidney BEAM	171	42.6 (21.4)	53.1 (23.1)	15.5 (12.6-18.4)	<0.0001	1.00
Waitlist control	167	45.0 (23.3)	39.5 (22.6)			
EQ-5D-3L utility score						
Kidney BEAM	171	0.65 (0.25)	0.71 (0.25)	0.10 (0.07-0.13)	<0.0001	1.00
Waitlist control	167	0.73 (0.23)	0.68 (0.26)			

- 1 Data are mean (standard deviation), median (interquartile range), or mean {95% confidence interval} ANCOVA adjusted scores. Control, waitlist control group (usual care);
- 2 Kidney BEAM, Kidney BEAM intervention group (physical activity training and education plus usual care); KDQOL, Kidney Disease Quality of Life Short Form (KDQOL-SF 1.3); MCS, Mental Component Score; PCS, Physical Component Summary; AU, arbitrary units; EQ-5D-3L, EuroQol five-dimension descriptive system.
- 3

Table 3 Base case model (assumes intervention £15 per person per year)

	Base case model: LVCF for missing cost components adjusted for baseline costs and EQ-5D	Complete case analysis adjusted for baseline costs and EQ-5D
N: WL	132 ⁺	92*
N: KB	91 ⁺	66*
Mean difference in Cost	£93.03 (-£360.60 to £613.40)	£273.60 (-£323 to £996.7)
Mean difference in QALYs	0.027 (0.013 to 0.040)	0.026 (0.009 to 0.043)
Incremental Cost effectiveness ratio (ICER)	£3,445.56	£10,523.08
Probability CE @ £20,000 per QALY gained	0.93	0.75
Probability CE @ £30,000 per QALY gained	0.98	0.87

Calculated at the average baseline value of cost (£1850) and EQ-5D score (0.70)


⁺ Excludes individuals with missing EQ-5D and cost baseline data (3 WL, 1 KB)

* Excludes individuals with missing EQ-5D and cost baseline data (1 WL, 1 KB)

Table 4 Number of patients with at least one Serious Adverse Event by MedDRA system organ class during the Kidney BEAM Trial

	All	Kidney BEAM	Waitlist control
	n (%)	n (%)	n (%)
Number of randomised patients who attended baseline visit	340	173	167
Number of patients with any event	9 (2)	4 (3)	5 (3)
Gastrointestinal disorders	2 (1)	1 (1)	1 (1)
Infections and infestations	4 (1)	2 (1)	2 (1)
Injury, poisoning and procedural complications	2 (1)	1 (1)	1 (1)
Renal and urinary disorders	1 (0.3)	0 (0)	1 (1)

Journal Pre-proof

Fig. 1: Flowchart of participants through the trial.

Note only 338 participants were included when analysing the primary outcome as two participants were missing both baseline and 6-month data.

