



King's Research Portal

DOI:

[10.1016/S0140-6736\(24\)02035-X](https://doi.org/10.1016/S0140-6736(24)02035-X)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Liu, L., Marshall, I. J., Bhalla, A., Wolfe, C., O'Connell, M., & Wang, Y. (2024). Associations between depression and mortality up to 15-years after stroke: a population-based cohort study. *Lancet*, 404(Special Issue), Article S75. [https://doi.org/10.1016/S0140-6736\(24\)02035-X](https://doi.org/10.1016/S0140-6736(24)02035-X)

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Associations between depression and mortality up to 15-years after stroke: a population-based cohort study

Lu Liu, Iain J Marshall, Ajay Bhalla, Charles Wolfe, Matthew DL O'Connell*, Yanzhong Wang*

Abstract

Background Limited data are available on the long-term mortality of post-stroke depression (PSD). We aim to estimate the associations between PSD and mortality up to 15-years after stroke and assess the differences by ethnicity.

Methods This is a secondary analysis of data from a prospective, population-based cohort study (the South London Stroke Register). Depression was assessed using the Hospital Anxiety and Depression Scale at 3-months after stroke (scores >7 = depression). Associations between depression at 3-months and mortality up to 15 years after stroke were estimated with Cox regression models adjusted for age, sex, ethnicity, smoking, social support, stroke severity, treatment with antidepressants and comorbidities (hypertension, diabetes and heart disease).

Findings A total of 2574 survivors were assessed at 3-months after stroke and 913 (35.5%) had depression. The number of deaths was 651 within 5-years, 958 within 10-years and 1101 within 15-years after stroke. Compared to stroke survivors without depression, those with depression had higher risks of mortality (adjusted hazard ratio [aHR] 1.20 [95% confidence interval (CI) 1.01–1.42]) within 5-years after stroke, but similar risks of mortality over 10-years (aHR 1.10, 95% CI 0.95–1.26) and 15-years (aHR 1.08, 95% CI 0.95–1.24) after stroke. In White ethnic group, stroke survivors with and without PSD had similar rates of mortality (aHR 1.08, 95% CI 0.95–1.24) within 5-years, while in Black ethnic group, survivors with depression had greater risks of mortality (aHR 1.45, 95% CI 1.01–2.10) than patients without depression.

Interpretation PSD is associated with higher mortality within 5-years after stroke and the association was evident in Black group. Depression at 3-months is not associated with mortality beyond 5-years after stroke. Targeted interventions to depression could be facilitated on the Black ethnic group within the first 5-years after stroke. These findings were limited to patients completing the depression assessment, who tended to have less severe stroke than excluded patients, so may not be generalized to all stroke survivors.

Funding Authors IM, AB, CW, MOC and YW received funding from the National Institute for Health and Care Research (NIHR) (Grant Reference Number: NIHR202339). Author LL received financial support from China Scholarship Council PhD Scholarship (CSC No. 202108310074). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright © 2024 Elsevier Ltd. All rights reserved, including those for text and data mining, AI training, and similar technologies.

Contributors

LL: funding acquisition, conceptualization, formal analysis, writing—original draft; IJM: funding acquisition, writing—review & editing; AB: funding acquisition, writing—review & editing; CW: funding acquisition, writing—review & editing; MDLO'C: funding acquisition, conceptualization, supervision, writing—review & editing; and YW: funding acquisition, conceptualization, supervision, writing—review & editing.

Declaration of interests

The authors declare no conflict of interest.

Date availability

Because of the sensitive nature of the data collected for this study, requests to access the data set for academic use should be made to the South London Stroke Register (SLSR) team: <https://www.kcl.ac.uk/lsm/research/divisions/hscr/research/groups/stroke/index.aspx>.

Acknowledgements

We thank the patients, their families, and the fieldworkers who have collected data for the SLSR since 1995.

Published Online
November 28, 2024

*Joint senior authors

School of Life Course and Population Sciences, King's College London, London, United Kingdom (L Liu MM, IJ Marshall PhD, A Bhalla MD, Prof C Wolfe PhD, M D L O'Connell PhD, Prof Y Wang PhD); NIHR Applied Research Collaboration (ARC) South London, London, United Kingdom (IJ Marshall PhD, Prof C Wolfe PhD, Prof Y Wang PhD); Department of Ageing Health and Stroke, Guy's and St Thomas' National Health Service Foundation Trust and King's College London, United Kingdom (A Bhalla MD)

Correspondence to:
Lu Liu, School of Life Course and Population Sciences, King's College London, London, UK
lu.7.liu@kcl.ac.uk