



King's Research Portal

DOI: 10.1093/schbul/sbv186

Document Version Publisher's PDF, also known as Version of record

Link to publication record in King's Research Portal

Citation for published version (APA):

Kumari, V., Peters, E., Guinn, A., Fannon, D., Russell, T., Sumich, A., Kuipers, E., Williams, S. C. R., & Ffytche, D. H. (2015). Mapping Depression in Schizophrenia: A Functional Magnetic Resonance Imaging Study. *Schizophrenia Bulletin*. https://doi.org/10.1093/schbul/sbv186

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Schizophrenia Bulletin doi:10.1093/schbul/sbv186

Mapping Depression in Schizophrenia: A Functional Magnetic Resonance Imaging Study

Veena Kumari^{*,1,2}, Emmanuelle Peters¹, Ashley Guinn¹, Dominic Fannon¹, Tamara Russell⁴, Alexander Sumich¹, Elizabeth Kuipers^{1,2}, Steven C. R. Williams⁵, and Dominic H. ffytche³

¹Department of Psychology, King's College London, Institute of Psychiatry, Psychology and Neuroscience, London, UK; ²NIHR Biomedical Research Centre for Mental Health, South London and Maudsley NHS Foundation Trust, London, UK; ³Department of Old Age Psychiatry and Department of Neuroimaging Sciences, King's College London, Institute of Psychiatry, Psychology and Neuroscience, London, UK; ⁴Department of Psychosis Studies, King's College London, Institute of Psychiatry, Psychology and Neuroscience, London, UK; ⁵Department of Neuroimaging, King's College London, Institute of Psychiatry, Psychology and Neuroscience, London, UK; ⁵Department of Neuroimaging, King's College London, Institute of Psychiatry, Psychology and Neuroscience, London,

*To whom correspondence should be addressed; Department of Psychology (P077), Institute of Psychiatry, Psychology and Neuroscience, King's College London, De Crespigny Park, London SE5 8AF, UK; tel: 44-207-848-0233, fax: 44-207-848-5006, e-mail: veena.kumari@kcl.ac.uk

Depressive symptoms are common in schizophrenia, often left untreated, and associated with a high relapse rate, suicidal ideation, increased mortality, reduced social adjustment and poor quality of life. The neural mechanisms underlying depression in psychosis are poorly understood. Given reports of altered brain response to negative facial affect in depressive disorders, we examined brain response to emotive facial expressions in relation to levels of depression in people with psychosis. Seventy outpatients (final N = 63) and 20 healthy participants underwent functional magnetic resonance imaging during an implicit affect processing task involving presentation of facial expressions of fear, anger, happiness as well as neutral expressions and a (no face) control condition. All patients completed Beck Depression Inventory (BDI-II) and had their symptoms assessed on the Positive and Negative Syndrome Scale (PANSS). In patients, depression (BDI-II) scores associated positively with activation of the left thalamus, extending to the putamen-globus pallidus, insula, inferior-middle frontal and para-post-pre-central gyri during fearful expressions. Furthermore, patients with moderateto-severe depression had significantly higher activity in these brain regions during fearful expressions relative to patients with no, minimal, or mild depression and healthy participants. The study provides first evidence of enhanced brain response to fearful facial expressions, which signal an uncertain source of threat in the environment, in patients with psychosis and a high level of self-reported depression.

Key words: psychosis/depressive symptoms/ fMRI/fear/thalamus

Introduction

Depression has been regarded as a key feature of schizophrenia since it was first defined by Bleuler.¹ A large body of evidence confirms that depressive symptoms are a core component,^{2,3} and may precede the clinical manifestation^{4,5} of psychosis. Prevalence rates for major depressive disorder (MDD) comorbid with psychotic disorders commonly vary between 25% to 75%,^{6,7} and around 70% of chronic schizophrenia patients without a MDD may show signs of sub-syndromal depression.^{2,3} Although there is overlap between depressive and negative symptoms, a distinction between them can be made,^{8,9} with negative symptoms indicating the severity of illness and depressive symptoms reflecting motivational hopelessness about the future.8 Importantly, depressed mood in schizophrenia is known to have detrimental consequences, such as a high relapse rate, suicidal ideation, increased mortality, reduced social adjustment and poor quality of life.^{8,10-14} A key question, central to refining and developing effective treatments for depression in schizophrenia is: what are the psychobiological abnormalities associated with depressive symptoms in schizophrenia, and to what extent are they similar or different to those commonly observed in MDD?

A large number of functional magnetic resonance imaging (fMRI) studies have already examined the neurobiology of MDD, particularly using facial affect processing paradigms.¹⁵ MDD patients, relative to healthy participants, show hyper-activation to negative facial affect, in particular sadness and fear, in the amygdala, insula, parahippocampal gyrus, fusiform face area and putamen.¹⁶⁻¹⁹ Structural and functional alterations in these regions are

© The Author 2015. Published by Oxford University Press on behalf of the Maryland Psychiatric Research Center.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Downloaded from http://schizophreniabulletin.oxfordjournals.org/ at King's College London - Journals Dept on January 18, 2016

widely implicated in disturbed implicit/explicit processing of emotive stimuli and poor affect regulation in MDD.^{20–} ²² Schizophrenia patients, on the other hand, are reported to show, on average, reduced activation of the amygdala, parahippocampal gyrus, superior frontal gyrus, insula and basal ganglia to negative facial affect.^{23,24} There are no fMRI studies to our knowledge specifically examining the role of depression in brain responses to negative facial affect in schizophrenia. Amygdala activation to fearful faces, however, was found to correlate positively with the severity of flat affect, a negative symptom dimension, in a previous study.²⁵

There is indirect support from structural brain imaging and neuropsychological studies that depressed mood in schizophrenia and MDD could have at least partially overlapping neurobiology, such as volumetric abnormalities of the temporal lobe and anterior cingulate²⁶ and attentional impairment.²⁷ Depression in schizophrenia, however, differs markedly amongst patients depending on the stage of the illness, with more severe depressive symptoms in first episode patients, perhaps as a natural reaction to developing psychotic symptoms, relative to multi-episode chronic patients.^{28,29} Depression in later phases of the illness is found to be more subjective and best detected with self-report instruments,^{29,30} such as the Beck Depression Inventory.³¹

In this study, our aim was to elucidate the neural correlates of subjective depression, using whole brain fMRI during an established facial affect paradigm,³² in patients with schizophrenia or schizoaffective disorder. Assuming depression in chronic psychosis patients to share functional brain abnormalities associated with MDD,^{15–17} we expected to find a positive association between the level of self-reported depression and brain activity, particularly in the amygdala, insula, parahippocampal gyrus, fusiform face area and putamen, during negative facial affect processing. In addition, we studied a group of healthy participants to examine whether a positive association between depression and brain activity in patients, if found, reflected a hyperresponse, or a stronger response within the normal range, in patients with moderate-tosevere depression.

Methods and Materials

Participants and Design

The study involved 70 right-handed³³ outpatients with a DSM-IV diagnosis³⁴ of schizophrenia or schizoaffective disorder, recruited from the South London and Maudsley NHS Foundation Trust (SLaM), United Kingdom. Twenty right-handed³³ healthy participants (with no current or previous diagnosis of an Axis I disorder³⁵) matched, on average, to age and sex of the patient group were also studied.

All recruited patients met the following inclusion criteria: (1) aged 18-65 years, (2) on a stable dosage of

antipsychotic medication for at least 3 months prior to taking part, (3) no history of neurological impairments, systematic illness or head injury, (4) no current substance use disorder, and (5) able to provide written informed consent. The final study sample consisted of 63 patients (table 1), out of 70 recruited initially, for whom we had obtained usable fMRI data, symptom ratings using the Positive and Negative Syndrome Scale (PANSS)³⁶ and who had fully completed (within 1 wk of fMRI) the Beck Depression Inventory-II (BDI-II).³¹ PANSS assessments on patients were carried out by an experienced psychiatrist (D.F.) who was blind to patients' BDI-II scores.

The study procedures were approved by the ethics committee of the joint research ethics committee of the SLaM and the Institute of Psychiatry, London. All participants provided written informed consent prior to their participation and were compensated for their time and travel.

fMRI Paradigm and Procedure

The task involved presentation of black and white pictures of faces from Ekman and Friesen's set.³⁷ Three types of emotions (fear, anger, happy; 100% expression) and relatively neutral expressions were displayed in 30-second blocks of 8 pictures each, with each picture presented for 3.75 seconds. There were sixteen 30-second blocks with facial expressions (4 fear, 4 anger, 4 neutral, 4 happy), and each block was followed by a 15-second control block during which just an oval frame, as on facial expression trials and matched for luminance but without the face inside, appeared 4 times (order of presentation over the 12-minute experiment: neutral-control-fear-controlhappy-control-anger-control-fear-control-anger-controlneutral-control-happy-control-anger-control-happycontrol-fear-control-neutral-control-happy-control-neutral-control-anger-control-fear-control). The oval frame. rather than neutral expressions, was used as the control condition because it has been observed that people with schizophrenia^{38,39} or those at a high risk for psychosis⁴⁰ may activate brain regions to neutral faces that are normally associated with fear processing. On presentation of each facial expression, participants were required to indicate whether the face was a male/female by pressing the left (female)/right (male) button on a button box, simply to ensure they were attending to the stimuli. During the no face (oval frame) condition, attention was monitored by asking participants to press either button (left/ right) when the blank oval frame appeared. All participants underwent task familiarization in advance of the scan through practice of the gender discrimination task once on all identities used in the fMRI experiment on a laptop computer and an identical button box device. The task and administration procedures were identical to that used in a previous study.³² Participants were requested to abstain from alcohol for at least 24 hours prior to their scanning. They were provided with tea and decaffeinated

	Patients						
	All Patient With Schiz 7 With Sch Disorder; 7	s (N = 63; 56 ophrenia and izoaffective 3% Men)	No/Minimal Depression (N = 25; 22 Schizophrenia, 3 Schizoaffective Disorder, 80% Men)	Mild Depression (<i>N</i> = 17; 16 Schizophrenia, 1 Schizoaffective Disorder; 70.6% Men)	Moderate-to-Severe Depression (<i>N</i> = 21; 17 Schizophrenia, 4 Schizoaffective Disorder; 66.7% Men)	Healthy Partic $(n = 20;75\% \text{ M})$	ipants [en)
Clinical Measures	Mean (SD)	Range	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Range
Age Years in education Predicted IQ (ªNART) ^d bBDI-II	38.19 (9.77 13.87 (2.58 107.94 (9.6 16.79 (11.6) 19–61) 8–20 2) 86–124 8) 0–54	39.36 (9.82) 14.32 (2.72) 108.17 (10.08) 5.80 (4.07)	34.76 (8.23) 12.94 (2.70) 108.70 (10.29) 16.12 (1.49)	39.57 (10.61) 14.09 (2.23) 107.09 (8.92) 30.43 (7.38)	37.25 (12.49) 14.80 (2.98) 115.74 (7.46) 2.80 (2.91)	20-65 10-20 91-126 0-7
Age at illness onset (y) Duration of illness (y) Positive symptoms	25.59 (8.58 13.46 (9.56 16.08 (4.75 17.65 (4.82	$\begin{array}{c} 15-48 \\ 1-43 \\ 7-25 \\ 7-27 \\ 7$	24.64 (8.15) 14.72 (8.32) 15.80 (4.71) 16.52 (4.81)	25.12 (6.10) 9.71 (7.27) 16.64 (5.19) 18.94 (4.74)	24.57 (9.81) 15.00 (11.80) 19.95 (4.62) 17.95 (4.83)	n/a n/a n/a	
^c General psychopathology ^c PANSS: total symptoms Antipsychotic medication (mg)	32.22 (6.62 65.95 (13.4 449.51 (29:	$\begin{array}{cccc} & 18-56 \\ 8) & 37-108 \\ 5.21) & 100-1600 \end{array}$	29.76 (6.37) 62.08 (13.26) 430.80 (215.51)	34.18 (7.68) 69.76 (15.68) 495.58 (10.29)	33.57 (5.24) 67,48 (11.03) 433.50 (327.01)	n/a n/a n/a	
Antidepressant medication	N (%) 22 (34.9%)		N (%) 7 (28%), 2 on Fluoxetine, 2 Citalopram, 1 Sertraline, 1 Mirtazipine, 1 Clomipramine	N (%) 7 (41%), 2 on Fluoxetine, 1 Citalopram, 2 Venlafaxine, 1 Paroxetine, 1 Clomipramine	N (%) 8 (38%), 2 on Fluoxetine, 2 Citalopram, 1 Venlafaxine, 1 Mirtazapine, 1	$N_{(0,0)}^{(0,0)}(0,0)$	
Cigarette smoking Task performance	34 (54%) All patient		13 (52%) Low depression	9 (52.9%) Moderate depression	High depression	7 (35%) Healthy	
Accuracy for correct gender Con discrimination responses Neu (%) Ang	ndition Mean (SD) utral 89.14 (11.7 s9.14 (11.7 88.94 (15.4 gry 84.62 (15.7) (6) (6)	Mean (SD) 87.32 (11.71) 89.00 (10.32) 81.87 (16.36)	Mean (SD) 92.46 (9.18) 92.28 (14.07) 90.44 (12.47)	Mean (SD) 88.54 (13.45) 86.16 (20.91) 83.18 (16.92)	participants Mean (SD) 92.50 (18.29) 92.66 (15.84) 91.25 (19.38)	
Hal Detection (%) No Reaction time (RT) to Net correct responses (s) Ang	ppy 93.06 (9.23 face 93.03 (16.1 utral 1.05 (0.23) rful 1.04 (0.23) gry 1.03 (0.25)	0	$\begin{array}{c} 92.12\ (9.42)\ 94.37\ (10.57)\ 1.09\ (0.24)\ 1.12\ (0.27)\ 1.10\ (0.27)\ 1.10\ (0.27) \end{array}$	95.96 (4.91) 92.09 (15.47) 1.00 (0.18) 0.99 (0.20) 0.97 (0.22)	91.82 (11.33) 92.19 (21.91) 1.05 (0.26) 1.01 (0.19) 1.00 (0.22)	94.53 (18.66) 93.75 (22.25) 0.94 (0.32) 0.91 (0.33) 0.95 (0.32)	
Ha Detection RT (s) No	ppy 1.02 (0.24) face 0.79 (0.24)		1.06(0.28) 0.84(0.27)	0.97 (0.20) 0.79 (0.23)	1.01 (0.22) 0.72 (0.20)	$0.92\ (0.35)$ $0.62\ (0.17)$	
<i>Note</i> : ^a NART: National Adult Re ^b BDI-II: Beck Depression Inventi ^c PANSS: Positive and Negative Sy ^d Data missing for 1 patient and 1 ^e Reliable dose information missin,	ading Test. ³⁷ ory-II. ²⁸ ymptom Scale. ³² healthy participant. g for 4 (1 no/minim	al, 1 mild and 2	moderate-to-high depression.) patients.			

Downloaded from http://schizophreniabulletin.oxfordjournals.org/ at King's College London - Journals Dept on January 18, 2016

coffee, as requested, up until 40 minutes before fMRI. Smokers were allowed to smoke as usual to avoid a state of smoking/nicotine withdrawal but care was taken not to start fMRI scanning them within 40 minutes of smoking a cigarette.

Image Acquisition

Echoplanar MR brain images were acquired using a 1.5 T GE Signa system (General Electric). In each of 16 near-axial noncontiguous planes parallel to the intercommissural plane, 240 T2*-weighted MR images depicting blood-oxygen-level-dependent (BOLD) contrast (echo time [TE] 40 ms, repetition time [TR] 3 s, Flip 90°, field of view 240 mm, matrix 64×64 , in-plane resolution 3.1 mm, slice thickness 7.0 mm, interslice gap 0.7 mm) were acquired over the 12-minute experiment. In the same session, a high resolution 3-D inversion recovery prepared spoiled GRASS volume dataset was acquired with TE = 5.3 ms, TI = 300 ms, TR = 12.2 ms, in-plane resolution = $0.94 \,\mathrm{mm}$, slice thickness = $1.5 \,\mathrm{mm}$.

Data Analysis

All behavioral analyses were performed using Statistical Package for Social Sciences (version 22). Alpha level for testing significance of effects was maintained at P < .05unless stated otherwise.

Demographic, Behavioral and Clinical Measures. Patients and healthy participants were compared on age, education, predicted IQ⁴¹ and BDI-II scores using independent sample t tests. Patient subgroups with varying depression levels (no/minimal: BDI-II score 0–13; mild: score 14–19, moderate-to-severe: score 20 or above) were compared on age, education, predicted IQ⁴¹ and PANSS symptoms using (separate) 1-way ANOVA, with Depression (no/ minimal, mild, moderate-to-severe) as a between-subjects factor, followed by independent sample t tests where needed.

Group and emotion effects in performance accuracy (% correct gender discrimination in facial expressions) and latency (reaction time to correct responses) were analysed separately using a Group (patients, healthy participants) × Emotion type (fear, anger, neutral, happy) repeated-measures ANOVA with Group as the betweensubjects factor and Emotion type as the within-subjects factor, followed by post hoc mean comparisons. The effects of varying depression in performance accuracy and latency of patients were analyzed (separately) using Depression × Emotion type ANOVA with Depression as the between-subjects factor and Emotion type as the within-subjects factor, followed by lower order ANOVAs and post hoc mean comparisons. Finally, Pearson's correlations were used to examine relationship of BDI-II scores to performance (accuracy and latency), symptoms,

Page 4 of 12

age, illness duration and the doses of antipsychotic medication in the patient group.

fMRI

Preprocessing. For each participant, the 240 volume functional time series were realigned to the first volume, corrected for motion artefacts (x, y and z translation, pitch, roll, yaw; time series with translations >5 mm or rotations $> 5^{\circ}$ were excluded from the analysis), transformed into stereotactic space using the EPI template in SPM (affine transformation x, y and z, 16 nonlinear iterations, $7 \times 9 \times 7$ basis functions), spatially smoothed with a 10mm FWHM Gaussian filter and band pass filtered using statistical parametric mapping software (SPM5; http://www.fil.ion.ucl.ac.uk/spm).

Models and Statistical Inferences. Data were analyzed using a random effect procedure.⁴² Participant-specific activations were identified with a factorial model consisting of 4 facial expression conditions, and no face control condition serving as an implicit baseline. The boxcar for each 30-second epoch was convolved with the haemodynamic response function. Motion parameters were included as covariates at this stage. The second stage of the random effect model identified generic task-related activations in the patient and healthy groups (separately) for each facial expression condition (fear, anger, neutral, happy) vs no face control condition (height threshold P < .005; clustercorrected $P \leq .05$), using statistical parametric mapping software (SPM8; http://www.fil.ion.ucl.ac.uk/spm).

The relationship of BDI-II scores in patients with neural activity across the whole brain was first examined (height threshold P < .05; cluster-corrected $P \le .05$), using regression model within SPM8 with BDI-II scores entered as a covariate for each facial expression condition (fear, anger, neutral, happy) vs no face control condition; and also for each emotional expression (fear, anger, happy) vs the neutral expression following the observation of no relationship between BDI-II scores and activity to neutral expressions (see Results). In addition, all regression analyses were repeated with gender included as an additional covariate. Since the inclusion of gender revealed the same cluster associations, the results of these additional SPM regressions are not reported hereafter. The participant-specific activation values were then extracted from the clusters (peak voxel) that showed a relationship with BDI-II scores in the patient group, and examined (within the SPSS) for their possible relationships with symptoms, age, illness duration and the doses of antipsychotic medication (in chlorpromazine equivalents) using Pearson correlations.

Next, we extracted participant-specific mean BOLD signal for all patients and healthy participants from each of 2 activation clusters (thalamus and superior frontal gyrus [SFG]; 10mm sphere around the peak voxel) that showed a significant association with BDI-II scores during fearful expression in SPM regression analyses (table 2) using the MarsBaR toolbox (http://marsbar.sourceforge.net/projects/marsbar). These data were then analysed (within the SPSS) to confirm the influence of depression in patients' brain activity using a Region (thalamus, SFG) × Emotion type (fear, anger, happy, neutral) × Depression (no/ minimal, mild, moderate-to-severe) ANOVA, followed by lower order ANOVAs and mean comparisons as

needed. A further Region (thalamus, SFG) \times Group (healthy participants, no/minimal depression patients, mild depression patients, moderate-to-severe depression patients) ANOVA was conducted to establish whether higher activations during fearful expressions in patients with moderate-to-high depression, relative to those with no/minimal/mild depression (detected with the earlier ANOVA), reflected a hyperresponse or a stronger response within the normal range.

Table 2. Significant Associations Between Brain Activity to Fearful Expressions and Depression in Patients (Voxel Threshold P < .005Uncorrected)

Brain AreaBAVentral posterior medial nucleus (thalamus) Postcentral gyrus3Globus pallidus (lentiform nucleus) Supramarginal gyrus40	Voxels (<i>n</i>) 5422	Side Left Left	$\frac{\text{MNI}}{x}$	$\frac{\text{Coordina}}{y}$	z	Voxel T	Cluster Corrected I
Brain AreaBAVentral posterior medial nucleus (thalamus) Postcentral gyrus3Globus pallidus (lentiform nucleus) Supramarginal gyrus40	Voxels (<i>n</i>) 5422	Side Left Left	x -16	<i>y</i>	Z	Voxel T	Cluster Competed D
Ventral posterior medial nucleus (thalamus) Postcentral gyrus 3 Globus pallidus (lentiform nucleus) Supramarginal gyrus 40	5422	Left Left	-16	22		Voxel T	Cluster-Corrected P
Postcentral gyrus3Globus pallidus (lentiform nucleus)Supramarginal gyrus40		Left	= 0	-22	0	4.31	.018
Globus pallidus (lentiform nucleus) Supramarginal gyrus 40		D' 1 /	-50	-14	48	3.17	
Supramarginal gyrus 40		Right	18	-12	-4	3.10	
		Left	-32	-50	32	3.06	
Insula		Left	-36	-48	24	3.01	
Inferior frontal gyrus 9		Left	-48	6	28	2.95	
Postcentral gyrus 3		Left	-48	-20	48	2.93	
Precentral gyrus 4		Left	-32	-20	52	2.91	
Paracentral lobule 5		Left	-14	-44	62	2.89	
Postcentral gyrus 2		Left	-48	-24	52	2.86	
Supramarginal gyrus 40		Left	-36	-48	30	2.84	
Postcentral gyrus 2		Left	-48	-24	46	2.73	
Superior parietal lobule 7		Left	-34	-52	50	2.73	
Fearful > Neutral							
Superior frontal gyrus 6	9081	Right	4	10	60	4.49	<.001
6		Right	4	16	58	4.11	
Medial frontal gyrus 6		Right	14	-4	60	3.94	
6		Right	12	0	60	3.91	
Middle frontal gyrus 6		Right	42	-4	46	3.51	
Superior frontal gyrus 6		Right	8	22	56	3.50	
Precentral gyrus 6		Left	-38	-10	58	3.28	
Medial frontal gyrus 6		Left	-12	2	62	3.26	
Middle frontal gyrus 6		Left	-20	-20	62	3.24	
Precentral gyrus 4		Left	-32	-20	52	3.24	
6		Right	50	2	46	3.23	
Precentral gyrus 6		Left	-24	-22	62	3.17	
Superior frontal gyrus 8		Right	10	26	54	3.15	
Precentral gyrus 4		Left	-50	-10	46	3.08	
Anterior cingulate 24		Left	-12	0	52	3.08	
Ventral posterior medial nucleus (thalamus)	3805	Left	-18	-22	2	3.74	.009*
Insula		Left	-40	-12	10	3.30	
Hippocampus		Right	30	-22	-8	2.78	
Mammillary body (thalamus)		Right	10	-18	-2	2.72	
Thalamus		Right	10	-16	4	2.70	
Putamen (lentiform nucleus)		Right	28	-18	8	2.69	
Transverse temporal gyrus 41		Left	-52	-16	8	2.68	
Putamen (lentiform nucleus)		Right	30	-18	4	2.67	
Claustrum		Right	34	-22	2	2.64	
Ventral lateral nucleus (thalamus)		Right	20	-16	8	2.43	
Globus pallidus (lentiform nucleus)		Right	20	-12	Õ	2.35	
Caudate		Right	10	4	8	2.30	

Note: BA, Brodmann area; MNI, Montreal Neurological Institute. *Uncorrected *P*.

Results

Demographic, Behavioral and Clinical Measures

Patients and healthy participants were comparable (table 1) on age, sex distribution and education (*P* values > .18) but patients had significantly lower premorbid IQ ($t_{79} = 3.21$, P = .002). Patients also rated themselves much higher on BDI-II depression ($t_{81} = 5.46$, P < .001). Patients with varying levels of depression (table 1) did not differ in age, sex distribution, education, age at illness onset, illness duration, antipsychotic dose, and PANSS positive, negative or total symptoms (*P* values > .16) but differed marginally in general psychopathology scores ($F_{2,60} = 1.90$, P = .05). The no/minimal depression subgroups had lower general psychopathology scores (accounted for by the positive correlation, described below, between the BDI-II and general psychopathology scores).

Both healthy and patient groups showed a high level of gender discrimination accuracy across all conditions (table 1). This was expected given the easy task demands, implemented simply to ensure patients' attention to task stimuli and not place cognitive demands on them. For accuracy, there was a significant main effect of Emotion type ($F_{3,243} = 5.84$, P < .001) but there was no effect of Group ($F_{1,81} = 1.30$, P = .26) or a Group × Emotion type interaction ($F_{3,243} = 1.15$, P = .33). The main effect of Emotion type indicated a lower accuracy for angry expressions than for fearful ($t_{82} = 2.36$, P = .02), neutral ($t_{82} = 3.21$, P = .002) and happy expressions ($t_{82} = 6.32$, P < .001), and in addition, for fearful ($t_{82} = 0.89$, P = .005) and neutral ($t_{82} = 5.33$, P < .001), relative to happy expressions. There was no significant difference between fearful and neutral expressions ($t_{82} = 0.09$, P = .93). For latency, no significant effects were found (Emotion type: $F_{3,243} = 1.58$ P = .19; Group: $F_{1,81} = 2.83$, P = .10; Emotion type × Group: $F_{3,243} = 1.23$, P = .30).

Patient subgroups with varying depression levels also showed comparable accuracy across all conditions (table 1), with no significant effect of Depression $(F_{2,60} = 1.42, P = .25)$. There was a significant effect of Emotion type $(F_{3,180} = 9.14, P < .001;$ reflecting the same pattern as above) but no Depression × Emotion type interaction $(F_{6,180} = 0.56, P = .76)$. There was no effect of Depression $(F_{2,60} = 1.38, P = .26)$, or any other effect, in latency.

In patients, BDI-II (depression) scores had modestly positive correlation with PANSS general psychopathology (r = .309, P = .014; for PANSS-G6 depression item, $\rho = .353$, P < .01) but had no correlation with PANSS positive (r = .01) or negative symptoms (r = .11). BDI-II scores did not correlate with accuracy or latency during any task condition (all P > .10), age (r = .02, P = .94), illness duration (r = .006, P = .96) or doses of antipsychotic medication (r = .004, P = .98).

fMRI

Generic Task Related Activations in Patients and Healthy Participants. The areas showing significant activity changes for each facial expression condition vs no face control condition in patient and healthy groups are described in supplementary appendix 1, and displayed in figure 1 (the clusters which did not reach corrected cluster-level significance are also displayed in figure 1 for completeness).

In patients, robust activation for fearful expressions (>no face) was limited to the right inferior frontal gyrus, with weaker (uncorrected; displayed in figure 1) activity seen in the left inferior frontal gyrus (peak, x = -40, y = 8, z = 32, cluster size = 417 voxels), right striatumhippocampus-parahippocampal gyrus (x = 30, y = -6, z = -10; 401 voxels), and the left precentral gyrus (x = 36, y = -20, z = 62; 417 voxels; figure 1). Significant activations for angry expressions included the inferior frontal gyrus (bilateral), thalamus (bilateral), parahippocampal gyrus (right), globus pallidus (left) and middle occipital gyrus (right). For neutral expressions, robust activation was seen in the middle frontal gyrus (bilateral), precentral gyrus (left), thalamus (bilateral) and globus pallidus (left); in addition, there was weak (uncorrected level) activation of the left superior parietal cortex (x = 30, y = -60, z = 52; 721 voxels; figure 1). No area was activated for happy expressions at the corrected level but 2 clusters with uncorrected significance were present, 1 in the right inferior-middle frontal gyrus (x = 42, y = 10, z = 24; 447 voxels) and the other in the left precentral gyrus (x = -32, y = -22, z = 62; 461 voxels; figure 1).

In healthy participants, fearful expressions activated parts of the inferior occipital gyrus, inferior-middle frontal gyri, striatum, insula, amygdala, hippocampus, parahippocampal gyrus and the thalamus. Angry expressions mainly activated the inferior occipital gyrus, cuneus and the inferior-middle frontal gyrus. Neutral expressions activated the inferior occipital and fusiform gyri, cuneus, insula and the caudate. Happy expressions activated only the inferior-middle occipital and fusiform gyri and the cuneus (supplementary appendix 1; figure 1). In general, the magnitude of activations across all conditions in healthy participants was greater than in patients.

Depression and Brain Activity in Patients. Regression Analyses Depression (BDI-II) scores correlated significantly positively with activity during fearful expressions (>no face) in the left thalamus extending to the parapost-pre-central gyrus, putamen-globus pallidus, supramarginal gyrus, insula and inferior-middle frontal gyrus (table 2, figure 2). BDI-II scores also correlated significantly positively with activity for fearful expressions when compared with neutral (fearful > neutral) expressions



Fig. 1. Task related increases (facial expressions > no face control stimuli) in patient and healthy groups with associated Montreal Neurological Institute (MNI) x, y and z coordinates (maps thresholded at P = .005 uncorrected). For each coordinate, sagittal and axial slices are shown. The left hemisphere is shown at the top of the axial slices.

in the right SFG extending to middle/medial frontal, left precentral and left anterior cingulate gyri. Another cluster, very similar to the cluster found to correlate positively with BDI-II scores during fearful > nonface contrast, with near identical peak in the left thalamus, was also present at uncorrected significance (table 2, figure 2). BDI-II scores did not correlate significantly (positively or negatively) with activation of any area during angry, neutral or happy expressions.

Participant-specific activation values extracted from peak voxel of the above 3 clusters that associated positively with BDI-II scores (peak of cluster 1 in the thalamus, fearful > no face control contrast; of cluster 2 in the superior frontal gyrus, and of cluster 3 in the thalamus, both fearful > neutral face contrast) did not correlate significantly with PANSS total or subscale scores (all *r* values <.17). Participant-specific activation values associated positively with BDI-II scores also did not correlate with age (*r* values -.009 to .045; all P > .72), duration of illness (*r* values -.015 to .03; all P > .80) or antipsychotic medication dose (*r* values -.025; all P > .76), and their correlation with BDI-II scores remained significant with near identical *P* values (all $P \le .001$) after controlling (partial *r*) for age, illness duration or doses of antipsychotic medication.

Categorical Analyses The Region × Emotion type × Depression (2×4 × 3) ANOVA to examine differences between patient subgroups revealed a significant Region × Emotion type effect ($F_{3,180} = 2.82$, P = .04) and a trend for Emotion type × Depression effect ($F_{6,180} = 1.76$, P = .11) with a significant linear trend ($F_{2,603} = 3.81$, P = .026). Follow-up analyses of these effects using Region × Depression ANOVAs separately for fearful, angry, happy and neutral expressions revealed a significant Depression effect for fearful expressions ($F_{2,60} = 4.86$, P = .01). This effect was due to significantly higher activity across the thalamic and SFG clusters in moderate-to-severe depression subgroup, relative to no/minimal ($F_{1,44} = 8.97$, P = .004) and mild depression ($F_{1,36} = 5.32$, P = .03) subgroups; no/minimal and mild depression subgroups did not differ from each other ($F_{1,40} = 0.04$, P = .85; figure 3). Region × Depression ANOVAs (separately) for angry,



Fearful > No face

Fig. 2. Brain activity positively associated with Beck Depression Inventory (BDI-II) depression with associated Montreal Neurological Institute (MNI) *x*, *y* and *z* coordinates (maps thresholded at P = .05 uncorrected; displayed clusters significant at corrected $P \le .05$). For each coordinate, sagittal and axial slices are shown. The left hemisphere is shown at the top of the axial slices.

happy and neutral expressions did not reveal any significant effects (figure 3).

Further Region × Group (no/minimal depression, mild depression and moderate-to-severe depression patients, and healthy participants) ANOVA to specifically probe activity differences for fearful expressions in moderate-to-severe depression patients, relative to healthy participants, also revealed a significant Group effect ($F_{3,79} = 3.88$, P = .01). This effect was due to significantly higher activity in the moderate-to-severe depression patient subgroup, relative to the healthy group ($F_{1,39} = 5.89$, P = .02) and no/minimal and mild depression patient subgroups (effects already described). Patients with no/minimal

 $(F_{1,43} = 0.36, P = .55)$ or mild depression $(F_{1,35} = 0.34, P = .56)$ did not differ from healthy participants (or each other, as described earlier).

Discussion

Our primary aim was to determine the brain activity associated with depression, as indexed with BDI-II, in patients with chronic schizophrenia during implicit processing of facial affect. The findings partially confirmed our a priori hypothesis in that there was a significantly positive association between depression (BDI-II) scores and brain activity to facial expressions of fear. This association was



Fig. 3. Mean functional magnetic resonance imaging (fMRI) response in the thalamic and superior frontal gyrus clusters (correlated with Beck Depression Inventory [BDI-II] scores during fearful, but not other, expressions) in patients with varying depression levels and healthy participants (*indicates P < .05; higher than all other groups).

robustly present (fearful > no face control) in a cluster that peaked in the left thalamus and extended to include parts of the insula, globus pallidus, and inferior-middle frontal and pre-post-para-precentral gyri. Although the thalamic cluster was present at the uncorrected level in fearful > neutral expressions contrast, it was in the same location as detected with fear > no face control contrast, and extended to include parts of the caudate, putamen, insula and hippocampus. In addition, during fearful > neutral contrast, bilateral activity in the superior-middlemedial frontal gyrus, extending to the precentral gyrus and anterior cingulate, was associated significantly positively with BDI-II scores. Importantly, patients with moderate-to-severe depression had significantly greater brain activity in these regions during fearful expressions, relative to patients with no/minimal/mild depression and healthy participants.

In general, our fMRI observations fit with previous findings of enhanced brain activity seen in MDD patients in response to facial expressions of fear.^{15,43} However, unlike previous findings in MDD which often show a hyperactive amygdala response to fearful expressions,^{16–18} we did not find amygdala activation to associate with

depression in psychosis patients, more likely because of their long-term antipsychotic use⁴⁴⁻⁴⁶ or other diagnosisrelated issues,^{24,43} rather than amygdala response habituation with repeated exposure to facial stimuli during the practice and fMRI sessions,⁴⁷ since there was amygdala activation to fearful expressions in healthy participants (figure 1) with identical task and scanning parameters. Patients with moderate-to-severe depression in this study also showed the strongest activity in the regions that showed "fear expressions-depression relationship" but were not significantly different from patients with no/ minimum/mild depression or healthy participants during angry expressions. Importantly, Gur and colleagues²⁵ also reported enhanced brain response to fearful, but not angry, expressions to be significantly associated with a high severity of flat affect in schizophrenia (n = 16) and this association was present in the thalamus and hippocampus in addition to the amygdala. It is also relevant in this context that we³² previously found a more robust reduction, following cognitive behavior therapy for psychosis (CBTp), in activity of the inferior frontal, insula, thalamus and putamen areas during fearful, than during angry, expressions; and improvements in subjective depression are a consistent and important outcome of CBTp.^{48,49} Taken together, these finding indicate a consistent relationship between depression and altered neural response to fearful expressions, a signal of uncertain threat,⁵⁰ in schizophrenia. Interestingly, there is recent meta-analytic evidence that schizophrenia patients (as a group) show under-activation in the entire facial emotion processing network whereas those with bipolar disorder (BD) show greater thalamic engagement during facial emotion processing.²⁴ The present findings regarding depression in the context of psychosis, therefore, seem related to emotion processing abnormalities in BD, and may point towards the use of therapies known to be effective for treating depressed mood in BD for treating depression in schizophrenia.

Of the regions that showed a positive association with depression during the viewing of fearful facial expressions, the pulvinar thalamus, with projections to the amygdala,⁵¹ has been strongly implicated in the control of visual attention,⁵²⁻⁵⁴ specifically during rapid detection of threatening stimuli.^{55,56} A negative correlation between avoidance response to threatening stimuli and activation of the pulvinar thalamus, insula and striatum has also been previously demonstrated in healthy humans.⁵⁷ It is thus possible that patients with mild/severe depression attended fearful expressions to a greater extent and possibly engaged in explicit processing of threat-related information.⁵⁸

The strengths of this study include the use of a large sample to examine depression-facial affect processing relationship, ie, sufficient power to investigate the questions of interest. The main limitation of this study is that it is cross-sectional in nature and therefore there is no direct evidence of cause and effect. Further limitations include a lack of reliable information on the proportion of patients who met DSM criteria for current/past MDD and possible confounding effects of different types and dosages of medications. It would be valuable in future to conduct longitudinal investigations while monitoring depression and medications, and to directly compare patients with schizophrenia and varying levels of depression to MDD and BD patients with varying levels of psychotic symptoms in their responses to facial affect, relative to an appropriate baseline (eg, scrambled face)⁵⁹ within the same study.

In conclusion, this study provides the first evidence that psychosis patients with a high level of subjective depression show enhanced brain response to fearful facial expressions, particularly in the thalamus. Our findings suggest an overlap between schizophrenia patients with moderate-to-severe depression and BD patients, the latter of which reliably show greater thalamic involvement during facial affect processing. Unlike previous findings in MDD,¹⁶⁻¹⁸ we found no evidence of amygdala activation during fearful expression to associate with depression in the context of psychosis.

Supplementary Material

Supplementary material is available at http://schizophreniabulletin.oxfordjournals.org.

Funding

This research was supported by the Wellcome Trust (067427/z/02/z). V.K. is part funded by the Biomedical Research Centre for Mental Health at the Institute of Psychiatry, King's College London, and the South London and Maudsley NHS Foundation Trust, UK.

Acknowledgment

The authors report no biomedical financial interests or potential conflicts of interest.

References

- 1. Bleuler E. *Dementia Praecox or the Group of Schizophrenias*. New York, NY: International Universities Press Inc; 1911.
- 2. Zisook S, McAdams LA, Kuck J, et al. Depressive symptoms in schizophrenia. *Am J Psychiatry*. 1999;156:1736–1743.
- 3. Zisook S, Nyer M, Kasckow J, Golshan S, Lehman D, Montross L. Depressive symptom patterns in patients with chronic schizophrenia and subsyndromal depression. *Schizophr Res.* 2006;86:226–233.
- Rosen JL, Miller TJ, D'Andrea JT, McGlashan TH, Woods SW. Comorbid diagnoses in patients meeting criteria for the schizophrenia prodrome. *Schizophr Res.* 2006;85:124–131.
- Krabbendam L, Myin-Germeys I, Hanssen M, et al. Development of depressed mood predicts onset of psychotic disorder in individuals who report hallucinatory experiences. *Br J Clin Psychol.* 2005;44:113–125.
- Buckley PF, Miller BJ, Lehrer DS, Castle DJ. Psychiatric comorbidities and schizophrenia. *Schizophr Bull.* 2009;35:383–402.
- 7. Birchwood M, Iqbal Z, Upthegrove R. Psychological pathways to depression in schizophrenia: studies in acute psychosis, post psychotic depression and auditory hallucinations. *Eur Arch Psychiatry Clin Neurosci.* 2005;255:202–212.
- 8. Rocca P, Bellino S, Calvarese P, et al. Depressive and negative symptoms in schizophrenia: different effects on clinical features. *Compr Psychiatry*. 2005;46:304–310.
- 9. Blanchard JJ, Cohen AS. The structure of negative symptoms within schizophrenia: implications for assessment. *Schizophr Bull.* 2006;32:238–245.
- Addington DE, Addington JM. Attempted suicide and depression in schizophrenia. Acta Psychiatr Scand. 1992;85:288–291.
- Jones JS, Stein DJ, Stanley B, Guido JR, Winchel R, Stanley M. Negative and depressive symptoms in suicidal schizophrenics. *Acta Psychiatr Scand*. 1994;89:81–87.
- 12. Häfner H, Löffler W, Maurer K, Hambrecht M, an der Heiden W. Depression, negative symptoms, social stagnation and social decline in the early course of schizophrenia. *Acta Psychiatr Scand.* 1999;100:105–118.
- 13. Kelly DL, Shim JC, Feldman SM, Yu Y, Conley RR. Lifetime psychiatric symptoms in persons with schizophrenia who died

by suicide compared to other means of death. J Psychiatr Res. 2004;38:531-536.

- Fialko L, Freeman D, Bebbington PE, et al. Understanding suicidal ideation in psychosis: findings from the Psychological Prevention of Relapse in Psychosis (PRP) trial. *Acta Psychiat Scand.* 2006;114:177–186.
- 15. Stuhrmann A, Suslow T, Dannlowski U. Facial emotion processing in major depression: a systematic review of neuroimaging findings. *Biol Mood Anxiety Disord*. 2011;1:10.
- Sheline YI, Barch DM, Donnelly JM, Ollinger JM, Snyder AZ, Mintun MA. Increased amygdala response to masked emotional faces in depressed subjects resolves with antidepressant treatment: an fMRI study. *Biol Psychiatry*. 2001;50:651–658.
- Peluso MA, Glahn DC, Matsuo K, et al. Amygdala hyperactivation in untreated depressed individuals. *Psychiat Res.* 2009;173:158–161.
- 18. Zhong M, Wang X, Xiao J, et al. Amygdala hyperactivation and prefrontal hypoactivation in subjects with cognitive vulnerability to depression. *Biol Psychol*. 2011;88:233–242.
- Suslow T, Konrad C, Kugel H, et al. Automatic mood-congruent amygdala responses to masked facial expressions in major depression. *Biol Psychiatry*. 2010;67:155–160.
- Davidson RJ, Pizzagalli D, Nitschke JB, Putnam K. Depression: perspectives from affective neuroscience. *Annu Rev Psychol.* 2002;53:545–574.
- Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception II: Implications for major psychiatric disorders. *Biol Psychiatry*. 2003;54:515–528.
- 22. Jaworska N, Yang XR, Knott V, Macqueen G. A review of fMRI studies during visual emotive processing in major depressive disorder *World J Biol Psychiatry*. 2015;16:448–471.
- Li H, Chan RC, McAlonan GM, Gong QY. Facial emotion processing in schizophrenia: a meta-analysis of functional neuroimaging data. *Schizophr Bull*. 2010;36:1029–1039.
- 24. Delvecchio G, Sugranyes G, Frangou S. Evidence of diagnostic specificity in the neural correlates of facial affect processing in bipolar disorder and schizophrenia: a metaanalysis of functional imaging studies. *Psychol Med.* 2013;43:553–569.
- 25. Gur RE, Loughead J, Kohler CG, et al. Limbic activation associated with misidentification of fearful faces and flat affect in schizophrenia. *Arch Gen Psychiatry*. 2007;64:1356–1366.
- Kohler C, Swanson CL, Gur RC, Mozley LH, Gur RE. Depression in schizophrenia: II. MRI and PET findings. *Biol Psychiatry*. 1998;43:173–180.
- 27. Kohler C, Gur RC, Swanson CL, Petty R, Gur RE. Depression in schizophrenia: I. Association with neuropsychological deficits. *Biol Psychiatry*. 1998;43:165–172.
- Addington D, Addington J, Patten S. Depression in people with first-episode schizophrenia. *Brit J Psychiatry*. 1998;172:90–92.
- Heald A, Morris J, Soni SD. Characterisation of depression in patients with schizophrenia. *Indian J Med Res.* 2008;127:544–550.
- Halari R, Mehrotra R, Sharma T, Kumari V. Does self-perceived mood predict more variance in cognitive performance than clinician-rated symptoms in schizophrenia? *Schizophr Bull*. 2006;32:751–757.
- 31. Beck A, Steer R, Brown G. *Manual for the BDI-II*. San Antonio, TX: Psychological Corporation; 1996.

- 32. Kumari V, Fannon D, Peters ER, et al. Neural changes following cognitive behaviour therapy for psychosis: a longitudinal study. *Brain*. 2011;134:2396–2407.
- 33. Oldfield RC. The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia*. 1971;9:97–113.
- 34. First MB, Spitzer RL, Gibbon M, Williams JBW. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Patient Edition. (SCID-I/P). New York, NY: New York State Psychiatric Institute, Biometrics Research; 2002
- 35. First MB, Spitzer RL, Gibbon M, Williams JBW. Structured Clinical Interview for DSM-IV-TR Axis I Disorders, Research Version, Non-patient Edition. (SCID-I/NP). New York, NY: New York State Psychiatric Institute, Biometrics Research; 2002.
- Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull*. 1987;13:261–276.
- 37. Ekman P, Friesen W. *Pictures of Facial Affect*. Palo Alto, CA: Consulting Psychologists Press; 1976.
- Horley K, Gonsalvez C, Williams L, Lazzaro I, Bahramali H, Gordon E. Event-related potentials to threat-related faces in schizophrenia. *Int J Neurosci*. 2001;107:113–130.
- Hall J, Whalley HC, McKirdy JW, et al. Overactivation of fear systems to neutral faces in schizophrenia. *Biol Psychiatry*. 2008;64:70–73.
- Seiferth NY, Pauly K, Kellermann T, et al. Neuronal correlates of facial emotion discrimination in early onset schizophrenia. *Neuropsychopharmacology*. 2009;34:477–487.
- 41. Nelson H, Willison J. NART Test Manual (Part II). New York, NY: NFER-Nelson; 1991.
- 42. Friston KJ, Holmes AP, Worsley KJ. How many subjects constitute a study? *Neuroimage*. 1999;10:1–5.
- Savitz J, Drevets WC. Bipolar and major depressive disorder: neuroimaging the developmental-degenerative divide. *Neurosci Biobehav Rev.* 2009;33:699–771.
- 44. Blasi G, Popolizio T, Taurisano P, et al. Changes in prefrontal and amygdala activity during olanzapine treatment in schizo-phrenia. *Psychiat Res.* 2009;173:31–38.
- 45. Bergé D, Carmona S, Salgado P, Rovira M, Bulbena A, Vilarroya O. Limbic activity in antipsychotic naïve first-episode psychotic subjects during facial emotion discrimination. *Eur Arch Psychiatry Clin Neurosci.* 2014;264:271–283.
- 46. Surguladze SA, Chu EM, Marshall N, et al. Emotion processing in schizophrenia: fMRI study of patients treated with risperidone long-acting injections or conventional depot medication. *J Psychopharmacol* 2011;25:722–733.
- 47. Plichta MM, Grimm O, Morgen K, et al. Amygdala habituation: a reliable fMRI phenotype. *Neuroimage*. 2014;103:383–390.
- 48. Garety PA, Fowler DG, Freeman D, Bebbington P, Dunn G, Kuipers E. Cognitive-behavioural therapy and family intervention for relapse prevention and symptom reduction in psychosis: randomised controlled trial. *Br J Psychiatry*. 2008;192:412–423.
- 49. Peters E, Landau S, McCrone P, et al. A randomised controlled trial of cognitive behaviour therapy for psychosis in a routine clinical service. *Acta Psychiatr Scand*. 2010;122:302–318.
- 50. Fridlund AJ. *Human Facial Expression: An Evolutionary View*. New York, NY: Academic Press; 1994.
- 51. Jones EG, Burton H. A projection from the medial pulvinar to the amygdala in primates. *Brain Res.* 1976;104:142–147.
- 52. Adler CM, Sax KW, Holland SK, Schmithorst V, Rosenberg L, Strakowski SM. Changes in neuronal activation with

increasing attention demand in healthy volunteers: an fMRI study. *Synapse*. 2001;42:266–272.

- Komura Y, Nikkuni A, Hirashima N, Uetake T, Miyamoto A. Responses of pulvinar neurons reflect a subject's confidence in visual categorization. *Nat Neurosci.* 2013;16:749–755.
- 54. Purushothaman G, Marion R, Li K, Casagrande VA. Gating and control of primary visual cortex by pulvinar. *Nat Neurosci.* 2012;15:905–912.
- 55. Van Le Q, Isbell LA, Matsumoto J, et al. Pulvinar neurons reveal neurobiological evidence of past selection for rapid detection of snakes. *Proc Natl Acad Sci U S A*. 2013;110:19000–19005.
- 56. Ward R, Danziger S, Bamford S. Response to visual threat following damage to the pulvinar. *Curr Biol.* 2005;15:571–573.
- 57. Schlund MW, Siegle GJ, Ladouceur CD, et al. Nothing to fear? Neural systems supporting avoidance behavior in healthy youths. *Neuroimage*. 2010;52:710–719.
- Scheuerecker J, Frodl T, Koutsouleris N, et al. Cerebral differences in explicit and implicit emotional processing–an fMRI study. *Neuropsychobiology*. 2007;56:32–39.
- 59. Underwood R, Peters E, Kumari V. Psychobiology of threat appraisal in the context of psychotic experiences: a selective review. *Eur Psychiatry*. 2015;30:817–829.