



King's Research Portal

DOI: 10.1016/j.jpsychores.2016.11.001

Document Version Peer reviewed version

Link to publication record in King's Research Portal

Citation for published version (APA):

Himmerich, H., Wesemann, U., Dalton, B., Holdt, L. M., Teupser, D., & Willmund, G. D. (2016). Letter to the editor: Exploring an association between hostility and serum concentrations of TNF-α and its soluble receptors. *Journal of Psychosomatic Research*, *91*, 87-88. https://doi.org/10.1016/j.jpsychores.2016.11.001

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Exploring an association between hostility and serum concentrations of TNF- α and its soluble receptors

Hubertus Himmerich, Ulrich Wesemann, Bethan Dalton, Lesca M. Holdt, Daniel Teupser, Gerd D. Willmund

PII:	S0022-3999(16)30466-4
DOI:	doi:10.1016/j.jpsychores.2016.11.001
Reference:	PSR 9235
To appear in:	Journal of Psychosomatic Research
Received date:	21 October 2016
Accepted date:	3 November 2016



Please cite this article as: Himmerich Hubertus, Wesemann Ulrich, Dalton Bethan, Holdt Lesca M., Teupser Daniel, Willmund Gerd D., Exploring an association between hostility and serum concentrations of TNF- α and its soluble receptors, *Journal of Psychosomatic Research* (2016), doi:10.1016/j.jpsychores.2016.11.001

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Exploring an association between hostility and serum concentrations of TNF-α and its soluble receptors

(Short title: Hostility, TNF-α and sTNF-R levels)

Hubertus Himmerich^{1,2#}, Ulrich Wesemann², Bethan Dalton¹, Lesca M. Holdt³, Daniel Teupser³, Gerd D. Willmund²

- ¹ Department of Psychological Medicine, King's College London, London, UK
- ² Department of Psychiatry, Psychotherapy and Psychotraumatology, Bundeswehr Hospital, Berlin, Germany
- ³ Institute of Laboratory Medicine, Ludwig-Maximilians-University Munich, Munich, Germany

Letter to the Editors

Manuscript submitted to:

Journal of Psychosomatic Research

Corresponding author: Dr. Hubertus Himmerich, Department of Psychological Medicine, King's College London, 103 Denmark Hill, London SE5 8AF, UK, Fax: +44-2078480182, Email: hubertus.himmerich@kcl.ac.uk

Key words:

Cytokines, tumor necrosis factor (TNF)-alpha, soluble TNF receptors, hostility

Dear Editors,

With great interest and pleasure, we have read the article "The relation between hostility and concurrent levels of inflammation is sex, age, and measure dependent" by Boisclair Demarble et al. [1] in the Journal of Psychosomatic Research which reports an association between hostility and TNF- α serum levels in women, but not men. It also gives a comprehensive overview of the somewhat contradictory results regarding cytokine levels and hostility reported in the literature. As several of the cited studies measured TNF- α levels [e.g. 2, 3], we thought that it would be of value to investigate hostility and levels of TNF- α , as well as its soluble receptors p55 (sTNF-R p55) and p75 (sTNF-R p75), in a cross-sectional study examining 135 male German soldiers [4].

In this study [4], the participating soldiers were assessed using the Brief Symptom Inventory (BSI) [5], which is an instrument that evaluates psychological distress and symptoms of psychiatric disorders. It is a 53-item self-report scale that takes approximately 10 minutes to complete. It contains subscales that address the following areas: somatization, obsessive-compulsive, interpersonal sensitivity, depression, anxiety, hostility, phobia, paranoia and psychoticism.

Blood was also drawn from study participants. Blood probes were immediately centrifuged at 3000 rpm for 10 min. The supernatant was aliquoted and stored in non-absorbing polypropylene tubes of 300 μ l. Probes were shock-frozen in liquid nitrogen and stored in freezers at -80 °C until further measurement. TNF- α and its soluble receptors sTNF-R p55 and p75 were measured using a Bio-Plex ProTM human cytokine immunoassay from Bio Rad, Germany.

In the Pearson correlation analysis, serum levels of TNF- α and its soluble receptor sTNF-R p75 were not significantly correlated with any of the BSI subscales. This is consistent with the finding that there is no association between hostility and TNF- α serum levels in men, as reported by Boisclair Demarble et al. [1]. However, sTNF-R p55 levels correlated significantly (p<0.05) with all BSI subscales: somatization (r=0.203, p=0.021, N=129), obsessive-compulsive (r=0.233, p=0.008, N=129), interpersonal sensitivity (r=0.263, p=0.003, N=129), depression (r=0.187, p=0.033, N=129), anxiety (r=0.230, p=0.009, N=129), hostility (r=0.249, p=0.004, N=129), phobia (r=0.185, p=0.036, N=129), paranoia (r=0.285, p=0.001, N=129), and psychoticism (r=0.206, p=0.019, N=129). We also reported [4] that body mass index (BMI) had a significant effect (F[1, 100]=7.36; p=0.008) on sTNF-R p55

levels, but also on TNF- α and sTNF-R p75 levels in this sample. In a regression analysis, hostility contributed significantly to sTNF-R p55 levels (F(1, 127)=8.41; p=0.004; corrected R²=0.055) even when controlling for BMI.

Our results support the view of Boisclair Demarble et al. [1] that hostility and TNF- α serum levels are not related in men. In addition to the importance of sex and age highlighted by Boisclair Demarble et al. [1], we would like to emphasize the influence of BMI on the TNF- α system. Based on our data, with regards to TNF- α signaling, we recommend that TNF- α receptors should also be taken into account. To our knowledge, this is the first report on correlations between sTNF-R p55 levels and hostility.

As sTNF-R p55 levels are related to several BSI subscales, one may consider psychological associations between these subscales. For example, hostility could be a consequence of paranoid thoughts. The answers to these questions, however, are beyond the scope of our data. Elevated TNF-R p55 levels have also been identified in other conditions like depression [6, 7], therefore, suggesting they may be a general indicator of psychological problems, rather than a specific marker for hostility.

Conflict of interest

The authors state that they do not have any conflicts of interest with regards to this article.

Acknowledgement

The authors thank Peter Zimmermann, Jörg-Egbert Wolf, Antje H. Bühler and Wolfgang Wilfert for their help. This work was financially supported by the Claussen-Simon-Foundation.

References

- J. Boisclair Demarble, D.S. Moskowitz, J.C. Tardif, B. D'Antono, The relation between hostility and concurrent levels of inflammation is sex, age, and measure dependent, J. Psychosom. Res. 76 (2014) 384-393.
- 2) E.C. Suarez, J.G. Lewis, C. Kuhn, The relation of aggression, hostility, and anger to

lipopolysaccharide-stimulated tumor necrosis factor (TNF)-alpha by blood monocytes from normal men, Brain Behav. Immun. 16 (2002) 675–684.

- 3) D. Janicki-Deverts, S. Cohen, W.J. Doyle, Cynical hostility and stimulated Th1 and Th2 cytokine production, Brain Behav. Immun. 24 (2010) 58–63.
- H. Himmerich, G.D. Willmund, P. Zimmermann, J.E. Wolf, A.H. Bühler, L.M. Holdt, D. Teupser, K.C. Kirkby, U. Wesemann, Serum concentrations of TNF-α, sTNF-R p55 and p75 and post-traumatic stress in German soldiers, Eur. Cytokine Netw. 26 (2015) 57-60.
- L.R. Derogatis, N. Melisaratos, The Brief Symptom Inventory: an introductory report, Psychol. Med. 13 (1983) 595-605.
- H. Himmerich, S. Fulda, J. Linseisen, H. Seiler, G. Wolfram, S. Himmerich, K. Gedrich, S. Kloiber, S. Lucae, M. Ising, M. Uhr, F. Holsboer, T. Pollmächer, Depression, comorbidities and the TNF-alpha system, Eur. Psychiatry 23 (2008) 421-429.
- R. Grassi-Oliveira, E. Brietzke, J.C. Pezzi, R.P. Lopes, A.L. Teixeira, M.E. Bauer, Increased soluble tumor necrosis factor-alpha receptors in patients with major depressive disorder, Psychiatry Clin. Neurosci. 63 (2009) 202-208.