



King's Research Portal

DOI:

[10.1136/bmjresp-2016-000156](https://doi.org/10.1136/bmjresp-2016-000156)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Qureshi, N. R., Rintoul, R. C., Miles, K. A., George, S., Harris, S., Madden, J., Cozens, K., Little, L. A., Eichhorst, K., Jones, J., Moate, P., McClement, C., Pike, L. C., Sinclair, D., Wong, W. L., Shekhdar, J., Eaton, R., Shah, A., Brindle, L., ... Gilbert, F. J. (2016). Accuracy and cost-effectiveness of dynamic contrast-enhanced CT in the characterisation of solitary pulmonary nodules — The SPUtNIK study. *BMJ Open Respiratory Research*, 3(1), Article e000156. <https://doi.org/10.1136/bmjresp-2016-000156>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accuracy and cost-effectiveness of dynamic contrast-enhanced CT in the characterisation of solitary pulmonary nodules – the SPUtNik study

N R Qureshi,¹ R C Rintoul,² K A Miles,³ S George,⁴ S Harris,⁴ J Madden,⁵ K Cozens,⁵ L A Little,⁵ K Eichhorst,⁵ J Jones,⁶ P Moate,⁵ C McClement,⁵ L Pike,⁷ D Sinclair,⁷ W L Wong,⁸ J Shekhdar,⁹ R Eaton,⁹ A Shah,⁹ L Brindle,¹⁰ C Peebles,¹¹ A Banerjee,¹¹ S Dizdarevic,¹² S Han,¹³ F W Poon,¹³ A M Groves,³ L Kurban,¹⁴ A J Frew,¹² M E Callister,¹⁵ P Crosbie,¹⁶ F V Gleeson,¹⁷ K Karunasaagar,¹⁸ O Kankam,¹⁹ F J Gilbert,²⁰ on behalf of the SPUTNIK investigators

To cite: Qureshi NR, Rintoul RC, Miles KA, *et al*. Accuracy and cost-effectiveness of dynamic contrast-enhanced CT in the characterisation of solitary pulmonary nodules—the SPUtNik study. *BMJ Open Resp Res* 2016;**3**:e000156. doi:10.1136/bmjresp-2016-000156

► Additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjresp-2016-000156>)

Received 11 August 2016
Accepted 17 August 2016

ABSTRACT

Introduction: Solitary pulmonary nodules (SPNs) are common on CT. The most cost-effective investigation algorithm is still to be determined. Dynamic contrast-enhanced CT (DCE-CT) is an established diagnostic test not widely available in the UK currently.

Methods and analysis: The SPUtNik study will assess the diagnostic accuracy, clinical utility and cost-effectiveness of DCE-CT, alongside the current CT and 18-fluorodeoxyglucose-positron emission tomography (¹⁸FDG-PET)-CT nodule characterisation strategies in the National Health Service (NHS). Image acquisition and data analysis for ¹⁸FDG-PET-CT and DCE-CT will follow a standardised protocol with central review of 10% to ensure quality assurance. Decision analytic modelling will assess the likely costs and health outcomes resulting from incorporation of DCE-CT into management strategies for patients with SPNs.

Ethics and dissemination: Approval has been granted by the South West Research Ethics Committee. Ethics reference number 12/SW/0206. The results of the trial will be presented at national and international meetings and published in an Health Technology Assessment (HTA) Monograph and in peer-reviewed journals.

Trial registration number: ISRCTN30784948; Pre-results.

BACKGROUND

Solitary pulmonary nodule (SPN) is defined as a discrete well-defined intraparenchymal lesion <3 cm. SPNs are present in 20–50% of individuals considered to be at high risk for lung cancer and present an important diagnostic problem. A small proportion of patients with a SPN will have early stage lung cancer with a high 5-year survival rate

following surgical resection. In the National Lung Screening Trial (NLST) an SPN was identified in 24% of the 53 454 asymptomatic current or former smokers with a 30 pack year history and of these >95% were benign.

CT is the first-line investigation for SPN. Fleischner Society guidelines have been widely used for many years although recently published British Thoracic Society (BTS) guidelines using the prediction scores and volumetry are now being more widely used in the UK. In general, nodules of 8 mm or less in diameter are assessed for interval growth using serial CT surveillance. For nodules >8 mm a number of imaging options are available determined by the patient's performance status and the probability for malignancy. For patients that are potentially suitable for treatment with curative intent, positron emission tomography-CT (PET-CT) is the investigation of choice. PET-CT provides both anatomical (CT) and functional information (PET) following intravenous administration of small quantities of the radioactive glucose analogue 18-fluorodeoxyglucose (¹⁸FDG). ¹⁸FDG-PET characterises SPNs on the basis of increased glucose metabolism in malignant lesions. However, false positives can occur in certain benign conditions including granulomatous disease and infective/inflammatory lesions. False-negative scans are seen in tumours with low glucose metabolism including lung adenocarcinoma with a lepidic or mucinous component and carcinoid tumours with no atypia. In order to minimise patient anxiety, risk of overdiagnosis, cumulative radiation burden and cost of performing multiple tests



CrossMark

For numbered affiliations see end of article.

Correspondence to

Professor FJ Gilbert;
fjg28@medschl.cam.ac.uk



a streamlined diagnostic pathway that involves the least number of investigations for accurate nodule characterisation is essential. In Paul Barnett's study looking at the management of patients with SPNs for the Veterans Affairs Positron Emission Tomography Imaging Cooperative study group, the average US Medicare expenditure for clinical management of an incidental SPN was \$50 233 (£30 363) when the nodule was malignant and \$22 461 (£13 577) when benign.

Dynamic contrast-enhanced CT (DCE-CT) is an imaging technique that involves the acquisition of a series of CT images through the nodule before and at fixed time points after the injection of an iodinated contrast medium. A region of interest is placed within the nodule and the mean enhancement value (Hounsfield Units) is calculated at each time point. DCE-CT characterises SPNs on the basis of increased enhancement in malignant nodules reflecting the presence of tumour neovascularisation. Malignant nodules typically demonstrate increased contrast enhancement >15 HU compared with benign nodules.

Pooled analysis from 10 DCE-CT studies (incorporating 1167 nodules) reported a sensitivity, specificity, PPV and NPV of 93%, 76%, 80% and 95%, respectively, for differentiating malignant from benign nodules.¹ A single study that compared the relative cost-effectiveness of DCE-CT to conventional CT surveillance and PET-based strategies for nodule evaluation, demonstrated savings of up to £2000 per patient compared with routine nodule surveillance. Furthermore, a strategy whereby patients only underwent ¹⁸FDG-PET if DCE-CT suggested malignancy had similar effectiveness to ¹⁸FDG-PET alone but was consistently less expensive.²

To date, only three studies have directly compared the diagnostic performances of ¹⁸FDG-PET and DCE-CT in the same cohort of patients. However, pooled data from these three studies evaluating 217 SPNs demonstrated that ¹⁸FDG-PET and DCE-CT had sensitivities of 92% and 87% and specificities of 90% and 83%, respectively.³⁻⁵ As yet, no large comparative multicentre trial of DCE-CT as a standalone technique or in combination with ¹⁸FDG-PET or integrated PET-CT for nodule characterisation has been performed.

STUDY DESIGN

The SPUtNik trial is a multicentre prospective cohort observational study designed to (1) assess the diagnostic performances of DCE-CT and ¹⁸FDG-PET-CT for the characterisation of SPNs in the National Health Service (NHS) setting and (2) use decision analytic modelling to assess the likely costs and health outcomes resulting from incorporation of DCE-CT into management strategies for patients with SPNs. Secondary objectives include assessment of the incremental value of incorporating the CT appearances of a SPN into the interpretation of integrated PET-CT examinations and assessing whether combining DCE-CT with ¹⁸FDG-PET-CT is more

accurate and/or cost-effective for characterising SPNs than either test used alone or in series. The study consists of a cohort of 375 patients and the full protocol is available as an online supplementary material.

The primary inclusion criterion is the presence of a dominant indeterminate soft tissue SPN identified on CT that measures ≥ 8 and ≤ 30 mm on axial plane (with no ancillary evidence strongly indicative of malignancy) which is being considered for further evaluation with ¹⁸FDG-PET-CT. Potential participants will be identified either at local multidisciplinary meetings or at time of referral for investigation of a SPN. The study involves a single DCE-CT (radiation dose 25 mSv) being performed in addition to the patient's standard SPN management. The DCE-CT is performed within 21 days of the ¹⁸FDG-PET-CT examination. Exclusion criteria include history of malignancy within 2 years, confirmed aetiology of the SPN at the time of the qualifying CT, biopsy of the nodule before the DCE-CT, contraindication to potential surgical resection or radiotherapy and contraindication to any of the imaging investigations.

Site accreditation and quality assessment for ¹⁸FDG-PET-CT and DCE-CT will be performed using established procedures by the PET core laboratory at St Thomas' Hospital, London and Mount Vernon Hospital, London, respectively. The ¹⁸FDG-PET-CT and DCE-CT image acquisition and data analysis will follow a standardised protocol. The ¹⁸FDG-PET-CT images and attenuation correction CT images will initially be classified according to a five-point characterisation scale. Further quantitative analysis will consist of measurements of FDG uptake expressed as the maximum standardised uptake value. The diagnostic performance of ¹⁸FDG-PET will be assessed with and without incorporation of the CT appearances. The presence of incidental extrathoracic findings on PET-CT will also be recorded. DCE-CT interpretation will be performed by thoracic radiologists at each participating site. For both DCE-CT and PET-CT, central review of 10% of cases will be performed by an expert radiologist/nuclear medicine physician to ensure quality assurance.

Following PET-CT and DCE-CT examinations subsequent SPN management is determined by the local specialist lung multidisciplinary meeting. The reference standard will comprise pathological and/or imaging follow-up data at 24 months. All patients without definitive pathological findings will undergo repeat CT examinations of the chest at 3, 9 and 24 months (with or without biopsy) in accordance with Fleischner guidelines. Clinical information, including information relating to costs will be extracted using a standardised data collection form to inform the economic analysis. A subanalysis using the BTS guidelines algorithm will also be undertaken.

In parallel with the main study, a quantitative substudy (IPCARD-SPN: validated in a population of general practitioner referred chest radiograph attendees) will aim to (1) identify the positive and negative predictive values of

symptoms that distinguish between malignant and non-malignant SPN and (2) ascertain whether or not the inclusion of symptoms found to distinguish between malignant and non-malignant nodules increases the diagnostic value of DCE-CT and ^{18}F FDG-PET-CT.

TRIAL OUTCOME MEASURES

Primary outcome measures will include diagnostic test characteristics (sensitivity, specificity, accuracy) for ^{18}F FDG-PET-CT and DCE-CT in relation to a subsequent clinical diagnosis of lung cancer. The outcome measures used in the economic model will include accuracy, estimated life expectancy and quality-adjusted life years (QALYs). Costs will be estimated from an NHS perspective. Incremental cost-effectiveness ratios will compare management strategies with DCE-CT to strategies without DCE-CT, where DCE-CT is expected to cost less than half that of ^{18}F FDG-PET-CT.

Secondary outcome measures will include diagnostic test characteristics for ^{18}F FDG-PET-CT with incorporation of CT appearances and combined DCE-CT/ ^{18}F FDG-PET-CT. The incidence of incidental extrathoracic findings on ^{18}F FDG-PET/CT, subsequent investigations and costs will also be determined.

Recruitment will be terminated when 375 patients have been recruited and undergone DCE-CT. The study has two possible end points for each patient: either the diagnosis of lung cancer via biopsy or a diagnosis of benign or non-lung cancer via either biopsy or failure of the imaged nodule to progress (increase in size) during the 2-year follow-up period. The end of the study will be reached when the last study patient reaches either of these two end points or withdraws full consent for continuing in the study.

CONCLUSIONS

With the potential adoption of a CT-based lung cancer screening programme in the UK, the number of patients with a SPN requiring further investigation could increase substantially. Furthermore, SPNs are a common finding on CT examinations undertaken for diagnostic or staging purposes. Novel cost-effective approaches to the assessment of SPNs will be of value to the NHS.

This study will provide accurate data on the diagnostic performances of DCE-CT and ^{18}F FDG-PET/CT in the NHS for the characterisation of SPNs: the decision analytic modelling will assess the likely costs and health outcomes resulting from incorporation of DCE-CT into management strategies for patients with SPNs.

Author affiliations

¹Department of Radiology, Papworth Hospital, Cambridge, UK

²Department of Thoracic Oncology, Papworth Hospital, Cambridge, UK

³Institute of Nuclear Medicine, University College London, London, UK

⁴Public Health Sciences and Medical Statistics, University of Southampton, Southampton, UK

⁵Southampton Clinical Trials Unit, University of Southampton, Southampton, UK

⁶Centre for Innovation and Leadership in Health Sciences, University of Southampton, UK

⁷Division of Imaging Sciences and Biomedical Engineering, King's College London, London, UK

⁸Department of Medical Physics, Paul Strickland Scanner Centre, Mount Vernon Hospital, East and North Herts NHS Trust, Stevenage, UK

⁹Radiation Protection Department, East and North Hertfordshire NHS Trust, Stevenage, UK

¹⁰Faculty of Health Sciences, University of Southampton, Southampton, UK

¹¹Department of Radiology and Respiratory Medicine, Southampton University Hospitals NHS Foundation Trust, Southampton, UK

¹²Departments of Respiratory and Nuclear Medicine, Brighton and Sussex University Hospitals NHS Trust, Brighton, UK

¹³West of Scotland PET Centre, Gartnavel Hospital, Glasgow, UK

¹⁴Department of Radiology, Aberdeen Royal Hospitals NHS Trust, Aberdeen, UK

¹⁵Department of Respiratory Medicine, Leeds Teaching Hospitals NHS Trust, Leeds, UK

¹⁶North West Lung Centre, University Hospital of South Manchester, Manchester, UK

¹⁷Department of Radiology, Churchill Hospital and University of Oxford, Oxford, UK

¹⁸Department of Radiology, Worcestershire Royal Hospital, Worcester, UK

¹⁹Department of Thoracic Medicine, East Sussex Hospitals NHS Trust, Saint Leonards-on-Sea, UK

²⁰Department of Radiology, University of Cambridge School of Clinical Medicine, Biomedical research centre, University of Cambridge, Cambridge, UK

Correction notice This article has been corrected since it published Online First. The section heading has been changed to 'Lung cancer'.

Acknowledgements The trial is funded by the NIHR HTA Programme (grant no: 09/22/117) and is being run by Southampton Clinical Trials Unit, directed by Professor Gareth Griffiths and part funded by CRUK. NRQ and RCR are part funded by the Cambridge Biomedical Research Centre and the Cancer Research Network: Eastern.

Collaborators The SPUTNIK Group: KAM, FJG, SG, SH, JJ, Andrew Clegg, NRQ, RCR, Ashley Groves, FVG, FWP, WLW, Richard Riley, LB, PM, CM, CP, AB, Andrea Lodge, John Buscombe, Theresa Green, Amanda Stone, Neal Navani, Robert Shortman, Gabriella Azzopardi, JS, RE, AS, Lucy Pike, DS, SD, AJF, Sarah Doffman, Janice Bush, Jane Lyttle, Kenneth Jacob, SH, Joris van der Horst, Joseph Sarvesvaran, Barbara McLaren, Lesley Gomersall, Lutfi Kurban, Ravi Sharma, Kathleen Collie, KK, Steve O'Hickey, Jayne Tyler, Sue King, John O'Brien, Rajiv Srivastava, Hugh Lloyd-Jones, Sandra Beech, Matthew Callister, Andrew Scarsbrook, Victoria Ashford-Turner, PC, Elaine Smith, Susan Mbale, Nick Adams, Osei Kankam, Gail Pottinger.

Contributors KAM, FJG, SG, SH, JJ, Andrew Clegg, AMG, Tony Frew, FVG, FWP, WLW and Richard Riley designed the study and obtained funding. Andrea Lodge, Theresa Green, Amanda Stone, Rob Shortman, Gabriella Azzopardi, Kathleen Collie, Barbara McLaren, June Innes, Kenneth Jacob, Jane Lyttle, Victoria Ashford-Turner, Susan Mbale, Jayne Tyler, Hugh Lloyd-Jones, Sandra Beech and Gail Pottinger were involved in data acquisition. KAM, FJG, SH, SG, JJ, NRQ, RCR, Andrew Clegg, AMG, FVG, FWP and RR were involved in data analysis and interpretation. RCR, FJG, NRQ, KAM, SG, SH, JJ, JM were involved in manuscript draft and revision for intellectual content. All members of the SPUTNIK Group were involved in approval of the final version. FJG is the guarantor of the study.

Funding Health Technology Assessment Programme (09/22/117).

Competing interests None declared.

Patient consent Obtained.

Ethics approval South West-Exeter.

Provenance and peer review Not commissioned; internally peer reviewed.

Data sharing statement Details of the methods used are freely available to any researcher wishing to use them.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which



permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Cronin P, Dwamena BA, Kelly AM, *et al.* Solitary pulmonary nodules: meta-analytic comparison of cross-sectional imaging modalities for diagnosis of malignancy. *Radiology* 2008;246:772–82.
2. Comber LA, Keith CJ, Griffiths M, *et al.* Solitary pulmonary nodules: impact of quantitative contrast-enhanced CT on the cost-effectiveness of FDG-PET. *Clin Radiol* 2003; 58:706–11.
3. Yi CA, Lee KS, Kim BT, *et al.* Tissue characterization of solitary pulmonary nodule: comparative study between helical dynamic CT and integrated PET/CT. *J Nucl Med* 2006;47:443–50.
4. Christensen JA, Nathan MA, Mullan BP, *et al.* Characterization of the solitary pulmonary nodule: 18F-FDG PET versus nodule-enhancement CT. *Am J Roentgenol* 2006;187: 1361–7.
5. Orlacchio A, Schillaci O, Antonelli L, *et al.* Solitary pulmonary nodules: morphological and metabolic characterisation by FDG-PET-MDCT. *Radiol Med* 2007;112:157–73.

Accuracy and cost-effectiveness of dynamic contrast-enhanced CT in the characterisation of solitary pulmonary nodules —the SPUtNik study

N R Qureshi, R C Rintoul, K A Miles, S George, S Harris, J Madden, K Cozens, L A Little, K Eichhorst, J Jones, P Moate, C McClement, L Pike, D Sinclair, W L Wong, J Shekhdar, R Eaton, A Shah, L Brindle, C Peebles, A Banerjee, S Dizdarevic, S Han, F W Poon, A M Groves, L Kurban, A J Frew, M E Callister, P Crosbie, F V Gleeson, K Karunasaagarar, O Kankam and F J Gilbert

BMJ Open Resp Res 2016 3:
doi: [10.1136/bmjresp-2016-000156](https://doi.org/10.1136/bmjresp-2016-000156)

Updated information and services can be found at:
<http://bmjopenrespres.bmj.com/content/3/1/e000156>

These include:

References

This article cites 5 articles, 1 of which you can access for free at:
<http://bmjopenrespres.bmj.com/content/3/1/e000156#BIBL>

Open Access

This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See:
<http://creativecommons.org/licenses/by/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Critical care](#) (6)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>