



King's Research Portal

DOI: 10.1007/JHEP11(2016)173

Document Version Publisher's PDF, also known as Version of record

Link to publication record in King's Research Portal

Citation for published version (APA): Acharya, B. S., Bozek, K., Romão, M. C., King, S. F., & Pongkitivanichkul, C. (2016). Neutrino mass from M theory SO(10). *Journal of High Energy Physics*, *2016*(11), Article 173. Advance online publication. https://doi.org/10.1007/JHEP11(2016)173

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Published for SISSA by 🖄 Springer

RECEIVED: August 11, 2016 ACCEPTED: November 19, 2016 PUBLISHED: November 29, 2016

Neutrino mass from M theory SO(10)

Bobby S. Acharya,^{*a,b*} Krzysztof Bożek,^{*a*} Miguel Crispim Romão,^{*c*} Stephen F. King^{*c*} and Chakrit Pongkitivanichkul^{*a*}

^a Department of Physics, King's College, WC2R 2LS, London, U.K.
^b International Centre for Theoretical Physics, I-34151 Trieste, Italy
^c School of Physics and Astronomy, University of Southampton, SO17 1BJ, Southampton, U.K.

E-mail: bobby.acharya@kcl.ac.uk, krzystof.bozek@kcl.ac.uk,

m.crispim-romao@soton.ac.uk, s.f.king@soton.ac.uk,

chakrit.pongkitivanichkul@kcl.ac.uk

ABSTRACT: We study the origin of neutrino mass from SO(10) arising from M Theory compactified on a G_2 -manifold. This is linked to the problem of the breaking of the extra U(1) gauge group, in the SU(5) × U(1) subgroup of SO(10), which we show can achieved via a (generalised) Kolda-Martin mechanism. The resulting neutrino masses arise from a combination of the seesaw mechanism and induced R-parity breaking contributions. The rather complicated neutrino mass matrix is analysed for one neutrino family and it is shown how phenomenologically acceptable neutrino masses can emerge.

KEYWORDS: GUT, Neutrino Physics, Compactification and String Models, Supersymmetric Standard Model

ARXIV EPRINT: 1607.06741



Contents

I Introduction and motivation		
SO(10) SUSY GUTS from M theory on G_2 -manifolds	3	
2.1 The vector-like family splitting	6	
2.2 R-parity violation	6	
2.3 The see-saw mechanism	10	
2.4 Effective light families	11	
$\mathrm{U}(1)_X$ breaking scenarios and mechanisms	12	
Neutrino-neutralino mass matrix	16	
4.1 The mass matrix hierarchies	20	
Numerical results	22	
5.1 ν component of the lightest state	23	
5.2 Matter neutrino Yukawas and B-RPV couplings	25	
Conclusions and discussion	28	
	Introduction and motivation SO(10) SUSY GUTS from M theory on G_2 -manifolds 2.1 The vector-like family splitting 2.2 R-parity violation 2.3 The see-saw mechanism 2.4 Effective light families U(1) _X breaking scenarios and mechanisms Neutrino-neutralino mass matrix 4.1 The mass matrix hierarchies Numerical results 5.1 ν component of the lightest state 5.2 Matter neutrino Yukawas and B-RPV couplings Conclusions and discussion	

1 Introduction and motivation

The discovery of neutrino mass and lepton mixing provides key evidence for new physics beyond the Standard Model (SM) [1–5]. The seesaw mechanism [6–10] is an attractive possibility to account for the origin of neutrino mass and lepton mixing in terms of righthanded neutrinos with large Majorana masses. SO(10) Grand Unified Theories (GUTs) [11] predict such right-handed neutrinos which appear along with SM matter fields in a single 16 multiplet. When the SO(10) gauge group is broken to that of the SM, neutrino mass is an inevitable consequence. In order to satisfy the constraint of gauge coupling unification, we shall here assume low energy supersymmetry (SUSY) [12]. However to also account for gravity, one needs to go beyond gauge theories, and here we shall focus on an M theory version of string theory [13, 14].

Recently we showed how SO(10) SUSY GUTs could emerge from M Theory compactified on a G_2 -manifold [15]. In this framework, discrete symmetry and Wilson lines [16] were used to prevent proton decay while maintaining gauge unification. In contrast to the SU(5) version [17, 18], the Wilson line symmetry breaking mechanism in SO(10) requires additional matter at the TeV scale, with the quantum numbers of an extra $\mathbf{16}_X$ plus $\mathbf{\overline{16}}_X$ [15]. In addition, there were a number of unresolved issues in this approach, notably the mechanism for breaking the extra gauged U(1)_X which accompanies the SM gauge group after the Wilson line symmetry breaking mechanism in SO(10). This gauge group is the usual one in the maximal SO(10) subgroup SU(5) × U(1)_X,¹ where SU(5) embeds the SM gauge group. The key point is that, since Abelian Wilson line symmetry breaking preserves the rank of the gauge group, the U(1)_X gauge group needs to be broken by some other mechanism in the low energy effective field theory. Since right-handed Majorana neutrino masses can only arise once the U(1)_X is broken, the origin of neutrino mass is therefore linked to this symmetry breaking.

In this paper we address the problem of $U(1)_X$ breaking and neutrino masses arising from the SO(10) M theory, following the construction in [15], although our approach to solving these problems may be more general than the specific example studied. To break the $U(1)_X$ gauge symmetry, we employ a (generalised) Kolda-Martin mechanism [19], where higher order operators can break the symmetry, inducing vacuum expectation values (VEVs) in the scalar right-handed neutrino components of both the matter **16** and the extra **16**_X, as well as their conjugate partners. The subsequent induced R-parity violation [20] provides additional sources of neutrino mass, in addition to that arising from the seesaw mechanism [6–10]. The resulting 11×11 neutrino mass matrix is analysed for one neutrino family (nominally the third family) and it is shown how a phenomenologically acceptable neutrino mass can emerge. We defer any discussion of flavour mixing to a possible future study of flavour from M theory. Here we only show that symmetry breaking and viable neutrino masses can arise within the framework of M theory SO(10), which is a highly non-trivial result, given the constrained nature of M theory constructions.

It is worth remarking that there are other alternative ways that have been proposed to study neutrino masses in string theory, which are complementary to the approach followed here. For example, it is possible to obtain large Majorana mass terms from instanton effects [21–25], large volume compactification [26], or orbifold compactifications of the heterotic string [25]. However the origin of Majorana mass terms in SO(10) has been non-trivial to realise from the string theory point of view. In GUTs all matter fields are unified in **16** multiplets whereas Higgs fields and triplet scalars are unified in **10**. Since string theory does not predict light particles in representations larger than the adjoint, the traditional renormalisable terms involving **126**, **126**, **210**, e.g., $W \sim 126 \, 16 \, 16 \, 16 \, 16$. Assuming that the supersymmetric partner of the right handed neutrino singlet gets a VEV, the Majorana mass is given by $M \sim \frac{\langle \tilde{N} \rangle^2}{M_{PL}}$. However, the required values of neutrino mass imply $M > 10^{14} \, \text{GeV}$, which gives $\langle \tilde{N} \rangle \sim \sqrt{Mm_{Pl}} \sim 10^{16} \, \text{GeV}$. The implementation of the seesaw mechanism [6–10] in other corners of string compactification has also been discussed [27–30].

The layout of the remainder of the paper is as follows. In section 2, we will review the SO(10) construction from M Theory on G_2 -manifolds, expanding the discussion in [15]. In section 3, the mechanism for $U(1)_X$ breaking will be given. The neutrino mass matrix will be analysed in section 4, and the numerical results presented in section 5. Finally we conclude in section 6.

¹The U(1)_X is also commonly called U(1)_X in the literature.

2 SO(10) SUSY GUTS from M theory on G_2 -manifolds

M Theory compactified on a G_2 -manifold leads to a 4 dimensional theory with $\mathcal{N} = 1$ SUSY, where gauge fields and chiral fermions are supported by different types of singularities in the compactified space [31, 32]. Yang-Mills fields are supported on three dimensional subspaces of the extra dimensions, along which there is an orbifold singularity, while chiral fermions will be further localised on conical singularities localised on these three dimensional spaces and interact with the gauge fields.

One of the key features of M Theory compactified on G_2 -manifolds without fluxes is that it provides a framework for generating hierarchies of mass scales. To understand the reason behind this notice that in M Theory, the moduli fields, s_i , are paired with the axions, a_i , in order to form a complex scalar component of a superfield Φ_i

$$\Phi_i = s_i + ia_i + \text{fermionic terms} . \tag{2.1}$$

In the absence of fluxes, the axions enjoy an approximate shift-symmetry, which is remnant of the higher dimensional gauge symmetry, $a_i \rightarrow a_i + c_i$ where c_i is an arbitrary constant. This Peccei-Quinn symmetry, in conjunction with holomorphicity of the superpotential, severely constrains the superpotential for the moduli. As such, terms which are polynomial in the moduli and matter fields are forbidden at tree-level in superpotential, appearing only in the Kähler potential.

In general non-perturbative effects such as instantons break the above shift symmetry, and generate a non-perturbative superpotential involving moduli and matter. Interactions will be generated by membrane instantons, whose actions are given by exponentials of the moduli. As the moduli stabilise and acquire VEVs, these exponentials will turn out to be small, and the VEV of the hidden sector superpotential naturally leading to a generation of hierarchical masses at the GUT scale [33]. These ideas were used to construct the G_2 -MSSM [17, 18], an SU(5) SUSY GUT from M Theory on a G_2 manifold with the MSSM spectrum. Here, we discuss an extention of the program to the SO(10) GUT group [15], while referring to previous work on G_2 compactifications and consequent predictions for the parameters [34, 35].

In the remainder of this section, we focus on the SO(10) SUSY GUT from M Theory on G_2 manifolds which we proposed in [15]. The breaking patterns of an abelian Wilson line are the same as the ones of an adjoint Higgs. The simplest case of a surviving group that is the most resembling to the SM is

$$\mathrm{SO}(10) \to \mathrm{SU}(3)_c \times \mathrm{SU}(2)_L \times \mathrm{U}(1)_Y \times \mathrm{U}(1)_X,$$

$$(2.2)$$

under which the branching rules of the GUT irreps read

$$\mathbf{10}: \ H_{u} = (\mathbf{1}, \mathbf{2})_{\left(\frac{1}{2}, 2\right)} \oplus H_{d} = (\mathbf{1}, \mathbf{2})_{\left(-\frac{1}{2}, -2\right)} \oplus D = (\mathbf{3}, \mathbf{1})_{\left(-\frac{1}{3}, 2\right)} \oplus \overline{D} = (\overline{\mathbf{3}}, \mathbf{1})_{\left(\frac{1}{3}, -2\right)}, \quad (2.3)$$

$$\mathbf{16}: \ L = (\mathbf{1}, \mathbf{2})_{\left(-\frac{1}{2}, 3\right)} \oplus e^{c} = (\mathbf{1}, \mathbf{1})_{(1, -1)} \oplus N = (\mathbf{1}, \mathbf{1})_{(0, -5)} \oplus u^{c} = (\overline{\mathbf{3}}, \mathbf{1})_{\left(-\frac{2}{3}, -1\right)} \oplus d^{c} = (\overline{\mathbf{3}}, \mathbf{1})_{\left(\frac{1}{3}, 3\right)} \oplus Q = (\mathbf{3}, \mathbf{2})_{\left(\frac{1}{6}, -1\right)}, \quad (2.4)$$

and the subscripts are the charges under $U(1)_Y \times U(1)_X$, which are normalised as $Q_Y = \sqrt{\frac{5}{3}}Q_1$, $Q_X = \sqrt{40}\tilde{Q}_X$, where Q_1 , \tilde{Q}_X are SO(10) generators.

The Wilson line can be conveniently represented as

$$\mathcal{W} = \exp\left[\frac{i2\pi}{N}\left(aQ_Y + bQ_X\right)\right] = \sum_{m=0}^{\infty} \frac{1}{m!} \left(\frac{i2\pi}{N}\right)^m \left(aQ_Y + bQ_X\right)^m, \quad (2.5)$$

where the coefficients a, b are constrained by the requirement that $\mathcal{W}^N = 1$ and specify the parametrisation of the Wilson line. Under the linear transformation

$$\frac{1}{2}a + 2b \to \alpha \,, \tag{2.6}$$

$$\frac{1}{3}a - 2b \to \beta \,, \tag{2.7}$$

its action on the fundamental irrep then reads

$$\mathcal{W}10 = \eta^{\alpha} H_u \oplus \eta^{-\alpha} H_d \oplus \eta^{-\beta} D \oplus \eta^{\beta} \overline{D}, \qquad (2.8)$$

where η is the *N*th root of unity.

Likewise the Wilson line matrix acts on the 16 irrep as

$$\mathcal{W}16 = \eta^{-\frac{3}{2}\beta}L \oplus \eta^{\alpha+\frac{3}{2}\beta}e^c \oplus \eta^{-\alpha+\frac{3}{2}\beta}N \oplus \eta^{-\alpha-\frac{1}{2}\beta}u^c \oplus \eta^{\alpha-\frac{1}{2}\beta}d^c \oplus \eta^{\frac{1}{2}\beta}Q, \qquad (2.9)$$

which could be simplified a bit further by replacing $\beta \to 2\beta$ without loss of generality, in order for the parameters to read as integers.

The effective discrete charges — of different states on a chiral supermultiplet that absorbs Wilson line phases — will be the overall charge of the discrete symmetry (common to all states belonging to the same GUT irrep) in addition to the Wilson line phases (different for each state inside the GUT irrep).

Having all the ingredients required to employ Witten's discrete symmetry proposal, we would like to have a consistent implementation of a well-motivated doublet-triplet splitting mechanism as it was done for SU(5). Unfortunately the customary approach to the problem does not seem to work with SO(10), as shown in [15]. To understand this first notice that Witten's splitting mechanism can only work in order to split couplings between distinct GUT irreps. This is understood as \mathcal{W} has the form of a gauge transformation of the surviving group and so it will never be able to split self bilinear couplings of a GUT irrep. For example, if one takes a **10** with Wilson line phases to contain the MSSM Higgses, we can see from eq. (2.8) that both mass terms for the Higgses and coloured triplets are trivially allowed. We could consider that in order to split the Higgses, H_u and H_d , from the coloured triplets — D, \overline{D} — we would need to add another **10**, but it was shown that this cannot be achieved and so we are ultimately left with light coloured triplets.

In order to allow for light D, \overline{D} we need to guarantee that they are sufficiently decoupled from matter to prevent proton-decay. To accomplish this, we can use the discrete symmetry to forbid certain couplings, namely to *decouple* D and \overline{D} from matter. Such couplings arise from the SO(10) invariant operator **10 16 16**, with **16** denoting the three SO(10) multiplets, each containing a SM family plus right handed neutrino N. If **16** transforms as η^{κ} **16**, the couplings and charge constraints are

 $H_u \mathbf{1616} : 2\kappa + \alpha + \omega = 0 \mod N \tag{2.10}$

$$H_d \mathbf{1616} : 2\kappa - \alpha + \omega = 0 \mod N \tag{2.11}$$

 $D1616 : 2\kappa - \beta + \omega \neq 0 \mod N \tag{2.12}$

$$\overline{D}\mathbf{1616} : 2\kappa + \beta + \omega \neq 0 \mod N, \tag{2.13}$$

where we allow for up-type quark Yukawa couplings together with couplings to the righthanded neutrinos,

$$y_u^{ij} H_u^w \mathbf{16}_i \mathbf{16}_j \equiv y_u^{ij} H_u^w (Q_i u_j^c + L_i N_j + i \leftrightarrow j), \qquad (2.14)$$

and similarly for down-type quarks and charged leptons.

The couplings forbidden at a renormalizable tree-level by the discrete symmetry are generically regenerated from Kähler interactions through the Giudice-Masiero mechanism [36]. While this provides the Higgsinos a TeV scale μ -term mass, it also originates effective trilinear couplings with an $\mathcal{O}(10^{-15})$ coefficient. As these are generic, we need to systematically study their physical implications at low energies, such as proton-decay, R-parity violation, and flavour mixing.

For proton decay, effective superpotential will be generate by the following Kähler potential

$$K \supset \frac{s}{m_{Pl}^2} \overline{D} d^c u^c + \frac{s}{m_{Pl}^2} De^c u^c + \frac{s}{m_{Pl}^2} DQQ + \frac{s}{m_{Pl}^2} \overline{D} QL + \frac{s}{m_{Pl}^2} DNd^c + \text{h.c.}, \qquad (2.15)$$

where we assume $\mathcal{O}(1)$ coefficients. As the moduli acquire non-vanishing VEVs, these become

$$W_{\text{eff}} \supset \lambda DQQ + \lambda De^{c}u^{c} + \lambda DNd^{c} + \lambda \overline{D}d^{c}u^{c} + \lambda \overline{D}QL, \qquad (2.16)$$

where we considering all couplings to be similar and taking one family for illustrative purposes. Notice that contrary to SU(5) case, there is no extra contribution from rotation of L and H_u as the bilinear term $\kappa L H_u$ is not allowed by gauge invariance.

We estimate the scalar triplet mediated proton decay rate to be

$$\Gamma_p \simeq \frac{|\lambda^2|^2}{16\pi^2} \frac{m_p^5}{m_D^4} \simeq \left(10^{42} \,\mathrm{yrs}\right)^{-1},\tag{2.17}$$

where we took the mass of the colour triplets to be $m_D \simeq 10^3 \,\text{GeV}$.

Another limit for triplet scalar comes from the cosmological constraints on its decay. As we have seen from proton-decay operators, triplet scalars can decay into quarks. If they start to decay during the Big Bang Nucleosynthesis (BBN) then nucleons could be disassociated, spoiling the predictions for light element abundances. We can estimate another limit on the triplet scalar mass by calculating its lifetime as it decay through the processes $D \to e^c u^c, QQ, QL, d^c u^c$, and we get

$$\Gamma \simeq \lambda^2 m_D \simeq (0.1 \text{ sec})^{-1}, \qquad (2.18)$$

which is approximately consistent with BBN constraint. They will also give interesting collider signatures due to their long-lived nature.

2.1 The vector-like family splitting

Because the presence of a light vector-like pair coloured triplets spoils unification, we need a workaround that will preserve unification while keeping the presented doublet-triplet problem solution. We achieve this by considering the presence of extra matter that would form a complete GUT irrep with the coloured triplets, and hence restore unification. Unification constraints requires heavy states with equivalent SM gauge numbers, say d_X^c and $\overline{d^c}_X$, that have to be subtracted from the spectrum. This can be achieved by adding a vector-like family pair, $\mathbf{16}_X \mathbf{\overline{16}}_X$, and splitting its mass terms using Wilson line phases.

Furthermore, as the Wilson line breaking pattern is rank-preserving, we still need to break the extra abelian gauge factor $U(1)_X$. This can be achieved if a scalar component of the right-handed conjugated neutrino pair of an extra vector-like family $\mathbf{16}_X$, $\mathbf{\overline{16}}_X$ acquires VEVs. On top of this, this VEV can generate a Majorana mass for the matter right-handed conjugated neutrinos, providing a crucial ingredient for a type I see-saw mechanism.

In order to preserve gauge coupling unification, we notice that the down-type quarks $-d_X^c$, $\overline{d^c}_X$ — have the same SM quantum numbers as the coloured triplet pair — D, \overline{D} — coming from the **10**. We take **16**_X to be localised along a Wilson line, and find that it transforms under the discrete symmetry as

$$\mathbf{16}_X \to \eta^x \left(\eta^{-3\gamma} L \oplus \eta^{3\gamma+\delta} e^c \oplus \eta^{3\gamma-\delta} N \oplus \eta^{-\gamma-\delta} u^c \oplus \eta^{-\gamma+\delta} d^c \oplus \eta^{\gamma} Q \right).$$
(2.19)

On the other hand, we let $\overline{\mathbf{16}}_X$ transform without Wilson line phases, $\overline{\mathbf{16}}_X \to \eta^{\overline{x}} \overline{\mathbf{16}}_X$, and the condition for the mass term that will split the vector-like family is

$$\overline{d^c}_X d^c_X : x - \gamma + \delta + \overline{x} = 0 \mod N, \tag{2.20}$$

whilst forbidding all the other self couplings that would arise from $\mathbf{16}_X \overline{\mathbf{16}}_X$. The d_X^c , $\overline{d^c}_X$ quarks will then be naturally endowed a GUT scale mass through membrane instantons, provided that the singularities supporting $\mathbf{16}_X$, $\overline{\mathbf{16}}_X$ are close enough to each other in the compactified space. The remaining states of $\mathbf{16}_X$, $\overline{\mathbf{16}}_X$ will have a μ term of order TeV through the Giudice-Masiero mechanism. The coloured triplets — D, \overline{D} — and the light components of $\mathbf{16}_X$, $\overline{\mathbf{16}}_X$ will effectively account for a full vector-like family. The light spectrum is then the one of MSSM in addition to this vector-like family, which in turn preserves unification, with a larger unification coupling at the GUT scale.

2.2 R-parity violation

Despite the existence of an effective matter parity symmetry inside SO(10), the presence of a vector-like family will lead to R-parity violating (RPV) interactions though the VEV of the N_X , \overline{N}_X components in the presence of moduli generated interactions. Furthermore, as we will see in detail in section 3, the scalar component of the matter conjugate right-handed neutrino, N, will also acquire a VEV. These VEVs break SO(10) and will inevitably generate RPV. These interactions will mediate proton-decay, enable the lightest supersymmetric particle (LSP) to decay, and generate extra contributions to neutrino masses. In our framework RPV is generic, not only arising from allowed superpotential terms but as well from Kähler interactions involving moduli fields.

The interactions that break R-parity can either be trilinear or bilinear (B-RPV), and have different origins in our framework. The first contribution we can find comes from the tree-level renormalizable superpotential allowed by the discrete symmetry. Since we will encounter $\langle N \rangle \neq 0$, this means that even in a minimal setup, there will be an R-RPV contribution from matter Dirac mass coupling

$$W \supset y_{\nu} N H_u L \,, \tag{2.21}$$

reading

$$W \supset y_{\nu} \langle N \rangle H_u L . \tag{2.22}$$

Next we turn our attention to the Kähler potential, where interactions otherwise forbidden by the discrete symmetry might arise if there is a modulus with required charge. In such case, there is another contribution arising from the non-vanishing VEVs of N_X , $N \overline{N}_X$ in conjugation with moduli VEVs. To see this, notice that in the Kähler potential there are generically interactions of the form

$$K \supset \frac{1}{m_{Pl}} N H_u L + \frac{s}{m_{Pl}^2} \overline{N}_X H_u L + \frac{s}{m_{Pl}^2} \overline{N}_X^{\dagger} H_u L + \text{ h.c.}, \qquad (2.23)$$

where while the first term exists in zeroth order in moduli (otherwise there would be no neutrino Dirac mass in the superpotential), the last two are otherwise forbidden by the discrete symmetry, and s denotes a generic modulus for each coupling. These terms will generate contributions to B-RPV as N_X , $N \overline{N}_X$, s acquire VEVs.

There are two types of contribution arising from the terms above. The first is generates through the Giudice-Masiero mechanism. As the moduli acquire VEVs, new holomorphic couplings will appear in the superpotential

$$W_{\rm eff,1} = \frac{m_{3/2}}{m_{Pl}} \langle N \rangle H_u L + 0.1 \frac{m_{3/2}}{m_{Pl}} \langle N_X \rangle H_u L + 0.1 \frac{m_{3/2}}{m_{Pl}} \langle \overline{N}_X^{\dagger} \rangle H_u L \,, \tag{2.24}$$

where $m_{3/2} \simeq \mathcal{O}(10^4)$ GeV, and since $s/m_{Pl} \simeq 0.1$ in *M* Theory. Notice that in principle we would also have a term in the Kähler potential involving *N*, but this can be found to be subleading in comparison to the term arising from the Dirac mass eq. (2.22).

The second contribution arises if the F-terms of the fields N_X , N, \overline{N}_X are non-vanishing. In this case, we expect the appearance of the contributions

$$W_{\text{eff},2} = \frac{\langle F_N \rangle}{m_{Pl}} H_u L + 0.1 \frac{\langle F_{N_X} \rangle}{m_{Pl}} H_u L + 0.1 \frac{\langle F_{\overline{N}_X^\dagger} \rangle}{m_{Pl}} H_u L \,, \tag{2.25}$$

and its magnitude will depend on how much F-breaking provoked by our symmetry breaking mechanism. Here we are considering that the case where $\overline{N}_X^{\dagger} H_u L$ cannot exist in the Kähler potential in zeroth order in a modulus field.

Putting all together, the B-RPV interactions account to the B-RPV paramter

$$W \supset \kappa H_u L \tag{2.26}$$

with

$$\kappa = \left(y_{\nu} + \frac{m_{3/2}}{m_{Pl}}\right) \langle N \rangle + 0.1 \frac{m_{3/2}}{m_{Pl}} \langle N_X \rangle + 0.1 \frac{m_{3/2}}{m_{Pl}} \langle \overline{N}_X^{\dagger} \rangle + \frac{\langle F_N \rangle}{m_{Pl}} + 0.1 \frac{\langle F_{N_X} \rangle}{m_{Pl}} + 0.1 \frac{\langle F_{N_X} \rangle}{m_{Pl}},$$
(2.27)

and the relative strength of each contribution is model detail dependent, namely on neutrino Yukawa textures, symmetry breaking details, and F-flatness deviation.

In a similar manner, trilinear RPV couplings will be generated when N, N_X , \overline{N}_X , s acquire VEVs. In order to systematically study this, we notice that the trilinear RPV couplings come from the term

16 16 16 16, **16**_X**16 16 16**,
$$\overline{\mathbf{16}}_X^{\dagger}$$
16 16 16 (2.28)

as the scalar component of N_X , N acquires non-vanishing VEVs. Notice that the last term lives in the Kähler potential. These are made forbidden at tree-level using the discrete symmetry of the compactified space. However, just like the μ terms and the B-RPV terms shown above, these terms will in general be present in the Kähler potential and will effectively be generated as the moduli acquire VEVs. This happens again through the Giudice-Masiero mechanism and we will find

$$\mathcal{O}\left(\frac{m_{3/2}}{m_{Pl}^2}(\langle N \rangle + \langle N_X \rangle + \langle \overline{N}_X^{\dagger} \rangle)\right) \{LLe^c, \ LQd^c, \ u^c d^c d^c\},$$
(2.29)

where $m_{3/2}/m_{Pl} \simeq \mathcal{O}(10^{-14})$. The apparent suppression of trilinear RPV is understood as these terms can only be generated by non-renormalizable terms in an SO(10) context.

Similarly to the B-RPV case, there will be further contributions if the F-terms of N_X , N, \overline{N}_X are non-vanishing. Namely we find

$$\mathcal{O}\left(\frac{\langle F_N \rangle + \langle F_{N_X} \rangle + \langle F_{\overline{N}_X^{\dagger}} \rangle}{m_{Pl}^2}\right) \{LLe^c, \ LQd^c, \ u^c d^c d^c\},$$
(2.30)

and again we expect these to be sub-leading even if the F-terms are not vanishing.

We see then that the values of all RPV coupling are strictly related to the details of the breaking mechanism employed to break the extra $U(1)_X$. This will be studied in great detail in section 3. Furthermore, the bilinear B-RPV term generates a contribution to the physical neutrino masses [20, 37]. The complete picture of neutrino masses, including B-RPV operators, will be discussed in section 4.

We can study now some direct effects of RPV in the dynamics of our class of models. Under the assumption that $\kappa \ll \mu$, performing a small rotation, of $\mathcal{O}(\kappa/\mu)$, in (H_d, L) space, the last term can be absorbed $\mu H_d H_u$. As a consequence, the first two terms will be enhanced by the Yukawa couplings $y_e H_d Le^c$, etc., leading to

$$W \supset y_e \frac{\kappa}{\mu} LLe^c + y_d \frac{\kappa}{\mu} LQd^c + \lambda \frac{v}{m_{Pl}} u^c d^c d^c, \qquad (2.31)$$

and we have dropped the $\mathcal{O}(1/m_{Pl})$ contributions to the first two terms since now the Yukawa rotated contributions are much larger. Also, we kept the last term with the parametrization v describing all contributions. These will be very small, for example in the case the VEVs are high-scale, $\langle N_X \rangle \simeq 10^{16}$ GeV, the trilinear RPV coupling strength is of $\mathcal{O}(10^{-16})$. A direct consequence of this result is that proton decay will be slow, even when the $\Delta L = 1$ terms are enhanced.

While the proton is relatively stable, the enhanced terms will provide a decay channel for the LSP, which is now unstable. In the limit that we can take the final states to be massless, and considering that the LSP is a neutralino mainly composed of neutral gauginos, the LSP lifetime through the decay $\tilde{\chi}^0 \to d^c QL$ can be estimated from a tree-level diagram involving a virtual \tilde{d}^c with mass $m_{0,2}^2$

$$\tau_{LSP} \simeq \left(3.9 \times 10^{-15}\right) \left(\frac{\mu}{g_w y_d \kappa}\right)^2 \left(\frac{m_0}{10 \text{ TeV}}\right)^4 \left(\frac{100 \text{ GeV}}{m_{LSP}}\right)^5 \text{sec}, \qquad (2.32)$$

where g_w is a weak gauge coupling. The LSP lifetime is bounded to be either $\tau_{LSP} \lesssim 1$ sec or $\tau_{LSP} \gtrsim 10^{25}$ sec [37, 38], from Big Bang Nucleosythesis (BBN) and indirect Dark Matter (DM) experiments, respectively. If we take $m_{LSP} \simeq 100$ GeV, $m_0 \simeq 10$ TeV, $y_d = y_b \simeq 10^{-2}, g_w \simeq 0.1$, we find that the VEV v_X is constrained to be either

$$\kappa \gtrsim 6 \times 10^{-2} \text{ GeV}$$
 (2.33)

or
$$\kappa \lesssim 2 \times 10^{-14} \text{ GeV},$$
 (2.34)

for a short- and long-lived LSP, respectively. In the above estimate we used the fact that the decay involving the bottom Yukawa is the largest contribution to the decay width.

We can use the above result to infer some parametric dependence on the scale of the U(1)_X breaking. If we have the leading contribution to the B-RPV coupling to be $\kappa \simeq \langle N_X \rangle \lambda \Rightarrow \langle N_X \rangle \gtrsim 10^{12} \,\text{GeV}$. In this case, the LSP is too short lived to be a good DM candidate, but decays quickly enough to not spoil BBN predictions. On the otherhand, a low-scale VEV is bound to be $\langle N_X \rangle \lesssim 1 \,\text{GeV}$ in order to allow for a long-lived LSP. This would imply the abelian gauge boson associated with extra U(1)_X to be light, $m_{Z'} < \mathcal{O}(1) \,\text{GeV}$. This last scenario is completely excluded from experimental searches.

The lack of a good DM candidate in the visible sector indicates us that DM is realised elsewhere. For instance, it has been recently suggested that in the context of String/MTheory, the generic occurrence of hidden sectors could account for the required DM mechanics [39].

²See, for example, the diagrams in [12].

2.3 The see-saw mechanism

The relevance of the bounds on the rank-breaking VEV is only fully understood when studying the details of symmetry breaking mechanism and neutrino masses. For example, if we start with an SO(10) invariant theory the Yukawas are unified for each family leading to at least one very heavy Dirac neutrino mass, m_{ν}^{D} . However, if the right-handed conjugated neutrino has a heavy Majorana mass, then the physical left-handed neutrino mass will be small through a type I see-saw mechanism. In order to accomplish this, one has to allow the following terms in the superpotential

$$W \supset y_{\nu}H_{u}LN + MNN, \tag{2.35}$$

where y_{ν} are the neutrino Yukawas, L the matter lepton doublets, N the right-handed conjugated neutrino, and M its Majorana mass, which we take $M \gg m_{\nu}^{D} = y_{\nu} \langle H_{u} \rangle$. With the above ingredients, a mostly left-handed light neutrino will have a physical mass

$$m_{phy}^{\nu} \simeq -\frac{(m_{\nu}^D)^2}{M}.$$
 (2.36)

One of the most appealing features of SO(10) models is that each family is in a **16** which includes a natural candidate for the right-handed conjugated neutrino, the N. In order to employ a type I see-saw mechanism, we need to generate a Majorana mass term for the matter right-handed conjugated neutrino through the operator $W \supset \overline{\mathbf{16}}_X \overline{\mathbf{16}}_X \mathbf{16} \mathbf{16}^3$ leading to the operator

$$\frac{1}{m_{Pl}}\overline{N}_X\overline{N}_XNN\,,\qquad(2.37)$$

from which the Majorana mass for the (CP conjugated) right-handed neutrino field N is emerges as

$$M \simeq \frac{\langle \overline{N}_X \rangle^2}{m_{Pl}} \ . \tag{2.38}$$

We can now relate the bounds on the value of the D-flat VEVs $\langle \overline{N}_X \rangle = \langle N_X \rangle$ from both RPV and the requirement of a realistic see-saw mechanism. Since the physical neutrino mass in type I see-saw mechanism is given by

$$m_{\nu}^{phy} \simeq \frac{(m_{\nu}^D)^2}{M},$$
 (2.39)

assuming $m_{\nu}^D \simeq \mathcal{O}(100 \text{ GeV})$, and knowing that the upper bound on the neutrino masses $m_{\nu}^{phy} \lesssim 0.1 \text{ eV}$, one finds

$$M \gtrsim 10^{14} \text{ GeV} \Rightarrow \langle N_X \rangle \gtrsim 10^{16} \text{ GeV}.$$
 (2.40)

The above argument suggests that we need to break the $U(1)_X$ close to the GUT scale. Since the Wilson line breaking mechanism is rank-preserving, we need to look for an alternative solution. Although the neutral fermion mass matrix will be considerably more intricate, obscuring the relations and hierarchies amongst different contributions to the neutrino masses, the above estimate motivates the need for a high-scale $U(1)_X$ breaking mechanism.

 $^{^{3}}$ Given that in M Theory one does not account for irreps larger than the adjoint, this is the lowest order term that can generate a right-handed neutrino Majorana mass.

2.4 Effective light families

For a simple SUSY SO(10) model where each family is unified into a single irrep with universal soft masses, it is well known that electroweak symmetry is difficult to break [40– 45]. Since the two Higgs soft masses are unified at GUT scale and have similar beta function due to Yukawa unification, either both masses are positive at electroweak scale and symmetry is not broken or both masses are negative and the potential becomes unbounded from below. Another aspect of Yukawa unification problem lies in the fact that low energy spectrum of quarks and leptons requires some degree of tuning in parameter space when their RG runnings are considered. A customary solution to these problems it the use of higher dimensional representations [46], which are not present in our framework.

The EWSB and Yukawa textures issues are naturally solved if each family is not contained in one single complete 16, but is instead formed of states from different Ultra Violet (UV) complete 16s. In order to implement this in our framework, first we assume the existence of multiple 16 with independent and different Wilson Line phases, alongside the existence of multiple 16. Second, we employ Witten's proposal to turn on some vector-like masses such that three effective light 16 survive. Since in M Theory the strength of the Yukawa couplings is given by membrane instantons, and are therefore related to distances between the singularities supporting the respective superfields, by constructing effective families from different UV 16s one can obtain different Yukawa couplings within each family.

Such solution can be achieved if one considers M complete $\overline{\mathbf{16}}_j$ and M + 3 complete $\mathbf{16}_i$ UV irreps. Allowing for masses between different states of these UV irreps to appear, one has schematically the mass terms in the superpotential

$$\mathbf{16}_{i}\mu_{ij}\overline{\mathbf{16}}_{j},\tag{2.41}$$

but since i = 1, ..., M while j = 1, ..., M + 3 the mass matrix μ_{ji} can only have at most rank M and hence there will be three linear combinations composing three **16** that will remain massless. If these masses are truly SO(10) invariant, i.e.

$$\mathbf{16}_{i}\mu_{ij}\overline{\mathbf{16}}_{j} = \mu_{ij}\left(Q_{i}\overline{Q}_{j} + L_{i}\overline{L}_{j} + \ldots\right), \qquad (2.42)$$

each effective light family will be SO(10) invariant. Consequently each family will retain unified Yukawa textures, and so this does not solve our problem of splitting the Yukawa couplings within each family.

However, Witten's proposal endows our framework with a GUT breaking discrete symmetry which can be employed to ensure that the superpotential mass matrices between the UV states

$$\mu_{ij}^Q Q_i \overline{Q}_j + \mu_{ij}^L L_i \overline{L}_j + \dots, \qquad (2.43)$$

are not the same, leading to different diagonalisations of Q, L, etc which in turn break the Yukawa SO(10) invariance. In order to accomplish that, take for example that the $\mathbf{16}_i$ absorb distinct and independent Wilson line phases, while $\mathbf{\overline{16}}_j$ do not, i.e. the UV irreps will transform under the discrete symmetry as

$$\mathbf{16}_{i} \to \eta^{m_{i}} \left(\eta^{-3\gamma_{i}} L_{i} \oplus \eta^{3\gamma_{i}+\delta_{i}} e_{i}^{c} \oplus \eta^{3\gamma_{i}-\delta_{i}} N_{i} \oplus \eta^{-\gamma_{i}-\delta_{i}} u_{i}^{c} \oplus \eta^{-\gamma_{i}+\delta_{i}} d_{i}^{c} \oplus \eta^{\gamma_{i}} Q_{i} \right)$$
(2.44)

$$\overline{\mathbf{16}}_{j} \to \eta^{\overline{m}_{i}} \overline{\mathbf{16}}_{j}, \tag{2.45}$$

and look for solutions for the discrete charges where different states have different mass matrices. Since explicit examples can only be given by solving extensive modular linear systems, which are computationally prohibitive, a fully working example with three lightfamilies is not provided.

3 $U(1)_X$ breaking scenarios and mechanisms

In this section we are interested in implementing a symmetry breaking mechanism for the extra $U(1)_X$ in which the breaking VEV is stabilised at high values, more or less close to the GUT scale. In order to do so, we will look into the D-flat direction of the potential that breaks the extra $U(1)_X$. It was shown [19, 47] that in the D-flat direction, non-renormalisable operators can provide such scenario. In its simplest inception, the Kolda-Martin mechanism [19] relies on a vector-like pair which lowest order term allowed in the superpotential is non-renormalizable

$$W = \frac{c}{m_{Pl}} (\Phi \bar{\Phi})^2 \tag{3.1}$$

and alongside the soft-term Lagrangian

$$-\mathcal{L}_{\text{soft}} = m_{\Phi}^2 |\Phi|^2 + m_{\bar{\Phi}}^2 |\bar{\Phi}|^2, \qquad (3.2)$$

it is immediate to find that along the D-flat direction the potential has a non-trivial minimum which fixes the VEVs at a high scale

$$\Phi^2 = \sqrt{-\frac{(m_{\Phi}^2 + m_{\bar{\Phi}}^2)m_{Pl}^2}{12c}},$$
(3.3)

where if we take $m \simeq 10^4$ the VEVs are estimated at $\Phi \simeq 10^{11}$ GeV.

There are some caveats to this mechanism as presented above. First, there is significant F-breaking as $\langle F \rangle \simeq \mathcal{O}(10^{15})$ GeV. While this is not a problem if the vector-like family does not share gauge interactions with ordinary matter, in our case non-vanishing F-terms will originate undesirable interactions, cf. section 2.2. We shall therefore focus on F-flat solutions.

Second, the mechanism is not complete in the absence of the full soft-terms Lagrangian, which has to include

$$-\mathcal{L}_{\text{soft}} \supset C \frac{1}{m_{Pl}} \Phi^2 \bar{\Phi}^2 + \text{ h.c..}$$
(3.4)

As we estimate $C \simeq \mathcal{O}(m_{3/2})$ at the GUT scale from the SUGRA [48], at the VEV scale this term is competing with the non-renormalisable terms in the potential arising from the superpotential, and therefore cannot be ignored.

Finally the model presented differs from ours as μ -terms are generically generated by moduli VEVs even if they are disallowed by the discrete symmetry of the compactified space.

In order to proceed, we turn to a more complete version of the mechanism. To do so, we include the μ -term

$$W = \mu \Phi \bar{\Phi} + \frac{c}{m_{Pl}} (\Phi \bar{\Phi})^2 \tag{3.5}$$

and the more complete soft Lagrangian,

$$-\mathcal{L}_{\text{soft}} = m_{\Phi}^2 |\Phi|^2 + m_{\bar{\Phi}}^2 |\bar{\Phi}|^2 - (B\mu \Phi \bar{\Phi} + \text{ h.c.}) + \frac{C}{m_{Pl}} \Phi^2 \bar{\Phi}^2 + \text{ h.c.}$$
(3.6)

Due to the presence of the μ -term, the F-term

$$F_{\Phi} = \mu \bar{\Phi} + \frac{2c}{m_{Pl}} \Phi \bar{\Phi}^2 \tag{3.7}$$

can be set to zero for two different field configurations

$$F_{\Phi} = 0 \Rightarrow \begin{cases} \bar{\Phi} = 0\\ \Phi \bar{\Phi} = -\frac{\mu m_{Pl}}{2c} \end{cases}$$
(3.8)

and the non-trivial VEV can be estimated. Taking $\mu \simeq \mathcal{O}(10^3)$ GeV, this leads to $|\Phi| = 10^{10.5}$ GeV. This looks very similar to the original Kolda-Martin case, with the exception being that the F-term can vanish, and the parametric dependence on the VEV is now on μ instead of a soft-mass. In general there might be a non-SUSY preserving vacuum elsewhere in field space, but we will work under the assumption that the SUSY vacua discovered with this approach are at least stable enough to host phenomenologically viable models.

We wish to assess if we can minimise the potential in this SUSY-preserving field configuration. For that, we need to check if the above field configuration will also extremise the soft-term Lagrangian. To see this we take

$$-\partial_{\Phi}\mathcal{L}_{\text{soft}} = m_{\Phi}^2 \Phi^* - B\mu \bar{\Phi} + \frac{2C}{m_{Pl}} \Phi \bar{\Phi}^2 = 0$$
(3.9)

and, in the limit the VEVs are real, we find a trivial and a non-trivial solutions

$$-\partial_{\Phi}\mathcal{L}_{\text{soft}} = 0 \Rightarrow \begin{cases} \Phi = 0\\ \Phi^2 = -\frac{(m_{\Phi}^2 - B\mu)m_{Pl}}{2C} \end{cases}$$
(3.10)

and the second one seems very similar to the non-trivial configuration derived through the F-term. In fact, both conditions can be met. To see this, we re-parametrise the soft-terms by factoring out their dimensionful dependence on $m_{3/2}$

$$B\mu = m_{3/2}\mu b \tag{3.11}$$

$$C = m_{3/2}\tilde{c} \tag{3.12}$$

$$m_{\Phi} = m_{3/2}a,$$
 (3.13)

where a, b, \tilde{c} are dimensionless, and from SUGRA formulae they are $\mathcal{O}(1)$ at the GUT scale. Of course they will evolve with the scale through RGE evolution, so they need not to be always of the same order. The condition that both the F-flatness and soft-term stabilisation are jointly achieved boils down to be a relation between parameters

$$\frac{\tilde{c}}{c} = \frac{2am_{\Phi} - \mu b}{\mu},\tag{3.14}$$

which is generically valid.

In order for the above non-trivial VEV be a minimum, we need the trivial VEV solution to account for a maximum. This is to say that the mass matrix for the system $(\Phi, \bar{\Phi}^*)$ evaluated at the origin has a negative eigen-value. In our case this accounts for allowing its determinant to be negative

$$(|\mu|^2 + m_{\Phi}^2)(|\mu|^2 + m_{\bar{\Phi}}^2) - B\mu^2 < 0.$$
(3.15)

We notice as well that the above discussion can be immediately extended for the case that the lowest order non-renormalisable term allowed by the discrete symmetry

$$W \supset \frac{c}{m_{Pl}^{2n-3}} (\Phi \bar{\Phi})^n \Rightarrow \Phi \simeq (\mu m_{Pl}^{2n-3})^{\frac{1}{2n-2}}$$
(3.16)

happens for $n \ge 2$, and not only for n = 2. Even so, the presented implementation of the Kolda-Martin mechanism only accounts for a vector-like pair of superfields, while in our case the system breaking the extra $U(1)_X$ is composed of N, N_X , \overline{N}_X states.

Therefore, we want to find similar solutions starting with the superpotential

$$W = \mu_{Xm}^N N \overline{N}_X + \mu_X^N N_X \overline{N}_X + \frac{c_{2,2}}{m_{Pl}} (N \overline{N}_X)^2 + \frac{c_{n,k}}{m_{Pl}^{2n-3}} (N_X \overline{N}_X)^{n-k} (N \overline{N}_X)^k \qquad (3.17)$$

where $n \ge 2$ and k < n. The third term generates a Majorana mass for the matter right-handed conjugated neutrino, N. The full soft-term Lagrangian for this theory is

$$-\mathcal{L}_{\text{soft}} = m_N^2 |N|^2 + m_{N_X}^2 |N_X|^2 + m_{\overline{N}_X}^2 |\overline{N}_X|^2 - (B\mu_{Xm}^N N \overline{N}_X + \text{h.c.}) - (B\mu_X^N N_X \overline{N}_X + \text{h.c.}) + \left(\frac{C_{2,2}}{m_{Pl}} (N \overline{N}_X)^2 + \text{h.c.}\right) + \left(\frac{C_{n,k}}{m_{Pl}^{2n-3}} (N_X \overline{N}_X)^{n-k} (N \overline{N}_X)^k + \text{h.c.}\right)$$
(3.18)

where again $C_{i,j}$ coefficients are $\mathcal{O}(m_{3/2})$ at the GUT scale.

The F-terms now read

$$F_N = \mu_{Xm}^N \overline{N}_X + \frac{2c_{2,2}}{m_{Pl}} N \overline{N}_X^2 + \frac{kc_{n,k}}{m_{Pl}^{2n-3}} N_X^{n-k} N^{k-1} \overline{N}_X^n$$
(3.19)

$$F_{N_X} = \mu_X^N \overline{N}_X + \frac{(n-k)c_{n,k}}{m_{Pl}^{2n-3}} N_X^{n-k-1} N^k \overline{N}_X^n$$
(3.20)

$$F_{\overline{N}_X} = \mu_{Xm}^N N + \mu_X^N N_X + \frac{2c_{2,2}}{m_{Pl}} N^2 \overline{N}_X + \frac{nc_{n,k}}{m_{Pl}^{2n-3}} N_X^{n-k} N^k \overline{N}_X^{n-1}$$
(3.21)

which have a significantly more challenging look than the simplified version presented above. Nonetheless, the same conclusions hold. The above F-terms become more tractable for the k = 0 and k = n - 1 cases. In these cases it is possible to get algebraic expressions for the VEVs estimates. For the k = 0, the F-flatness conditions alone give us

$$N\overline{N}_X = -\frac{\mu_{Xm}^N m_{Pl}}{2c_{2,2}} \tag{3.22}$$

$$N_X \overline{N}_X = \left(-\frac{\mu_X^N m_{Pl}^{2n-3}}{nc_{n,o}}\right)^{\frac{1}{n-1}}$$
(3.23)

while for k = n - 1, analogous expressions can be obtained

$$|N\overline{N}_X| \simeq (\mu_{Xm}^N m_{Pl}^{2n-3})^{\frac{1}{n-1}}$$
(3.24)

$$|N_X \overline{N}_X| \simeq ((\mu_X^N)^{3-n} m_{Pl}^{3n-5})^{\frac{1}{n-1}}$$
(3.25)

where the approximations mean we dropped $\mathcal{O}(1)$ parameters and took all μ -terms to be of the same order, which is expected.

In both cases, the ratio between the N_X and N VEV is follows the same dependency on n

$$\left|\frac{N_X}{N}\right| \simeq \left(\frac{m_{Pl}}{\mu}\right)^{\frac{n-2}{n-1}} \simeq \begin{cases} 1 & n=2\\ 10^{7.5} & n=3\\ 10^{10} & n=4 \end{cases}$$
(3.26)

where we μ is an $\mathcal{O}(\mu_X^N, \mu_{Xm}^N)$ parameter. This result shows that there is a hierarchy between N_X and N VEVs, which is very desirable as N VEVs can generate large B-RPV couplings, cf. section 2.2.

Just like before, we use the D-flat direction

$$\left|\frac{\overline{N}_X}{N_X}\right|^2 = \left|\frac{N}{N_X}\right|^2 + 1,\tag{3.27}$$

which sets the magnitude of the three VEVs. The results for k = 0 and k = n - 1 can be immediately estimated algebraically, in contrast to the other cases. The full result of SUSY preserving configurations can be seen in table 1. It is important to note that for n = 4, the only viable scenario is for k = 0, while for n = 3 the k = 2 is not viable as there are super-GUT VEVs. In the end we are only interested in the sensible cases, where the VEVs are below the GUT scale and therefore the mechanism is self-consistent.

The SUSY configurations above are expected stabilise the soft-terms Lagrangian just before. The stabilisation conditions are

$$m_{\Phi_1}^2 \Phi_1^* - B\mu_1 \bar{\Phi} + \frac{2C_{2,2}}{m_{Pl}} \Phi_1 \bar{\Phi}^2 = 0$$
$$m_{\Phi_2}^2 \Phi_2^* - B\mu_2 \bar{\Phi} + \frac{nC_{n,0}}{m_{Pl}^{2n-3}} \Phi_2^{n-1} \bar{\Phi}^n = 0$$
$$m_{\bar{\Phi}}^2 \bar{\Phi}^* - B\mu_1 \Phi_1 - B\mu_2 \Phi_2 + \frac{2C_{2,2}}{m_{Pl}} \Phi_1^2 \bar{\Phi} + \frac{nC_{n,0}}{m_{Pl}^{2n-3}} \Phi_2^n \bar{\Phi}^{n-1} = 0$$

n	k	$N \; ({\rm GeV})$	N_X (GeV)	\overline{N}_X (GeV)
2	0	$10^{10.5}$	$10^{10.5}$	$10^{10.5}$
	1	$10^{10.5}$	$10^{10.5}$	$10^{10.5}$
3	0	$10^{6.5}$	$10^{14.25}$	$10^{14.25}$
	1	$10^{10.2}$	$10^{15.5}$	$10^{15.5}$
	2	$10^{10.5}$	10^{18}	10^{18}
4	0	$10^{5.5}$	$10^{15.5}$	$10^{15.5}$
	1	$10^{10.1}$	$10^{16.5}$	$10^{16.5}$
	2	$10^{10.3}$	10^{18}	10^{18}
	3	$10^{10.5}$	$10^{20.5}$	$10^{20.5}$

Table 1. Estimate of the magnitude of the VEVs in SUSY vacua for different implementations of the modified Kolda-Martin mechanism. In all cases the scalar component of the (CP conjugated) right-handed neutrino field N develops a VEV, breaking R-parity, in addition to the N_X and \overline{N}_X VEVs.

and re-parametrising the dimensionful soft-terms just as before, the above conditions will resemble the F-flatness conditions in form and so they'll be jointly respected taken the parameters of the theory respect relations between them.

As before, the condition that the above extrema are minima is that the potential has a runaway direction around the origin. This is the same to say that, when close to the origin the potential takes the form

$$V \simeq \mathbf{N}^* \cdot M_N \cdot \mathbf{N} \tag{3.28}$$

with $\mathbf{N} = (N, N_X, \overline{N}_X^*)$, such that M_N at least one negative eigenvalue to account for a run-away behaviour at the trivial extremum. Boundness of the potential in the D-flat direction is achieved by noticing that — for each field direction — at least a quadratic term from the non-renormalisable interactions becomes the leading contribution, while keeping a run-away behaviour at the origin.

4 Neutrino-neutralino mass matrix

The different breaking scenarios discussed in the previous section rely on different superpotential terms, which are either present or suppressed depending the discrete symmetry of the compactified G_2 space. Furthermore, the generic presence of a matter field VEV, $\langle N \rangle$, will generate B-RPV terms, as seen in section 2.2. In turn, these provide a new source of neutrino masses which has to be taken into account. To be more precise we enumerate all the interactions that contribute to neutrino masses. First, we let the matter neutrino to have a Yukawa coupling at tree-level, of the form

$$W_{\text{tree}} \supset y_{\nu} N L H_u . \tag{4.1}$$

Next we have to consider the non-renormalizable terms that employ the KM mechanism for each scenario. Alongside this, we also keep a term that can generate a Majorana mass for the matter right-handed conjugated neutrino, N. On top of these, we include a set of non-renormalizable terms involving the Higgses or *L*-type fields, in first order of $1/m_{Pl}$. The non-renormalizable terms that will affect the neutral fermion mass matrix are then

$$W_{\text{non.ren.}} \supset \frac{c_{2,2}}{m_{Pl}} (NN) \left(\overline{N}_X \overline{N}_X \right) + \frac{c_{n,k}}{m_{Pl}^{2n-3}} \left(N_X \overline{N}_X \right)^{n-k} \left(N \overline{N}_X \right)^k + \frac{1}{m_{Pl}} \left(b_1 H_d H_u L \overline{L}_X + b_2 L L \overline{L}_X \overline{L}_X + b_3 H_d H_u L_X \overline{L}_X + b_4 L L_X \overline{L}_X \overline{L}_X \right) + b_5 L_X L_X \overline{L}_X \overline{L}_X + b_6 H_d H_u N \overline{N}_X + b_7 L \overline{L}_X N \overline{N}_X + b_8 L_X \overline{L}_X N \overline{N}_X + b_9 H_d H_u N_X \overline{N}_X + b_{10} L \overline{L}_X N_X \overline{N}_X + b_{11} L_X \overline{L}_X N_X \overline{N}_X \right).$$
(4.2)

The terms that are disallowed by discrete symmetry are generically re-generated as the moduli acquire VEVs. As such, the following Kähler potential terms will have an important contribution for neutrino masses

$$K \supset \frac{s}{m_{Pl}}\overline{L}_X L_X + \frac{s}{m_{Pl}}\overline{L}_X L + \frac{s}{m_{Pl}}\overline{N}_X N_X + \frac{s}{m_{Pl}}\overline{N}_X N + \frac{s}{m_{Pl}}\overline{H}_u H_d \tag{4.3}$$

$$+\frac{s}{m_{Pl}^{2}}N_{X}L_{X}H_{u}+\frac{s}{m_{Pl}^{2}}NLH_{u}+\frac{s}{m_{Pl}^{2}}N_{X}LH_{u}+\frac{s}{m_{Pl}^{2}}NL_{X}H_{u}+\frac{s}{m_{Pl}^{2}}\overline{N}_{X}\overline{L}_{X}H_{d},$$

where s denotes a generic modulus fields that counterbalances the discrete charge. This modulus field needs not to be the same for each coupling. As the moduli acquire VEVs as they are stabilised, the above terms will generate the effective superpotential

$$W_{\text{eff}} \supset \mu_{XX}^L \overline{L}_X L_X + \mu_{Xm}^L \overline{L}_X L + \mu_{XX}^N \overline{N}_X N_X + \mu_{Xm}^N \overline{N}_X N + \mu H_u H_d$$

$$+ \lambda_{\overline{XX}} H_d \overline{L}_X \overline{N}_X + \lambda_{\nu} H_u L N + \lambda_{mX} H_u L N_X + \lambda_{Xm} H_u L_X N + \lambda_{XX} H_u L_X N_X$$

$$(4.4)$$

where the parameters can be estimated to lie inside the orders of magnitude

$$\mu \simeq m_{3/2} \frac{s}{m_{Pl}} \simeq \mathcal{O}(10^3) \text{ GeV}$$
(4.5)

$$\lambda \simeq m_{3/2} \frac{s}{m_{Pl}^2} \simeq \mathcal{O}(10^{-15}).$$
 (4.6)

Therefore, the total superpotential, which includes all the interactions that contribute to the neutral fermion mass matrix is give by

$$W_{\text{total}} \supset W_{\text{tree}} + W_{\text{non.ren.}} + W_{\text{eff}}.$$
 (4.7)

In our framework we have VEVs of the *N*-type fields that can be significantly large, depending on which implementation of the KM mechanism we assume. As such, B-RPV couplings, mixing Higgses superfields with *L*-type superfields, appear in the superpotential as

$$\kappa_m H_u L + \kappa_X H_u L_X + \kappa_{\overline{X}} H_d \overline{L}_X \tag{4.8}$$

where the κ -parameters read

$$\kappa_m \simeq (y_\nu + \lambda_\nu) \langle N \rangle + \lambda_{mX} \langle N_X \rangle \tag{4.9}$$

$$\kappa_X \simeq \lambda_{Xm} \langle N \rangle + \lambda_{XX} \langle N_X \rangle \tag{4.10}$$

$$\kappa_{\overline{X}} \simeq \lambda_{\overline{X}\overline{X}} \langle \overline{N}_X \rangle \tag{4.11}$$

where we are dropping the *F*-terms contribution as the solutions for our KM mechanism presented in section 3 are aligned in the *D* and *F* directions. We also note that we are assuming no tree-level Yukawa couplings involving extra vector-like N_X , \overline{N}_X for the KM scenarios.

Furthermore, the presence of B-RPV induces a sub-EWS VEV on the scalar components of the ν -type fields. In our case, below the EWS, we expect all ν -type scalars to acquire a non-vanishing VEV, generating a mixing between N-type fermions and Higgsinos through

$$\epsilon_m H_u^0 N + \epsilon_X H_u^0 N_X + \epsilon_{\overline{X}} H_d^0 N_{\overline{X}} \tag{4.12}$$

where the coefficients read

$$\epsilon_m \simeq (y_\nu + \lambda_\nu) \langle \nu \rangle + \lambda_{mX} \langle \nu_X \rangle \tag{4.13}$$

$$\epsilon_X \simeq \lambda_{Xm} \langle \nu \rangle + \lambda_{XX} \langle \nu_X \rangle \tag{4.14}$$

$$\epsilon_{\overline{X}} \simeq \lambda_{\overline{XX}} \langle \nu_{\overline{X}} \rangle \tag{4.15}$$

and, as expected, they have the same generic form as the κ -parameters since both set of parameters arise from trilinear, Yukawa, couplings in the superpotential.

Finally, as in the MSSM, the presence of VEVs will mix some fermions with gauginos through kinetic terms, namely the Higgsinos with \tilde{B}_1 , \tilde{W}^0 due to the Higgses VEVs. In our case we also have *N*-type and ν -type scalar VEVs, which will mix gauginos with matter fermions through kinetic terms. We have, for the SU(2) states,

$$g'\widetilde{B}\langle\widetilde{\nu}_i\rangle\nu_i, \quad g\widetilde{W}^0\langle\widetilde{\nu}_i\rangle\nu_i, \quad g''\widetilde{B}_X\langle\widetilde{\nu}_i\rangle\nu_i \tag{4.16}$$

while for the N-states, which are singlets under the SM gauge group, the mixing with the gaugino of the extra $U(1)_X$ gauge group

$$g''\widetilde{B}_X\langle\widetilde{N}_i\rangle N_i$$
 (4.17)

where, in both expressions, we used the shorthand $g' = \sqrt{\frac{5}{3}}g_1$ and $g'' = \frac{1}{2\sqrt{10}}g_X$. With all the above considerations, we can now construct the 11×11 mass matrix for

With all the above considerations, we can now construct the 11×11 mass matrix for neutral fermions of our model. We define this matrix in the basis

$$\psi = (\widetilde{B}, \widetilde{W}^0, \widetilde{B}_X, \widetilde{H}^0_d, \widetilde{H}^0_u, \nu, \nu_X, \nu_{\overline{X}}, N, N_X, N_{\overline{X}}),$$
(4.18)

and it has the schematic form

$$\mathbf{M}_{\chi-\nu} = \begin{pmatrix} \mathbf{M}_{\chi^0}^{5\times 5} & \mathbf{M}_{\chi\nu}^{5\times 6} \\ (\mathbf{M}_{\chi\nu}^{5\times 6})^T & \mathbf{M}_{\nu}^{6\times 6} \end{pmatrix}.$$
 (4.19)

The usually called neutralino part of the matrix includes only mass terms involving gauginos and Higgsinos, and its form is very similar to the MSSM, except we have an extended gauge group with one more $U(1)_X$ factor. It reads

$$\mathbf{M}_{\chi^0}^{5\times5} = \begin{pmatrix} M_1 & 0 & 0 & -\frac{1}{\sqrt{2}}g'v_d & \frac{1}{\sqrt{2}}g'v_u\\ 0 & M_2 & 0 & \frac{1}{\sqrt{2}}gv_d & -\frac{1}{\sqrt{2}}gv_u\\ 0 & 0 & M_X & -2\sqrt{2}g''v_d & 2\sqrt{2}g''v_u\\ -\frac{1}{\sqrt{2}}g'v_d & \frac{1}{\sqrt{2}}gv_d & -2\sqrt{2}g''v_d & 0 & -\mu\\ \frac{1}{\sqrt{2}}g'v_u & -\frac{1}{\sqrt{2}}gv_u & 2\sqrt{2}g''v_u & -\mu & 0 \end{pmatrix}$$
(4.20)

The next block is the one involving terms mixing the usual neutralino states with matter states. As such, they include B-RPV masses that mix matter with higgses. The matrix reads

$$\mathbf{M}_{\chi\nu}^{5\times6} = \begin{pmatrix} -\frac{1}{\sqrt{2}}g\nu & -\frac{1}{\sqrt{2}}g'\nu_X & \frac{1}{\sqrt{2}}g'\overline{\nu}_X & 0 & 0 & 0\\ \frac{1}{\sqrt{2}}g\nu & \frac{1}{\sqrt{2}}g\nu_X & -\frac{1}{\sqrt{2}}g\overline{\nu}_X & 0 & 0 & 0\\ 3\sqrt{2}g''\nu & 3\sqrt{2}g''\nu_X & -3\sqrt{2}g''\overline{\nu}_X & -5\sqrt{2}g''N & -5\sqrt{2}g''N_X & 5\sqrt{2}g''\overline{N}_X\\ 0 & 0 & \kappa_{\overline{X}} & 0 & 0 & \epsilon_{\overline{X}}\\ \kappa_m & \kappa_X & 0 & \epsilon_m & \epsilon_X & 0 \end{pmatrix}$$
(4.21)

where, in order to de-clutter notation, we are taking the fields names as to represent the VEVs. We notice that the B-RPV couplings κ and ϵ are superpotential terms, while the top three rows is generated by kinetic terms only.

The lower-right 6×6 block is purely from the superpotential, and includes only the masses involving ν -type and/or N-type fermions. To obtain the mass, one performs the usual SUSY rule for fermionic masses

$$\left(\mathbf{M}_{\nu}^{6\times 6}\right)_{ij} = -\frac{1}{2} \frac{\partial^2}{\partial \psi_i \partial \psi_j} W_{\text{total}}$$

$$(4.22)$$

where $i, j = \{\nu, \nu_X, \overline{\nu}_X, N, N_X, \overline{N}_X\}.$

This 6×6 matrix has three main blocks: the $\nu\nu$ block, νN block, and NN block. Schematically they are arranged, in our basis, as

$$\mathbf{M}_{\nu}^{6\times6} = -\frac{1}{2} \left(\frac{M_{\nu\nu}}{M_{\nu N}^{T}} \frac{M_{\nu N}}{M_{NN}} \right)$$
(4.23)

The actual form of the matrix is obtained using the full superpotential in eq. (4.7). Doing so, one gets the following sub-blocks. First we have the $\nu\nu$ block that has mixing between $\overline{\nu}_X$ and ν , ν_X . In the sub-basis $(\nu, \nu_X, \overline{\nu}_X)$ this reads

$$M_{\nu\nu} = \begin{pmatrix} 0 \ 0 \ \frac{b_7 \overline{N}_X N}{m_{Pl}} + \frac{b_{10} \overline{N}_X N_X}{m_{Pl}} + \mu_{Xm}^L \\ 0 \ \frac{b_8 \overline{N}_X N}{m_{Pl}} + \frac{b_{11} \overline{N}_X N_X}{m_{Pl}} + \mu_X^L \\ 0 \end{pmatrix}$$
(4.24)

where we dropped the terms ν^2/m_{Pl} , $v_{u/d}^2/m_{Pl}$ as they are irrelevant and to de-clutter, and since this block is symmetric we omit the lower left triangular part. But notice that the terms with coefficients b_7 , b_8 , b_{10} , b_{11} can play an important role as they can generate heavy Dirac masses, depending on the KM mechanism.

Next we have the νN block, where one can find the neutrino Dirac masses generated by the Higgses VEV at the EWS. Taking the rows to be along the basis $(\nu, \nu_X, \overline{\nu}_X)$, while the columns along (N, N_X, \overline{N}_X) , this block reads

$$M_{\nu N} = \begin{pmatrix} v_{u} y_{\nu} + \frac{b_{7} \overline{N}_{X} \overline{\nu}_{X}}{m_{Pl}} & \frac{b_{10} \overline{N}_{X} \overline{\nu}_{X}}{m_{Pl}} & \frac{b_{7} N \overline{\nu}_{X}}{m_{Pl}} + \frac{b_{10} N_{X} \overline{\nu}_{X}}{m_{Pl}} \\ \frac{b_{8} \overline{N}_{X} \overline{\nu}_{X}}{m_{Pl}} & \frac{b_{11} \overline{N}_{X} \overline{\nu}_{X}}{m_{Pl}} & \frac{b_{8} N \overline{\nu}_{X}}{m_{Pl}} + \frac{b_{11} N_{X} \overline{\nu}_{X}}{m_{Pl}} \\ \frac{b_{7} \overline{N}_{X} \nu}{m_{Pl}} + \frac{b_{8} \overline{N}_{X} \nu_{X}}{m_{Pl}} & \frac{b_{10} \overline{N}_{X} \nu}{m_{Pl}} + \frac{b_{11} \overline{N}_{X} \nu_{X}}{m_{Pl}} & \frac{b_{7} N \nu}{m_{Pl}} + \frac{b_{10} N_{X} \overline{\nu}_{X}}{m_{Pl}} \\ \end{pmatrix}$$

$$(4.25)$$

where we dropped the sub-leading terms $v_{u/d}\lambda \simeq \mathcal{O}(10^{-12})$ GeV.

Finally we have the NN block, that involves Dirac and Majorana masses generated through the first two terms in equation (4.2). Ignoring the terms generated by Higgses and sneutrino VEVs, in the sub-basis (N, N_X, \overline{N}_X) this block reads

$$M_{NN} =$$

$$\begin{pmatrix} \frac{c_{n,k}(k-1)k}{m_{Pl}^{2n-3}} \overline{N}_{X}^{n} N_{X}^{n-k} N^{k-2} + \frac{2c_{2,2}}{m_{Pl}} \overline{N}_{X}^{2} & \frac{c_{n,k}k(n-k)}{m_{Pl}^{2n-3}} \overline{N}_{X}^{n} N^{k-1} N_{X}^{-k+n-1} & \frac{c_{n,k}kn}{m_{Pl}^{2n-3}} \overline{N}_{X}^{n-1} N_{X}^{n-k} N^{k-1} + \mu_{Xm}^{N} + \frac{4c_{2,2}}{m_{Pl}} \overline{N}_{X} N \\ & \frac{c_{n,k}(-k+n-1)(n-k)}{m_{Pl}^{2n-3}} \overline{N}_{X}^{n} N^{k} N_{X}^{n-k-2} & \frac{c_{n,k}n(n-k)}{m_{Pl}^{2n-3}} \overline{N}_{X}^{n-1} N_{X}^{-k+n-1} N^{k} + \mu_{Xx}^{N} \\ & \frac{c_{n,k}(-k+n-1)(n-k)}{m_{Pl}^{2n-3}} \overline{N}_{X}^{n-k} N^{k-1} + \frac{c_{n,k}n(n-k)}{m_{Pl}^{2n-3}} \overline{N}_{X}^{n-1} N_{X}^{-k+n-1} N^{k} + \mu_{Xx}^{N} \\ & \frac{c_{n,k}(n-1)n}{m_{Pl}^{2n-3}} \overline{N}_{X}^{n-2} N_{X}^{n-k} N^{k} + \frac{2c_{2,2}}{m_{Pl}} N^{2} \end{pmatrix}$$

where the orders of magnitude of each entry will largely depend on which KM scenario is being considered. The matrix is symmetric so only the upper diagonal entries are displayed.

4.1 The mass matrix hierarchies

Following the description of the mass matrix above, we will now try to infer the hierarchies between the entries of the matrix. First we notice that, regardless of the case (i.e. the allowed Kolda-Martin operators), the biggest entry in the mass matrix is always in the Gaugino-N mixing block.⁴ This result is understandable as we expect the breaking of the extra U(1)_X to transform a chiral superfield and a massless vector superfield into a single massive vector superfield. The degrees of freedom add up correctly, and would mean that below the U(1)_X breaking scale we can take \tilde{B}_X and the linear combination of N-states that break the U(1)_X to be integrated out jointly. The linear combination that breaks the extra U(1)_X depends on the exact values of the VEVs, but we can highlight some characteristics and how the mass-matrix will look like after this is integrated out.

In order to single out the correct liner combination that breaks the extra $U(1)_X$, one can perform a rotation in the last three states — N, N_X , \overline{N}_X — in order to retain only one mixing mass between these states and the \tilde{B}_X . In order to do so, in the limit the mass

⁴The caveat to this statement is if we allow for an order 1 Neutrino Yukawa, in that case the κ entry originated from $y_{\nu}\langle N \rangle LH_u$, will have the same order of magnitude. But since the B-RPV coupling above does not involve \tilde{B}_X , N, N_X , or \overline{N}_X , the magnitude of this coupling does not change the following discussion. We will return to B-RPV couplings further below.

matrix is real, the rotation is

$$U = \begin{pmatrix} 1 & 0 \\ 0 & 1 & \cdots \\ \vdots & \ddots & \vdots \\ & \cos(\theta) - \sin(\theta)\cos(\phi) & \sin(\theta)\sin(\phi) \\ & \sin(\theta) & \cos(\theta)\cos(\phi) & -\cos(\theta)\sin(\phi) \\ & 0 & \sin(\phi) & \cos(\phi) \end{pmatrix}$$

where the angles are determined by the strength of the mixing mass parameters. For instance, in the n = 2, k = 0 Kolda-Martin mechanism presented before, the VEVs of the scalar components of N, N_X , \overline{N}_X are all of same order. In such case, taking $\theta \simeq 3\pi/4$ and $\phi \simeq \arctan \sqrt{2}$ will leave only one state mixing with \tilde{B}_X . For the other Kolda-Martin implementations, the N_X , \overline{N}_X VEVs are much larger than N VEV and so we can take $\theta \simeq 0$ with $\phi \simeq 3\pi/4$ to accomplish the same.

The rotation above affects only the last three columns and rows. Since the matrix is unitary (orthogonal in the case the masses are real), the entries of last three columns of a given row will be mixed with at-most order 1 coefficients, and whilst there might be cancellations there will be no order of magnitude enhancements. Once the rotation is performed one can then integrate out \tilde{B}_X jointly with its Dirac partner. This in turn will affect all the remainder of the matrix. For example, the entry i, j will receive a contribution from integrating out a Dirac mass at position a, b of order

$$-\frac{M_{i3}M_{bj}}{M_{3b}}$$

with some order one coefficients from the rotation. In this case we are setting one of the indices to 3 as this is the position of \tilde{B}_X in our basis. The remaining index, b, refers to the position of the linear combination that breaks the extra $U(1)_X$. If, for example, the breaking linear combination that breaks the extra $U(1)_X$ is mostly composed of N_X , \overline{N}_X states, the main contribution to the ν Majorana mass is given by

$$\frac{b_{10}}{m_{Pl}} \langle \nu \rangle \langle \overline{\nu}_X \rangle \ll 10^{-10} \text{ GeV}$$

even if we let the respective coupling on, i.e. $b_{10} \simeq \mathcal{O}(1)$. Therefore, after the above rotation and integrating out, the mass matrix remains schematically the same, but with the absence of \tilde{B}_X and a linear combination composed of N, N_X , \overline{N}_X .

After integrating out the Dirac fermion originated by the breaking, one can see that the Majorana and Dirac masses — generated at the $U(1)_X$ breaking scale — involving only the surviving terms of the N, N_X, \overline{N}_X system are the leading entries of the mass matrix. These are present in the bottom-right-most 2×2 block. These states will then be responsible for a type of see-saw mechanism involving the lighter SU(2) doublet states $\nu, \nu_X, \overline{\nu}_X$, with EW scale Dirac mass terms. In order to make sense of this see-saw mechanism, the ν -states need to be protected from too much mixing with the remaining gauginos and higgsinos,

such that the lightest mass eigenstate is dominantly composed of ν . Actually the mixing between the ν -type states with gauginos is negligible since it is generated by ν -type VEVs and are therefore sub-EWS. But the mixing with Higgsinos is parametrically dependent on N-type VEVs through B-RPV terms, the so called κ mass parameters.

The κ parameters defined in equations (4.9), (4.10) and (4.10) can have other potentially undesirable consequences as they can spoil Higgs physics. Take for example the matter B-RPV interaction, with κ_m significantly larger than any other mass involving H_u . If were to happens, then L and H_u superfields would pair up to produce a heavy vectorlike pair. Then H_u would be much heavier than the EWS physics and would spoil Higgs physics, where H_u and H_d are identified as a vector-like pair. In order to preserve viable Higgs physics, we need all κ -parameters to be much smaller than the remaining masses appearing in the Higgs potential.

Finally, there is risk that ν , ν_X , $\overline{\nu}_X$ states will mix with each other too much. To see this consider the 3 × 3 sub-block of the matrix as shown in eq. (4.24). If all b_i couplings are suppressed, this matrix will maximally mix ν and ν_X through the μ -terms. But it is important to note that while most of the b_i interactions will be generated by Higgses and ν -type VEVs (making them naturally sub-leading even if they are allowed by discrete symmetry) there are two terms that can have important contributions

$$\frac{b_{10}}{m_{Pl}}N_X\overline{N}_X\nu\overline{\nu}_X, \ \frac{b_{11}}{m_{Pl}}N_X\overline{N}_X\nu_X\overline{\nu}_X, \tag{4.27}$$

which for the KM cases can generate Dirac masses much greater than μ -terms if the respective b_i coefficients are unsuppressed. This can then provide a natural mechanism to split ν from ν_X , $\overline{\nu}_X$, if the coupling b_{10} is forbidden while b_{11} is allowed. In this case, we define

$$\mu_{11} = \frac{b_{11}}{m_{Pl}} N_X \overline{N}_X \tag{4.28}$$

and the leading entries for eq. (4.24) will take the form

$$\begin{pmatrix} 0 & 0 & \mu_{Xm}^L \\ 0 & 0 & \mu_{11} \\ \mu_{Xm}^L & \mu_{11} & 0 \end{pmatrix},$$
(4.29)

which will then lead to ν_X , $\overline{\nu}_X$ to pair up and decouple from ν .

5 Numerical results

As the full mass matrix presents an intricate structure of relations and hierarchies between different states, it is ultimately impossible to obtain a simple and revealing analytic expression that describes how one should obtain good neutrino physics. Instead, we perform a numerical scan over space, ensuring that the above constraints are satisfied. In so doing, we divided the analysis into different realisations of the Kolda-Martin mechanism, parametrised by different values of (n, k), corresponding to the scenarios in table 1. In all the cases, we considered a point of the parameter space to be good if the mass of the lightest eigenstate of the mass matrix, identified as a physical neutrino, has a mass in the range

$$[50, 100] \text{ meV},$$
 (5.1)

and in addition that the corresponding eigenstate is mostly composed of the left-handed doublet component ν (i.e. the state arising from $(\nu \ e)^T$). In order to do so, we compute the decomposition of the eigenstate in the original basis

$$|\nu_{\text{light}}\rangle = \alpha |\nu\rangle + \dots \tag{5.2}$$

and impose α to be the largest of the coefficients. As discussed in the previous section, the prevalence of ν as the largest component of ν_{light} will depend greatly on the parameters of the mass matrix that mix different states, i.e. Dirac masses. For definiteness, we shall also require that the second lightest mass eigenstate (essentially the lightest non-neutrino-like neutralino) to be at least 100 GeV.

For each example, we only allow the particular desired Kolda-Martin operator while preventing all tree-level Yukawas involving states of the extra vector-like family. Furthermore, unless otherwise stated we assume that all quadratic terms in eq. (4.2) involving large VEVs are turned off. As expected within the M Theory framework, the disallowed tree-level couplings are regenerated through moduli VEVs, and so the respective coupling strength was set to be of $\mathcal{O}(10^{-15})$. Along the same line, the μ -terms generated by moduli VEVs were set to $\mathcal{O}(1)$ TeV.

Below we will show our findings for the only promising cases, which are (n, k) = (2, 0), (2, 1), (3, 0). The other (n, k) assignments either returned to little points or no viable correlation to enhance α . This happens as for the (3, 1), (4, 0) cases, since $N_X, \overline{N}_X \simeq 10^{15.5}$ GeV, the B-RPV coupling is generically greater than 1 GeV. As we will see below, the only viable regions of the parameter space coincide with a naturally suppressed B-RPV parameter.

5.1 ν component of the lightest state

From the discussion above, we expect the value of α to be correlated with some parameters of the theory. Namely, we expect α to be enhanced if b_{11} is not suppressed and if the B-RPV coupling κ_m is much smaller than any other mass involving Higgsinos. Since any disallowed tree-level coupling can be regenerated through moduli VEVs with a $\lambda \simeq 10^{-15}$ suppression, we started our numerical study by looking at the behaviour of α as we let b_{11} vary in the range

$$b_{11} \in [10^{-15}, 1], \tag{5.3}$$

which, in conjugation with a non-vanishing N_X , \overline{N}_X VEVs will lead to non-vanishing μ_{11} as defined in eq. (4.28).

In order to assess the strength of the B-RPV term, κ_m , allowed in the regions of the parameter space that return good neutrinos, we also registered the value of κ_m at each point which returned the mass inside the bounds stated.



Figure 1. Scatter plots showing the amplitude α of the left-handed doublet state ν in the lightest mass eigenstate ν_{light} as b_{11} varies for the (2,0) and (2,1) cases. The points are fairly evenly distributed with a slight clustering near the desired value of $\alpha \approx 1$ for $b_{11} \approx 1$.

(2,0) and (2,1) cases. For these two Kolda-Martin implementation cases, the three (N, N_X, \overline{N}_X) VEVs are all of order $\mathcal{O}(10^{10.5})$ GeV. As such, we allowed these VEVs to take values around

$$N, N_X, \overline{N}_X \in [10^{9.5}, 10^{11.5}] \text{ GeV}$$
(5.4)

to cover the range of expected values. Since with these values the mass matrix is very similar for both (2,0) and (2,1) cases, we present them together.

As a consequence of the values of the VEVs above, the μ_{11} Dirac mass between ν_X , $\overline{\nu}_X$, defined in eq. (4.28), will take values spanning

$$\mu_{11} = b_{11} \frac{N_X \overline{N}_X}{m_{Pl}} = b_{11} [10, 10^5] \text{ GeV}$$
(5.5)

which means that, only for non-suppressed b_{11} we expect

$$\mu_{11} > \mu_{Xm}^L \tag{5.6}$$

as required to split ν from ν_X , as discussed in section 4.1.

The above considerations indicate us that the mechanism to split ν from ν_X will only work for large values of b_{11} . This can be seen in figures 1a and 1b, where a slight agglomeration of points around $(\alpha, b_{11}) \simeq (1, 1)$ can be identified.

On the other hand, we find that the κ_m parameter is mostly bounded to be smaller than 1 GeV, as is shown in figures 2a and 2b. Although such small values of κ_m are welcome, the fact that there is no clear preference for $\kappa_m \gtrsim 10^{-2}$ GeV suggests this class of models is challenged by BBN constraints, cf. eq. (2.33).

(3,0) case. For the (3,0) Kolda-Martin realisation, we found much promising results. Since the N_X , \overline{N}_X VEVs are expected to be around $\mathcal{O}(10^{14.25})$ GeV, if we allow them to be in the range

$$N_X, \ \overline{N}_X \in [10^{13.25}, 10^{15.25}] \text{ GeV}$$
 (5.7)



Figure 2. Scatter plots showing the amplitude α of the left-handed doublet state ν in the lightest mass eigenstate ν_{light} as κ_m varies for the (2,0) and (2,1) cases. The points are fairly evenly distributed with a slight clustering near the desired value of $\alpha \approx 1$. The horizontal dashed line represents the bound on the LSP lifetime, cf. eq. (2.33).

we find

$$\mu_{11} \in b_{11}[10^{8.25}, 10^{12.25}] \text{ GeV}$$
(5.8)

which implies that it is natural to achieve

$$\mu_{11} \gg \mu_{Xm}^L \tag{5.9}$$

and consequently ν will decouple easily from the other ν -type states.

The above expectations are confirmed by the numerical results, and the lightest state will be mostly composed of ν even for values of b_{11} below $\mathcal{O}(1)$. This behaviour can be seen in figure 3a.

Interestingly, in the (α, κ_m) plane, shown in figure 3b we can see again that the mass matrix prefers $\kappa_m < 1 \text{ GeV}$ in order to reproduce a mostly- ν lightest state. This is a nice result which ensures that whenever we have good physical neutrinos, we also find sufficiently suppressed B-RPV. Furthermore, all the good points also suggest $\kappa_m \gtrsim 10^{-2} \text{ GeV}$, satisfying the requirement for successful BBN physics, cf. eq. (2.33).

5.2 Matter neutrino Yukawas and B-RPV couplings

From the above analysis we learned that for the (2,0), (2,1) and (3,0) cases we expect a non-suppressed b_{11} to enhance the component of ν in the lightest state. As such, we will now consider this coupling to be of order 1 and re-run the analysis for these cases, with the goal being to assess what typical values κ_m and y_{ν} should take for a successful implementation of the proposed Kolda-Martin mechanism.

(2,0) and (2,1) cases. In figures 4a and 4b we see that the preferred points are those with $y_{\nu} \leq 10^{-10}$. This suggests that for theses cases, the see-saw mechanism does not take a great role in explaining the light neutrino masses.



Figure 3. Scatter plots showing the amplitude α of the left-handed doublet state ν in the lightest mass eigenstate ν_{light} as κ_m varies for the (3,0) case. The points are fairly evenly distributed except for a significant clustering near the desired value of $\alpha \approx 1$ for larger values of b_{11} . The horizontal dashed line represents the bound on the LSP lifetime, cf. eq. (2.33). The right panel shows that nearly all the points satisfy $\kappa_m \gtrsim 10^{-2} \text{ GeV}$.



Figure 4. Histograms for the values of y_{ν} for the (2,0) and (2,1) cases with unsuppressed b_{11} .

In figures 5a and 5b we see that for these cases, the B-RPV parameter κ_m is naturally very small. This result is easy to understand, considering the main contribution to κ_m to be

$$\kappa_m \simeq y_\nu v_m,$$

and given the range of values that we are allowing the VEVs to take, κ_m is expected to be small. Unfortunately, all points returning good neutrino physics also return $\kappa_m > 10^{-2}$ GeV, which means that these classes of models spoil BBN, cf. (2.33). Although not shown here one can also find that κ_X , $\kappa_{\overline{X}}$ parameters, which mix L_X , \overline{L}_X with H_u , H_d respectively, are also constrained to be smaller than 1 GeV.

(3,0) case. For this realisation of the Kolda-Martin mechanism, the results are slightly different but in line with our expectations. In figure 6a we can see that the matter Yukawa



Figure 5. Histograms for the values of y_{ν} and κ_m for the (2, 1) case with unsuppressed b_{11} . The vertical dashed line represents the bound on the LSP lifetime, cf. eq. (2.33).



Figure 6. Histograms for the values of y_{ν} and κ_m for the (3,0) case with unsuppressed b_{11} .

coupling is allowed to take values larger than in the previous case. This indicates that the see-saw mechanism is having an effect on reducing the contribution of the matter neutrino Dirac mass to the lightest eigenstate.

In figure 6b we see that κ_m is bound to be smaller than 1 GeV. The fact that κ_m takes larger values for (3,0) case than for the n = 2 cases is easily understandable. The main contributions to κ_m are

$$\kappa_m \simeq y_\nu N + \lambda N_X \tag{5.10}$$

where the VEVs are expected as in table 1. These contributions are in general greater than those in n = 2 cases, but they are still bounded to be smaller than 1 GeV. This is fortunate, as $\kappa_m \gtrsim 10^{-2}$ GeV and hence this class of models retain the successful predictions of BBN, cf. (2.33). As before, although not shown here also finds that κ_X , $\kappa_{\overline{X}}$ parameter are also constrained to be smaller than 1 GeV.

6 Conclusions and discussion

In this paper we have studied the origin of neutrino mass from SO(10) SUSY GUTs arising from M Theory compactified on a G_2 -manifold. We have seen that this problem is linked to the problem of U(1)_X gauge symmetry breaking, which appears in the SU(5) × U(1)_X subgroup of SO(10), and remains unbroken by the Abelian Wilson line breaking mechanism. In order to break the U(1)_X gauge symmetry, we considered a (generalised) Kolda-Martin mechanism. Our results show that it is possible to break the U(1)_X gauge symmetry without further SUSY breaking while achieving high-scale VEVs that play a crucial role in achieving the desired value of neutrino mass.

The subsequent induced R-parity violation provides an additional source of neutrino mass, in addition to that arising from the seesaw mechanism from non-renormalisable terms. The resulting 11×11 neutrino mass matrix was analysed for one neutrino family and it was shown how a phenomenologically acceptable neutrino mass can emerge. This happens easily for the (n, k) = (3, 0) case of the Kolda-Martin mechanism we developed. For this class of models, not only is the neutrino masses phenomenologically viable, but also the physical light neutrino eigenstate is almost entirely composed of the left-handed (weakly charged) state ν in the same doublet as the electron (ν, e) , as desired. Furthermore, our analysis showed that the B-RPV parameters, which play an important role in neutrino masses and low-energy dynamics, are in the required range, being smaller than 1 GeV. Finally, we notice that contrary to the n = 2 cases, the n = 3 type of Kolda-Martin mechanism immediately preserves the successful predictions of BBN by allowing the LSP to decay quickly in early universe.

In conclusion, we have shown that SO(10) SUSY GUTs from M Theory on G_2 manifolds provides a phenomenologically viable framework, in which the rank can be broken in the effective theory below the compactification scale, leading to acceptable values of neutrino mass, arising from a combination of the seesaw mechanism and induced R-parity breaking contributions. In principle the mechanism presented here could be extended to three neutrino families and eventually could be incorporated into a complete theory of flavour, based on M Theory SO(10), however such questions are beyond the scope of the present paper.

Acknowledgments

The work of BSA is supported by U.K. STFC via the research grant ST/J002798/1. SFK acknowledges support from the STFC Consolidated grant ST/L000296/1 and the European Union Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreements InvisiblesPlus RISE No. 690575 and Elusives ITN No. 674896. MCR acknowledges support from the FCT under the grant SFRH/BD/84234/2012. CP is supported by the KCL NMS graduate school and ICTP Trieste. The work of KB is supported by a KCL GTA studentship.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- S.F. King, Models of neutrino mass, mixing and CP-violation, J. Phys. G 42 (2015) 123001 [arXiv:1510.02091] [INSPIRE].
- [2] S.F. King, Neutrino mass models, Rept. Prog. Phys. 67 (2004) 107 [hep-ph/0310204]
 [INSPIRE].
- [3] G. Altarelli and F. Feruglio, Discrete flavor symmetries and models of neutrino mixing, Rev. Mod. Phys. 82 (2010) 2701 [arXiv:1002.0211] [INSPIRE].
- S.F. King and C. Luhn, Neutrino mass and mixing with discrete symmetry, Rept. Prog. Phys. 76 (2013) 056201 [arXiv:1301.1340] [INSPIRE].
- [5] S.F. King, A. Merle, S. Morisi, Y. Shimizu and M. Tanimoto, Neutrino mass and mixing: from theory to experiment, New J. Phys. 16 (2014) 045018 [arXiv:1402.4271] [INSPIRE].
- [6] P. Minkowski, $\mu \to e\gamma$ at a rate of one out of 10⁹ muon decays?, Phys. Lett. **B** 67 (1977) 421 [INSPIRE].
- M. Gell-Mann, P. Ramond and R. Slansky, Complex spinors and unified theories, Conf. Proc. C 790927 (1979) 315 [arXiv:1306.4669] [INSPIRE].
- [8] T. Yanagida, Horizontal symmetry and masses of neutrinos, Conf. Proc. C 7902131 (1979) 95 [INSPIRE].
- R.N. Mohapatra and G. Senjanović, Neutrino mass and spontaneous parity violation, Phys. Rev. Lett. 44 (1980) 912 [INSPIRE].
- [10] J. Schechter and J.W.F. Valle, Neutrino masses in $SU(2) \times U(1)$ theories, Phys. Rev. D 22 (1980) 2227 [INSPIRE].
- [11] H. Fritzsch and P. Minkowski, Unified interactions of leptons and hadrons, Annals Phys. 93 (1975) 193 [INSPIRE].
- S.P. Martin, A supersymmetry primer, Adv. Ser. Direct. High Energy Phys. 18 (1998) 1
 [Adv. Ser. Direct. High Energy Phys. 21 (2010) 1] [hep-ph/9709356] [INSPIRE].
- [13] E. Witten, String theory dynamics in various dimensions, Nucl. Phys. B 443 (1995) 85 [hep-th/9503124] [INSPIRE].
- P. Hořava and E. Witten, Heterotic and type-I string dynamics from eleven-dimensions, Nucl. Phys. B 460 (1996) 506 [hep-th/9510209] [INSPIRE].
- B.S. Acharya, K. Bożek, M. Crispim Romão, S.F. King and C. Pongkitivanichkul, SO(10) grand unification in M-theory on a G₂ manifold, Phys. Rev. D 92 (2015) 055011
 [arXiv:1502.01727] [INSPIRE].
- [16] E. Witten, Deconstruction, G₂ holonomy and doublet triplet splitting, hep-ph/0201018
 [INSPIRE].
- B.S. Acharya, K. Bobkov, G.L. Kane, J. Shao and P. Kumar, *The G₂-MSSM: an M-theory motivated model of particle physics*, *Phys. Rev.* D 78 (2008) 065038 [arXiv:0801.0478]
 [INSPIRE].
- [18] B.S. Acharya, G. Kane, E. Kuflik and R. Lu, Theory and phenomenology of μ in M-theory, JHEP 05 (2011) 033 [arXiv:1102.0556] [INSPIRE].
- [19] C.F. Kolda and S.P. Martin, Low-energy supersymmetry with D term contributions to scalar masses, Phys. Rev. D 53 (1996) 3871 [hep-ph/9503445] [INSPIRE].

- [20] R. Barbier et al., *R-parity violating supersymmetry*, *Phys. Rept.* 420 (2005) 1
 [hep-ph/0406039] [INSPIRE].
- [21] B.S. Acharya, K. Bobkov, G. Kane, P. Kumar and D. Vaman, An M-theory solution to the hierarchy problem, Phys. Rev. Lett. 97 (2006) 191601 [hep-th/0606262] [INSPIRE].
- [22] R. Blumenhagen, M. Cvetič and T. Weigand, Spacetime instanton corrections in 4D string vacua: the seesaw mechanism for D-brane models, Nucl. Phys. B 771 (2007) 113
 [hep-th/0609191] [INSPIRE].
- [23] L.E. Ibáñez and A.M. Uranga, Neutrino Majorana masses from string theory instanton effects, JHEP 03 (2007) 052 [hep-th/0609213] [INSPIRE].
- [24] M. Cvetič, R. Richter and T. Weigand, Computation of D-brane instanton induced superpotential couplings: Majorana masses from string theory, Phys. Rev. D 76 (2007) 086002 [hep-th/0703028] [INSPIRE].
- [25] W. Buchmüller, K. Hamaguchi, O. Lebedev, S. Ramos-Sanchez and M. Ratz, Seesaw neutrinos from the heterotic string, Phys. Rev. Lett. 99 (2007) 021601 [hep-ph/0703078]
 [INSPIRE].
- [26] J.P. Conlon and D. Cremades, The neutrino suppression scale from large volumes, Phys. Rev. Lett. 99 (2007) 041803 [hep-ph/0611144] [INSPIRE].
- [27] A.E. Faraggi, ν_{τ} mass as possible evidence for a superstring inspired standard like model, *Phys. Lett.* **B 245** (1990) 435 [INSPIRE].
- [28] A.E. Faraggi and E. Halyo, Neutrino masses in superstring derived standard-like models, Phys. Lett. B 307 (1993) 311 [hep-th/9303060] [INSPIRE].
- [29] C. Corianò and A.E. Faraggi, String inspired neutrino mass textures in light of KamLAND and WMAP, Phys. Lett. B 581 (2004) 99 [hep-ph/0306186] [INSPIRE].
- [30] D.M. Ghilencea, L.E. Ibáñez, N. Irges and F. Quevedo, TeV scale Z' bosons from D-branes, JHEP 08 (2002) 016 [hep-ph/0205083] [INSPIRE].
- [31] B.S. Acharya and E. Witten, Chiral fermions from manifolds of G₂ holonomy, hep-th/0109152 [INSPIRE].
- [32] B.S. Acharya and S. Gukov, M theory and singularities of exceptional holonomy manifolds, Phys. Rept. 392 (2004) 121 [hep-th/0409191] [INSPIRE].
- [33] B.S. Acharya, K. Bobkov, G.L. Kane, P. Kumar and J. Shao, Explaining the electroweak scale and stabilizing moduli in M-theory, Phys. Rev. D 76 (2007) 126010 [hep-th/0701034]
 [INSPIRE].
- [34] B.S. Acharya and K. Bobkov, Kähler independence of the G_2 -MSSM, JHEP **09** (2010) 001 [arXiv:0810.3285] [INSPIRE].
- [35] B.S. Acharya, G. Kane and P. Kumar, Compactified string theories generic predictions for particle physics, Int. J. Mod. Phys. A 27 (2012) 1230012 [arXiv:1204.2795] [INSPIRE].
- [36] G.F. Giudice and A. Masiero, A natural solution to the μ problem in supergravity theories, Phys. Lett. **B 206** (1988) 480 [INSPIRE].
- [37] T. Banks, Y. Grossman, E. Nardi and Y. Nir, Supersymmetry without R-parity and without lepton number, Phys. Rev. D 52 (1995) 5319 [hep-ph/9505248] [INSPIRE].
- [38] H.K. Dreiner, An introduction to explicit R-parity violation, Pramana 51 (1997) 123 [Adv. Ser. Direct. High Energy Phys. 21 (2010) 565] [hep-ph/9707435] [INSPIRE].

- [39] B.S. Acharya, S.A.R. Ellis, G.L. Kane, B.D. Nelson and M.J. Perry, *The lightest visible-sector supersymmetric particle is likely to be unstable*, *Phys. Rev. Lett.* **117** (2016) 181802 [arXiv:1604.05320] [INSPIRE].
- [40] L.J. Hall, R. Rattazzi and U. Sarid, The top quark mass in supersymmetric SO(10) unification, Phys. Rev. D 50 (1994) 7048 [hep-ph/9306309] [INSPIRE].
- [41] R. Rattazzi, U. Sarid and L.J. Hall, Yukawa unification: the good, the bad and the ugly, in Yukawa couplings and the origins of mass. Proceedings, 2nd IFT Workshop, Gainesville U.S.A. February 11–13 1994 [hep-ph/9405313] [INSPIRE].
- [42] H. Murayama, M. Olechowski and S. Pokorski, Viable t-b-τ Yukawa unification in SUSY SO(10), Phys. Lett. B 371 (1996) 57 [hep-ph/9510327] [INSPIRE].
- [43] H. Baer, M.A. Diaz, J. Ferrandis and X. Tata, Sparticle mass spectra from SO(10) grand unified models with Yukawa coupling unification, Phys. Rev. D 61 (2000) 111701
 [hep-ph/9907211] [INSPIRE].
- [44] D. Auto, H. Baer, C. Balázs, A. Belyaev, J. Ferrandis and X. Tata, Yukawa coupling unification in supersymmetric models, JHEP 06 (2003) 023 [hep-ph/0302155] [INSPIRE].
- [45] H. Baer, S. Kraml and S. Sekmen, Is 'just-so' Higgs splitting needed for t-b-τ Yukawa unified SUSY GUTs?, JHEP 09 (2009) 005 [arXiv:0908.0134] [INSPIRE].
- [46] A.S. Joshipura and K.M. Patel, Yukawa coupling unification in SO(10) with positive μ and a heavier gluino, Phys. Rev. D 86 (2012) 035019 [arXiv:1206.3910] [INSPIRE].
- [47] M. Drees, Intermediate scale symmetry breaking and the spectrum of super partners in superstring inspired supergravity models, Phys. Lett. B 181 (1986) 279 [INSPIRE].
- [48] A. Brignole, L.E. Ibáñez and C. Muñoz, Soft supersymmetry breaking terms from supergravity and superstring models, Adv. Ser. Direct. High Energy Phys. 18 (1998) 125
 [hep-ph/9707209] [INSPIRE].