



King's Research Portal

DOI: 10.1186/s13033-016-0115-1

Document Version Publisher's PDF, also known as Version of record

Link to publication record in King's Research Portal

Citation for published version (APA):

Docherty, M., Shaw, K., Goulding, L., Parke, H., Eassom, E., Ali, F., & Thornicroft, G. (2017). Evidence-based guideline implementation in low and middle income countries: Lessons for mental health care. *International Journal Of Mental Health Systems*, *11*(1), Article 8. https://doi.org/10.1186/s13033-016-0115-1

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

REVIEW

Open Access



Evidence-based guideline implementation in low and middle income countries: lessons for mental health care

Mary Docherty¹, Kate Shaw², Lucy Goulding², Hannah Parke², Erica Eassom², Farnoosh Ali² and Graham Thornicroft^{3*}

Abstract

Background: There is a significant treatment gap in provision of effective treatment for people with mental disorders globally. In some Low and Middle Income Countries (LMICs) this gap is 90% or more in terms of untreated cases. Clinical practice guidelines (CPGs) are one tool to improve health care provision. The aim of this review is to examine studies of the effectiveness of evidence-based CPG implementation across physical and mental health care, to inform mental healthcare provision in low and middle income countries (LMICs), and to identify transferable lessons from other non-communicable diseases to mental health.

Methods: A systematic literature review employing narrative synthesis and utilising the tools developed by the Cochrane Effective Practice and Organisation of Care (EPOC) group was conducted. Experimental studies of CPG implementation relating to non-communicable diseases, including mental disorders, in LMICs were retrieved and synthesised.

Results: Few (six) studies were identified. Four cluster randomised controlled trials (RCTs) related to the introduction of CPGs for non-communicable diseases in physical health; one cluster-RCT included CPGs for both a non-communicable disease in physical health and mental health, and one uncontrolled before and after study described the introduction of a CPG for mental health. All of the included studies adopted multi-faceted CPG implementation strategies and used education as part of this strategy. Components of the multi-faceted strategies were sometimes poorly described. Results of the studies included generally show statistically significant improvement on some, but not all, outcomes.

Conclusion: Evidence for the effectiveness of interventions to improve uptake of, and compliance with, evidencebased CPGs in LMICs for mental disorders and for other non-communicable diseases is at present limited. The sparse literature does, however, suggest that multifaceted CPG implementation strategies that involve an educational component may be an effective way of improving guideline adherence and therefore of improving clinical outcomes. Further work is needed to examine cost-effectiveness of CPG implementation strategies in LMICs and to draw conclusions on the transferability of implementation experience in physical health care to mental health practice settings. Strategies to ensure that CPGs are developed with clear guidance for implementation, and with explicit, methods to evaluate them should be a priority for mental health researchers and for international agencies.

Keywords: Guideline implementation, Clinical practice guideline, Evidence-based practice, Implementation science, Implementation strategies, Low income countries, Middle income countries, Mental health, Physical health, Systematic literature review

Institute of Psychiatry, Psychology and Neuroscience, King's College

London, London, UK

Full list of author information is available at the end of the article



© The Author(s) 2017. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/ publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

^{*}Correspondence: graham.thornicroft@kcl.ac.uk

³ Centre for Global Mental Health and Centre for Implementation Science,

Background

The global burden of mental, neurological and substance use (MNS) disorders is relentlessly high, resulting in long-term disability combined with premature mortality [1, 2]. Untreated MNS disorders also have a negative impact on global health priorities [3], and may be associated with human rights abuses [4]. The majority of people with MNS disorders in LMICs are unable to access effective mental health care, with the treatment gap higher than 90% in many such countries [5–7]. The most recent estimates of the global burden of mental and neurological disorders suggest that these may be considerably greater than previously thought [8].

Issues related to quality improvement and implementation science are central to these challenges [9, 10]. Health system constraints are recognised to be potent threats to the scale-up of access to evidence-based mental health care for people affected by MNS disorders in low- and middle-income countries (LMICs) [11]. Policy makers and planners play a critical role in the successful strengthening of mental health systems, but may not be appropriately equipped for the task. For example, in a mixed-methods study of the challenges faced during implementation of national mental health policy in South Africa, the key barriers included the low priority given to mental health care by planners, provincial bureaucracy around service coordination, insufficient staff for policymaking and service planning, and disinclination by some local authorities to lead mental health policy implementation [12].

In a qualitative study involving national and regional stakeholders in Ghana, South Africa, Uganda and Zambia, low perceived legitimacy of the problem of scaling up mental health services and inadequate government support were identified as factors perpetuating the low priority accorded to mental health care [13]. A survey of leaders and specialists in international mental health specifically identified a need for a more over-arching public health perspective among mental health policy-makers [14]. The lack of training and experience of clinicians to fulfil leadership roles in policy making and planning was particularly emphasised.

There is international consensus on the need for mental health system strengthening and for a specific focus on building the capacity of key stakeholders, including policy makers and planners and service users [15–18].

These issues speak to the *quantity* of mental health care available in LMICs. Indeed, the key mental health focus of the WHO is the *mental health gap*, namely the difference between true prevalence rates of mental disorders and treated prevalence rates. Such treated prevalence rates are sometimes expressed as *treatment coverage*. Nevertheless the usual way in which coverage is conceptualised and measured refers to *contact coverage*, i.e. the occurrence of any treatment encounter, whether or not this confers benefit to the patient. A more satisfactory definition refers to quality of care as expressed in terms of *effectiveness coverage*, which means the proportion of people with mental disorders, at a time point or over a time period, who receive effective treatment and care (see Fig. 1) [19].

Regarding implementation, it is clear in terms of the provision of mental health care that there are two central problems: weak or absent (1) national mental health plans, and (2) mechanisms to implement these national plans, including patient and practitioner level interventions such as evidence-based clinical practice guidelines, which are simply not put into practice. One can therefore speak of an *implementation gap*, which is a complex set of barriers standing in the way of better mental health care across most countries of the world [20]. Among the reasons for non-implementation of carefully constructed national level strategic plans, and local level treatment guidelines is the paucity of leadership skills for general health systems strengthening [17, 21–23].

Turning now to the *quality of care*, clinical practice guidelines (CPGs) are an established tool to support reduction in variation, improvement in quality and efficiency, and delivery of evidence-based care [24]. Over the last two decades there has been a relatively large production of mental health guidelines in high-income countries, particularly in the UK, Canada, New Zealand and Australia. With respect to these efforts however, only a small corollary body of research focused on their implementation [25]. Recognition of this knowledge gap and lack of clarity as to the benefits of implementing specific psychiatric guidelines in routine primary care or mental health specialist teams [26] has led to calls for more formal evaluations in this field [27].

A recent systematic review concluded that there was sufficient evidence from high income countries to view CPGs in mental healthcare as an essential asset if appropriately developed and implemented [28]. They showed trends towards improvement in process and patient outcomes following guideline implementation [28]. This supports the legitimacy of CPG development and implementation in LMICs as one of a number of possible tools to approach scale up of mental health care in these settings. On-going difficulties in methodology to evaluate implementation techniques has however been identified as a significant barrier to optimising the potential value of evidence-based guidelines as tools for population level improvements in care [28].

The context for this paper is therefore an appreciation that little has so far been published within the mental health field about implementation of evidence-based



policies and practices in LMICs [29–31]. At the same time we wish to learn from other health sectors of how such progress in LMICs can be made. The aims of this paper are therefore to: (1) review evidence for CPG implementation in physical and mental health care settings in LMICs; (2) consider the transferability of lessons learned from implementation of CPGs in physical health to mental health in LMICs.

Methods

Design of the review

The systematic review was conducted following guidelines produced by the Cochrane Collaboration [32]. The PRISMA checklist was used to inform reporting of the review [33].

Searches

The following electronic databases were searched: MED-LINE, EMBASE, PsychInfo, Global Health and LILACS (Latin American and Caribbean Health Sciences Literature). The search strategy was prepared for MEDLINE (see Additional file 1: Web Appendix) and then translated to other databases following their requirements. Literature published since the earliest date indexed in each database up to the search date (October 2014) was retrieved. Duplicates were removed prior to screening.

Reference list checks of studies found during the electronic search were made. General searches were conducted using internet search engines and key authors in the field were asked to indicate potentially relevant studies. The review team did not have any funds to provide translation; however, Google Translate was used to translate the full text of three studies that were not reported in English. None of these papers met the inclusion criteria.

Study inclusion and exclusion criteria Desian

Studies relating to a non-communicable physical health condition using any of the following designs were included: randomised controlled trials (RCT), cluster-RCTs, controlled clinical trials (CCT), controlled-beforeand-after studies (CBA) and interrupted time series (ITS) studies with concurrent controls.

Studies relating to a mental health condition using any of the following designs were included: randomised controlled trials (RCT), cluster-RCTs, controlled clinical trials (CCT), controlled-before-and-after studies (CBA), interrupted time series (ITS) studies with concurrent controls, and uncontrolled before and after studies.

Included studies were permitted to have any number of sites. Studies with non-experimental designs were excluded. However, findings from process evaluation or qualitative studies conducted alongside experimental studies were included in the data extraction and synthesis of results.

Countries and populations included

Included studies were conducted in a low or middle income country. The World Bank Atlas method for 2014 defined low-income economies as those with a GNI per capita of \$1045 or less; middle-income economies had a GNI per capita of more than \$1045 but less than \$12,746. Studies conducted in high-income countries were excluded.

Eligible participants in included studies were children or adults of any age with a non-communicable disease—including mental and physical health conditions. Examples include: depression, dementia, coronary heart disease, cancer. Studies solely involving participants with communicable diseases (e.g. malaria, HIV, influenza) were excluded. Studies involving outcomes related to surgical site infection and hand hygiene were also excluded. Studies that included participants with both non-communicable and communicable diseases were included with outcomes for people with non-communicable diseases being the focus during synthesis and reporting of the results.

Interventions

Included studies described the introduction of a CPG relating to management of people who have a noncommunicable disease (including mental and physical health). The Institute of Medicine's definition of a CPG was adopted. This states that "clinical practice guidelines are statements that include recommendations intended to optimize patient care that are informed by a systematic review of evidence and an assessment of the benefit and harms of alternative care options" [34].

Studies that assessed the implementation of more than one CPG were included if at least one of the guidelines was in relation to a non-communicable disease (physical or mental health); and results were drawn from that part of the study only. CPGs produced for a health system, a group of healthcare professionals, a country, state, or province were included. Included studies described and assessed the guideline implementation strategy adopted. No restrictions were placed upon the type or number of strategies used. Studies that did not describe a guideline implementation strategy were excluded.

Comparisons

For studies targeting physical health conditions, included studies could compare a group where one or more CPG implementation strategies was used to a control group where no specific implementation strategy was adopted, or groups using different implementation strategies could be directly compared.

For studies targeting mental health conditions, included studies did not have to include a control group. In such cases, the comparison of interest was outcomes before and after the introduction of the CPG and supporting implementation strategy. For study designs employing control groups, included studies could compare a group where one or more CPG implementation strategies was used to a control group where no specific implementation strategy was adopted, or groups using different implementation strategies could be directly compared.

Outcomes

Included studies demonstrated pre and post measurement of processes or outcomes targeted by the CPG. Outcome measures could therefore concern the effectiveness of the implementation strategy measured by compliance with the CPG (for example prescribing behaviours), changes in the knowledge, attitudes, beliefs of behaviours of healthcare professionals, or changes in patients' health. Outcomes could be clinician or patient reported. Post-implementation measurement could take place at any point following the introduction of the CPG and accompanying implementation strategies. Studies that did not include pre and post CPG implementation measurement were excluded.

Screening

Screening of titles and abstracts and full-texts was conducted by four authors (MD, KS, HP and EE). Authors met to agree on final inclusion. Uncertainties experienced during the screening and extraction process were resolved with input from an additional author (LG).

Data extraction and quality assessment

Three authors (MD, KS and FA) independently extracted data from included studies and appraised the potential risk of bias in each study using tools developed by the Cochrane Effective Practice and Organisation of Care (EPOC) group [35]. An additional author (LG) double checked all data extractions and risk of bias assessments. The Cochrane Collaboration's 'EPOC taxonomy' was used to classify the implementation strategies adopted in included papers [36].

Data synthesis

Due to the nature of the review question there was heterogeneity in the participants, interventions, comparisons, outcome measures and outcomes of included studies. A descriptive data synthesis was consequently undertaken to summarise the characteristics and results from included studies in table form and to address the review questions.

Results

Results of the search

The electronic searches yielded 18,060 citations. 7950 remained after duplicates were removed. An additional 38 citations were uncovered by hand searching reference lists of key papers and through contact with experts in the field. This was reduced to 159 papers for full text screening, of which 6 met the inclusion criteria for the review. Details of included studies are provided in Table 1 and Fig. 2.

Included studies

Six papers met the inclusion criteria for the review. Four of these papers related to the introduction of CPGs for non-communicable diseases in physical health [37–40],

Lead author (year of publication) Country Mental or physical health	Study design Risk of bias assessment	Health condition or behav- iour targeted by Clinical Practice Guideline (GPG) Setting and participants	Brief description of interven- tion / control conditions and implementation strate- gies adopted (classified according to EPOC tax- onomy)	Primary outcomes	Narrative summary of results
Berwanger et al. [37] Brazil Physical health (acute coronary syndromes)	Cluster randomised trial Low-moderate risk of bias	<i>CPG</i> The intervention included posters containing evidence- based recommendations for the treatment of acute coronary syndromes (ACS) <i>Setting and participants</i> 36 general public hospitals with emergency centres in major urban areas in Brazil. 19 hos- pitals were randomised to receive the intervention and 17 to control. The analysis represented 602 patients with a diagnosis of ACS in the intervention group and 548 in the control group and 548 in the control group and 548 in the control of the number and type of health professionals exposed to the intervention is not provided	Intervention A multifaceted educational quality improve- ment intervention for man- agement of acute coronary syndromes Implementation strategies adopted were: reminders; case management; educa- tional materials, educational outreach; local opinion leaders Checklists were also used Control: routine care	The percentage of eligible patients who received all appropriate evidence-based therapies (aspirin, clopi- dogrel, anticoagulants and statins) within the first 24 h of admission to hospital without contraindications	A multifaceted educational intervention resulted in statisti- cally significant improvement in the use of evidence-based therapies both within 24 h of admission and at discharge in comparison to routine care There were no statistically sig- ificant differences in the rate of major cardiovascular events, mortality at 30 days, new myo- cardial infarction or incidence of major bleeding—however, the study was not powered for evaluation of clinical outcomes Adherence to the reminders and checklists by healthcare professionals in the interven- tion group was 82.7%

Table 1 Description of studies meeting the inclusion criteria

Lead author (year of publication) Country Mental or physical health	Study design Risk of blas assessment	Health condition or behav- iour targeted by Clinical Practice Guideline (GPG) Setting and participants	Brief description of interven- tion / control conditions and implementation strate- gies adopted (classified according to EPOC tax- onomy)	Primary outcomes	Narrative summary of results
Du et al. [52] China Physical health (acute coronary syndromes)	Cluster randomised controlled trial Moderate-high risk of bias	<i>CPG</i> Clinical pathways for acute coronary syndromes based on the American Heart Association / American College of Cardiology guidelines Setting and participants 70 urban hospitals throughout China routinely admitting >100 patients annually with suspected acute coronary syndromes were randomly allocated to receive the intervention (n = 32 hospitals, n = 8049 patients) or control (n = 38 hospitals, n = 5731 patients) Detail regarding the number and type of health professionals exposed to the intervention is not provided to the in	Intervention group Hospitals implemented clinical path- ways (and accompanying multifaceted implementation strategies) for acute coronary syndromes early' Implementation strategies adopted: audit and feedback; monitoring the performance of the delivery of healthcare; educational meetings <i>Control group</i> Usual care for first year. Hospitals imple- mented clinical pathways (and accompanying imple- mentation strategies) for acute coronary syndromes 'late'(1 year after intervention sites) Key performance indicators in intervention sites were collected 12 months after the introduction of the inter- vention and compared with baseline data in control sites The guideline was tailored to the local context. Teams led by senior cardiologists took responsibility for implemen- tation	Proportion of patients with final diagnosis consistent with biomarker findings; pro-portion of patients with ST-segment-elevation myocardial infarction (STEMI) receiving thrombolysis or primary percutaneous coronary intervention (PCI) among those arriving within 12 h of symptom onset; door-to-needle time for patients with STEMI undergo- ing thrombolysis; door-to- balloon time for patients with STEMI undergoing primary PCI; proportion of high-risk patients undergoing primary PCI; proportion of low-risk patients (no on- going symptoms, persistently normal electrocardiogram, and persistently normal biomarkers) undergoing functional testing; proportion of patients discharged on combination medical therapy (including any antiplatelet therapy, B-blocker, and statin); lendth of stav	The use of a clinical pathway for treatment of acute coronary syndromes compared with usual care statistically significantly improved use of secondary prevention treatments (an increase in the number of patients who were discharged on appropriate medical therapy) but there were no statistically significant differences on other measures of quality of care A survey of 556 health profes- sionals found that 98.2% had head of the intervention, and >80% had attended training sessions, used the pathway in clinical practice and were aware of study reports.>90% completely or strongly agreed that the pathway was valuable. A qualitative process evaluation aids understanding of findings

Table 1 continued

Lead author Stu					
(year of publication) Ris Country ass Mental or physical health	ıdy design ik of bias sessment	Health condition or behav- iour targeted by Clinical Practice Guideline (GPG) Setting and participants	Brief description of interven- tion / control conditions and implementation strate- gies adopted (classified according to EPOC tax- onomy)	Primary outcomes	Narrative summary of results
Okasha et al. [42] Unc Egypt Hig Mental health Hig	controlled before and after tudy gh risk of bias	<i>CPG</i> The tenth revision of the International Statistical Classification of Mental and Behavioural disorders Primary Health Care (ICD-10 PHC) Setting and participants: 20 GPs working in 6 primary health care centres (rural and urban) in Egypt	Intervention: Training was con- ducted by an experienced psychiatrist. The ICD-10 training kit, an educational training programme and materials produced by the WHO Implementation strate- gies therefore comprised: educational materials	GP's attitudes, knowledge, interview skills and ability for diagnosing psychiatric disorders in a primary care setting	There was statistically significant improvement in GP's optimism about helping patients with mental disorders and in their confidence in their ability to diagnose mental disorders fol- lowing the intervention. There was no statistically significant change in GP's interest in patients with mental disorders or in their reporting of the importance of diagnosing mental disorders.
Pagaiya and Gamer [41] Clu Thailand tr Physical health (diabetes Mo mellitus) and mental health (anxiety and panic disorder)	uster randomised controlled rial oderate risk of bias	<i>CPG</i> Four clinical guidelines introduced: two for children (acute respiratory infection and diarrhoea); two for adults (diazepam prescribing for anxiety and panic disorder and management of diabe- tes mellitus) <i>Setting and participants</i> Health centres in Thailand, mainly in rural areas. 9 health centres were randomised to receive the intervention and 9 to control. 110 patients for diazepam prescribing and all diabetes patients at each centre were included	Intervention group introduction of four clinical guidelines (two of which meet the inclusion criteria for the present review) plus the following implementation strategies: educational meet- ings; educational out-reach visits; audit and feedback <i>Control group</i> usual care. The PRECEDE-PROCEED imple- mentation framework was adopted	Diazepam prescribing; pre- scribing costs per patient; management of diabetes	Clinical guidelines implemented with educational meetings and outreach visits and audit and feedback improved some aspects of prescribing but not others in the short-term Diazepam prescribing was statis- tically significantly reduced in the intervention group. There was no statistically significant difference in the management of diabetes between the intervention and control groups. For all four CPGs combined, average drug costs per patient in the intervention group were statistically significantly less than in the control group

Table 1 continued					
Lead author (year of publication) Country Mental or physical health	Study design Risk of bias assessment	Health condition or behav- iour targeted by Clinical Practice Guideline (GPG) Setting and participants	Brief description of interven- tion / control conditions and implementation strate- gies adopted (classified according to EPOC tax- onomy)	Primary outcomes	Narrative summary of results
Shrestha et al. [3 9] Nepal Physical health (asthma, COPD)	Cluster randomised trial High risk of bias	<i>CPG Practical</i> Approach to Lung health (PAL) guidelines on prescription behaviour and total cost of prescription for patients with asthma, pneumonia and chronic obstructive pulmonary disease (COPD) Setting and participants: 0f 76 health facilities in Nawalpar- asi district, 40 were included in the study on the basis of highest patient attendance. Facilities were stratified by type and subsequently randomized to receive the intervention [19]. Detail regarding the number of patients in each group is not provided, rather number of patients in each group is not provided, rather number of prescriptions is given. Detail regarding the number and type of health professionals exposed to the intervention is not provided	Intervention Introduction of the PAL guideline plus the following accompanying implementation strategies: educational materials, educa- tional meetings, educational outreach (train-the trainer) <i>Control</i> Usual practice The guideline was adapted to the Nepalese context. Differ- ent professionals, organisa- tions and health workers were involved in this process	Healthcare professional pre- scribing practices for asthma and COPD	The guidelines and accompany- ing implementation strategies led to a statistically significant reduction in poly-pharmacy (multiple prescriptions) in comparison to the control condition. There were no sta- tistically significant differences in the prescription of generic and essential drugs, average prescription of antibi- otics between the intervention and control groups

Lead author (year of publication) Country Mental or physical health	Study design Risk of bias assessment	Health condition or behav- iour targeted by Clinical Practice Guideline (GPG) Setting and participants	Brief description of interven- tion / control conditions and implementation strate- gies adopted (classified according to EPOC tax- onomy)	Primary outcomes	Narrative summary of results
Steyn et al. [40] South Africa Physical health (diabetes mel- litus and hypertension)	Cluster randomised controlled trial Low -moderate risk of bias	<i>CPG</i> A structured record which incorporated the national guidelines for the manage- ment of patients with type 2 Diabetes mellitus or hyperten- sion or both conditions <i>Setting and participants</i> 18 randomly selected public sector primary healthcare clinics known as Community Health Centres (CHCs) in Cape Town in 1999 and 2000. 9 clinics were randomised to deliver the intervention and 9 to control. At baseline, in the intervention with diabetes and n = 461 patients with hypertranion. At baseline, in the control clusters there were n = 227 patients with diabetes and n = 459 patients with hypertranion. The interven- tion was targeted at General Practitioners and nurses who consult patients with diabetes ing the number of health professionals exposed to the intervention is not provided	Intervention a structured record that included algorithms for diagnosis and guideline- based management of diabetes and hypertension, including educational top- ics to be covered with the patient Implementation strategies adopted: case management; reminders; educational out- reach from a local diabetes and hypertension expert <i>Control group</i> usual care, which included the guidelines passively disseminated by the National Department of Health	Mean level of glycated hae- moglobin in all patients with diabetes Mean systolic and mean diastolic BP in all those with hypertension as measured at the end of the 1 year intervention period	The intervention had no statisti- cally significant impact on either diabetes or hyperten- sion control No differences were observed in the recording of process measures, e.g. visual acuity or foot examinations The process measures collected implied that the structured record was not widely adopted by healthcare professionals in the primary care clinics. At follow-up, the structured record was found in the folders of only 58% of patients with diabetes and 47% of patients with hypertension



one paper included CPGs for both a non-communicable disease in physical health and mental health [41] and one paper examined the introduction of a CPG for mental health [42].

Design

Five cluster randomised controlled trials (cluster RCTs) [37–41] and one uncontrolled before and after study (in mental health) [42] were included in the review.

Population

Due to the nature of the interventions being reviewed (implementation of CPGs); all interventions were delivered to healthcare professionals. Where specified, participants receiving the intervention were typically physicians and nurses.

Setting

Included studies were conducted in Brazil [37], China [38], Thailand [41], Nepal [39], South Africa [40] and Egypt [42]. Two studies were set in general hospitals in urban areas [37, 38]. Four studies were set in primary health care centres: one of these was conducted in an urban area [40]; two in rural areas [39, 41] and one in a mix of urban and rural areas [42].

Behaviour/clinical condition targeted by the CPG

The non-communicable clinical conditions and behaviours targeted by the CPGs were as follows: use of evidence-based therapies for acute coronary syndromes [37, 38]; diazepam prescribing practices for anxiety or panic disorder and management of diabetes mellitus [41]; prescribing practices for asthma and COPD [39]; management of diabetes mellitus and hypertension [40]; and management of mental and behavioural disorders in primary healthcare through implementation of the ICD-10 [42].

Format of clinical practice guidelines

Evidence-based guidelines were presented by: poster [37]; a clinical pathway document containing structured algorithms to be filled in by the healthcare professional [38]; a structured record to be filled in by the healthcare professional which was a three sided, folded A3 sheet of coloured paper placed with each patient's notes [40]; and A4 laminated documents printed on coloured paper [41]. In two studies, the format of the CPG was unclear [39, 42]. Four studies report that the CPG was either originally developed for the local context or tailored to the local context [38–41].

CPG implementation strategies adopted

All of the included studies adopted multi-faceted CPG implementation strategies. The components of the multifaceted interventions were sometimes poorly described. All studies employed implementation strategies that targeted healthcare professionals' use of the CPG. Implementation strategies adopted to facilitate uptake of CPGs (classified according to the EPOC taxonomy) were: reminders, case management, educational materials and educational outreach [37]; audit and feedback, monitoring the performance of the delivery of healthcare and educational meetings [38]; educational meetings, educational out-reach and audit and feedback [41]; educational materials, educational meetings and educational outreach [39]; case management, reminders and educational outreach [40]; educational meetings and educational materials [42].

Only one study [41] explicitly described adoption of a theoretical framework to guide CPG implementation. One paper reported the lack of adaptation of the CPG implementation strategies to the local context as a limitation [40]—this finding was pronounced in qualitative interviews undertaken with healthcare professionals as part of a process evaluation.

Control groups

In four of the cluster-RCTs, the intervention (CPG and accompanying implementation strategies) was compared to routine care (no CPG or accompanying implementation strategies) [37–39, 41]. In one cluster-RCT, the intervention (CPG and accompanying implementation strategies) was compared to passive diffusion of the CPG (no accompanying implementation strategies) [40]. The before and after study in mental health [42] did not include a control group.

Primary outcomes

Primary outcomes relating to processes undertaken by healthcare professionals were: the proportion of patients who received evidence-based treatments for acute coronary syndromes [37, 38] and prescribing practices [39, 41]. Primary outcomes relating to healthcare professionals were: attitudes, knowledge and skills [42]. Primary outcomes relating to patients were level of glycated haemoglobin in patients with diabetes and systolic and diastolic blood pressure in patients with hypertension [40].

Risk of bias in included studies

The risk of bias in included studies was mixed. In the case of the cluster-RCTs, the nature of the intervention (CPG) and accompanying implementation strategies (e.g. educational sessions) often meant that it was not feasible to blind participants or personnel to the allocation of each cluster. In some cases, staff from different groups met occasionally leading to opportunity for contamination. Randomisation processes were judged to be low risk in four studies [37, 38, 40, 41] and unclear in one study [39]. Risk of bias in the mental health paper that used the ICD-10 [42] was judged to be very high due to the use of an uncontrolled before and after study design.

Effects of interventions

Results of the included studies generally show statistically significant improvement on some but not all outcomes with the exception of one study [40] where the CPG intervention did not demonstrate any statistically significant improvement in comparison to control. The authors of this paper proposed that poor implementation of their intervention and accompanying implementation strategies was the major contributor to ineffectiveness.

Multifaceted implementation strategies were generally deemed to be effective in encouraging use of guidelines and creating positive change in outcomes. However, it is not possible to infer which approach to guideline implementation is most effective. Furthermore, the small number of included studies means that there are no observable trends according to clinical setting or topic addressed by the CPG (e.g. physical versus mental health).

Implementation outcomes

Implementation outcomes, as defined by Proctor et al. [43], were infrequently studied. Cost effectiveness of the interventions (e.g. taking into account the cost of delivering the CPG implementation strategies) was not assessed in any of the included studies. However, the effect of the intervention on prescription costs was calculated in two studies [39, 41]. Fidelity and acceptability of the intervention was explored via qualitative process evaluation

in two studies [38, 40]. The length of time to follow-up in included studies was generally short: 30 days [37]; 6 months [41]; 8 months [39]; 1 year [38, 40] and unclear but assumed to be immediately following the intervention [42]. This limits the ability to draw inferences regarding sustainability of any improved practice.

Discussion

The aims of this review were to examine studies of the effectiveness of evidence-based clinical practice guideline implementation for non-communicable diseases in LMICs, and to learn transferable lessons from other non-communicable diseases to mental health. The review revealed a significant paucity of good quality controlled studies not only in the field of mental health but across other non-communicable diseases in LMICs.

The small number of included papers made it difficult to draw clear conclusions about the role of different implementation strategies to support CPGs in mental health or in other non-communicable disease areas. Our inclusion criteria for mental health studies permitted uncontrolled designs in order to capture any existing work in this area. For other non-communicable disease areas, only high quality studies including control groups were included to support better validity of any conclusions being drawn about transferability of lessons learned across health sectors. As a result of this design, the relatively stringent inclusion criteria led to a large number of non-controlled studies in physical health being excluded from review. Future reviews may therefore wish to relax the inclusions criteria to gather a wider range of information which can create a conceptual map of this type of evidence.

This larger volume of excluded non-controlled studies in physical health had significant methodological limitations in addition to the absence of a control group. A number of studies described interventions with adequate study design for inclusion, but did not detail or refer to an evidence-base in the development of the CPG (e.g. [44]). Other common omissions included failure to clearly describe the implementation strategy used, to consider reproducibility of results, or to consider a theoretical framework in which to seat the implementation study and evaluation. Many excluded papers relied on pre and post analysis without clear attention to confounders that may have affected whether the implementation strategy itself, or other variables, impacted upon any observed change in practice. This difficulty is noted ten years after this problem had already been highlighted in a previous literature review [45]. Overall, the methodological limitations of the excluded studies (see Additional file 1: Web Appendix) mirrored similar problems reported within current literature on guideline implementation in

high-income countries suggesting that these study design difficulties are consistent across these different health care settings [24, 27, 47].

Most of the included studies did not explore the relative efficacy of one implementation strategy compared to another in supporting the uptake of and adherence to CPGs, leaving important knowledge gaps with regards to the most effective and efficient strategies to support change within low resource settings. Additionally, there was a frequent lack of delineation between the effect of a strategy to improve uptake of guidelines distinct from whether improved implementation actually improved clinical outcomes targeted by the CPG (e.g. [40, 46]).

Multifaceted interventions with an educational component appear to be effective at supporting change, and attention to their delivery alongside CPG implementation is important to ensure optimum impact. Development of the CPG with consideration of local context, including staff attitudes and available resources, appear to be very important. The one study that showed no improvement in any targeted outcomes had developed CPGs specifically for the local context but had not sufficiently developed a feasible implementation strategy. Qualitative interviews with the professionals involved in implementing CPGs in this study revealed not only an ambivalence towards the perceived utility of the intervention, but also a fundamental failure of the guideline to accommodate resource limitations meaning that were the guideline to be fully implemented it would simply not be affordable [40]. In comparison, the studies that tailored both CPG development and implementation to the local context appeared to be relatively more efficacious in respect of achieving targeted outcomes [38–40].

Known barriers to improving mental health provision in LMICS includes the perceived lack of importance of this field relative to other clinical areas. These observations together support the impression that guideline implementers need to engage those using or impacted upon by the CPGS prior to implementing other strategies focused directly on the guideline use. In mental health this could for example include preliminary anti-stigma interventions. Pagaiya recommends that a "well-planned stepwise process be adopted which takes account of both theoretical and empirical evidence, as well as obstacles to change in relation to the individual staff and the local context" in order to approach successful implementation [41].

Prescribing featured in several of the included studies including one of those addressing mental health guidelines (the ICD-10), which may reflect the relatively more straightforward task of monitoring this sort of intervention. It could be argued that other mental health interventions such as the delivery of psychological therapy Page 12 of 16

or case management approach are more challenging to formulate into guidelines and associated guideline adherence monitoring. Medication-based interventions in mental health CPGs implemented in high income countries may be a feasible place for future researchers to gain insight. Consideration of local availability of specific medications, facilities to store them and resource constraints on formulations offered will be essential in developing and implementing such guidelines in lower income settings.

There was an insufficient number of studies to be able to draw conclusions about transferability of findings to specific diseases, populations or care settings. However, the diversity of the included studies illustrates that CPGs can be implemented and evaluated across a range of populations and care settings within LMICs. Asthma, COPD, hypertension, acute coronary syndrome (ACS) and diabetes were targeted in the included studies on physical health conditions and primary care mental health provision and anxiety disorders targeted in the mental health paper. The transferability of lessons from most of the included physical illnesses to mental health seems feasible in principle due to the parallels between these disease courses and treatment requirements. Most conditions studied were chronic diseases which present with a risk of fluctuations and requirement of a series of interventions to support recovery and stability, in keeping with the clinical course of many mental health disorders (e.g. schizophrenia). ACS is an acute presentation with the requirement of treatment intervention and systematic follow up for a period in keeping with most mental health crises (e.g. suicide plans). There were insufficient studies to explore the question of whether similar pathways and protocols can be constructed for mental health conditions but the findings did not exclude the transferability of these approaches.

Despite these similarities in principle supporting the transferability of implementation experience across mental health and other non-communicable disease areas, there are also important differences in these fields. Mental health workers employ different diagnostic processes, with less reliance on technology and more reliance on human resources to deliver both assessment and treatment. These features highlight a risk for even greater variation in practices and the need to learn from experience in other non-communicable disease areas of the importance of embedding systems for standardisation and measurement of interventions within guidelines. Outcomes in mental health are frequently qualitatively different to those in other non-communicable diseases and less amenable to conventional measurements. In the absence of biomarker outcome measures the importance of developing and incorporating simple and tractable

measures such as quality of life, ability to sustain employment, activities of daily living (ADLs) or relationships in order to validate and monitor the efficacy of CPGs in this field is essential.

Local context including prevalent knowledge, behaviours and attitudes towards mental health conditions has predicted larger impact on the potential success of a guideline than in many of the non-communicable diseases considered in this review. Despite some literature acknowledging the context in which a guideline was to be implemented, few gave consideration to the range of barriers that would need to be considered in mental health before implementing and evaluating change. Although beyond the scope of this paper, an important area to consider in future reviews is how impediments to changes in practice due to stigma in some communicable disease areas, for example HIV, have been overcome and addressed in guideline development and implementation.

A consistent difficulty observed in included studies was the limited follow up to evaluate longer-term impact and interventions needed to embed changes in clinical practice in the medium and long term. Longer follow up periods are needed to understand the requirements for sustained change such as on-going interventions (e.g. educational updates) or wider system changes (e.g. changes in job roles, informatics or care pathways and checklists). There is evidence that active on-going efforts to support changes in practice may be required beyond the initial implementation period (e.g. [45, 48, 49]. This has important resource implications for health planners looking to embed improvements into routine clinical care. None of the included studies conducted cost effectiveness analyses of CPG implementation which is a very important omission given the resource constraints in these settings. This should be considered a clear priority in future study designs.

Despite most of the included studies showing improvements in a selection of outcomes, it is not possible to reach conclusions regarding the sustainability, feasibility or practicality of these approaches for mental health planners. For example, all included studies used educational implementation strategies and showed them to be broadly efficacious but they can be labour intensive. One difficulty in providing such strategies in LMICs is the available modes and associated resources for delivering the educational intervention. Furthermore, education is an important tool to support changes in clinical practice but rarely a one off solution to altering knowledge and behaviours. Multiple sessions and top ups are usually required alongside rolling programmes to support staff turnover. Implementing interventions based around education usually require additional strategies such as rigorous systems for monitoring to support change. Outreach approaches were used in some of the studies but again these can be costly and if practitioners are geographically very far apart, this would make educational meetings difficult. Educational sessions also reduce time to engage in clinical activity in already resource stretched services, this places an onus on organisers to ensure high quality

sessions or material to support perceived utility and buy

in towards the CPG. Another implementation strategy commonly used in low resource settings, known as task shifting, was not adopted in the studies included within this review. Task shifting refers to the process of transferring a task usually delivered by a scare resource such as a physician to a more rapidly trained and less scare resource such as a health care worker. Evidence has shown that this strategy has been effective in increasing use of treatment guides and protocols in the management of a range of noncommunicable conditions such as asthma, hypertension, epilepsy, diabetes and depression [50, 51]. Integrated protocols that involved strategies such as care bundles or care pathways that supported whole systems change seem potentially more promising than focus on just the guideline alone (e.g. [52]). Directed studies to explore the use of task shifting alongside other CPG implementation strategies may be an important area for future research in LMICs considering the need for CPGs not only to improve patient outcomes but also to address efficiency.

Despite the emergence of some lessons for mental health planners, the volume of literature excluded from this review (see Additional file 1: Web Appendix) reveals significant limitations in both study methodology and reporting practices and a need to increase the volume of good quality research. This highlights the importance of a concerted effort within LMICs to improve the rigor of CPG implementation studies. Efforts to increase the availability and existence of context adapted evidencebased mental health guidelines are in process but it is essential that mental health care planners and researchers learn from the limitations encountered in other non-communicable disease guideline implementation research. The World Health Organization mhGAP Intervention Guide, for example, is now in use in over 90 countries worldwide, where the guidelines are intended to be locally adapted for each country and each context, but as yet few evaluations of its use have been published [29, 30, 53-55].

This raises the more fundamental question of whether the scope of this review was too narrow. Clearly this field is not at the stage where many RCTs have been published, from which strong summary findings can be draw. Indeed, given the relative infancy of this field, we did consider whether to conduct a broader narrative review of the literature, for example to summarise what is known of barrier and facilitator factors in guideline implementation in LMICs. Similarly we considered including process evaluation papers in this review. On balance we decided to take a narrower focus on the better quality papers for this paper. The results indicate that future reviews may need to adopt somewhat broader criteria, for example including non-experimental studies, until such time as the quality of the available evidence improves sufficiently.

The overall disconnection which was observed between the volume of literature on guideline development in LMICs, and that on guideline implementation raises important questions for those developing and adapting guidelines in mental health about with whom the responsibility lies for ensuring validation and implementation studies of CPGs are conducted. It has been argued that sound validation studies should be considered a prerequisite for conferring the label of an evidence-based mental health guideline [26]. The importance of this approach in the field of mental health in lower resource settings has particular resonance due to the highly variable existing health care resources and infrastructures in which mental health care can be delivered and the requirement of a local approach to map knowledge, attitudes and behaviour before developing an implementation strategy. In resource limited settings with such significant diagnosis and treatment gaps, efforts to increase effective coverage of mental health care must consider implementation strategies and attention to resource constraints as necessary components of the guideline development process.

This review demonstrates the salience of the implementation gap across medical specialities in LMICs and the real risks for mental health in not responding to these lessons from CPG development over the last few decades. Those looking to scale up mental health in LMICs must prioritise implementation research. Those facilitating the development of CPGs, including professional bodies contributing to them, have a responsibility to ensure that their efforts and the money invested in them lead to tangible improvements in care. Unless agreement is made as to how to take forward this requirement for mental health provision, a real opportunity to learn from lessons in other sectors and settings will be missed.

Conclusions

Current evidence for the effectiveness of interventions to improve uptake of and compliance with evidence based guidelines in LMICs for mental disorders and for other non-communicable diseases is very limited. The literature suggests that multifaceted CPG implementation strategies that involve an educational component may be an effective way of improving guideline adherence and therefore improving clinical outcomes. Further work is needed to examine cost effectiveness of CPG implementation strategies in LMICs and to draw conclusions on the transferability of implementation experience in other non-communicable disease areas to mental health. Strategies to ensure that CPGs are developed with clear guidance for implementation and methods to evaluate them should be a priority for mental health researchers and for international agencies.

Additional file

Additional file 1: Web Appendix. Excluded papers.

Abbreviations

ACS: acute coronary syndrome; COPD: chronic obstructive pulmonary disorder; CPGs: clinical practice guidelines; EPOC: Cochrane Effective Practice and Organisation of Care; CBA: controlled-before-and-after studies; CCT: controlled clinical trials; GNI: gross national income; ITS: interrupted time series; LMICS: Low and Middle Income Countries; PRISMA: preferred reporting items for systematic reviews and meta-analyses; RCTs: randomised controlled trials.

Authors' contributions

MD helped to develop the design of the review, conducted screening and data extraction, contributed to exclusion and exclusion criteria, drafted the discussion section of the manuscript and co-ordinated production of the manuscript. KS conducted screening and data extraction. LG helped to develop the design of the review, checked inclusions and data extraction, and drafted the methods and results sections of the manuscript. HP helped to develop the design of the review, wrote the search strategy, conducted the search, and conducted screening. EE conducted screening and reviewed inclusion and exclusion criteria. FA conducted data extraction. GT conceived the review, helped to develop the design of the review, and drafted the introduction. All authors provided comments on drafts of the manuscript and approved the final version to be published. All authors read and approved the final manuscript.

Author details

¹ South London and Maudsley NHS Foundation Trust, The Maudsley Hospital, Denmark Hill, London SE5 8AF, UK. ² Centre for Implementation Science, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK. ³ Centre for Global Mental Health and Centre for Implementation Science, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK.

Acknowledgements

GT is supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care South London at King's College London Foundation Trust. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. GT acknowledges financial support from the Department of Health via the National Institute for Health Research (NIHR) Biomedical Research Centre and Dementia Unit awarded to South London and Maudsley NHS Foundation Trust in partnership with King's College London and King's College Hospital NHS Foundation Trust. GT is supported by the European Union Seventh Framework Programme (FP7/2007-2013) Emerald project.

Competing interests

The authors declare that they have no competing interests.

Availability of data and supporting materials

There is no additional data associated with this work. A Additional file of the review search strategy used has been submitted.

Consent for publication

Not applicable. This is a review article and does not contain any individual person's data.

Ethics approval and consent to participate

Not applicable. This is a review article and does not report on or involve the use of any animal or human data or tissue.

Funding

KS, LG, HP and EE were supported by King's Improvement Science. King's Improvement Science is part of the NIHR CLAHRC South London and comprises a specialist team of improvement scientists and senior researchers based at King's College London. It's work is funded by King's Health Partners (Guy's and St Thomas' NHS Foundation Trust, King's College Hospital NHS Foundation Trust, King's College London and South London and Maudsley NHS Foundation Trust), Guy's and St Thomas' Charity, the Maudsley Charity and the Health Foundation. There was no direct individual funding for this work. MD did not receive funding for this work. There was no distinct role of any funding body in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript as detailed in contributions.

Received: 8 November 2016 Accepted: 21 December 2016 Published online: 05 January 2017

References

- Fekadu A, Medhin G, Kebede D, Alem A, Cleare A, Prince M, et al. Excess mortality in severe mental disorders: a 10-year population-based cohort study in rural Ethiopia. Br J Psychiatry. 2015;206(4):289–96. doi:10.1192/ bjp.bp.114.149112.
- Whiteford HA, Degenhardt L, Rehm J, Baxter AJ, Ferrari AJ, Erskine HE, et al. Global burden of disease attributable to mental and substance use disorders: findings from the Global Burden of Disease Study 2010. Lancet. 2013;382(9904):1575–86.
- Prince M, Patel V, Saxena S, Maj M, Maselko J, Phillips MR, et al. Global mental health 1: no health without mental health. Lancet. 2007;370:859–77.
- Drew N, Funk M, Tang S, Lamichhane J, Chávez E, Katontoka S, et al. Human rights violations of people with mental and psychosocial disabilities: an unresolved global crisis. Lancet. 2011;378(9803):1664–75.
- Alem A, Kebede D, Fekadu A, Shibre T, Fekadu D, Beyero T, et al. Clinical course and outcome of schizophrenia in a predominantly treatmentnaive cohort in rural Ethiopia. Schizophr Bull. 2009;35(3):646–54.
- Wang PS, Aguilar-Gaxiola S, Alonso J, Angermeyer MC, Borges G, Bromet EJ, et al. Use of mental health services for anxiety, mood, and substance disorders in 17 countries in the WHO world mental health surveys. Lancet. 2007;370(9590):841–50.
- Thornicroft G. Most people with mental illness are not treated. Lancet. 2007;370(9590):807–8.
- Vigo D, Thornicroft G, Atun R. Estimating the true global burden of mental illness. Lancet Psychiatry. 2016;3(2):171–8.
- 9. Peters DH, Adam T, Alonge O, Agyepong IA, Tran N. Implementation research: what it is and how to do it. BMJ. 2013;347:f6753.
- Thornicroft G. Evidence-based mental health care and implementation science in low- and middle-income countries. Epidemiol Psychiatr Sci. 2012;21(3):241–4.
- Jacob KS, Sharan P, Mirza I, Garrido-Cumbrera M, Seedat S, Mari JJ, et al. Mental health systems in countries: where are we now? Lancet. 2007;370(9592):1061–77.
- Draper CE, Lund C, Kleintjes S, Funk M, Omar M, Flisher AJ. Mental health policy in South Africa: development process and content. Health Policy Plan. 2009;24(5):342–56.
- 13. Bird P, Omar M, Doku V, Lund C, Nsereko JR, Mwanza J. Increasing the priority of mental health in Africa: findings from qualitative research in Ghana, South Africa, Uganda and Zambia. Health Policy Plan 2010.
- Saraceno B, Van OM, Batniji R, Cohen A, Gureje O, Mahoney J, et al. Barriers to improvement of mental health services in low-income and middleincome countries. Lancet. 2007;370(9593):1164–74.
- Petersen I, Lund C, Stein DJ. Optimizing mental health services in low-income and middle-income countries. Curr Opin Psychiatry. 2011;24(4):318–23.

- 16. Saraceno B, van Ommeren M, Batniji R, Cohen A, Gureje O, Mahoney J, et al. Barriers to improvement of mental health services in low-income and middle-income countries. Lancet. 2007;370(9593):1164–74.
- Swanson RC, Cattaneo A, Bradley E, Chunharas S, Atun R, Abbas KM, et al. Rethinking health systems strengthening: key systems thinking tools and strategies for transformational change. Health Policy Plan. 2012;27(Suppl 4):iv54–61.
- WHO. Health systems: improving performance. Geneva: World Health Organization;2000.
- Da Silva MCA, Patel V. Evaluations of effectiveness in the real world. In: Thornicroft GPV, editor. Global mental health trials. Oxford: Oxford University Press; 2014.
- 20. Hanlon C, Fekadu A, Jordans M, Kigozi F, Petersen I, Shidhaye R, et al. District mental healthcare plans for five low- and middle-income countries: commonalities, variations and evidence gaps. Br J Psychiatry. 2015.
- 21. Rowe LA, Brillant SB, Cleveland E, Dahn BT, Ramanadhan S, Podesta M, et al. Building capacity in health facility management: guiding principles for skills transfer in Liberia. Hum Resour Health. 2010;8:5.
- 22. World Health Organization. Mental health systems in selected low- and middle-income countries: a WHO-AIMS cross-national analysis: WHO; 2009.
- Curry LA, Alpern R, Webster TR, Byam P, Zerihun A, Tarakeshwar N, et al. Community perspectives on roles and responsibilities for strengthening primary health care in rural Ethiopia. Glob Public Health. 2012;7(9):961–73.
- 24. Grimshaw J, Eccles M, Thomas R, MacLennan G, Ramsay C, Fraser C, et al. Toward evidence-based quality improvement. Evidence (and its limitations) of the effectiveness of guideline dissemination and implementation strategies 1966–1998. J Gen Intern Med. 2006;21(Suppl 2):S14–20.
- 25. Bauer MS. A review of quantitative studies of adherence to mental health clinical practice guidelines. Harv Rev Psychiatry. 2002;10(3):138–53.
- Weinmann S, Koesters M, Becker T. Effects of implementation of psychiatric guidelines on provider performance and patient outcome: systematic review. Acta Psychiatr Scand. 2007;115(6):420–33.
- 27. Barbui C, Girlanda F, Ay E, Cipriani A, Becker T, Koesters M. Implementation of treatment guidelines for specialist mental health care. Cochrane Database Syst Rev. 2014;1:CD009780.
- Girlanda F, Fiedler I, Ay E, Barbui C, Koesters M. Guideline implementation strategies for specialist mental healthcare. Curr Opin Psychiatry. 2013;26(4):369–75.
- Barbui C, Dua T, van Ommeren M, Yasamy MT, Fleischmann A, Clark N, et al. Challenges in developing evidence-based recommendations using the GRADE approach: the case of mental, neurological, and substance use disorders. Plos Med. 2010;7(8):e1000322.
- Dua T, Barbui C, Clark N, Fleischmann A, Poznyak V, van Ommeren M, et al. Evidence-based guidelines for mental, neurological, and substance use disorders in low- and middle-income countries: summary of WHO recommendations. Plos Med. 2011;8(11):e1001122.
- 31. World Health Organization. WHO mental health atlas. Geneva: World Health Organization; 2011.
- 32. HJPTG S, editor. Cochrane Handbook for Systematic Reviews of Interventions. 2011.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Reprint—preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. Phys Ther. 2009;89(9):873–80.
- Medicine Io. Clinical practice guidelines we can trust. Institute of Medicine; 2011.
- EPOC. Effective practice and organisation of care (EPOC) resources for review authors. Oslo: Norwegian Knowledge Centre for the Health Services; 2015. http://epoc.cochrane.org/ epoc-specific-resources-review-authors.
- 36. EPOC. EPOC taxonomy. 2015. https://epoc.cochrane.org/epoc-taxonomy.
- Berwanger O, Guimaraes HP, Laranjeira LN, Cavalcanti AB, Kodama AA, Zazula AD, et al. Effect of a multifaceted intervention on use of evidencebased therapies in patients with acute coronary syndromes in Brazil: the BRIDGE-ACS randomized trial. JAMA. 2012;307(19):2041–9.
- Du X, Gao R, Turnbull F, Wu Y, Rong Y, Lo S, et al. Hospital quality improvement initiative for patients with acute coronary syndromes in China: a cluster randomized, controlled trial. Circ Cardiovasc Qual Outcomes. 2014;7(2):217–26.

- Shrestha N, Samir KC, Baltussen R, Kafle KK, Bishai D, Niessen L. Practical approach to lung health in Nepal: better prescribing and reduction of cost. Trop Med Int Health. 2006;11(5):765–72.
- Steyn K, Lombard C, Gwebushe N, Fourie JM, Everett-Murphy K, Zwarenstein M, et al. Implementation of national guidelines, incorporated within structured diabetes and hypertension records at primary level care in Cape Town, South Africa: a randomised controlled trial. Glob Health Action. 2013;6:20796.
- Pagaiya N, Garner P. Primary care nurses using guidelines in Thailand: a randomized controlled trial. Trop Med Int Health. 2005;10(5):471–7.
- Okasha AFM, Haggag W, et al. A psychiatric training programme for general practitioners in primary health care in Egypt. Prim Care Psychiatry. 2002;8(1):9–16.
- Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. Adm Policy Ment Health. 2011;38(2):65–76.
- Deng Y, Liang L, Liu J, Jiang H. Analysis of monitoring results of iodine deficiency disorders in Ankang, Shaanxi province in 2012. Chin J Control Endemic Dis. 2013;28(4):284–6.
- Siddiqi K, Newell J, Robinson M. Getting evidence into practice: what works in developing countries? Int J Qual Health Care. 2005;17(5):447–54.
- Okasha A, Fahmy M, Haggag W, Awad M, Okasha T, Moez K. A psychiatric training programme for general practitioners in primary health care in Egypt. Primary Care Psychiatry. 2002;8(1):9–16.
- Brusamento S, Legido-Quigley H, Panteli D, Turk E, Knai C, Saliba V, et al. Assessing the effectiveness of strategies to implement clinical guidelines for the management of chronic diseases at primary care level in EU Member States: a systematic review. Health Policy. 2012;107(2/3):168–83.

- Rowe AK, Onikpo F, Lama M, Osterholt DM, Rowe SY, Deming MS. A multifaceted intervention to improve health worker adherence to integrated management of childhood illness guidelines in Benin. Am J Public Health. 2009;99(5):837–46.
- Rowe AK, Osterholt DM, Kouame J, Piercefield E, Herman KM, Onikpo F, et al. Trends in health worker performance after implementing the integrated management of childhood illness strategy in Benin. Trop Med Int Health. 2012;17(4):438–46.
- Joshi R, Alim M, Kengne AP, Jan S, Maulik PK, Peiris D, et al. Task shifting for non-communicable disease management in low and middle income countries—a systematic review. Plos ONE. 2014;9(8):e103754.
- Petersen I, Lund C, Bhana A, Flisher AJ. A task shifting approach to primary mental health care for adults in South Africa: human resource requirements and costs for rural settings. Health Policy Plan. 2012;27(1):42–51.
- Du X, Gao R, Turnbull F, Wu Y, Rong Y, Lo S, et al. Hospital quality improvement initiative for patients with acute coronary syndromes in China: a cluster randomized, controlled trial. Circ Cardiovasc Qual Outcomes. 2014;7(2):217–26.
- 53. World Health Organization. mhGAP intervention guide for mental, neurological and substance use disorders in non-specialized health settings: mental health Gap Action Programme (mhGAP). Geneva: WHO; 2010.
- 54. Siriwardhana C, Adikari A, Van Bortel T, McCrone P, Sumathipala A. An intervention to improve mental health care for conflict-affected forced migrants in low-resource primary care settings: a WHO MhGAP-based pilot study in Sri Lanka (COM-GAP study). Trials. 2013;14:423.
- Abdulmalik J, Kola L, Fadahunsi W, Adebayo K, Yasamy MT, Musa E, et al. Country contextualization of the mental health gap action programme intervention guide: a case study from Nigeria. PLoS Med. 2013;10(8):e1001501.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Submit your manuscript at www.biomedcentral.com/submit

• Maximum visibility for your research

