



King's Research Portal

DOI: 10.1016/j.scr.2015.12.047

Document Version Publisher's PDF, also known as Version of record

Link to publication record in King's Research Portal

Citation for published version (APA):

Devito, L., Petrova, A., Wood, V., Kadeva, N., Cornwell, G., Codognotto, S., Stephenson, E., & Ilic, D. (2016). Generation of KCL033 clinical grade human embryonic stem cell line. *Stem Cell Research*, *16*(2), 296-299. https://doi.org/10.1016/j.scr.2015.12.047

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

•Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research. •You may not further distribute the material or use it for any profit-making activity or commercial gain •You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Contents lists available at ScienceDirect

Stem Cell Research



Lab Resource: Stem Cell Line

Generation of KCL033 clinical grade human embryonic stem cell line



Liani Devito, Anastasia Petrova, Victoria Wood, Neli Kadeva, Glenda Cornwell, Stefano Codognotto, Emma Stephenson, Dusko Ilic*

Stem Cell Laboratories, Division of Women's Health, Faculty of Life Sciences and Medicine, King's College London and Assisted Conception Unit, Guys' Hospital, London, United Kingdom

ARTICLE INFO

Article history: Received 28 December 2015 Accepted 29 December 2015 Available online 15 January 2016

ABSTRACT

The KCL033 human embryonic stem cell line was derived from a normal healthy blastocyst donated for research. The ICM was isolated using laser microsurgery and plated on γ -irradiated human foreskin fibroblasts. Both the derivation and cell line propagation were performed in an animal product-free environment and under current Good Manufacturing Practice (cGMP) standards. Pluripotent state and differentiation potential were confirmed by in vitro assays. The line was also validated for sterility and specific and non-specific human pathogens. © 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

(continued)

Resource table

Name of stem cell line KCL033 Institution Derivation team King's College London, London, UK Neli Kadeva, Victoria Wood, Glenda C Codognotto, Emma Stephenson Contact person and email Dusko Ilic, email: dusko.ilic@kcl.ac.uk archived/stock date Type of resource Sub-type Biological reagent: cell line Human pluripotent stem cell line Pluripotent stem cell markers: NANO Reverssion Key marker expression Pluripotent stem cell markers: NANO Identity and purity of line confirmed 1) Jacquet, L., Stephenson, E., Collins, Al-Bedaery, R., Renwick, P., Ogilvi (direct URL links and full references) Jacquet, L., Stephenson, E., Collins, Al-Bedaery, R., Renwick, P., Ogilvi banks that HLA-match a target po Med. 5 (1), 10–17. doi: 10.1002/emmm.201201973 http://www.ncbi.nlm.nih.gov/pub 2) Canham, A., Van Deusen, A., Briso Downie, J., Devito, L., Hewitt, Z.A., Moore, H.D., Murray, H., Kunath,	KCI 033		 karyotype of 25 clinical-grade human embryonic stem cells lines. <i>Sci. Rep.</i> 5, 17258. doi: 10.1038/srep17258 http://www.ncbi.nlm.nih.gov/pubmed/26607962 3) Ilic, D., Stephenson, E., Wood, V., Jacquet, L., Stevenson, D., Petrova, A., Kadeva, N., Codognotto, S., Patel, H., Semple, M., Cornwell, G., Ogilvie, C., Braude, P., 2012. Derivation and feeder-free propagation of human embryonic stem cells under xeno-free conditions. Cytotherapy. 14 (1), 122–128. doi: 10.3109/14653249.2011.623692 http://www.ncbi.nlm.nih.gov/pubmed/22029654
	King's College London, London, UK Neli Kadeva, Victoria Wood, Glenda Cornwell, Stefano Codognotto, Emma Stephenson Dusko Ilic, email: dusko.ilic@kcl.ac.uk Aug 17, 2011		
	 Biological reagent: cell line Human pluripotent stem cell line Human embryo Pluripotent stem cell markers: NANOG, OCT4, TRA-1-60, TRA-1-81, alkaline phosphatase (AP) activity Identity and purity of line confirmed 1) Jacquet, L., Stephenson, E., Collins, R., Patel, H., Trussler, J., Al-Bedaery, R., Renwick, P., Ogilvie, C., Vaughan, R., Ilic, D., 2013. Strategy for the creation of clinical grade hESC line banks that HLA-match a target population. EMBO Mol. Med. 5 (1), 10–17. doi: 10.1002/emmm.201201973 http://www.ncbi.nlm.nih.gov/pubmed/23161805 		 4) Stephenson, E., Jacquet, L., Miere, C., Wood, V., Kadeva, N., Cornwell, G., Codognotto, S., Dajani, Y., Braude, P., Ilic, D., 2012. Derivation and propagation of human embryonic stem cell lines from frozen embryos in an animal product-free environment. Nat. Protoc. 7 (7), 1366–1381. doi: 10.1038/nprot.2012.080 http://www.ncbi.nlm.nih.gov/pubmed/22722371 5) Devito, L., Petrova, A., Miere, C., Codognottom S., Blakely, N., Lovatt, A., Ogilvie, C., Khalaf, Y., Ilic, D., 2014. Cost-effective master cell bank validation of multiple clinical-grade human pluripotent stem cell lines from a single donor. Stem Cells Transl. Med. 3(10), 1116–1124. doi: 10.5966/sctm 2014-0015
	 Canham, A., Van Deusen, A., Brison, D.R., De Sousa, P., Downie, J., Devito, L., Hewitt, Z.A., Ilic, D., Kimber, S.J., Moore, H.D., Murray, H., Kunath, T., 2015. The molecular 	Information in public databases	http://www.ncbi.nlm.nih.gov/pubmed/25122690 KCL033 is a National Institutes of Health (NIH) registered hESC line NIH Registration Number: NIHhESC-14-0267
		Ethics	http://grants.nih.gov/stem_cells/registry/current.htm?id=653 The hESC line KCL033 is derived under license from the UK Human Fertilisation and Embryology Authority (research

(continued on next page)

* Corresponding author.

http://dx.doi.org/10.1016/j.scr.2015.12.047

1873-5061/© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

(continued)

license numbers: R0075 and R0133) and also has local ethical approval (UK National Health Service Research Ethics Committee Reference: 06/Q0702/90). Informed consent was obtained from all subjects and the experiments conformed to the principles set out in the WMA Declaration of Helsinki and the NIH Belmont Report. No financial inducements are offered for donation.

Resource details

Consent signed	May 26, 2009
Embryo thawed	Jul 11, 2011
UK stem cell bank deposit approval	Mar 08, 2012
	Reference: SCSC12-54
Sex	Female 46, XX
Grade	Clinical
Disease status	Healthy/Unaffected
Karyotype (aCGH)	No copy number changes detected.
SNP array	Gain in regions 5p14.3 and 12p11.21
	(Canham et al., 2015)
DNA fingerprint	Allele sizes (in bp) of 16 microsatellite
	markers specific for chromosomes 13,
	18 and 21 (Jacquet et al., 2013)
HLA typing	HLA-A 11,29; B 44,51; Bw 4; C 04,16;
	DRB1 04,07; DRB4 01; DQB1 02,03
	(Jacquet et al., 2013; Canham et al.,
	2015)
Viability testing	Pass
Mycoplasma	Negative
Sterility	Pass
Pluripotent markers	NANOG, OCT4, TRA-1-60, TRA-1-81
(immunostaining)	
(Fig. 1)	
Three germ layers differentiation	Endoderm: AFP
in vitro (immunostaining)	Ectoderm: TUBB3 (tubulin, beta 3 class
(Fig. 2)	III)
	Mesoderm: ACTA2 (actin, alpha 2,
	smooth muscle)
Sibling lines available	KCL032, KCL034

We generated KCL033 clinical grade hESC line following protocols, established previously (Ilic et al., 2012; Stephenson et al., 2012), and now adapted to cGMP conditions. The expression of the pluripotency markers was tested after freeze/thaw cycle (Fig. 1). Differentiation potential into three germ layers was verified in vitro (Fig. 2).

Molecular karyotyping identified a novel 2.4 Mb gain on chromosome 5p14.3 and a gain on chromosome 12p11.21, which was also found in KCL040.

The gain on chromosome 5p14.3 containing a single gene, *CDH18* (Cadherin-18), was also present in one of two sibling cell lines, KCL032, but not in KCL034, a third sibling. A duplication of this size has not been reported to date, but its presence in two sibling hESC lines strongly suggests that it was inherited from one of the parents rather than by acquisition during hESC derivation and culture (Canham et al., 2015). The 2498.8 bp gain starts at bp 19086546 and ends at bp 21585311 as referred to Human Genome Build 38.

The gain on chromosome 12p11.21 contains no genes and it has been also reported in at least 14 submissions at Database of Genomic Variants (DGV; http://dgv.tcag.ca), which has collected structural variations in more than 14,000 healthy individuals from worldwide population (MacDonald et al., 2014). Estimated frequency in the human population is 4.70% (Canham et al., 2015).

Validation for sterility and specific and non-specific human pathogens (Devito et al., 2014) conformed that the cells in the Master Bank were sterile, mycoplasma-free, and negative for *Treponema pallidum*, chlamydia, *Neisseria gonorrhoeae*, human immunodeficiency virushuman cytomegalovirus (hCMV), human parvovirus B19, SV40, JCV, BKV, enterovirus, HAV, HCV, nonspecific viral and other adventitious contaminants.

We also generated research grade of KCL033 line that is adapted to feeder-free conditions.

Materials and methods

Consenting process

We distribute Patient Information Sheet (PIS) and consent form to the in vitro fertilization (IVF) patients if they opted to donate to research embryos that were stored for 5 or 10 years. They mail signed consent back to us and that might be months after the PIS and consent were mailed to them. If in meantime new versions of PIS/consent are implemented, we do not send these to the patients or ask them to resign; the whole process is done with the version that was given them initially. The PIS/consent documents (FRO-V.6) were created on Dec. 18, 2008. HFEA Code of Practice that was in effect at the time of document creation: Edition 7 – R.4 (http://www.hfea.gov.uk/2999.html). The donor couple signed the consent on May 26, 2009. HFEA Code of Practice that was in effect at the time of donor signature: Edition 7 – R.4. HFEA Code of Practice Edition 7 – R.4 was in effect: 02 Oct. 2008– 30 Sep. 2009.

Embryo culture and micromanipulation

Embryo culture and laser-assisted dissection of inner cell mass (ICM) were carried out as previously described in detail (Ilic et al., 2012; Stephenson et al., 2012). The cellular area containing the ICM was then washed and transferred to plates containing mitotically inactivated human neonatal foreskin fibroblasts (HFF).

Cell culture

ICM plated on mitotically inactivated HFF were cultured as described (llic et al., 2012; Stephenson et al., 2012). TE cells were removed mechanically from the outgrowth (llic et al., 2007; llic et al., 2010). hES colonies were expanded and cryopreserved at the third passage.



Fig. 1. Expression of pluripotency markers. Pluripotency is confirmed by immunostaining (Oct. 4, Nanog, TRA-1-60, TRA-1-81). Scale bar, 50 μm.



Fig. 2. Differentiation of three germ layers in vitro is confirmed by detection of markers: smooth muscle actin (ACTA2, red) for mesoderm, β-III tubulin (TUBB3, red) for ectoderm and α-fetoprotein (AFP, red) for endoderm. Nuclei are visualized with Hoechst 33342 (blue). Scale bar, 50 µm.

Viability test

Straws with the earliest frozen passage (p. 2–3) are thawed and new colonies are counted three days later. These colonies are then expanded up to passage 8, at which point cells were part frozen and part subjected to standard battery of tests (pluripotency markers, in vitro and in vivo differentiation capability, genetics, sterility, mycoplasma).

Pluripotency markers

Pluripotency was assessed with immunostaining for pluripotency markers as described (llic et al., 2012; Stephenson et al., 2012).

Differentiation

Spontaneous differentiation into three germ layers was assessed in vitro and in vivo as described (Petrova et al., 2014; Stephenson et al., 2012).

Genotyping

DNA was extracted from hES cell cultures using a Chemagen DNA extraction robot according to the manufacturer's instructions. Amplification of polymorphic microsatellite markers was carried out as described (llic et al., 2012). Allele sizes were recorded to give a unique fingerprint of each cell line.

Array comparative genomic hybridization (aCGH)

aCGH was performed as described in detail (Ilic et al., 2012).

Whole-genome single nucleotide polymorphism (SNP) array

SNP array was performed as described in detail (Canham et al., 2015).

HLA typing

HLA-A, -B and -DRB1 typing was performed with a PCR sequencespecific oligonucleotide probe (SSOP; Luminex, Austin, TX, USA) hybridization protocol at the certified Clinical Transplantation Laboratory, Guy's and St Thomas' NHS Foundation Trust and Serco Plc. (GSTS) Pathology (Guy's Hospital, London, UK) as described (Jacquet et al., 2013). HLA typing was also performed independently by other group (Canham et al., 2015). Validation for sterility and specific and non-specific human pathogens

Validation for sterility and specific and non-specific human pathogens was performed as described (Devito et al., 2014). All validation studies were conducted by SGS Vitrology (Glasgow, U.K., http:// www.sgs.com), in compliance with the principles of GMP as set out in Directive 2003/94/EC for medicinal products for human use (Directive 2003/94/EC, 2003) and 91/412/EEC for veterinary medicinal products (Directive 91/412/EEC, 1991).

Sterility testing was performed in accordance with the current requirements of the European Pharmacopoeia, Section 2.6.1 Sterility, U.S. Pharmacopeia, 71. Sterility Tests, and International Conference on Harmonisation Topic Q5D guidelines.

Mycoplasma testing was performed in accordance with the current requirements of the European Pharmacopoeia, Section 2.6.7, Mycoplasmas.

All PCR-based assays used were compliant with the current edition of the European Pharmacopoeia, 2.6.21, Nucleic Acid Amplification Techniques.

Author disclosure statement

There are no competing financial interests in this study.

Acknowledgments

This work was supported by the UK Medical Research Council grants G0701172 and G0801061. We thank Dr. Yacoub Khalaf, Director of the Assisted Conception Unit of Guy's and St Thomas' NHS Foundation Trust and his staff for supporting the research program. We are especially indebted to Prof Peter Braude and patients who donated the embryos.

References

- Canham, A., Van Deusen, A., Brison, D.R., De Sousa, P., Downie, J., Devito, L., Hewitt, Z.A., Ilic, D., Kimber, S.J., Moore, H.D., Murray, H., Kunath, T., 2015. The molecular karyotype of 25 clinical-grade human embryonic stem cells lines. Sci. Rep. 5, 17258.
- Devito, L., Petrova, A., Miere, C., Codognottom S., Blakely, N., Lovatt, A., Ógilvie, C., Khalaf, Y., Ilic, D., 2014. Cost-effective master cell bank validation of multiple clinical-grade human pluripotent stem cell lines from a single donor. Stem Cells Transl Med. 3(10), 1116–1124.
- Directive 2002/94/EC, 2003. Laying down the principles and guidelines of good manufacturing practice in respect of medical products for human use and investigational medicinal products for human use. Off. J. Eur. Union L262, 22–26 (of the European Parliament and of the Council of October 08, October 14; 2003; Available at http://ec.europa.eu/halth/files/eudralex/vol-1/dir_2003_94/dir_2003_94_en.pdf).
- Directive 91/412/EEC, 1991. Laying down the principles and guidelines of good manufacturing practice for veterinary medicinal products. Off. J. Eur. Union L228, 70–73 (of July 23, August 17, 1991; Available at: http://ec.europa.eu/health/files/ eudralex/vol-5/dir_1991_412/dir_1991_412_en.pdf).
- Ilic, D., Genbacev, O., Krtolica, A., 2007. Derivation of hESC from intact blastocysts. Curr. Protoc. Stem Cell Biol. (Chapter 1: Unit 1 A.2).
- Ilic, D., Caceres, E., Lu, S., Julian, P., Foulk, R., Krtolica, A., 2010. Effect of karyotype on successful human embryonic stem cell derivation. Stem Cells Dev. 19 (1), 39–46.

- Ilic, D., Stephenson, E., Wood, V., Jacquet, L., Stevenson, D., Petrova, A., Kadeva, N., Codognotto, S., Patel, H., Semple, M., Cornwell, G., Ogilvie, C., Braude, P., 2012. Derivation and feeder-free propagation of human embryonic stem cells under xeno-free conditions. Cytotherapy 14 (1), 122–128.
- conditions. Cytotherapy 14 (1), 122–128.
 Jacquet, L., Stephenson, E., Collins, R., Patel, H., Trussler, J., Al-Bedaery, R., Renwick, P., Ogilvie, C., Vaughan, R., Ilic, D., 2013. Strategy for the creation of clinical grade hESC line banks that HLA-match a target population. EMBO Mol. Med. 5 (1), 10–17.
- Line banks that HLA-match a target population. EMBO Mol. Med. 5 (1), 10–17.
 MacDonald, J.R., Ziman, R., Yuen, R.K.C., Feuk, L., Scherer, S.W., 2014. The database of genomic variants: a curated collection of structural variation in the human genome. Nucleic Acids Res. 42, D986–D992.
- Petrova, A., Celli, A., Jacquet, L., Dafou, D., Crumrine, D., Hupe, M., Arno, M., Hobbs, C., Cvoro, A., Karagiannis, P., Devito, L., Sun, R., Adame, L.C., Vaughan, R., McGrath, J.A., Mauro, T.M., Ilic, D., 2014. 3D In vitro model of a functional epidermal permeability barrier from human embryonic stem cells and induced pluripotent stem cells. Stem Cell Rep. 2 (5), 675–689.
- Stephenson, E., Jacquet, L., Miere, C., Wood, V., Kadeva, N., Cornwell, G., Codognotto, S., Dajani, Y., Braude, P., Ilic, D., 2012. Derivation and propagation of human embryonic stem cell lines from frozen embryos in an animal product-free environment. Nat. Protoc. 7 (7), 1366–1381.