



King's Research Portal

DOI:

[10.1016/j.jgo.2017.02.006](https://doi.org/10.1016/j.jgo.2017.02.006)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Raimi-Abraham, B. T., de Orbe Izquierdo, M. S., Collignon, O., & Cerreta, F. (2017). Regulatory Considerations on the Enrollment of Older Adults in Oncology Clinical Trials. *Journal of Geriatric Oncology*, 151-153. Advance online publication. <https://doi.org/10.1016/j.jgo.2017.02.006>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1 **Regulatory Considerations on the Enrollment of Older Adults in Oncology**
2 **Clinical Trials**

3 Bahijja Tolulope Raimi-Abraham^{1,4±}, Maria Silvia de Orbe Izquierdo^{2,4}, Olivier Collignon^{3,4},
4 Francesca Cerreta^{4*}

5 ¹ University College London School of Pharmacy, 29-39 Brunswick Square, London WC1N
6 1AX.

7 [±]Present address: King's College London, Faculty of Life Sciences and Medicine, Institute of
8 Pharmaceutical Science, Drug Delivery Group, Franklin-Wilkins Building, 150 Stamford
9 Street, London SE1 9NH, United Kingdom.

10

11 ² Fundación para la Investigación Biomédica del Hospital Universitario Ramón y Cajal.
12 Instituto Ramón y Cajal de Investigación Sanitaria (IRYCIS).Ctra. de Colmenar Viejo, Km.
13 9,100. 28034 Madrid.

14 ³ Luxembourg Institute of Health, 1A rue Thomas Edison, L1445, Strassen, Luxembourg.

15 ⁴ European Medicines Agency, 30 Churchill Place, London E14 5E.

16 *Corresponding author: Francesca.Cerreta@ema.europa.eu

17

1 **Introduction**

2 The International Conference of Harmonisation (ICH) E7 efficacy guideline “Studies in
3 support of special populations: geriatrics” was adopted in 1993 by the regulatory bodies of
4 Europe, USA and Japan, stating that “*a minimum of 100 patients would usually allow detection*
5 *of clinically important differences*” and that “*it is also important not to exclude unnecessarily*
6 *patients with concomitant illness*”. [1] Since then, the world’s population has been aging
7 rapidly: in the United States, the percentages of patients with cancer who are older than 65, 70
8 and 75 years are, respectively, 60%, 46% and 31%. [2]. Cancer incidence in the over 65 years
9 category is expected to increase dramatically by 67% from 2010 to 2030. [3] Restrictive
10 eligibility criteria such as arbitrary upper age limits or exclusion criteria based on
11 comorbidities, polypharmacy or reduced life expectancy, have resulted in the exclusion of older
12 people from clinical trials. [4]

13 In 2006, the European Commission requested the European Medicines Agency (EMA) to
14 provide their opinion on the ‘adequacy of guidance on the elderly regarding medicinal products
15 for human use’: [5] 56 drug development guidelines were reviewed and analyzed together with
16 10 recently approved medicine dossiers in indications relevant to older patients.
17 Recommendations in the report included an update of ICH E7 for better representation of the
18 target population. [5]

19 This resulted in the adoption in 2010 of the addendum to ICH E7 in the form of a questions
20 and answers (Q&A) document underlining that the number of geriatric patients to be included
21 in a marketing application should be influenced by the target patient population and expected
22 use: [6] “... *it is very important to ensure, to the extent possible, that the population included*
23 *in the clinical development program is representative of the target patient population and that*
24 *in the marketing application, depending on the numbers of patients, data should be presented*

1 *for various age groups (i.e.<65, 65-74, 75-84 and > 85) to assess the consistency of the*
2 *treatment effect and safety profile in these patients with the non-geriatric patient population.”*

3 However, recruiting in a Phase 3 trial a patient sample whose age structure is similar to the
4 target population can be challenging, and even so, the resulting smaller number of older
5 patients would not necessarily guarantee sufficient power to show significant efficacy in that
6 subgroup: this is compounded by the greater variability inherent to the older population.

7 Therefore in 2011 the EMA adopted its Geriatric Medicines Strategy, [7] which aims, when
8 granting a marketing authorization for a new medicine, to consider its benefit/risk balance in
9 relation to the expected use, and to clearly communicate the conclusions (in terms of findings
10 and/or request for post-authorization data) in the EMA approval documents.

11 As the world’s population is aging, the relevance of this approach has been underlined in a
12 recent paper [8] where survival gains demonstrated in a clinical trial setting have not been
13 verified in an older real-world population with more comorbidities.


14 **Representation of older patients in clinical trials: two recently approved drugs**

15 We have considered the proportion of older people included in clinical trials for two recently
16 approved oncology medicines in indications highly prevalent in the older population: Xofigo®
17 (radium-223 dichloride) for prostate cancer, and Perjeta® (pertuzumab) for breast cancer. Both
18 drugs were approved in 2013 by the EMA. In terms of age groups, there was good
19 representation of older patients with lower representation only of patients older than 85 years.


20 This appears reasonable for the clinical trial population of a Phase 3 trial, where the aim is to
21 characterize the benefit/risk of a medicine without exposing potentially frailer patients in the
22 initial phase of development. [9] These data satisfy the requirements of ICH E7 Q&A. A more
23 in depth-look at the characteristics of these older patients shows that the protocol of Xofigo®

1 pivotal study specified ECOG PS ≥ 3 as exclusion criterion (i.e. “capable of only limited self-
 2 care, confined to bed, more than 50% of walking hours”), while the Perjeta® pivotal study
 3 excluded patients with ECOG PS ≥ 2 (i.e. “ambulatory and capable of all self-care but unable
 4 to carry out any work activities, up and about more than 50% of waking hours”) [10]: this might
 5 have contributed to the fact that for Perjeta® enrolment in the oldest age bracket was lower,
 6 reflecting a known enrolment trend for older-old women in breast cancer trials. [9]

7



8



1 **Figure 1.** a) Prostate cancer and b) breast cancer drug % (1dp) of elderly patients (i.e. 65 to
2 85+ years) in PK, PD, efficacy and safety studies.

3

4 An analysis of the exclusion criteria in 25 phase II and III clinical trials for breast cancer (i.e.
5 Perjeta®, Halaven®, Tyverb®, Kadcylla®, Afinitor®, Herceptin®, Avastin®, Abraxane®)
6 showed that age limits are, however, still present as an exclusion criterion in many protocols,
7 and only one of these pivotal clinical trials considered patients with ECOG PS 3 to be eligible.

8 It is important that the information on frailty or ECOG status of the studied population is clearly
9 communicated: achieving “real evidence based medicine” for the ageing population will
10 constitute the challenge for the coming years.[8, 10]

11 **Who can help?**

12 Companies should take into account the recommendations in ICH E7 Q&A when designing
13 clinical trials. While every effort should be made to include geriatric patients using concomitant
14 therapies and with co-morbidities in the premarketing clinical development program, it is
15 recognized that enrolment of these patients can be challenging, particularly in the initial phases
16 of drug development: if this is the case, a specific plan to collect data post-marketing should
17 be discussed with the regulators during development, and presented in the marketing
18 application. To assist characterization of these patients beyond chronological age, the EMA is
19 developing guidance on frailty assessment tools which can be used as demographic
20 stratification parameters for older patients included in clinical trials. [11]

21 Patient preferences play an increasingly important part in drug development, and the relevance
22 of additional trial endpoints, such as QoL, could be considered for the older population. [12]

1 Lastly, to support physicians and patients in their therapeutic decisions, there should be clear
2 communication of the available evidence, on the conclusions of the benefit risk in the
3 population of expected use, and of any post-authorization data collection required.

4

5 **Acknowledgements**

6 Dr Bahijja Tolulope Raimi-Abraham would like to thank and acknowledge the financial
7 support from University College London (UCL) EPSRC IAA account and policy engagement
8 support from UCL Public Policy towards her secondment at the European Medicines Agency
9 (EMA).

10

11 **Disclaimer**

12 The views presented in this article are those of the authors and should not be understood or
13 quoted as being made on behalf of the European Medicines Agency and/or its scientific
14 committees.

15

16 **References**

- 17 1. *ICH Topic E7. Studies in support of special populations: geriatrics*. 1994 [20/12/2016];
18 Available from: Available at
19 [www.ema.europa.eu/ema/index.jsp?curl=pages/includes/document/document_detail.jsp?web](http://www.ema.europa.eu/ema/index.jsp?curl=pages/includes/document/document_detail.jsp?webContentId=WC500002875&mid=WC0b01ac058009a3dc)
20 [ContentId=WC500002875&mid=WC0b01ac058009a3dc](http://www.ema.europa.eu/ema/index.jsp?curl=pages/includes/document/document_detail.jsp?webContentId=WC500002875&mid=WC0b01ac058009a3dc).
- 21 2. Scher, K.S. and A. Hurria, *Under-Representation of Older Adults in Cancer Registration*
22 *Trials: Known Problem, Little Progress*. *Journal of Clinical Oncology*, 2012. **30**(17): p. 2036-
23 2038.
- 24 3. Smith, B.D., et al., *Future of Cancer Incidence in the United States: Burdens Upon an Aging,*
25 *Changing Nation*. *Journal of Clinical Oncology*, 2009. **27**(17): p. 2758-2765.

- 1 4. Crome, P., A. Cherubini, and J. Oristrell, *The PREDICT (increasing the participation of the*
2 *elderly in clinical trials) study: the charter and beyond*. Expert Review of Clinical
3 Pharmacology, 2014. **7**(4): p. 457-468.
- 4 5. *European Medicines Agency, Committee for Medicinal Products for Human Use. Adequacy*
5 *of guidance on the elderly regarding medicinal products for human use.* . 2006 20/12/2016];
6 Available from: Available at
7 [http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2010/01/WC5](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2010/01/WC500049541.pdf)
8 [00049541.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2010/01/WC500049541.pdf).
- 9 6. *ICH topic E7. Studies in support of special populations: geriatrics questions and answers.*
10 2010 20/12/2016]; Available from: Available from:
11 [www.ema.europa.eu/ema/index.jsp?curl=pages/includes/document/document_detail.jsp?web](http://www.ema.europa.eu/ema/index.jsp?curl=pages/includes/document/document_detail.jsp?webContentId=WC500005218&mid=WC0b01ac058009a3dc)
12 [ContentId=WC500005218&mid=WC0b01ac058009a3dc](http://www.ema.europa.eu/ema/index.jsp?curl=pages/includes/document/document_detail.jsp?webContentId=WC500005218&mid=WC0b01ac058009a3dc). .
- 13 7. *The European Medicines Agency's Geriatric Medicines Strategy.* 2011 20/12/2016];
14 Available from: Available from
15 http://www.ema.europa.eu/docs/en_GB/document_library/Other/2011/02/WC500102291.pdf
- 16 8. Mailankody, S. and V. Prasad, *Overall survival in cancer drug trials as a new surrogate end*
17 *point for overall survival in the real world*. JAMA Oncology, 2016.
- 18 9. Kemeny, M.M., et al., *Barriers to Clinical Trial Participation by Older Women With Breast*
19 *Cancer*. Journal of Clinical Oncology, 2003. **21**(12): p. 2268-2275.
- 20 10. Tracey E Howe, S.M., Vivian Welch *Introducing Cochrane Global Ageing: towards a new*
21 *era of evidence[editorial]*. Cochrane Database of Systematic Reviews, 2016. **9**.
- 22 11. Maggioni, A.P., et al., *The real-world evidence of heart failure: findings from 41 413 patients*
23 *of the ARNO database*. European Journal of Heart Failure, 2016. **18**(4): p. 402-410.
- 24 12. Wildiers, H., et al., *Pharmacology of Anticancer Drugs in the Elderly Population*. Clinical
25 Pharmacokinetics, 2003. **42**(14): p. 1213-1242.

26