



## King's Research Portal

DOI:

[10.1002/oby.21922](https://doi.org/10.1002/oby.21922)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Menni, C., Migaud, M., Kastenmüller, G., Pallister, T., Zierer, J., Peters, A., Mohny, R. P., Spector, T. D., Bagnardi, V., Gieger, C., Moore, S. C., & Valdes, A. M. (2017). Metabolomic Profiling of Long-Term Weight Change: Role of Oxidative Stress and Urate Levels in Weight Gain. *Obesity*, 25(9), 1618-1624. Advance online publication. <https://doi.org/10.1002/oby.21922>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Metabolomic Profiling of Long-Term Weight Change: Role of Oxidative Stress and Urate Levels in Weight Gain

Cristina Menni<sup>1</sup>, Marie Migaud<sup>2</sup>, Gabi Kastenmüller<sup>1,3,4</sup>, Tess Pallister<sup>1</sup>, Jonas Zierer<sup>1,3</sup>, Annette Peters<sup>5</sup>, Robert P. Mohny<sup>6</sup>, Tim D. Spector<sup>1</sup>, Vincenzo Bagnardi<sup>7</sup>, Christian Gieger<sup>5</sup>, Steve C. Moore<sup>8</sup>, and Ana M. Valdes<sup>1,9,10</sup>

**Objective:** To investigate the association between long-term weight change and blood metabolites.

**Methods:** Change in BMI over  $8.6 \pm 3.79$  years was assessed in 3,176 females from the TwinsUK cohort (age range: 18.3–79.6, baseline BMI:  $25.11 \pm 4.35$ ) measured for 280 metabolites at follow-up. Statistically significant metabolites (adjusting for covariates) were included in a multivariable least absolute shrinkage and selection operator (LASSO) model. Findings were replicated in the Cooperative Health Research in the Region of Augsburg (KORA) study ( $n = 1,760$ ; age range: 25–70, baseline BMI:  $27.72 \pm 4.53$ ). The study examined whether the metabolites identified could prospectively predict weight change in KORA and in the Prostate, Lung, Colorectal, and Ovarian Cancer (PLCO) study ( $n = 471$ ; age range: 55–74, baseline BMI:  $27.24 \pm 5.37$ ).

**Results:** Thirty metabolites were significantly associated with change in BMI per year in TwinsUK using Bonferroni correction. Four were independently associated with weight change in the multivariable LASSO model and replicated in KORA: namely, urate (meta-analysis  $\beta$  [95% CI] = 0.05 [0.040 to 0.063];  $P = 1.37 \times 10^{-19}$ ), gamma-glutamyl valine ( $\beta$  [95% CI] = 0.06 [0.046 to 0.070];  $P = 1.23 \times 10^{-20}$ ), butyryl-carnitine ( $\beta$  [95% CI] = 0.04 [0.028 to 0.051];  $P = 6.72 \times 10^{-12}$ ), and 3-phenylpropionate ( $\beta$  [95% CI] =  $-0.03$  [ $-0.041$  to  $-0.019$ ];  $P = 9.8 \times 10^{-8}$ ), all involved in oxidative stress. Higher levels of urate at baseline were associated with weight gain in KORA and PLCO.

**Conclusions:** Metabolites linked to higher oxidative stress are associated with increased long-term weight gain.

Obesity (2017) 00, 00–00. doi:10.1002/oby.21922

<sup>1</sup> Department of Twin Research & Genetic Epidemiology, King's College London, London, UK <sup>2</sup> Mitchell Cancer Institute, University of South Alabama, Mobile, Alabama, USA <sup>3</sup> Institute of Bioinformatics and Systems Biology, Helmholtz Zentrum München, Neuherberg, Germany <sup>4</sup> German Center for Diabetes Research, Neuherberg, Germany <sup>5</sup> Institute of Epidemiology II, Helmholtz Zentrum München, Neuherberg, Germany <sup>6</sup> Metabolon, Inc., Durham, North Carolina, USA <sup>7</sup> Department of Statistics and Quantitative Methods, University of Milano-Bicocca, Milan, Italy <sup>8</sup> Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, Maryland, USA <sup>9</sup> School of Medicine, University of Nottingham, Nottingham, UK <sup>10</sup> National Institute for Health Research, Nottingham Biomedical Research Centre, Nottingham, UK. Correspondence: Ana M. Valdes (ana.valdes@nottingham.ac.uk)

**Funding agencies:** This work was supported by the FP7 project HEALS (Health and Environment-wide Associations based on Large population Surveys) Project No. 603946 of the European Union's Seventh Framework Programme and by the Medical Research Council Ancestry and Biological Informative Markers for Stratification of Hypertension grant (MR/M016560/1). Metabolomic analysis was funded by Pfizer. The TwinsUK research was funded by the Wellcome Trust and the European Community's Seventh Framework Programme (FP7/2007–2013). The study also received support from the National Institute for Health Research (NIHR) Clinical Research Facility at Guy's & St Thomas' National Health Service (NHS) Foundation Trust and NIHR Biomedical Research Centre, based at Guy's and St Thomas' NHS Foundation Trust and King's College London. TDS is an NIHR senior investigator and is holder of a European Research Council Advanced Principal Investigator award. AMV is funded by the NIHR Nottingham Biomedical Research Centre.

The Cooperative Health Research in the Region of Augsburg (KORA) research platform was initiated and financed by the Helmholtz Zentrum München German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research and by the State of Bavaria. Furthermore, KORA research was supported within the Munich Center of Health Sciences, Ludwig-Maximilians-Universität, as part of LMUinnovativ.

This work was also supported by the Intramural Research Program of the National Cancer Institute, National Institutes of Health, Department of Health and Human Services, and by the Breast Cancer Research Stamp Fund, awarded through competitive review.

**Disclosure:** RPH is an employee of Metabolon, Inc. All other authors declare no conflict of interest.

**Author contributions:** CM, TDS, and AMV conceived and designed the experiments; RPM performed the experiments; CM analyzed the data; GK and SCM analyzed the replication samples; TP, JZ, AP, CG, and VB contributed reagents/materials/analysis tools; and CM, MM, and AMV wrote the manuscript. All authors were involved in revising the manuscript and had final approval of the submitted version.

Additional Supporting Information may be found in the online version of this article.

**Received:** 23 January 2017; **Accepted:** 31 May 2017; **Published online** 00 Month 2017. doi:10.1002/oby.21922

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

## Introduction

Obesity is a major public health concern resulting from a positive energy balance for an extended period of time (1). Although most people in industrialized nations have overweight or obesity, significant numbers of people with normal weight never develop overweight or obesity, partly reflecting the large interindividual variation in response to excess caloric intake (1). A clear understanding of the mechanisms of obesity and development of effective strategies for its prevention and treatment require consideration of biochemical changes resulting from weight gain and loss of metabolic homeostasis (2).

Comprehensive metabolic profiling provides an opportunity to gain new insights into biochemical differences involved in weight loss or gain in humans. Several metabolomic studies have been carried out to date, although most of them have focused on the overlap between obesity and type 2 diabetes mellitus (3). Dozens of metabolites fall into that category, including branched-chain amino acids (BCAAs), cysteine, glutamine, phenylalanine, proline, tyrosine, threonine, tryptophan, choline, and pantothenic acid, which are increased in both obesity and diabetes. In contrast, asparagine, glycine, methionine, and citrulline are decreased in both obesity and diabetes (4). Metabolomic profiling of BMI and obesity independent of the link with diabetes has also been performed. In a cross-sectional study of 947 participants, 37 metabolites were significantly associated with BMI, including 19 lipids, 12 amino acids, and 6 others. Eighteen of these associated metabolites had not been previously reported on, including histidine and butyrylcarnitine, a lipid marker of whole-body fatty acid oxidation (5).

Similar results have been derived from smaller studies focused on child obesity. A study profiling serum samples of 80 children with obesity and 40 children with normal weight identified 14 metabolites (glutamine, methionine, proline, 9 phospholipids, and 2 acylcarnitines) that were significantly different when comparing children with normal weight and children with obesity (6).

Less research has been conducted to investigate the metabolites related not to high or low BMI but to change in BMI per year. One study investigated the metabolic adaptations seen after bariatric surgery. The metabolic footprint of bariatric procedures appears to be specifically characterized by an increase of bile acid circulating pools and a decrease of ceramide levels, a greater perioperative decline in BCAA, and the rise of circulating serine and glycine, mirroring glycemic control and inflammation improvement (7).

A recent study from the Cooperative Health Research in the Region of Augsburg (KORA) cohort defined metabolite modules or clusters of related molecules (using weighted correlation network) and then identified four metabolite modules significantly and stably associated with body weight gain. These included major branches of metabolism, with very low-density lipoprotein, low-density lipoprotein (LDL), and large high-density lipoprotein subclasses, triglycerides, BCAAs, and markers of energy metabolism, among others (8).

However, a key question is “What are the molecular pathways that are associated with greater or lower weight change over time?” This is relevant if these factors are to be addressed and eventually modified for therapeutic purposes.

In this study, we used longitudinal measures of BMI in 3,176 women from the TwinsUK cohort assessed 8.6 years apart. Having adjusted

for age and BMI at baseline, smoking, and family relatedness, we carried out metabolomic profiling of change in BMI per year. We replicated our results in 1,760 individuals from KORA and in an elderly cohort from the United States. Having identified metabolites whose levels are significantly and reproducibly altered by weight change, we then tested in a subanalysis whether the levels of any of these at baseline (before weight change) were also associated with weight change.

## Methods

### Discovery cohort

Study subjects were twins enrolled in the TwinsUK registry, a national register of adult twins recruited as volunteers without selecting for any particular disease or trait (9). All recruited twins were of the same sex. In this study, we analyzed data from 3,176 females (age range: 18.3-79.6) from the TwinsUK cohort with BMI assessed on average  $8.6 \pm 3.79$  years apart and metabolomic profiling conducted at follow-up.

The study was approved by St Thomas' Hospital Research Ethics Committee, and all twins provided informed written consent.

### Replication cohorts

The replication cohort consisted of individuals from the follow-up study (KORA F4) drawn from the general population of the region of Augsburg, Germany (10). The average follow-up time from the baseline study (KORA S4) was  $7.11 \pm 0.22$  years. In total, we analyzed 1,760 individuals (age range: 25-70) with fasting serum metabolomic profiles made available using the Metabolon, Inc. (Durham, North Carolina) platform and with the measure of longitudinal BMI.

We tested whether the identified and replicated metabolites could also predict weight change on a different set of fasting and nonfasting samples from the KORA S4 study, which comprised 1,069 subjects (age range: 32-77) with nontargeted metabolomic data at baseline and longitudinal BMI.

We also included 471 female controls (age range: 55-74) from the Prostate, Lung, Colorectal, and Ovarian Cancer (PLCO) study (11). Eligibility criteria were that participants had to be controls, had to have a follow-up weight measurement, and had to have metabolomic profiling available. For PLCO, metabolomic profiling was available only at baseline.

The PLCO trial is registered at ClinicalTrials.gov as NCT00002540, and it was approved by the institutional review boards of the US National Cancer Institute and the 10 screening centers. Only controls from this trial are included in this study. The KORA F4/S4 study was approved by the ethics committee of the Bavarian Medical Association.

### Assessment of weight gain, weight loss, and covariates

Height and weight were measured once at baseline and once at follow-up, which was on average  $8.6 \pm 3.79$  years after the baseline assessment. BMI was calculated by dividing weight (in kilograms) by the square of height (in meters).

For the subanalysis and for computing box plots of the metabolites' distribution, the change in BMI per year was calculated by adjusting for age, BMI at baseline, smoking, metabolite batch, and family relatedness. Subjects were then categorized based on these tertiles. The weight gain group was defined as the top tertile, while the low weight gain (weight loss) group was defined as the lower tertile (12). The middle tertile was excluded from the subanalysis.

Metabolomics measurements

Nontargeted metabolite detection and quantification were conducted by the metabolomics provider Metabolon on 3,176 fasting samples from participants in the TwinsUK study, as described previously (13) (for details, see the Supporting Information). The metabolomic data set measured by Metabolon includes 280 metabolites of known chemical identity containing the following broad categories: amino acids, acylcarnitines, sphingomyelins, glycerophospholipids, carbohydrates, vitamins, lipids, nucleotides, peptides, xenobiotics, and steroids.

Circulating levels of saturated fatty acids (SFAs), total fatty acids (TotFAs), monounsaturated fatty acids, and polyunsaturated fatty acids (PUFAs) were measured by Nightingale Health, previously known as Brainshake Ltd. (Vantaa, Finland; <https://nightingale-health.com/services>), from fasting serum samples using proton nuclear magnetic resonance spectroscopy at 500 MHz and 600 MHz, as previously described (14). Traits were natural log-transformed and then scaled to standard deviation (SD) units, as previously proposed by Würtz et al (15). For the metabolites containing zeros, one was added to all values of that metabolite before natural log-transformation.

Statistical analysis

Statistical analysis was carried out using the data analysis and statistical software Stata version 11 (StataCorp LLC, College Station, Texas) and R version 3.3.1 (R Foundation for Statistical Computing, Vienna, Austria). We inverse normalized the metabolite data, as the metabolite concentrations were not normally distributed. We excluded metabolic traits with more than 20% of values missing. We imputed individual metabolites not detected in a sample using the minimum run-day measures.

Linear regression analysis adjusting for age, BMI at baseline, smoking, metabolite batch, and family relatedness was undertaken to identify associations between biomarkers and weight change. We corrected for multiple testing using Bonferroni correction ( $P < 1.2 \times 10^{-4}$ ). To identify a set of metabolites independently associated with weight change, we fitted the least absolute shrinkage and selection operator method (LASSO), incorporating all significant metabolites together with age, smoking, and metabolite batch. LASSO is a regression method that performs both variable selection and regularization. It involves penalizing the absolute size of the regression coefficients with some parameter estimates shrunk to zero (16). The penalty parameter was chosen as the largest  $\lambda$  such that error is within one standard error of the minimum as determined by cross validation using the glmnet package in R (17).

Replication analysis

The metabolites independently associated with weight change were replicated in 1,760 individuals from the KORA cohort. Results were

TABLE 1 Descriptive characteristics of the study population

	TwinsUK	KORA	PLCO
Females (%)	100	51	100
Age at baseline	50.26 ± 10.24	53.72 ± 8.79	62.65 ± 5.12
Age at follow-up	58.86 ± 10.54	60.82 ± 8.79	71.74 ± 5.89
BMI at baseline	25.11 ± 4.35	27.72 ± 4.53	27.24 ± 5.37
BMI at follow-up	26.38 ± 4.76	28.17 ± 4.81	27.60 ± 5.71
Current smoker (%)	8.09	14.77	5.94

Values represent the mean (SD) unless otherwise specified.

then combined using an inverse variance fixed-effect meta-analysis. The fixed-effect model provides a weighted average of the study estimates, the weights being the inverse of the variance of the study estimate.

We computed the area under the curve (AUC) and 95% CIs for a receiver operating characteristic curve comparing the top and bottom tertiles of BMI change per year in the TwinsUK cohort. Two models were fitted: (1) demographics and lifestyle, including age, baseline BMI, and smoking and (2) all variables in (1) plus the inverse normal concentrations of the four successfully replicated metabolites. This allowed us to evaluate the extent to which the significant metabolites contribute to the AUC in addition to the demographic and lifestyle variables. As metabolites may also be predictive of BMI change per year, we finally tested the association of weight change and metabolites at baseline in both KORA and PLCO. Metabolomic data at baseline in twins were not available.

Results

The descriptive characteristics of the study participants are presented in Table 1. Thirty metabolites were associated with weight change after adjusting for covariates and multiple testing using Bonferroni correction. We investigated whether these associations were sensitive to the various adjustments and found that results were consistent (Supporting Information Tables S1-S2).

We then used a LASSO model to analyze these 30 metabolites, identifying 7 that contributed independently to weight change (Table 2). Of those, 4 were successfully replicated in the KORA cohort and were therefore meta-analyzed using inverse variance fixed-effect meta-analysis. Urate (meta-analysis  $\beta$  [95% CI] = 0.05 [0.040 to 0.063];  $P = 1.37 \times 10^{-19}$ ), gamma-glutamyl valine ( $\beta$  [95% CI] = 0.06 [0.046 to 0.070];  $P = 1.23 \times 10^{-20}$ ), and butyrylcarnitine ( $\beta$  [95% CI] = 0.04 [0.028 to 0.051];  $P = 6.72 \times 10^{-12}$ ) were positively associated with weight gain, while 3-phenylpropionate (hydrocinnamate) ( $\beta$  [95% CI] = -0.03 [-0.041 to -0.019];  $P = 9.8 \times 10^{-8}$ ) was inversely associated with weight change. Supporting Information Figure S1 shows the box plot of the distribution of the four metabolite concentrations (in the inverse normal scale) for each of the three tertiles of change in BMI per year. Tertile 1 refers to individuals with the lowest BMI change per year, while tertile 3 refers to the highest BMI gain over time.



TABLE 2 List of metabolites associated with weight change in the LASSO model										
Super	Sub	Metabolite	TwinsUK			KORA			Fixed-effect meta-analysis	
			$\beta^a$	SE	P	$\beta^a$	SE	P	$\beta$ (95% CI)	P
Amino acid	Phenylalanine and tyrosine metabolism	3-Phenylpropionate (hydrocinnamate)	−0.03	0.006	$3.01 \times 10^{-6}$	−0.03	0.007	$3.33 \times 10^{-6}$	−0.03 (−0.041 to −0.019)	$9.8 \times 10^{-8}$
Cofactors and vitamins	Hemoglobin and porphyrin metabolism	Bilirubin (Z,Z isomer)	−0.03	0.006	$2.30 \times 10^{-7}$	−0.09	0.007	$1.91 \times 10^{-1}$		
Lipid	Fatty acid metabolism (also BCAA metabolism)	Butyrylcarnitine	0.04	0.007	$7.88 \times 10^{-9}$	0.04	0.007	$8.63 \times 10^{-9}$	0.04 (0.028 to 0.051)	$6.72 \times 10^{-12}$
Lipid	Lysolipid	1-Docosahexaenoyl-GPC (22:6)	−0.04	0.007	$5.19 \times 10^{-9}$	−0.02	0.007	$1.20 \times 10^{-2}$		
Nucleotide	Purine and urate metabolism	Urate	0.05	0.007	$3.04 \times 10^{-13}$	0.06	0.008	$2.27 \times 10^{-15}$	0.05 (0.040 to 0.063)	$1.37 \times 10^{-19}$
Peptide	Gamma-glutamyl	Gamma-glutamylvaline	0.06	0.008	$8.28 \times 10^{-17}$	0.04	0.008	$1.62 \times 10^{-8}$	0.06 (0.046 to 0.070)	$1.23 \times 10^{-20}$
Peptide	Polypeptide	HWESASXX	0.04	0.008	$2.86 \times 10^{-9}$	−0.01	0.007	$1.48 \times 10^{-1}$		

$\beta$  values (SEs) are estimated with linear regressions adjusting for age, BMI at baseline, smoking, batch, and familiar relatedness. Results are combined (if replicated in the KORA study) using fixed-effect meta-analysis.  $\beta^a$  represents the yearly change in BMI per 1 SD increase of metabolite concentration. Analyses in KORA are additionally adjusted for sex.

To quantify how much of the change in BMI per year was explained by these four metabolites, we computed the AUC in a receiver operating characteristic analysis. We found that the AUC obtained by adding these four metabolites was significantly greater than for the lifestyle and demographic variables alone. Demographics plus smoking gave an AUC of 0.563 (95% CI: 0.539-0.587), whereas the addition of the metabolite data resulted in an AUC of 0.637 (95% CI: 0.614-0.660) (Supporting Information Figure S2).

The full metabolome has been previously investigated for its association with intake of 71 food groups (18). (The full information can be searched online at <http://www.twinsuk.ac.uk/dietmetab-data/>.) Of the metabolites identified here as related to weight change, urate, gamma-glutamyl valine, and butyrylcarnitine were not associated with any of the 71 food groups that we considered. On the other hand, 3-phenylpropionate (hydrocinnamate), which was associated with negative BMI change, was associated with higher consumption of apples and pears ( $\beta$  [SE] = 0.02 [0.004];  $P = 1.24 \times 10^{-8}$ ) and lower consumption of fried fish ( $\beta$  [SE] = −0.17 [0.05];  $P = 4.12 \times 10^{-11}$ ) and savory pies ( $\beta$  [SE] = −0.23 [0.04];  $P = 3.72 \times 10^{-10}$ ) (18).


We further tested whether baseline levels of these metabolites were also predictive of weight change and found that circulating levels of urate at baseline were associated with higher weight gain at follow-up in both KORA ( $\beta$  [SE] = 0.02 [0.009];  $P = 0.04$ ) and PLCO ( $\beta$  [SE] = 0.04 [0.020];  $P = 0.008$ ) and in the meta-analysis ( $\beta$  [95% CI] = 0.023 [0.007-0.039];  $P = 0.004$ ).

We hypothesized that urate may be a surrogate marker of the metabolic change or changes that result in weight gain. If this were true, we would expect levels of urate at baseline to be positively correlated with levels of fatty acids in blood in individuals who gain the most weight. If that were the case, there should be no such metabolic change in people who lose weight. Given the link to oxidative stress, we further hypothesized that this might affect SFAs and PUFAs differently. We therefore tested the interactions between urate and circulating levels of SFA, TotFA, monounsaturated fatty acid, and PUFA in the weight gain and weight loss groups (top and bottom tertiles of the weight change distribution, which had changes of  $-0.17 \pm 0.26$  and  $0.39 \pm 0.22$  kg/m<sup>2</sup> per year, respectively) and found that the association between urate and both SFAs and TotFAs was only present in those who gained weight (SFA:  $\beta$  [SE] = 0.09 [0.04];  $P = 0.04$ ; TotFA:  $\beta$  [SE] = 0.1 [0.04];  $P = 0.02$ ), while lower PUFAs were associated with higher circulating urate levels in those who lost weight (PUFA:  $\beta$  [SE] = −0.1 [0.03];  $P = 0.002$ ) (Figure 1).

## Discussion

In this study, we profiled the metabolic signatures of long-term weight change. Having assessed the association between circulating metabolites and weight change, we identified four metabolites—urate, butyrylcarnitine, gamma-glutamylvaline and 3-phenylpropionate (hydrocinnamate), all of which have been previously linked to higher oxidative stress—that were independently associated with long-term weight gain and successfully replicated in an independent cohort.

Previous studies reported that several metabolites—including BCAAs, butyrylcarnitine cysteine, glutamine, phenylalanine, proline,



**Figure 1** Association between circulating urate and fatty acids among participants with weight gain and those with weight loss. Linear regression coefficient  $\beta$  values (95% CIs) are depicted.

tyrosine, histidine, threonine, tryptophan, choline, pantothenic acid, LDL, very low-density lipoprotein, asparagine, glycine, methionine, and citrulline, among others—were associated with obesity (3-5). Some of the metabolites we identified as related to weight change have been previously linked to obesity, specifically butyrylcarnitine, which was reported by Moore et al. in individuals of both Asian and European descent (5). We did not find a correlation with BCAAs, which are well-known markers of visceral fat (19), obesity in general (7), and type 2 diabetes (3). Also, unlike the authors of previous studies (8), we did not find a strong correlation between weight change and lipids such as LDL or other forms of cholesterol. These findings are in line with those of studies of high fat and high sucrose intake with regards to the generation of reactive oxygen species (ROS). Treatment with high glucose or high fatty acid concentrations in cultured skeletal muscle cells has been shown to induce ROS production and mitochondrial damage in myotubes (20). In both rodents and humans, a high-fat diet increased the hydrogen peroxide-emitting potential of mitochondria without any changes in oxidative capacity (21). Not only does high-fat, high-sucrose feeding in mice result in increased ROS production, but this ROS production also results in mitochondrial dysfunction (20,21). In light of this finding, a possible mechanism linking the metabolites identified in our study and previous research could be the strong link between a dysfunction in energy metabolism and ROS generation. **O**

### Butyrylcarnitine

Butyrylcarnitine is an acylcarnitine, a type of compound derived from mitochondrial acyl-CoA metabolism, and it has been associated with diet-induced insulin resistance (22). Acylcarnitines comprise an acyl group esterified to L-carnitine, which enables them to cross the mitochondrial membrane. Several recent studies show increased levels of multiple acylcarnitines in subjects with obesity and insulin resistance (23). Acylcarnitines are generated during incomplete fatty acid beta-oxidation, and treatment of cultured murine and human myotubes with these compounds induces insulin resistance and oxidative stress (24). High levels of butyrylcarnitine are therefore indicative of incomplete beta-oxidation and are likely causative of increased ROS.

### Gamma-glutamylvaline

Gamma-glutamylvaline, like other gamma-glutamylated amino acids, is an established marker of oxidative stress and reflective of glutathione (GSH) turnover (25). GSH is the major redox buffer in the cell (26). It protects cellular components from oxidative damage of ROS such as hydrogen peroxide and organic peroxides via GSH peroxidases and also detoxifies the electrophilic metabolites of toxins (27). The intracellular redox environment can be measured through

the redox state of the GSH disulfide/GSH couple (28). Under conditions of oxidative stress, intracellular GSH turnover is increased, activating the enzyme  $\gamma$ -glutamyl transpeptidase to cleave extracellular GSH to release gamma-glutamylated amino acids, which are taken up for intracellular GSH reconstitution. A drop in such amino acids has been reported in response to vitamin E treatment (25).

### 3-phenylpropionate (hydrocinnamate)

Lower levels of the metabolite 3-phenylpropionate (hydrocinnamate) were also reproducibly associated with weight gain. Hydrocinnamate is a phenolic compound (29) with well-characterized antioxidant activity (30). Moreover, derivatives of this compound are known to have agonistic effects on peroxisome proliferator-activated receptors with antidiabetic and lipid-lowering activity (31).

### Urate

In humans, the final compound of purine catabolism is uric acid (32). Urate is a well-known marker of intracellular prooxidant and inflammatory status. Although it has antioxidant activity extracellularly, it has been widely linked to increased risk of metabolic syndrome (33). We found urate to be associated with weight change in all cohorts (Table 2), with urate levels at baseline being correlated with higher weight gain at follow-up.

Urate, despite being a major antioxidant in the human plasma, both correlates and predicts development of obesity, hypertension, and cardiovascular disease, conditions associated with oxidative stress (34). Urate is a prooxidant when forming radicals in reactions with other oxidants, and these radicals predominantly target lipids (LDL and membranes) rather than other cellular components. However, the hydrophobic environment created by lipids is unfavorable for the antioxidant effects of uric acid, and oxidized lipids thus convert uric acid into an oxidant (34). There is a substantial body of evidence indicating that uric acid has a direct effect on adipose tissue, and this effect has a redox-dependent component (34–36). Moreover, hyperuricemia was previously found to be a mediator linking obesity with kidney disease and might, thus, increase oxidative stress due to decreasing renal function (37).

### Proposed biological mechanism

The metabolic signatures we saw therefore could be interpreted as follows: The increase in fatty acids and carbohydrates produces increased ROS (as witnessed by the accumulation of gamma glutamyl amino acids) and incomplete beta-oxidation (accumulation of acylcarnitines), which in turn causes mitochondrial dysfunction linked to a dysregulation of the tricarboxylic acid cycle. This would result in decreased adenosine triphosphate production, and eventually mitochondrial DNA degradation, leading to the release of nucleotides, nucleosides, and nucleobases, which are metabolized to urate by oxidation.

In a subanalysis of TwinsUK cohort data we found that urate is correlated with increased levels of SFAs and TotFAs, but only in individuals who gain weight (Figure 1). On the other hand, PUFAs, which have an antioxidant effect themselves (38), are correlated with lower levels of urate, but only in individuals who lose weight. Our data therefore suggest that increases (even at normal levels) of urate may potentially be markers of the start of metabolic changes

that result in long-term weight gain. It also suggests that the metabolites identified in our study might be used to monitor the efficacy of therapies aimed at restoring mitochondrial function, such as ongoing clinical trials using nicotinamide adenine dinucleotide precursors (39). Urate is not only an easily measured marker that can be monitored in therapies aimed at reducing weight gain, but, importantly, may also be modified through the use of urate-lowering therapy. We note, however, that this is merely one interpretation of the data, and we were unable to prove causality; therefore, other interpretations are possible.

The current study has several strengths. It used a nontargeted metabolomics approach that identified a wide range of biochemicals in addition to lipids. TwinsUK is a very large and extensively phenotyped cohort, and we replicated our results in two independent cohorts.

We also note several study limitations. Our discovery sample consisted of females only. Longitudinal metabolomic data were not available in all of the three cohorts. Also, we did not have a direct measure of oxidative phosphorylation and redox function in these subjects. The lack of metabolomic data at baseline in the discovery cohort did not allow us to test the predictive value of these metabolites on weight loss or gain.

Another important limitation is the possibility of confounding with total energy intake and expenditure. Adjustment using self-reported measures from questionnaires on diet and physical activity did not make a difference to our findings, but these self-report values are not objectively measured, are subject to bias, and have wide margins of error (40,41). In this study, we were unable to adjust for objective measures of energy intake and expenditure, and therefore, the effects that we reported may be affected if adjusted by such measures. On the other hand, the four metabolites that we focused on were replicated in two independent studies, lending further confidence to the robustness of the four metabolomic associations reported here.

In conclusion, our study shows that metabolites linked to higher oxidative stress define a metabolic profile correlated with long-term weight gain and that urate at baseline is associated with weight gain over time.

© 2017 The Authors. *Obesity* published by Wiley Periodicals, Inc. on behalf of The Obesity Society (TOS)

### References

1. Bray GA, Smith SR, de Jonge L, et al. Effect of dietary protein content on weight gain, energy expenditure, and body composition during overeating: a randomized controlled trial. *JAMA* 2012;307:47–55.
2. Haqq AM, Lien LF, Boan J, et al. The Study of the Effects of Diet on Metabolism and Nutrition (STEDMAN) weight loss project: rationale and design. *Contemp Clin Trials* 2005;26:616–625.
3. Abu Bakar MH, Sarmidi MR, Cheng KK, et al. Metabolomics - the complementary field in systems biology: a review on obesity and type 2 diabetes. *Mol Biosyst* 2015;11:1742–1774.
4. Park S, Sadanala KC, Kim EK. A metabolomic approach to understanding the metabolic link between obesity and diabetes. *Mol Cells* 2015;38:587–596.
5. Moore SC, Matthews CE, Sampson JN, et al. Human metabolic correlates of body mass index. *Metabolomics* 2014;10:259–269.
6. Wahl S, Yu Z, Kleber M, et al. Childhood obesity is associated with changes in the serum metabolite profile. *Obes Facts* 2012;5:660–670.
7. Tulipani S, Griffin J, Palau-Rodriguez M, et al. Metabolomics-guided insights on bariatric surgery versus behavioral interventions for weight loss. *Obesity (Silver Spring)* 2016;24:2451–2466.
8. Wahl S, Vogt S, Stuckler F, et al. Multi-omic signature of body weight change: results from a population-based cohort study. *BMC Med* 2015;13:48. doi:10.1186/s129160150282y

9. Moayyeri A, Hammond CJ, Valdes AM, Spector TD. Cohort Profile: TwinsUK and healthy ageing twin study. *Int J Epidemiol* 2013;42:76-85.
10. Holle R, Happich M, Lowel H, Wichmann HE. KORA—a research platform for population based health research. *Gesundheitswesen* 2005;67(suppl 1):S19-S25.
11. Ahn J, Moore SC, Albanes D, Huang WY, Leitzmann MF, Hayes RB; Prostate, Lung, Colorectal, and Ovarian (PLCO) Cancer Screening Trial Project Team. Height and risk of prostate cancer in the prostate, lung, colorectal, and ovarian cancer screening trial. *Br J Cancer* 2009;101:522–525.
12. Menni C, Jackson MA, Pallister T, Steves C, Spector T, Valdes A. Gut microbiome diversity and high fibre intake are related to lower long term weight gain. *Int J Obes (Lond)* 2017;41:1099-1105.
13. Menni C, Kastenmuller G, Petersen AK, et al. Metabolomic markers reveal novel pathways of ageing and early development in human populations. *Int J Epidemiol* 2013;42:1111-1119.
14. Soininen P, Kangas AJ, Wurtz P, Suna T, Ala-Korpela M. Quantitative serum nuclear magnetic resonance metabolomics in cardiovascular epidemiology and genetics. *Circ Cardiovasc Genet* 2015;8:192-206.
15. Wurtz P, Havulinna AS, Soininen P, et al. Metabolite profiling and cardiovascular event risk: a prospective study of 3 population-based cohorts. *Circulation* 2015;131:774-785.
16. Tibshirani R. Regression shrinkage and selection via the lasso: a retrospective. *J R Stat Soc B* 2011;73:273-282.
17. Friedman J, Hastie T, Tibshirani R. Regularization paths for generalized linear models via coordinate descent. *J Stat Softw* 2010;33:1-22.
18. Pallister T, Jennings A, Mohny RP, et al. Characterizing blood metabolomics profiles associated with self-reported food intakes in female twins. *PLoS One* 2016;11:e0158568. doi:10.1371/journal.pone.0158568
19. Menni C, Migaud M, Glastonbury CA, et al. Metabolomic profiling to dissect the role of visceral fat in cardiometabolic health. *Obesity (Silver Spring)* 2016;24:1380-1388.
20. Bonnard C, Durand A, Peyrol S, et al. Mitochondrial dysfunction results from oxidative stress in the skeletal muscle of diet-induced insulin-resistant mice. *J Clin Invest* 2008;118:789-800.
21. Anderson EJ, Lustig ME, Boyle KE, et al. Mitochondrial H<sub>2</sub>O<sub>2</sub> emission and cellular redox state link excess fat intake to insulin resistance in both rodents and humans. *J Clin Invest* 2009;119:573-581.
22. Schooneman MG, Vaz FM, Houten SM, Soeters MR. Acylcarnitines: reflecting or inflicting insulin resistance? *Diabetes* 2013;62:1-8.
23. Mihalik SJ, Goodpaster BH, Kelley DE, et al. Increased levels of plasma acylcarnitines in obesity and type 2 diabetes and identification of a marker of glucolipotoxicity. *Obesity (Silver Spring)* 2010;18:1695-1700.
24. Aguer C, McCain CS, Knotts TA, et al. Acylcarnitines: potential implications for skeletal muscle insulin resistance. *FASEB J* 2015;29:336-345.
25. Cheng J, Joyce A, Yates K, Aouizerat B, Sanyal AJ. Metabolomic profiling to identify predictors of response to vitamin E for non-alcoholic steatohepatitis (NASH). *PLoS One* 2012;7:e44106. doi:10.1371/journal.pone.0044106
26. Moriarty-Craige SE, Jones DP. Extracellular thiols and thiol/disulfide redox in metabolism. *Annu Rev Nutr* 2004;24:481-509.
27. Hanigan MH. Gamma-glutamyl transpeptidase: redox regulation and drug resistance. *Adv Cancer Res* 2014;122:103-141.
28. Schafer FQ, Buettner GR. Redox environment of the cell as viewed through the redox state of the glutathione disulfide/glutathione couple. *Free Radic Biol Med* 2001;30:1191-1212.
29. Jenner AM, Rafter J, Halliwell B. Human fecal water content of phenolics: the extent of colonic exposure to aromatic compounds. *Free Radic Biol Med* 2005;38:763-772.
30. Roleira FM, Siquet C, Orru E, et al. Lipophilic phenolic antioxidants: correlation between antioxidant profile, partition coefficients, and redox properties. *Bioorg Med Chem* 2010;18:5816-5825.
31. Raval P, Jain M, Goswami A, et al. Revisiting glitazars: thiophene substituted oxazole containing alpha-ethoxy phenylpropanoic acid derivatives as highly potent PPARalpha/gamma dual agonists devoid of adverse effects in rodents. *Bioorg Med Chem Lett* 2011;21:3103-3109.
32. Grassi D, Ferri L, Desideri G, et al. Chronic hyperuricemia, uric acid deposit and cardiovascular risk. *Curr Pharm Des* 2013;19:2432-2438.
33. Srikanthan K, Feyh A, Visweshwar H, Shapiro JI, Sodhi K. Systematic review of metabolic syndrome biomarkers: a panel for early detection, management, and risk stratification in the West Virginian population. *Int J Med Sci* 2016;13:25-38.
34. Sautin YY, Johnson RJ. Uric acid: the oxidant-antioxidant paradox. *Nucleosides Nucleotides Nucleic Acids* 2008;27:608-619.
35. Moriya C, Satoh H. Teneligliptin decreases uric acid levels by reducing xanthine dehydrogenase expression in white adipose tissue of male wistar rats. *J Diabetes Res* 2016;2016:3201534. doi:10.1155/2016/3201534
36. Tsushima Y, Nishizawa H, Tochino Y, et al. Uric acid secretion from adipose tissue and its increase in obesity. *J Biol Chem* 2013;288:27138-27149.
37. Zierer J, Pallister T, Tsai PC, et al. Exploring the molecular basis of age-related disease comorbidities using a multi-omics graphical model. *Sci Rep* 2016;6:37646. doi:10.1038/srep37646
38. Richard D, Kefi K, Barbe U, Bausero P, Visioli F. Polyunsaturated fatty acids as antioxidants. *J Biochem Pharmacol Res* 2008;57:451-455.
39. Trammell SA, Schmidt MS, Weidemann BJ, et al. Nicotinamide riboside is uniquely and orally bioavailable in mice and humans. *Nat Commun* 2016;7:12948. doi:10.1038/ncomms12948
40. Kromhout D. Energy and macronutrient intake in lean and obese middle-aged men (the Zutphen study). *Am J Clin Nutr* 1983;37:295-299.
41. Freedman LS, Commins JM, Moler JE, et al. Pooled results from 5 validation studies of dietary self-report instruments using recovery biomarkers for potassium and sodium intake. *Am J Epidemiol* 2015;181:473-487.